

# PROBABILITY & STATISTICS

Lecture 11 – Confidence Intervals

# CONFIDENCE INTERVALS

# MOTIVATION

- Point estimation provides a single “best guess”.

# MOTIVATION

- Point estimation provides a single “best guess”.
- How confident can we be in our estimation?

# MOTIVATION

- Point estimation provides a single “best guess”.
- How confident can we be in our estimation?
- CI proposes a range of plausible values.

# DEFINITION

A  $1 - \alpha$  confidence interval for a parameter  $\theta$  is an interval  $C_n = (a, b)$  such that  $T_1 = t_1(X_1, \dots, X_n)$ ,  $T_2 = t_2(X_1, \dots, X_n)$  and

$$P(T_1 < \theta < T_2) \geq 1 - \alpha$$

# DEFINITION

A  $1 - \alpha$  confidence interval for a parameter  $\theta$  is an interval  $C_n = (T_1, T_2)$  such that  $T_1 = t_1(X_1, \dots, X_n)$ ,  $T_2 = t_2(X_1, \dots, X_n)$  and

$$P(T_1 < \theta < T_2) \geq 1 - \alpha$$

- **Random** intervals:  $T_1$  and  $T_2$  are functions of random samples.

# DEFINITION

A  $1 - \alpha$  confidence interval for a parameter  $\theta$  is an interval  $C_n = (T_1, T_2)$  such that  $T_1 = t_1(X_1, \dots, X_n)$ ,  $T_2 = t_2(X_1, \dots, X_n)$  and

$$P(T_1 < \theta < T_2) \geq 1 - \alpha$$

- **Random** intervals:  $T_1$  and  $T_2$  are functions of random samples.
- $\theta$  is unknown, but fixed  
 $T_1$  and  $T_2$  are random



# INTERPRETATION

$$C_n = (T_1, T_2): P(T_1 < \theta < T_2) \geq 1 - \alpha$$

# INTERPRETATION

$$C_n = (T_1, T_2): P(T_1 < \theta < T_2) \geq 1 - \alpha$$

- **Not** a probability statement about  $\theta$  since it's fixed.

# INTERPRETATION

$$C_n = (T_1, T_2): P(T_1 < \theta < T_2) \geq 1 - \alpha$$

- **Not** a probability statement about  $\theta$  since it's fixed.
- Common interpretation:

# INTERPRETATION

$$C_n = (T_1, T_2): P(T_1 < \theta < T_2) \geq 1 - \alpha$$

- **Not** a probability statement about  $\theta$  since it's fixed.
- Common interpretation:

*If I repeat the experiment many times, the interval will contain the true value of  $\theta$  95% of the time ( $\alpha=0.05$ ).*

# CI FOR NORMAL DATA

CI for  $\mu$ , *known*  $\sigma$

# CI FOR $\mu$ , KNOWN $\sigma$

$X_1, X_2, \dots, X_n$  — samples from  $N(\mu, \sigma^2)$ ,  $\sigma$  is known.

How to construct a CI for  $\mu$ ?

# CI FOR $\mu$ , KNOWN $\sigma$

$X_1, X_2, \dots, X_n$  – samples from  $N(\mu, \sigma^2)$ ,  $\sigma$  is known.

How to construct a CI for  $\mu$ ? Consider  $\bar{X}$ :

# CI FOR $\mu$ , KNOWN $\sigma$

$X_1, X_2, \dots, X_n$  – samples from  $N(\mu, \sigma^2)$ ,  $\sigma$  is known.

How to construct a CI for  $\mu$ ? Consider  $\bar{X}$ :

$$\bar{X} = \frac{1}{n} \sum_{i=1}^n X_i \sim$$



# CI FOR $\mu$ , KNOWN $\sigma$

$X_1, X_2, \dots, X_n$  – samples from  $N(\mu, \sigma^2)$ ,  $\sigma$  is known.

How to construct a CI for  $\mu$ ? Consider  $\bar{X}$ :

$$\bar{X} = \frac{1}{n} \sum_{i=1}^n X_i \sim N( \quad ) \Rightarrow$$

# CI FOR $\mu$ , KNOWN $\sigma$

$X_1, X_2, \dots, X_n$  – samples from  $N(\mu, \sigma^2)$ ,  $\sigma$  is known.

How to construct a CI for  $\mu$ ? Consider  $\bar{X}$ :

$$\bar{X} = \frac{1}{n} \sum_{i=1}^n X_i \sim N(\mu, \quad ) \Rightarrow$$

# CI FOR $\mu$ , KNOWN $\sigma$

$X_1, X_2, \dots, X_n$  – samples from  $N(\mu, \sigma^2)$ ,  $\sigma$  is known.

How to construct a CI for  $\mu$ ? Consider  $\bar{X}$ :

$$\bar{X} = \frac{1}{n} \sum_{i=1}^n X_i \sim N(\mu, \sigma^2/n) \quad \Rightarrow$$

# CI FOR $\mu$ , KNOWN $\sigma$

$X_1, X_2, \dots, X_n$  – samples from  $N(\mu, \sigma^2)$ ,  $\sigma$  is known.

How to construct a CI for  $\mu$ ? Consider  $\bar{X}$ :

$$\bar{X} = \frac{1}{n} \sum_{i=1}^n X_i \sim N(\mu, \sigma^2/n) \quad \Rightarrow \quad \frac{\bar{X} - \mu}{\sigma/\sqrt{n}} \sim N(0,1)$$

# CI FOR $\mu$ , KNOWN $\sigma$

$X_1, X_2, \dots, X_n$  – samples from  $N(\mu, \sigma^2)$ ,  $\sigma$  is known.

How to construct a CI for  $\mu$ ? Consider  $\bar{X}$ :

$$\bar{X} = \frac{1}{n} \sum_{i=1}^n X_i \sim N(\mu, \sigma^2/n) \Rightarrow \frac{(\bar{X} - \mu)\sqrt{n}}{\sigma} \sim N(0,1)$$

# CI FOR $\mu$ , KNOWN $\sigma$


$X_1, X_2, \dots, X_n$  – samples from  $N(\mu, \sigma^2)$ ,  $\sigma$  is known.

How to construct a CI for  $\mu$ ? Consider  $\bar{X}$ :


$$\bar{X} = \frac{1}{n} \sum_{i=1}^n X_i \sim N(\mu, \sigma^2/n) \quad \Rightarrow \quad \frac{(\bar{X} - \mu)\sqrt{n}}{\sigma} \sim N(0,1)$$

$$P\left( -z_{\alpha/2} < \frac{(\bar{X} - \mu)\sqrt{n}}{\sigma} < z_{\alpha/2} \right) = 1 - \alpha$$

# CI FOR $\mu$ , KNOWN $\sigma$




# CI FOR $\mu$ , KNOWN $\sigma$







# CI FOR $\mu$ , KNOWN $\sigma$



# CI FOR $\mu$ , KNOWN $\sigma$



# CI FOR $\mu$ , KNOWN $\sigma$



# CI FOR $\mu$ , KNOWN $\sigma$

$X_1, X_2, \dots, X_n$  – samples from  $N(\mu, \sigma^2)$ ,  $\sigma$  is known.

How to construct a CI for  $\mu$ ? Consider  $\bar{X}$ :

$$\bar{X} = \frac{1}{n} \sum_{i=1}^n X_i \sim N(\mu, \sigma^2/n) \quad \Rightarrow \quad \frac{(\bar{X} - \mu)\sqrt{n}}{\sigma} \sim N(0,1)$$

$$P\left(-z_{1-\frac{\alpha}{2}} < \frac{(\bar{X} - \mu)\sqrt{n}}{\sigma} < z_{1-\frac{\alpha}{2}}\right) = 1 - \alpha$$

# CI FOR $\mu$ , KNOWN $\sigma$

$X_1, X_2, \dots, X_n$  – samples from  $N(\mu, \sigma^2)$ ,  $\sigma$  is known.

How to construct a CI for  $\mu$ ? Consider  $\bar{X}$ :

$$\bar{X} = \frac{1}{n} \sum_{i=1}^n X_i \sim N(\mu, \sigma^2/n) \quad \Rightarrow \quad \frac{(\bar{X} - \mu)\sqrt{n}}{\sigma} \sim N(0,1)$$

$$P\left(-z_{1-\frac{\alpha}{2}} < \frac{(\bar{X} - \mu)\sqrt{n}}{\sigma} < z_{1-\frac{\alpha}{2}}\right) = 1 - \alpha$$

$$P\left(\bar{X} - \frac{\sigma}{\sqrt{n}} z_{1-\frac{\alpha}{2}} < \mu < \bar{X} + \frac{\sigma}{\sqrt{n}} z_{1-\frac{\alpha}{2}}\right) = 1 - \alpha$$

# CI FOR $\mu$ , KNOWN $\sigma$ : EXAMPLE 1

- $X_1, X_2, \dots, X_{100}$  – samples from  $N(\mu, 3^2)$ ,  $\bar{X} = 12$ .
- Give the 95%-CI for  $\mu$ .

# CI FOR $\mu$ , KNOWN $\sigma$ : EXAMPLE 1

- $X_1, X_2, \dots, X_{100}$  – samples from  $N(\mu, 3^2)$ ,  $\bar{X} = 12$ .
- Give the 95%-CI for  $\mu$ .

- Interval:

$$\bar{X} \pm \frac{\sigma}{\sqrt{n}} \cdot z_{1-\alpha/2}$$

# CI FOR $\mu$ , KNOWN $\sigma$ : EXAMPLE 1

- $X_1, X_2, \dots, X_{100}$  – samples from  $N(\mu, 3^2)$ ,  $\bar{X} = 12$ .
- Give the 95%-CI for  $\mu$ .

- Interval:

$$\bar{X} \pm \frac{\sigma}{\sqrt{n}} \cdot z_{1-\alpha/2}$$

$$1 - \alpha = 0.95 \Rightarrow \alpha = 0.05,$$



# CI FOR $\mu$ , KNOWN $\sigma$ : EXAMPLE 1

- $X_1, X_2, \dots, X_{100}$  – samples from  $N(\mu, 3^2)$ ,
- Give the 95%-CI for  $\mu$ .

- Interval:

$$\bar{X} \pm \frac{\sigma}{\sqrt{n}} \cdot z_{1-\alpha/2}$$

$$1 - \alpha = 0.95 \Rightarrow \alpha = 0.05, \quad z_{1-0.025} =$$

Quantile (p)	$\Phi^{-1}(p, 0, 1)$
0.995	2.58
0.99	2.33
0.975	1.96
0.95	1.64
0.9	1.28

# CI FOR $\mu$ , KNOWN $\sigma$ : EXAMPLE 1

- $X_1, X_2, \dots, X_{100}$  – samples from  $N(\mu, 3^2)$ ,
- Give the 95%-CI for  $\mu$ .
- Interval:

$$\bar{X} \pm \frac{\sigma}{\sqrt{n}} \cdot z_{1-\alpha/2}$$

$$1 - \alpha = 0.95 \Rightarrow \alpha = 0.05, \quad z_{1-0.025} =$$

Quantile (p)	$\Phi^{-1}(p, 0, 1)$
0.995	2.58
0.99	2.33
0.975	1.96
0.95	1.64
0.9	1.28

# CI FOR $\mu$ , KNOWN $\sigma$ : EXAMPLE 1

- $X_1, X_2, \dots, X_{100}$  – samples from  $N(\mu, 3^2)$ ,
- Give the 95%-CI for  $\mu$ .
- Interval:

$$\bar{X} \pm \frac{\sigma}{\sqrt{n}} \cdot z_{1-\alpha/2}$$

$$1 - \alpha = 0.95 \Rightarrow \alpha = 0.05, \quad z_{1-0.025} = 1.96$$

Quantile (p)	$\Phi^{-1}(p, 0, 1)$
0.995	2.58
0.99	2.33
0.975	1.96
0.95	1.64
0.9	1.28

# CI FOR $\mu$ , KNOWN $\sigma$ : EXAMPLE 1

- $X_1, X_2, \dots, X_{100}$  – samples from  $N(\mu, 3^2)$ ,  $\bar{X} = 12$ .
- Give the 95%-CI for  $\mu$ .

- Interval:

$$\bar{X} \pm \frac{\sigma}{\sqrt{n}} \cdot z_{1-\alpha/2}$$

$$1 - \alpha = 0.95 \Rightarrow \alpha = 0.05, \quad z_{1-0.025} = 1.96$$

$$12 \pm \frac{3}{\sqrt{100}} \cdot 1.96$$

# CI FOR $\mu$ , KNOWN $\sigma$ : EXAMPLE 2

- $X_1, X_2, \dots, X_{100}$  — samples from  $N(\mu, 1^2)$ ,  $\bar{X} = 5$ .
- Give the 90%-CI for  $\mu$ .

# CI FOR $\mu$ , KNOWN $\sigma$ : EXAMPLE 2

- $X_1, X_2, \dots, X_{100}$  – samples from  $N(\mu, 1^2)$ ,  $\bar{X} = 5$ .
- Give the 90%-CI for  $\mu$ .

- Interval:

$$\bar{X} \pm \frac{\sigma}{\sqrt{n}} \cdot z_{1-\alpha/2}$$

Quantile (p)	$\Phi^{-1}(p, 0, 1)$
0.995	2.58
0.99	2.33
0.975	1.96
0.95	1.64
0.9	1.28

# CI FOR $\mu$ , KNOWN $\sigma$ : EXAMPLE 2

- $X_1, X_2, \dots, X_{100}$  – samples from  $N(\mu, 1^2)$ ,  $\bar{X} = 5$ .
- Give the 90%-CI for  $\mu$ .

- Interval:

$$\bar{X} \pm \frac{\sigma}{\sqrt{n}} \cdot z_{1-\alpha/2}$$

Quantile (p)	$\Phi^{-1}(p, 0, 1)$
0.995	2.58
0.99	2.33
0.975	1.96
0.95	1.64
0.9	1.28

# CI FOR $\mu$ , KNOWN $\sigma$ : EXAMPLE 2

- $X_1, X_2, \dots, X_{100}$  – samples from  $N(\mu, 1^2)$ ,  $\bar{X} = 5$ .
- Give the 90%-CI for  $\mu$ .

- Interval:

$$\bar{X} \pm \frac{\sigma}{\sqrt{n}} \cdot z_{1-\alpha/2}$$

$$1 - \alpha = 0.90 \Rightarrow \alpha = 0.1, \quad z_{1-0.05} = 1.64$$



# CI FOR $\mu$ , KNOWN $\sigma$ : EXAMPLE 2

- $X_1, X_2, \dots, X_{100}$  – samples from  $N(\mu, 1^2)$ ,  $\bar{X} = 5$ .
- Give the 90%-CI for  $\mu$ .

- Interval:

$$\bar{X} \pm \frac{\sigma}{\sqrt{n}} \cdot z_{1-\alpha/2}$$

$$1 - \alpha = 0.90 \Rightarrow \alpha = 0.1, \quad z_{1-0.05} = 1.64$$

$$5 \pm \frac{1}{\sqrt{100}} \cdot 1.64$$

# PRACTICE!

Google Classroom -> z-intervals

# CONFIDENCE INTERVALS: RECAP

- NORMAL DISTRIBUTION:

$X_1, X_2, \dots, X_n$  – i.i.d. samples

- CI for  $\mu$ ,  $\sigma$  is known: **z-interval**

$$\bar{X} \pm \frac{\sigma}{\sqrt{n}} z_{1-\alpha/2}, \quad z_{1-\alpha/2} \text{ – quantile from } N(0,1)$$

# CI FOR NORMAL DATA

CI for  $\mu$ , *unknown*  $\sigma$

# CI FOR $\mu$ , UNKNOWN $\sigma$

$X_1, X_2, \dots, X_n$  – samples from  $N(\mu, \sigma^2)$ ,  $\sigma$  is unknown.

How to construct a CI for  $\mu$ ?

# CI FOR $\mu$ , UNKNOWN $\sigma$

$X_1, X_2, \dots, X_n$  – samples from  $N(\mu, \sigma^2)$ ,  $\sigma$  is unknown.

How to construct a CI for  $\mu$ ? Consider  $\bar{X}$ :

# CI FOR $\mu$ , UNKNOWN $\sigma$

$X_1, X_2, \dots, X_n$  – samples from  $N(\mu, \sigma^2)$ ,  $\sigma$  is unknown.

How to construct a CI for  $\mu$ ? Consider  $\bar{X}$ :

$$\bar{X} = \frac{1}{n} \sum_{i=1}^n X_i \sim N(\mu, \sigma^2/n)$$

# CI FOR $\mu$ , UNKNOWN $\sigma$

$X_1, X_2, \dots, X_n$  – samples from  $N(\mu, \sigma^2)$ ,  $\sigma$  is unknown.

How to construct a CI for  $\mu$ ? Consider  $\bar{X}$ :

$$\bar{X} = \frac{1}{n} \sum_{i=1}^n X_i \sim N(\mu, \sigma^2/n) \Rightarrow \frac{(\bar{X} - \mu)\sqrt{n}}{\sigma} \sim N(0,1)$$



# CI FOR $\mu$ , UNKNOWN $\sigma$

$X_1, X_2, \dots, X_n$  – samples from  $N(\mu, \sigma^2)$ ,  $\sigma$  is unknown.

How to construct a CI for  $\mu$ ? Consider  $\bar{X}$ :

$$\bar{X} = \frac{1}{n} \sum_{i=1}^n X_i \sim N(\mu, \sigma^2/n) \Rightarrow \frac{(\bar{X} - \mu)\sqrt{n}}{\sigma} \sim N(0,1)$$

# CI FOR $\mu$ , UNKNOWN $\sigma$

$X_1, X_2, \dots, X_n$  – samples from  $N(\mu, \sigma^2)$ ,  $\sigma$  is unknown.

How to construct a CI for  $\mu$ ? Consider  $\bar{X}$ :

$$\bar{X} = \frac{1}{n} \sum_{i=1}^n X_i \sim N(\mu, \sigma^2/n) \Rightarrow \frac{(\bar{X} - \mu)\sqrt{n}}{\sigma} \sim N(0,1)$$

Approximate with sample variance  $s^2 = \frac{1}{n-1} \sum_{i=1}^n (X_i - \bar{X})^2$ :

# CI FOR $\mu$ , UNKNOWN $\sigma$

$X_1, X_2, \dots, X_n$  – samples from  $N(\mu, \sigma^2)$ ,  $\sigma$  is unknown.

How to construct a CI for  $\mu$ ? Consider  $\bar{X}$ :

$$\bar{X} = \frac{1}{n} \sum_{i=1}^n X_i \sim N(\mu, \sigma^2/n) \Rightarrow \frac{(\bar{X} - \mu)\sqrt{n}}{\sigma} \sim N(0,1)$$

Approximate with sample variance  $s^2 = \frac{1}{n-1} \sum_{i=1}^n (X_i - \bar{X})^2$ :

$$\frac{(\bar{X} - \mu)\sqrt{n}}{s}$$

# CI FOR $\mu$ , UNKNOWN $\sigma$

$X_1, X_2, \dots, X_n$  – samples from  $N(\mu, \sigma^2)$ ,  $\sigma$  is unknown.

How to construct a CI for  $\mu$ ? Consider  $\bar{X}$ :

$$\bar{X} = \frac{1}{n} \sum_{i=1}^n X_i \sim N(\mu, \sigma^2/n) \Rightarrow \frac{(\bar{X} - \mu)\sqrt{n}}{\sigma} \sim N(0,1)$$

Approximate with sample variance  $s^2 = \frac{1}{n-1} \sum_{i=1}^n (X_i - \bar{X})^2$ :

$$\frac{(\bar{X} - \mu)\sqrt{n}}{s} \sim t(n-1) \text{ — Student distribution}$$

# STUDENT DISTRIBUTION



William Sealy Gosset

# STUDENT DISTRIBUTION


$$\frac{(\bar{X}_n - \mu)\sqrt{n}}{s} \sim t(n - 1)$$



William Sealy Gosset

# STUDENT DISTRIBUTION

$$\frac{(\bar{X}_n - \mu)\sqrt{n}}{s} \sim t(n - 1)$$



William Sealy Gosset

# CI FOR $\mu$ , UNKNOWN $\sigma$

$X_1, X_2, \dots, X_n$  – samples from  $N(\mu, \sigma^2)$ ,  $\sigma$  is unknown.

How to construct a CI for  $\mu$ ? Consider  $\bar{X}$ :

$$\frac{(\bar{X} - \mu)\sqrt{n}}{s} \sim t(n - 1) \text{ – Student distribution}$$



# CI FOR $\mu$ , UNKNOWN $\sigma$

$X_1, X_2, \dots, X_n$  – samples from  $N(\mu, \sigma^2)$ ,  $\sigma$  is unknown.

How to construct a CI for  $\mu$ ? Consider  $\bar{X}$ :

$$\frac{(\bar{X} - \mu)\sqrt{n}}{s} \sim t(n - 1) - \text{Student distribution}$$

Before: 
$$P\left(-z_{1-\alpha/2} < \frac{(\bar{X} - \mu)\sqrt{n}}{\sigma} < z_{1-\alpha/2}\right) = 1 - \alpha$$

# CI FOR $\mu$ , UNKNOWN $\sigma$

$X_1, X_2, \dots, X_n$  – samples from  $N(\mu, \sigma^2)$ ,  $\sigma$  is unknown.

How to construct a CI for  $\mu$ ? Consider  $\bar{X}$ :

$$\frac{(\bar{X} - \mu)\sqrt{n}}{s} \sim t(n - 1) - \text{Student distribution}$$

$$\text{Before: } P\left(-z_{1-\alpha/2} < \frac{(\bar{X} - \mu)\sqrt{n}}{\sigma} < z_{1-\alpha/2}\right) = 1 - \alpha$$

Now:

$$P\left(-t_{n-1, 1-\alpha/2} < \frac{(\bar{X} - \mu)\sqrt{n}}{s} < t_{n-1, 1-\alpha/2}\right) = 1 - \alpha$$

# CI FOR $\mu$ , UNKNOWN $\sigma$

$X_1, X_2, \dots, X_n$  – samples from  $N(\mu, \sigma^2)$ ,  $\sigma$  is unknown.

How to construct a CI for  $\mu$ ? Consider  $\bar{X}$ :

$$\frac{(\bar{X} - \mu)\sqrt{n}}{s} \sim t(n - 1) - \text{Student distribution}$$

$$\text{Before: } P\left(-z_{1-\alpha/2} < \frac{(\bar{X} - \mu)\sqrt{n}}{\sigma} < z_{1-\alpha/2}\right) = 1 - \alpha$$

Now: same logic, replace  $z_{1-\alpha/2}$  with  $t_{1-\alpha/2}$ :

$$P\left(-t_{1-\alpha/2} < \frac{(\bar{X} - \mu)\sqrt{n}}{s} < t_{1-\alpha/2}\right) = 1 - \alpha$$

# CI FOR $\mu$ , UNKNOWN $\sigma$

$X_1, X_2, \dots, X_n$  – samples from  $N(\mu, \sigma^2)$ ,  $\sigma$  is unknown.

How to construct a CI for  $\mu$ ? Consider  $\bar{X}$ :

$$\frac{(\bar{X} - \mu)\sqrt{n}}{s} \sim t(n - 1) - \text{Student distribution}$$

$$\text{Before: } P\left(-z_{1-\alpha/2} < \frac{(\bar{X} - \mu)\sqrt{n}}{\sigma} < z_{1-\alpha/2}\right) = 1 - \alpha$$

Now: same logic, replace  $z_{1-\alpha/2}$  with  $t_{1-\alpha/2}$ :

$$P\left(-t_{1-\alpha/2} < \frac{(\bar{X} - \mu)\sqrt{n}}{s} < t_{1-\alpha/2}\right) = 1 - \alpha$$

# CI FOR $\mu$ , UNKNOWN $\sigma$

$X_1, X_2, \dots, X_n$  – samples from  $N(\mu, \sigma^2)$ ,  $\sigma$  is unknown.

How to construct a CI for  $\mu$ ? Consider  $\bar{X}$ :

$$\frac{(\bar{X} - \mu)\sqrt{n}}{s} \sim t(n - 1) - \text{Student distribution}$$

$$\text{Before: } P\left(-z_{1-\alpha/2} < \frac{(\bar{X} - \mu)\sqrt{n}}{\sigma} < z_{1-\alpha/2}\right) = 1 - \alpha$$

Now: same logic, replace  $z_{1-\alpha/2}$  with  $t_{1-\alpha/2}$ :

$$P\left(\bar{X} - \frac{s}{\sqrt{n}} t_{1-\alpha/2} < \mu < \bar{X} + \frac{s}{\sqrt{n}} t_{1-\alpha/2}\right) = 1 - \alpha$$

# CI FOR $\mu$ , UNKNOWN $\sigma$ : EXAMPLE

- $X_1, X_2, \dots, X_{20}$  – samples from  $N(\mu, \sigma^2)$ ,  $\sigma$  is unknown.
- $\bar{X} = 42, \quad s^2 = 36$
- Give the 95%-CI for  $\mu$ .

# CI FOR $\mu$ , UNKNOWN $\sigma$ : EXAMPLE

- $X_1, X_2, \dots, X_{20}$  – samples from  $N(\mu, \sigma^2)$ ,  $\sigma$  is unknown.
- $\bar{X} = 42, \quad s^2 = 36$
- Give the 95%-CI for  $\mu$ .

$$\mu = \bar{X} \pm \frac{s}{\sqrt{n}} t_{1-\alpha/2}$$

# CI FOR $\mu$ , UNKNOWN $\sigma$ : EXAMPLE

- $X_1, X_2, \dots, X_{20}$  – samples from  $N(\mu, \sigma^2)$ ,  $\sigma$  is unknown.
- $\bar{X} = 42, \quad s^2 = 36$
- Give the 95%-CI for  $\mu$ .

$$\mu = \bar{X} \pm \frac{s}{\sqrt{n}} t_{1-\alpha/2}$$

$$1 - \alpha = 0.95 \Rightarrow \alpha = 0.05,$$



# CI FOR $\mu$ , UNKNOWN $\sigma$ : EXAMPLE

- $X_1, X_2, \dots, X_{20}$  – samples from  $N(\mu, \sigma^2)$ ,  $\sigma$  is unknown.
- $\bar{X} = 42, \quad s^2 = 36$
- Give the 95%-CI for  $\mu$ .

$$\mu = \bar{X} \pm \frac{s}{\sqrt{n}} t_{1-\alpha/2}$$

$$1 - \alpha = 0.95 \Rightarrow \alpha = 0.05, \quad t_{1-0.025} =$$

# CI FOR $\mu$ , UNKNOWN $\sigma$ : EXAMPLE

- $X_1, X_2, \dots, X_{20}$  – samples from  $N(\mu, \sigma^2)$ ,  $\sigma$  is unknown.
- $\bar{X} = 42, \quad s^2 = 36$
- Give the 95%-CI for  $\mu$ .

$$\mu = \bar{X} \pm \frac{s}{\sqrt{n}} t_{1-\alpha/2}$$

$$1 - \alpha = 0.95 \Rightarrow \alpha = 0.05, \quad t_{1-0.025} = 2.093$$

# CI FOR $\mu$ , UNKNOWN $\sigma$ : EXAMPLE

- $X_1, X_2, \dots, X_{20}$  – samples from  $N(\mu, \sigma^2)$ ,  $\sigma$  is unknown.
- $\bar{X} = 42, \quad s^2 = 36$
- Give the 95%-CI for  $\mu$ .

$$\mu = \bar{X} \pm \frac{s}{\sqrt{n}} t_{1-\alpha/2}$$

$$1 - \alpha = 0.95 \Rightarrow \alpha = 0.05, \quad t_{1-0.025} = 2.093$$

$$\mu = 42 \pm \frac{6}{\sqrt{20}} \cdot 2.093$$

# CI FOR NORMAL DATA

CI for  $\sigma$ , unknown  $\mu$ ,  $\sigma$

# CI FOR $\sigma$ , UNKNOWN $\mu, \sigma$

- $X_1, X_2, \dots, X_{100}$  — samples from  $N(\mu, \sigma^2)$ ,  $\sigma$  is unknown.
- How to construct a CI for  $\sigma$ ?

# CI FOR $\sigma$ , UNKNOWN $\mu, \sigma$

- $X_1, X_2, \dots, X_{100}$  — samples from  $N(\mu, \sigma^2)$ ,  $\sigma$  is unknown.
- How to construct a CI for  $\sigma$ ? Consider  $S^2$ :

# CI FOR $\sigma$ , UNKNOWN $\mu, \sigma$

- $X_1, X_2, \dots, X_{100}$  – samples from  $N(\mu, \sigma^2)$ ,  $\sigma$  is unknown.
- How to construct a CI for  $\sigma$ ? Consider  $S^2$ :

$$s^2 = \frac{1}{n-1} \sum_{i=1}^n (X_i - \bar{X})^2$$

# CI FOR $\sigma$ , UNKNOWN $\mu, \sigma$

- $X_1, X_2, \dots, X_{100}$  – samples from  $N(\mu, \sigma^2)$ ,  $\sigma$  is unknown.
- How to construct a CI for  $\mu$ ? Consider  $S^2$ :

$$s^2 = \frac{1}{n-1} \sum_{i=1}^n (X_i - \bar{X})^2$$

$$\frac{(n-1)s^2}{\sigma^2}$$



# CI FOR $\sigma$ , UNKNOWN $\mu, \sigma$

- $X_1, X_2, \dots, X_{100}$  – samples from  $N(\mu, \sigma^2)$ ,  $\sigma$  is unknown.
- How to construct a CI for  $\mu$ ? Consider  $S^2$ :

$$s^2 = \frac{1}{n-1} \sum_{i=1}^n (X_i - \bar{X})^2$$

$$\frac{(n-1)s^2}{\sigma^2} \sim \chi^2(n-1) -$$

Chi-squared distribution

# CHI-SQUARED DISTRIBUTION

$$Z_1, Z_2, \dots, Z_n \text{ — i.i.d. } N(0,1)$$

# CHI-SQUARED DISTRIBUTION

$$Z_1, Z_2, \dots, Z_n \text{ — i. i. d. } N(0,1)$$

$\Rightarrow$


$$Q = Z_1^2 + \dots + Z_n^2 \sim \chi^2(n - 1)$$

# CHI-SQUARED DISTRIBUTION

$$Z_1, Z_2, \dots, Z_n \text{ — i.i.d. } N(0,1)$$

$\Rightarrow$

$$Q = Z_1^2 + \dots + Z_n^2 \sim \chi^2(n-1)$$



# CI FOR $\sigma$ , UNKNOWN $\mu, \sigma$

- $X_1, X_2, \dots, X_{100}$  – samples from  $N(\mu, \sigma^2)$ ,  $\sigma$  is unknown.
- How to construct a CI for  $\sigma$ ? Consider  $S^2$ :

$$\frac{(n-1)s^2}{\sigma^2} \sim \chi^2(n-1)$$


# CI FOR $\sigma$ , UNKNOWN $\mu, \sigma$

- $X_1, X_2, \dots, X_{100}$  – samples from  $N(\mu, \sigma^2)$ ,  $\sigma$  is unknown.
- How to construct a CI for  $\sigma$ ? Consider  $S^2$ :

$$\frac{(n-1)s^2}{\sigma^2} \sim \chi^2(n-1)$$

$$P\left( < \frac{(n-1)s^2}{\sigma^2} < \right) = 1 - \alpha$$

# CI FOR $\sigma$ , UNKNOWN $\mu, \sigma$



# CI FOR $\sigma$ , UNKNOWN $\mu, \sigma$

- $X_1, X_2, \dots, X_{100}$  – samples from  $N(\mu, \sigma^2)$ ,  $\sigma$  is unknown.
- How to construct a CI for  $\sigma$ ? Consider  $S^2$ :

$$\frac{(n-1)s^2}{\sigma^2} \sim \chi^2(n-1)$$

$$P\left(c_{\alpha/2} < \frac{(n-1)s^2}{\sigma^2} < c_{1-\alpha/2}\right) = 1 - \alpha$$



# CI FOR $\sigma$ , UNKNOWN $\mu, \sigma$

- $X_1, X_2, \dots, X_{100}$  – samples from  $N(\mu, \sigma^2)$ ,  $\sigma$  is unknown.
- How to construct a CI for  $\sigma$ ? Consider  $S^2$ :

$$\frac{(n-1)s^2}{\sigma^2} \sim \chi^2(n-1)$$

$$P\left(c_{\alpha/2} < \frac{(n-1)s^2}{\sigma^2} < c_{1-\alpha/2}\right) = 1 - \alpha$$

$$P\left(\frac{(n-1)s^2}{c_{1-\alpha/2}} < \sigma^2 < \frac{(n-1)s^2}{c_{\alpha/2}}\right) = 1 - \alpha$$

# CI FOR $\sigma$ , UNKNOWN $\mu, \sigma$ : EXAMPLE

- $X_1, X_2, \dots, X_{20}$  — samples from  $N(\mu, \sigma^2)$ ,  $\sigma$  is unknown.
- $\bar{X} = 42, \quad s^2 = 36$

# CI FOR $\sigma$ , UNKNOWN $\mu, \sigma$ : EXAMPLE

- $X_1, X_2, \dots, X_{20}$  — samples from  $N(\mu, \sigma^2)$ ,  $\sigma$  is unknown.
- $\bar{X} = 42, \quad s^2 = 36$
- Give the 95%-CI for  $\sigma$ .

# CI FOR $\sigma$ , UNKNOWN $\mu$ , $\sigma$ : EXAMPLE

- $X_1, X_2, \dots, X_{20}$  – samples from  $N(\mu, \sigma^2)$ ,  $\sigma$  is unknown.
- $\bar{X} = 42, \quad s^2 = 36$
- Give the 95%-CI for  $\sigma$ .

$$\left[ \frac{(n-1)s^2}{c_{1-\alpha/2}}; \frac{(n-1)s^2}{c_{\alpha/2}} \right]$$

# CI FOR $\sigma$ , UNKNOWN $\mu$ , $\sigma$ : EXAMPLE

- $X_1, X_2, \dots, X_{20}$  – samples from  $N(\mu, \sigma^2)$ ,  $\sigma$  is unknown.
- $\bar{X} = 42, \quad s^2 = 36$
- Give the 95%-CI for  $\sigma$ .

$$\left[ \frac{(n-1)s^2}{c_{1-\alpha/2}}; \frac{(n-1)s^2}{c_{\alpha/2}} \right]$$

$$\left[ \frac{19 \cdot 36}{32.85}; \frac{19 \cdot 36}{8.91} \right]$$

# LARGE SAMPLES

Using Central Limit Theorem

# CI FOR LARGE SAMPLES

- Typical task: estimating the mean of a distribution.
- Suppose  $X_1, \dots, X_n$  is drawn from an unknown distribution.
- How to construct a CI?

- CLT:

If  $\mu, \sigma^2 < \infty$  and if  $n$  is sufficiently large, then:

$$\frac{(\bar{X} - \mu)\sqrt{n}}{s} \approx N(0,1)$$

# CI FOR LARGE SAMPLES

- Typical task: estimating the mean of a distribution.
- Suppose  $X_1, \dots, X_n$  is drawn from an unknown distribution.
- How to construct a CI?

- CLT:

If  $\mu, \sigma^2 < \infty$  and if  $n$  is sufficiently large, then:

$$\frac{(\bar{X} - \mu)\sqrt{n}}{s} \approx N(0,1) \Rightarrow \mu \approx$$



# CI FOR LARGE SAMPLES

- Typical task: estimating the mean of a distribution.
- Suppose  $X_1, \dots, X_n$  is drawn from an unknown distribution.
- How to construct a CI?

- CLT:

If  $\mu, \sigma^2 < \infty$  and if  $n$  is sufficiently large, then:

$$\frac{(\bar{X} - \mu)\sqrt{n}}{s} \approx N(0,1) \Rightarrow \mu \approx \bar{X} \pm \frac{s}{\sqrt{n}} z_{1-\alpha/2}$$


# CI FOR BERNOULLI DISTRIBUTION

# POLITICAL POLLS

- Political polls are often reported as a value with a margin-of-error.

*52% favor candidate A with a margin-of-error of  $\pm 5\%$ .*

# Presidential Election Preview: How Clinton, Trump Voters from 2016 Plan to Vote in 2020



A total of 1,510 eligible voters, who are adult members of the USC Dornsife Center for Economic and Social Research's Understanding America Study internet panel, participated from August 11 – 16, 2020. Margin of sampling error for this preliminary sample is +/-3 percentage points. Tracking graphs will be updated every day. For full question text, methodology, and other information, visit <https://uacdata.usc.edu/index.php>.

# POLITICAL POLLS

- Political polls are often reported as a value with a margin-of-error.

*52% favor candidate A with a margin-of-error of  $\pm 5\%$ .*

- The actual precise meaning:

# POLITICAL POLLS

- Political polls are often reported as a value with a margin-of-error.

*52% favor candidate A with a margin-of-error of  $\pm 5\%$ .*

- The actual precise meaning:

*If  $p$  is the proportion of the population that supports A, then*

# POLITICAL POLLS

- Political polls are often reported as a value with a margin-of-error.

*52% favor candidate A with a margin-of-error of  $\pm 5\%$ .*

- The actual precise meaning:

*If  $p$  is the proportion of the population that supports A, then  
the point estimate for  $p$  is  $\hat{p} = 0.52$*

# POLITICAL POLLS

- Political polls are often reported as a value with a margin-of-error.

*52% favor candidate A with a margin-of-error of  $\pm 5\%$ .*

- The actual precise meaning:

*If  $p$  is the proportion of the population that supports A, then  
the point estimate for  $p$  is  $\hat{p} = 0.52$   
and the 95% - CI is  $52\% \pm 5\%$*



# CI FOR P (BERNOULLI)

- Construct a confidence interval for the parameter  $p$  of the Bernoulli distribution based on samples  $X_1, \dots, X_n$ .

# CI FOR P (BERNOULLI)

- Construct a confidence interval for the parameter  $p$  of the Bernoulli distribution based on samples  $X_1, \dots, X_n$ .

$$\hat{p} = \bar{X}$$

# CI FOR P (BERNOULLI)

- Construct a confidence interval for the parameter  $p$  of the Bernoulli distribution based on samples  $X_1, \dots, X_n$ .

$$\text{CLT: } \hat{p} = \bar{X} \approx N\left( \quad , \quad \right)$$

# CI FOR P (BERNOULLI)

- Construct a confidence interval for the parameter  $p$  of the Bernoulli distribution based on samples  $X_1, \dots, X_n$ .

$$\text{CLT: } \hat{p} = \bar{X} \approx N\left(p, \right)$$

# CI FOR P (BERNOULLI)

- Construct a confidence interval for the parameter  $p$  of the Bernoulli distribution based on samples  $X_1, \dots, X_n$ .

$$\text{CLT: } \hat{p} = \bar{X} \approx N\left(p, \frac{p(1-p)}{n}\right)$$

# CI FOR P (BERNOULLI)

- Construct a confidence interval for the parameter  $p$  of the Bernoulli distribution based on samples  $X_1, \dots, X_n$ .

$$\text{CLT: } \hat{p} = \bar{X} \approx N\left(p, \frac{p(1-p)}{n}\right)$$

$$\text{Approximation: } \frac{p(1-p)}{n} \approx \frac{\hat{p}(1-\hat{p})}{n} \text{ (not valid for small samples)}$$

# CI FOR P (BERNOULLI)

- Construct a confidence interval for the parameter  $p$  of the Bernoulli distribution based on samples  $X_1, \dots, X_n$ .

$$\text{CLT: } \hat{p} = \bar{X} \approx N\left(p, \frac{p(1-p)}{n}\right)$$

$$\text{Approximation: } \frac{p(1-p)}{n} \approx \frac{\hat{p}(1-\hat{p})}{n} \text{ (not valid for small samples)}$$

$$\approx N(0,1)$$

# CI FOR P (BERNOULLI)

- Construct a confidence interval for the parameter  $p$  of the Bernoulli distribution based on samples  $X_1, \dots, X_n$ .

$$\text{CLT: } \hat{p} = \bar{X} \approx N\left(p, \frac{p(1-p)}{n}\right)$$

$$\text{Approximation: } \frac{p(1-p)}{n} \approx \frac{\hat{p}(1-\hat{p})}{n} \text{ (not valid for small samples)}$$

$$\frac{(\hat{p} - p)\sqrt{n}}{\sqrt{\hat{p}(1 - \hat{p})}} \approx N(0,1)$$



# CI FOR P (BERNOULLI)

- Construct a confidence interval for the parameter  $p$  of the Bernoulli distribution based on samples  $X_1, \dots, X_n$ .

$$\frac{(\hat{p} - p)\sqrt{n}}{\sqrt{\hat{p}(1 - \hat{p})}} \approx N(0, 1)$$

# CI FOR P (BERNOULLI)

- Construct a confidence interval for the parameter  $p$  of the Bernoulli distribution based on samples  $X_1, \dots, X_n$ .

$$\frac{(\hat{p} - p)\sqrt{n}}{\sqrt{\hat{p}(1 - \hat{p})}} \approx N(0, 1)$$

$$P\left(-z_{1-\alpha/2} < \frac{(\hat{p} - p)\sqrt{n}}{\sqrt{\hat{p}(1 - \hat{p})}} < z_{1-\alpha/2}\right) = 1 - \alpha$$

# CI FOR P (BERNOULLI)

- Construct a confidence interval for the parameter  $p$  of the Bernoulli distribution based on samples  $X_1, \dots, X_n$ .

$$\frac{(\hat{p} - p)\sqrt{n}}{\sqrt{\hat{p}(1 - \hat{p})}} \approx N(0, 1)$$

$$P\left(-z_{1-\alpha/2} < \frac{(\hat{p} - p)\sqrt{n}}{\sqrt{\hat{p}(1 - \hat{p})}} < z_{1-\alpha/2}\right) = 1 - \alpha$$

$$P\left(\hat{p} - z_{1-\alpha/2} \sqrt{\frac{\hat{p}(1 - \hat{p})}{n}} < p < \hat{p} + z_{1-\alpha/2} \sqrt{\frac{\hat{p}(1 - \hat{p})}{n}}\right) = 1 - \alpha$$

# CI FOR P (BERNOULLI)

- Construct a confidence interval for the parameter  $p$  of the Bernoulli distribution based on samples  $X_1, \dots, X_n$ .

$$P\left(\hat{p} - z_{1-\alpha/2} \sqrt{\frac{\hat{p}(1-\hat{p})}{n}} < p < \hat{p} + z_{1-\alpha/2} \sqrt{\frac{\hat{p}(1-\hat{p})}{n}}\right) = 1 - \alpha$$

# CI FOR P (BERNOULLI)

- Construct a confidence interval for the parameter  $p$  of the Bernoulli distribution based on samples  $X_1, \dots, X_n$ .

$$P\left(\hat{p} - z_{1-\alpha/2} \sqrt{\frac{\hat{p}(1-\hat{p})}{n}} < p < \hat{p} + z_{1-\alpha/2} \sqrt{\frac{\hat{p}(1-\hat{p})}{n}}\right) = 1 - \alpha$$

- Example: toss a coin 100 times, 60 heads. CI for  $p$  ( $\alpha = 0.05$ ):

# CI FOR P (BERNOULLI)

- Construct a confidence interval for the parameter  $p$  of the Bernoulli distribution based on samples  $X_1, \dots, X_n$ .

$$P\left(\hat{p} - z_{1-\alpha/2} \sqrt{\frac{\hat{p}(1-\hat{p})}{n}} < p < \hat{p} + z_{1-\alpha/2} \sqrt{\frac{\hat{p}(1-\hat{p})}{n}}\right) = 1 - \alpha$$

- Example: toss a coin 100 times, 60 heads. CI for  $p$  ( $\alpha = 0.05$ ):

$$p \in (0.6 - 1.96 \cdot 0.049 ; 0.6 + 1.96 \cdot 0.049)$$

# CI FOR P (BERNOULLI)

- Construct a confidence interval for the parameter  $p$  of the Bernoulli distribution based on samples  $X_1, \dots, X_n$ .

$$P\left(\hat{p} - z_{1-\alpha/2} \sqrt{\frac{\hat{p}(1-\hat{p})}{n}} < p < \hat{p} + z_{1-\alpha/2} \sqrt{\frac{\hat{p}(1-\hat{p})}{n}}\right) = 1 - \alpha$$

- Example: toss a coin 100 times, 60 heads. CI for  $p$  ( $\alpha = 0.05$ ):

$$p \in (0.6 - 1.96 \cdot 0.049 ; 0.6 + 1.96 \cdot 0.049)$$

$$p \in (0.504; 0.696)$$

# THE 95%-CI: RULE-OF-THUMB

$$p \approx \hat{p} \pm z_{1-\frac{\alpha}{2}} \sqrt{\frac{\hat{p}(1-\hat{p})}{n}}$$



# THE 95%-CI: RULE-OF-THUMB

$$p \approx \hat{p} \pm z_{1-\frac{\alpha}{2}} \sqrt{\frac{\hat{p}(1 - \hat{p})}{n}}$$

What is the maximum possible value of  $\hat{p}(1 - \hat{p})$ ?

# THE 95%-CI: RULE-OF-THUMB

$$p \approx \hat{p} \pm z_{1-\frac{\alpha}{2}} \sqrt{\frac{\hat{p}(1 - \hat{p})}{n}}$$

What is the maximum possible value of  $\hat{p}(1 - \hat{p})$ ? 1/4!

# THE 95%-CI: RULE-OF-THUMB

$$p \approx \hat{p} \pm z_{1-\frac{\alpha}{2}} \sqrt{\frac{\hat{p}(1-\hat{p})}{n}}$$

What is the maximum possible value of  $\hat{p}(1-\hat{p})$ ? 1/4!

$$z_{1-\frac{\alpha}{2}} \sqrt{\frac{\hat{p}(1-\hat{p})}{n}} \leq$$

# THE 95%-CI: RULE-OF-THUMB

$$p \approx \hat{p} \pm z_{1-\frac{\alpha}{2}} \sqrt{\frac{\hat{p}(1-\hat{p})}{n}}$$

What is the maximum possible value of  $\hat{p}(1-\hat{p})$ ? 1/4!

$$z_{1-\frac{\alpha}{2}} \sqrt{\frac{\hat{p}(1-\hat{p})}{n}} \leq z_{1-\frac{\alpha}{2}} \cdot \frac{1}{2\sqrt{n}} =$$

# THE 95%-CI: RULE-OF-THUMB

$$p \approx \hat{p} \pm z_{1-\frac{\alpha}{2}} \sqrt{\frac{\hat{p}(1-\hat{p})}{n}}$$

What is the maximum possible value of  $\hat{p}(1-\hat{p})$ ? 1/4!

$$z_{1-\frac{\alpha}{2}} \sqrt{\frac{\hat{p}(1-\hat{p})}{n}} \leq z_{1-\frac{\alpha}{2}} \cdot \frac{1}{2\sqrt{n}} = 1.96 \cdot \frac{1}{2\sqrt{n}} \approx$$

# THE 95%-CI: RULE-OF-THUMB

$$p \approx \hat{p} \pm z_{1-\frac{\alpha}{2}} \sqrt{\frac{\hat{p}(1-\hat{p})}{n}}$$

What is the maximum possible value of  $\hat{p}(1-\hat{p})$ ? 1/4!

$$z_{1-\frac{\alpha}{2}} \sqrt{\frac{\hat{p}(1-\hat{p})}{n}} \leq z_{1-\frac{\alpha}{2}} \cdot \frac{1}{2\sqrt{n}} = 1.96 \cdot \frac{1}{2\sqrt{n}} \approx \frac{1}{\sqrt{n}}$$

# THE 95%-CI: RULE-OF-THUMB

$$p \approx \hat{p} \pm z_{1-\frac{\alpha}{2}} \sqrt{\frac{\hat{p}(1-\hat{p})}{n}}$$

What is the maximum possible value of  $\hat{p}(1-\hat{p})$ ? 1/4!

$$z_{1-\frac{\alpha}{2}} \sqrt{\frac{\hat{p}(1-\hat{p})}{n}} \leq z_{1-\frac{\alpha}{2}} \cdot \frac{1}{2\sqrt{n}} = 1.96 \cdot \frac{1}{2\sqrt{n}} \approx \frac{1}{\sqrt{n}}$$

$$p \approx$$

# THE 95%-CI: RULE-OF-THUMB

$$p \approx \hat{p} \pm z_{1-\frac{\alpha}{2}} \sqrt{\frac{\hat{p}(1-\hat{p})}{n}}$$

What is the maximum possible value of  $\hat{p}(1-\hat{p})$ ? 1/4!

$$z_{1-\frac{\alpha}{2}} \sqrt{\frac{\hat{p}(1-\hat{p})}{n}} \leq z_{1-\frac{\alpha}{2}} \cdot \frac{1}{2\sqrt{n}} = 1.96 \cdot \frac{1}{2\sqrt{n}} \approx \frac{1}{\sqrt{n}}$$

$$p \approx \hat{p} \pm \frac{1}{\sqrt{n}}$$



# EXAMPLE: POLITICAL POLLS

- Two polls:
  - Fast and First:  
polls 40 random voters and finds 22 support A.
  - Quick but Cautious:  
polls 400 random voters and finds 190 support A.

# EXAMPLE: POLITICAL POLLS

- Two polls:
  - Fast and First:  
polls 40 random voters and finds 22 support A.
  - Quick but Cautious:  
polls 400 random voters and finds 190 support A.
- Let's find the point estimates and 95% rule-of-thumb confidence intervals for each poll:

# EXAMPLE: POLITICAL POLLS

- Two polls:
  - Fast and First:  
polls 40 random voters and finds 22 support A.
  - Quick but Cautious:  
polls 400 random voters and finds 190 support A.
- Let's find the point estimates and 95% rule-of-thumb confidence intervals for each poll:

$$p = \hat{p} \pm \frac{1}{\sqrt{n}}$$

# EXAMPLE: POLITICAL POLLS

- Two polls:
  - Fast and First:  
polls 40 random voters and finds 22 support A.
  - Quick but Cautious:  
polls 400 random voters and finds 190 support A.

- Poll 1:

$$\hat{p} =$$

- Poll 2:

$$\hat{p} =$$

# EXAMPLE: POLITICAL POLLS

- Two polls:
  - Fast and First:  
polls 40 random voters and finds 22 support A.
  - Quick but Cautious:  
polls 400 random voters and finds 190 support A.

- Poll 1:

$$\hat{p} = 22/40$$

- Poll 2:

$$\hat{p} =$$

# EXAMPLE: POLITICAL POLLS

- Two polls:
  - Fast and First:  
polls 40 random voters and finds 22 support A.
  - Quick but Cautious:  
polls 400 random voters and finds 190 support A.

- Poll 1:

$$\hat{p} = 22/40$$

- Poll 2:

$$\hat{p} = 190/400$$

# EXAMPLE: POLITICAL POLLS

- Two polls:
  - Fast and First:  
polls 40 random voters and finds 22 support A.
  - Quick but Cautious:  
polls 400 random voters and finds 190 support A.

- Poll 1:

$$\hat{p} = 22/40 \qquad p = \hat{p} \pm \frac{1}{\sqrt{n}} =$$

- Poll 2:

$$\hat{p} = 190/400 \qquad p = \hat{p} \pm \frac{1}{\sqrt{n}} =$$

# EXAMPLE: POLITICAL POLLS

- Two polls:
  - Fast and First:  
polls 40 random voters and finds 22 support A.
  - Quick but Cautious:  
polls 400 random voters and finds 190 support A.

- Poll 1:

$$\hat{p} = 22/40 \qquad p = \hat{p} \pm \frac{1}{\sqrt{n}} = 0.55 \pm 0.16$$

- Poll 2:

$$\hat{p} = 190/400 \qquad p = \hat{p} \pm \frac{1}{\sqrt{n}} =$$



# EXAMPLE: POLITICAL POLLS

- Two polls:
  - Fast and First:  
polls 40 random voters and finds 22 support A.
  - Quick but Cautious:  
polls 400 random voters and finds 190 support A.

- Poll 1:

$$\hat{p} = 22/40 \qquad p = \hat{p} \pm \frac{1}{\sqrt{n}} = 0.55 \pm 0.16$$

- Poll 2:

$$\hat{p} = 190/400 \qquad p = \hat{p} \pm \frac{1}{\sqrt{n}} = 0.475 \pm 0.05$$

# EXAMPLE: POLITICAL POLLS

- Two polls:
  - Fast and First:  
polls 40 random voters and finds 22 support A.
  - Quick but Cautious:  
polls 400 random voters and finds 190 support A.

- Poll 1:

$$\hat{p} = 22/40 \qquad p = \hat{p} \pm \frac{1}{\sqrt{n}} = 0.55 \pm 0.16$$

- Poll 2:

$$\hat{p} = 190/400 \qquad p = \hat{p} \pm \frac{1}{\sqrt{n}} = 0.475 \pm 0.05$$

# SAMPLE SIZE

*How large should my sample be?*

# SAMPLE SIZE DETERMINATION

- Data collection is difficult.
- How much is 'just enough'?
- Example: estimating CI for the mean,  $\sigma$  is know.

# SAMPLE SIZE DETERMINATION

- Data collection is difficult.
- How much is 'just enough'?
- Example: estimating CI for the mean,  $\sigma$  is know.

$$\mu \in \bar{X} \pm \frac{\sigma}{\sqrt{n}} z_{1-\frac{\alpha}{2}}$$

# SAMPLE SIZE DETERMINATION

- Data collection is difficult.
- How much is 'just enough'?
- Example: estimating CI for the mean,  $\sigma$  is know.

$$\mu \in \bar{X} \pm \frac{\sigma}{\sqrt{n}} z_{1-\frac{\alpha}{2}}$$

# SAMPLE SIZE DETERMINATION

- Data collection is difficult.
- How much is 'just enough'?
- Example: estimating CI for the mean,  $\sigma$  is know.

$$\mu \in \bar{X} \pm \frac{\sigma}{\sqrt{n}} Z_{1-\frac{\alpha}{2}}$$

Limit the width of the interval:  $\frac{\sigma}{\sqrt{n}} Z_{1-\alpha/2} \leq \epsilon$

# SAMPLE SIZE DETERMINATION

- Data collection is difficult.
- How much is 'just enough'?
- Example: estimating CI for the mean,  $\sigma$  is know.

$$\mu \in \bar{X} \pm \frac{\sigma}{\sqrt{n}} Z_{1-\frac{\alpha}{2}}$$

Limit the width of the interval:  $\frac{\sigma}{\sqrt{n}} Z_{1-\alpha/2} \leq \epsilon$

$$n \geq$$



# SAMPLE SIZE DETERMINATION

- Data collection is difficult.
- How much is 'just enough'?
- Example: estimating CI for the mean,  $\sigma$  is know.

$$\mu \in \bar{X} \pm \frac{\sigma}{\sqrt{n}} Z_{1-\frac{\alpha}{2}}$$

Limit the width of the interval:  $\frac{\sigma}{\sqrt{n}} Z_{1-\alpha/2} \leq \epsilon$

$$n \geq \frac{\sigma^2 Z_{1-\alpha/2}^2}{\epsilon^2}$$

# SAMPLE SIZE DETERMINATION

- At a juice factory machine is set up as follows:
  - the average content of juice per bottle equals  $\mu$ ;
  - $\sigma = 5\text{cl}$ .

# SAMPLE SIZE DETERMINATION

- At a juice factory machine is set up as follows:
  - the average content of juice per bottle equals  $\mu$ ;
  - $\sigma = 5\text{cl}$ .
- What sample size is required to estimate the average contents to within 0.5cl at the 95% confidence level?

# SAMPLE SIZE DETERMINATION

- At a juice factory machine is set up as follows:
  - the average content of juice per bottle equals  $\mu$ ;
  - $\sigma = 5\text{cl}$ .
- What sample size is required to estimate the average contents to within 0.5cl at the 95% confidence level?

$$n \geq \frac{\sigma^2 z_{1-\frac{\alpha}{2}}^2}{\epsilon^2} =$$

# SAMPLE SIZE DETERMINATION

- At a juice factory machine is set up as follows:
  - the average content of juice per bottle equals  $\mu$ ;
  - $\sigma = 5\text{cl}$ .
- What sample size is required to estimate the average contents to within  $0.5\text{cl}$  at the 95% confidence level?

$$n \geq \frac{\sigma^2 z_{1-\frac{\alpha}{2}}^2}{\epsilon^2} = \frac{5^2 \cdot 1.96^2}{0.5^2} \approx$$

# SAMPLE SIZE DETERMINATION

- At a juice factory machine is set up as follows:
  - the average content of juice per bottle equals  $\mu$ ;
  - $\sigma = 5\text{cl}$ .
- What sample size is required to estimate the average contents to within  $0.5\text{cl}$  at the 95% confidence level?

$$n \geq \frac{\sigma^2 z_{1-\frac{\alpha}{2}}^2}{\epsilon^2} = \frac{5^2 \cdot 1.96^2}{0.5^2} \approx 384.16$$

# SAMPLE SIZE DETERMINATION

- At a juice factory machine is set up as follows:
  - the average content of juice per bottle equals  $\mu$ ;
  - $\sigma = 5\text{cl}$ .
- What sample size is required to estimate the average contents to within 0.5cl at the 95% confidence level?

$$n \geq \frac{\sigma^2 z_{1-\frac{\alpha}{2}}^2}{\epsilon^2} = \frac{5^2 \cdot 1.96^2}{0.5^2} \approx 384.16$$

$$n^* = 385$$

# SAMPLE SIZE DETERMINATION

- Similar reasoning applies to the CI for Bernoulli.
- *How many customers should be surveyed in order to estimate the share of satisfied customers to within 3% at the 95% confidence level?*



# SAMPLE SIZE DETERMINATION

- Similar reasoning applies to the CI for Bernoulli.
- *How many customers should be surveyed in order to estimate the share of satisfied customers to within 3% at the 95% confidence level?*

$$\hat{p} \pm z_{1-\frac{\alpha}{2}} \sqrt{\frac{\hat{p}(1-\hat{p})}{n}}$$

# SAMPLE SIZE DETERMINATION

- Similar reasoning applies to the CI for Bernoulli.
- *How many customers should be surveyed in order to estimate the share of satisfied customers to within 3% at the 95% confidence level?*

$$\hat{p} \pm z_{1-\frac{\alpha}{2}} \sqrt{\frac{p_0(1-p_0)}{n}}$$

# SAMPLE SIZE DETERMINATION

- Similar reasoning applies to the CI for Bernoulli.
- *How many customers should be surveyed in order to estimate the share of satisfied customers to within 3% at the 95% confidence level?*

$$\hat{p} \pm z_{1-\frac{\alpha}{2}} \sqrt{\frac{p_0(1-p_0)}{n}}$$

Let  $p_0 = 0.5$  – the worst possible variance

# SAMPLE SIZE DETERMINATION

- Similar reasoning applies to the CI for Bernoulli.
- *How many customers should be surveyed in order to estimate the share of satisfied customers to within 3% at the 95% confidence level?*

$$\hat{p} \pm z_{1-\frac{\alpha}{2}} \sqrt{\frac{p_0(1-p_0)}{n}}$$

Let  $p_0 = 0.5$  – the worst possible variance

$$1.96 \sqrt{\frac{0.25}{n}} \leq 0.03 \Leftrightarrow$$

# SAMPLE SIZE DETERMINATION

- Similar reasoning applies to the CI for Bernoulli.
- *How many customers should be surveyed in order to estimate the share of satisfied customers to within 3% at the 95% confidence level?*

$$\hat{p} \pm z_{1-\frac{\alpha}{2}} \sqrt{\frac{p_0(1-p_0)}{n}}$$

Let  $p_0 = 0.5$  – the worst possible variance

$$1.96 \sqrt{\frac{0.25}{n}} \leq 0.03 \Leftrightarrow n \geq 1067.1$$