

# PROBABILITY & STATISTICS

Lecture 3 – Conditional probability

# LAST TIME

- Frequentist interpretation of probability
  - Python
  - Practice problems

# TODAY

- Multiple events
- Conditional probability
  - The law of total probability
  - Bayes Rule

# FLIPPING A COIN 10 TIMES

- Which outcome you think is more likely?

HEADS, TAILS, HEADS, TAILS, TAILS,  
HEADS, TAILS, HEADS, HEADS, TAILS

HEADS, HEADS, HEADS, HEADS, HEADS,  
HEADS, HEADS, HEADS, HEADS, HEADS

HEADS, HEADS, HEADS, HEADS, HEADS,  
TAILS, TAILS, TAILS, TAILS, TAILS

# Independent events

# INDEPENDENT EVENTS

- Two events  $A$  and  $B$  from the same sample space  $S$  are independent if

$$P(A \& B) = P(A) \cdot P(B)$$

# ROLLING DICE

- Two dice are rolled. Let  $E$  be the event that sum 7 is rolled on the two dice and let  $F$  be the event that the first die is 1. Are  $E$  and  $F$  independent events?

# ROLLING DICE

- Two dice are rolled. Let  $E$  be the event that sum 7 is rolled on the two dice and let  $F$  be the event that the first die is 1. Are  $E$  and  $F$  independent events?

$$P(E) = \frac{6}{6 \cdot 6} = \frac{1}{6},$$



# ROLLING DICE

- Two dice are rolled. Let  $E$  be the event that sum 7 is rolled on the two dice and let  $F$  be the event that the first die is 1. Are  $E$  and  $F$  independent events?

$$P(E) = \frac{6}{6 \cdot 6} = \frac{1}{6}, \quad P(F) = \frac{1}{6},$$

# ROLLING DICE

- Two dice are rolled. Let  $E$  be the event that sum 7 is rolled on the two dice and let  $F$  be the event that the first die is 1. Are  $E$  and  $F$  independent events?

$$P(E) = \frac{6}{6 \cdot 6} = \frac{1}{6}, \quad P(F) = \frac{1}{6}, \quad P(EF) =$$

# ROLLING DICE

- Two dice are rolled. Let  $E$  be the event that sum 7 is rolled on the two dice and let  $F$  be the event that the first die is 1. Are  $E$  and  $F$  independent events?

$$P(E) = \frac{6}{6 \cdot 6} = \frac{1}{6}, \quad P(F) = \frac{1}{6}, \quad P(EF) = \frac{1}{6 \cdot 6} = \frac{1}{36}$$

# ROLLING DICE

- Two dice are rolled. Let  $E$  be the event that sum 7 is rolled on the two dice and let  $F$  be the event that the first die is 1. Are  $E$  and  $F$  independent events?

$$P(E) = \frac{6}{6 \cdot 6} = \frac{1}{6}, \quad P(F) = \frac{1}{6}, \quad P(EF) = \frac{1}{6 \cdot 6} = \frac{1}{36}$$

$$P(EF) = P(E) \cdot P(F) = \frac{1}{36}$$

# ROLLING DICE

- Two dice are rolled. Let  $E$  be the event that sum 7 is rolled on the two dice and let  $F$  be the event that the first die is 1. Are  $E$  and  $F$  independent events?

$$P(E) = \frac{6}{6 \cdot 6} = \frac{1}{6}, \quad P(F) = \frac{1}{6}, \quad P(EF) = \frac{1}{6 \cdot 6} = \frac{1}{36}$$

$$P(EF) = P(E) \cdot P(F) = \frac{1}{36} \rightarrow E \& F \text{ are independent!}$$

# Practice problems

Google classroom -> Day 3

# FLIPPING A COIN

- A coin is flipped 6 times. Let  $E$  be the event that heads and tails come up an equal number of times. Let  $F$  be the event that heads and tails come up once each during the first two flips of the coin. Are  $E$  and  $F$  independent events?

# FLIPPING A COIN

- A coin is flipped 6 times. Let  $E$  be the event that heads and tails come up an equal number of times. Let  $F$  be the event that heads and tails come up once each during the first two flips of the coin. Are  $E$  and  $F$  independent events?

$$P(E) =$$

$$P(F) =$$

$$P(EF) =$$



# FLIPPING A COIN

- A coin is flipped 6 times. Let  $E$  be the event that heads and tails come up an equal number of times. Let  $F$  be the event that heads and tails come up once each during the first two flips of the coin. Are  $E$  and  $F$  independent events?

$$P(E) = \frac{C(6,3)}{2^6} = \frac{5}{2^4}, \quad P(F) =$$

$$P(EF) =$$

# FLIPPING A COIN

- A coin is flipped 6 times. Let  $E$  be the event that heads and tails come up an equal number of times. Let  $F$  be the event that heads and tails come up once each during the first two flips of the coin. Are  $E$  and  $F$  independent events?

$$P(E) = \frac{C(6,3)}{2^6} = \frac{5}{2^4}, \quad P(F) = \frac{2 \cdot 2^4}{2^6} = \frac{1}{2},$$

$$P(EF) =$$

# FLIPPING A COIN

- A coin is flipped 6 times. Let  $E$  be the event that heads and tails come up an equal number of times. Let  $F$  be the event that heads and tails come up once each during the first two flips of the coin. Are  $E$  and  $F$  independent events?

$$P(E) = \frac{C(6,3)}{2^6} = \frac{5}{2^4}, \quad P(F) = \frac{2 \cdot 2^4}{2^6} = \frac{1}{2},$$

$$P(EF) = \frac{2 \cdot C(4,2)}{2^6} = \frac{3}{2^4}$$

# FLIPPING A COIN

- A coin is flipped 6 times. Let  $E$  be the event that heads and tails come up an equal number of times. Let  $F$  be the event that heads and tails come up once each during the first two flips of the coin. Are  $E$  and  $F$  independent events?

$$P(E) = \frac{C(6,3)}{2^6} = \frac{5}{2^4}, \quad P(F) = \frac{2 \cdot 2^4}{2^6} = \frac{1}{2},$$

$$P(EF) = \frac{2 \cdot C(4,2)}{2^6} = \frac{3}{2^4} \neq P(E) \cdot P(F) = \frac{5}{2^5}$$

# FLIPPING A COIN

- A coin is flipped 6 times. Let  $E$  be the event that heads and tails come up an equal number of times. Let  $F$  be the event that heads and tails come up once each during the first two flips of the coin. Are  $E$  and  $F$  independent events?

$$P(E) = \frac{C(6,3)}{2^6} = \frac{5}{2^4}, \quad P(F) = \frac{2 \cdot 2^4}{2^6} = \frac{1}{2},$$

$$P(EF) = \frac{2 \cdot C(4,2)}{2^6} = \frac{3}{2^4} \neq P(E) \cdot P(F) = \frac{5}{2^5}$$

→  $E$  &  $F$  aren't independent

# FLIPPING A COIN 2

- Let  $E$  be the event that when a coin is flipped three times, we don't get all heads or all tails. Let  $F$  be the event that when a coin is flipped three times, heads comes up at most once. Are  $E$  and  $F$  independent events?

$$P(E) =$$

$$P(F) =$$

$$P(EF) =$$

# FLIPPING A COIN 2

- Let  $E$  be the event that when a coin is flipped three times, we don't get all heads or all tails. Let  $F$  be the event that when a coin is flipped three times, heads comes up at most once. Are  $E$  and  $F$  independent events?

$$P(E) = 1 - 2 \cdot \frac{1}{2^3} = \frac{3}{4}$$

$$P(F) =$$

$$P(EF) =$$

# FLIPPING A COIN 2

- Let  $E$  be the event that when a coin is flipped three times, we don't get all heads or all tails. Let  $F$  be the event that when a coin is flipped three times, heads comes up at most once. Are  $E$  and  $F$  independent events?

$$P(E) = 1 - 2 \cdot \frac{1}{2^3} = \frac{3}{4}$$

$$P(F) = P(0H) + P(1H) =$$

$$P(EF) =$$



# FLIPPING A COIN 2

- Let  $E$  be the event that when a coin is flipped three times, we don't get all heads or all tails. Let  $F$  be the event that when a coin is flipped three times, heads comes up at most once. Are  $E$  and  $F$  independent events?

$$P(E) = 1 - 2 \cdot \frac{1}{2^3} = \frac{3}{4}$$

$$P(F) = P(0H) + P(1H) = \frac{1}{2^3} + \frac{3}{2^3} = \frac{1}{2}$$

$$P(EF) =$$

# FLIPPING A COIN 2

- Let  $E$  be the event that when a coin is flipped three times, we don't get all heads or all tails. Let  $F$  be the event that when a coin is flipped three times, heads comes up at most once. Are  $E$  and  $F$  independent events?

$$P(E) = 1 - 2 \cdot \frac{1}{2^3} = \frac{3}{4}$$

$$P(F) = P(0H) + P(1H) = \frac{1}{2^3} + \frac{3}{2^3} = \frac{1}{2}$$

$$P(EF) = P(1H) =$$

# FLIPPING A COIN 2

- Let  $E$  be the event that when a coin is flipped three times, we don't get all heads or all tails. Let  $F$  be the event that when a coin is flipped three times, heads comes up at most once. Are  $E$  and  $F$  independent events?

$$P(E) = 1 - 2 \cdot \frac{1}{2^3} = \frac{3}{4}$$

$$P(F) = P(0H) + P(1H) = \frac{1}{2^3} + \frac{3}{2^3} = \frac{1}{2}$$

$$P(EF) = P(1H) = \frac{3}{2^3} = P(E) \cdot P(F)$$

# FLIPPING A COIN 2

- Let  $E$  be the event that when a coin is flipped three times, we don't get all heads or all tails. Let  $F$  be the event that when a coin is flipped three times, heads comes up at most once. Are  $E$  and  $F$  independent events?

$$P(E) = 1 - 2 \cdot \frac{1}{2^3} = \frac{3}{4}$$

$$P(F) = P(0H) + P(1H) = \frac{1}{2^3} + \frac{3}{2^3} = \frac{1}{2}$$

$$P(EF) = P(1H) = \frac{3}{2^3} = P(E) \cdot P(F)$$

→  $E$  &  $F$  are independent!

# FACULTY BOARD

- Faculty board, consisting of 15 professors (10 men and 5 women) and 10 students (5 men and 5 women), is randomly choosing a candidate to represent the faculty at the visiting day.

Are the following events independent?

$$E_1 = \{man\}, \quad E_2 = \{professor\}$$

# FACULTY BOARD

- Faculty board, consisting of 15 professors (10 men and 5 women) and 10 students (5 men and 5 women), is randomly choosing a candidate to represent the faculty at the visiting day.

Are the following events independent?

$$E_1 = \{man\}, \quad E_2 = \{professor\}$$

$$P(E_1) =$$

# FACULTY BOARD

- Faculty board, consisting of 15 professors (10 men and 5 women) and 10 students (5 men and 5 women), is randomly choosing a candidate to represent the faculty at the visiting day.

Are the following events independent?

$$E_1 = \{man\}, \quad E_2 = \{professor\}$$

$$P(E_1) = \frac{15}{25} = \frac{3}{5}$$

# FACULTY BOARD

- Faculty board, consisting of 15 professors (10 men and 5 women) and 10 students (5 men and 5 women), is randomly choosing a candidate to represent the faculty at the visiting day.

Are the following events independent?

$$E_1 = \{man\}, \quad E_2 = \{professor\}$$

$$P(E_1) = \frac{15}{25} = \frac{3}{5}, \quad P(E_2) =$$



# FACULTY BOARD

- Faculty board, consisting of 15 professors (10 men and 5 women) and 10 students (5 men and 5 women), is randomly choosing a candidate to represent the faculty at the visiting day.

Are the following events independent?

$$E_1 = \{man\}, \quad E_2 = \{professor\}$$

$$P(E_1) = \frac{15}{25} = \frac{3}{5}, \quad P(E_2) = \frac{15}{25} = \frac{3}{5}$$

# FACULTY BOARD

- Faculty board, consisting of 15 professors (10 men and 5 women) and 10 students (5 men and 5 women), is randomly choosing a candidate to represent the faculty at the visiting day.

Are the following events independent?

$$E_1 = \{man\}, \quad E_2 = \{professor\}$$

$$P(E_1) = \frac{15}{25} = \frac{3}{5}, \quad P(E_2) = \frac{15}{25} = \frac{3}{5}, \quad P(E_1 \cap E_2) =$$

# FACULTY BOARD

- Faculty board, consisting of 15 professors (10 men and 5 women) and 10 students (5 men and 5 women), is randomly choosing a candidate to represent the faculty at the visiting day.

Are the following events independent?

$$E_1 = \{man\}, \quad E_2 = \{professor\}$$

$$P(E_1) = \frac{15}{25} = \frac{3}{5}, \quad P(E_2) = \frac{15}{25} = \frac{3}{5}, \quad P(E_1 \cap E_2) = \frac{10}{25} = \frac{2}{5}$$

# FACULTY BOARD

- Faculty board, consisting of 15 professors (10 men and 5 women) and 10 students (5 men and 5 women), is randomly choosing a candidate to represent the faculty at the visiting day.

Are the following events independent?

$$E_1 = \{man\}, \quad E_2 = \{professor\}$$

$$P(E_1) = \frac{15}{25} = \frac{3}{5}, \quad P(E_2) = \frac{15}{25} = \frac{3}{5}, \quad P(E_1 \cap E_2) = \frac{10}{25} = \frac{2}{5}$$

$$P(E_1 \cap E_2) \quad P(E_1) \cdot P(E_2)$$

# FACULTY BOARD

- Faculty board, consisting of 15 professors (10 men and 5 women) and 10 students (5 men and 5 women), is randomly choosing a candidate to represent the faculty at the visiting day.

Are the following events independent?

$$E_1 = \{man\}, \quad E_2 = \{professor\}$$

$$P(E_1) = \frac{15}{25} = \frac{3}{5}, \quad P(E_2) = \frac{15}{25} = \frac{3}{5}, \quad P(E_1 \cap E_2) = \frac{10}{25} = \frac{2}{5}$$

$$P(E_1 \cap E_2) \neq P(E_1) \cdot P(E_2)$$

# FACULTY BOARD

- Faculty board, consisting of 15 professors (10 men and 5 women) and 10 students (5 men and 5 women), is randomly choosing a candidate to represent the faculty at the visiting day.

Are the following events independent?

$$E_1 = \{man\}, \quad E_2 = \{professor\}$$

$$P(E_1) = \frac{15}{25} = \frac{3}{5}, \quad P(E_2) = \frac{10}{25} = \frac{2}{5}, \quad P(E_1 \cap E_2) = \frac{10}{25} = \frac{2}{5}$$

$P(E_1 \cap E_2) \neq P(E_1) \cdot P(E_2) \rightarrow$   
the events are not independent

# WEATHER FORECAST

- There is a 50% chance of rain on Saturday and a 50% chance of rain on Sunday. If having rain on Saturday and having rain on Sunday are independent events, then what is the probability of having no rain over the weekend?

# WEATHER FORECAST

- There is a 50% chance of rain on Saturday and a 50% chance of rain on Sunday. If having rain on Saturday and having rain on Sunday are independent events, then what is the probability of having no rain over the weekend?

$$P(Sat \& Sun) =$$



# WEATHER FORECAST

- There is a 50% chance of rain on Saturday and a 50% chance of rain on Sunday. If having rain on Saturday and having rain on Sunday are independent events, then what is the probability of having no rain over the weekend?

$$P(Sat \& Sun) = P(Sat) \cdot P(Sun) =$$

# WEATHER FORECAST

- There is a 50% chance of rain on Saturday and a 50% chance of rain on Sunday. If having rain on Saturday and having rain on Sunday are independent events, then what is the probability of having no rain over the weekend?

$$P(Sat \& Sun) = P(Sat) \cdot P(Sun) = \frac{1}{4}$$

# Conditional probability

# EXAMPLE: DEMOGRAPHICS

- In Czechia, 46% of the population is male, and 54% is female.
- 14% of the population aged 65+.
- Among them, 31% are male and 69% is female.

# EXAMPLE: DEMOGRAPHICS

- In Czechia, 46% of the population is male, and 54% is female.
- 14% of the population aged 65+.
- Among them, 31% are male and 69% is female.
- So, men aged 65+ are % of the population.

# EXAMPLE: DEMOGRAPHICS

- In Czechia, 46% of the population is male, and 54% is female.
- 14% of the population aged 65+.
- Among them, 31% are male and 69% is female.
- So, men aged 65+ are 4.3% ( $0.31 \cdot 0.14 = 0.0434$ ) of the population.

# EXAMPLE: DEMOGRAPHICS

- In Czechia, 46% of the population is male, and 54% is female.
- The probability to be male is 0.46.
- 14% of the population aged 65+.
- Among them, 31% are male and 69% is female.
- So, men aged 65+ are 4.3% ( $0.31 \cdot 0.14 = 0.0434$ ) of the population.

# EXAMPLE: DEMOGRAPHICS

- In Czechia, 46% of the population is male, and 54% is female.
- 14% of the population aged 65+.
- Among them, 31% are male and 69% is female.
- So, men aged 65+ are 4.3% ( $0.31 \cdot 0.14 = 0.0434$ ) of the population.
- The probability to be male is 0.46.
- The probability to be aged 65+ is 0.14.



# EXAMPLE: DEMOGRAPHICS

- In Czechia, 46% of the population is male, and 54% is female.
- 14% of the population aged 65+.
- Among them, 31% are male and 69% is female.
- So, men aged 65+ are 4.3% ( $0.31 \cdot 0.14 = 0.0434$ ) of the population.
- The probability to be male is 0.46.
- The probability to be aged 65+ is 0.14.
- *Joint probability* of being a man **and** aged 65+ is 0.0434.

# EXAMPLE: DEMOGRAPHICS

- In Czechia, 46% of the population is male, and 54% is female.
- 14% of the population aged 65+.
- Among them, 31% are male and 69% is female.
- So, men aged 65+ are 4.3% ( $0.31 \cdot 0.14 = 0.0434$ ) of the population.
- The probability to be male is 0.46.
- The probability to be aged 65+ is 0.14.
- The *conditional probability* of being a male **given** age 65+ is 0.31.
- *Joint probability* of being a man **and** aged 65+ is 0.0434.

# EXAMPLE: DEMOGRAPHICS

- The probability to be male is 0.46:
- The probability to be aged 65+ is 0.14:

# EXAMPLE: DEMOGRAPHICS

- The probability to be male is 0.46:  $P(M) = 0.46$
- The probability to be aged 65+ is 0.14:  $P(65+) = 0.14$

# EXAMPLE: DEMOGRAPHICS

- The probability to be male is 0.46:  $P(M) = 0.46$
- The probability to be aged 65+ is 0.14:  $P(65+) = 0.14$
- The *conditional probability* of being a male **given** age 65+ is 0.31:

# EXAMPLE: DEMOGRAPHICS

- The probability to be male is 0.46:  $P(M) = 0.46$
- The probability to be aged 65+ is 0.14:  $P(65+) = 0.14$
- The *conditional probability* of being a male **given** age 65+ is 0.31:

$$P(M|65+) = 0.31$$

# EXAMPLE: DEMOGRAPHICS

- The probability to be male is 0.46:  $P(M) = 0.46$
- The probability to be aged 65+ is 0.14:  $P(65+) = 0.14$
- The *conditional probability* of being a male **given** age 65+ is 0.31:

$$P(M|65+) = 0.31$$

- *Joint probability* of being a man **and** aged 65+ is 0.0434:

# EXAMPLE: DEMOGRAPHICS

- The probability to be male is 0.46:  $P(M) = 0.46$
- The probability to be aged 65+ is 0.14:  $P(65+) = 0.14$
- The *conditional probability* of being a male **given** age 65+ is 0.31:

$$P(M|65+) = 0.31$$

- *Joint probability* of being a man **and** aged 65+ is 0.0434:

$$P(M \& 65+) = 0.0434$$



# EXAMPLE: DEMOGRAPHICS

- The probability to be male is 0.46:  $P(M) = 0.46$
- The probability to be aged 65+ is 0.14:  $P(65+) = 0.14$
- The *conditional probability* of being a male **given** age 65+ is 0.31:

$$P(M|65+) = 0.31$$

- *Joint probability* of being a man **and** aged 65+ is 0.0434:

$$P(M \& 65+) = 0.0434$$

Note:  $P(M \& 65+) = P(M|65+) \cdot P(65+)$

# EXAMPLE: ROLLING A DIE

- A man rolls a pair of dice. For the game, it is important that he rolls a sum of 7. What's the probability of that?

# EXAMPLE: ROLLING A DIE

- A man rolls a pair of dice. For the game, it is important that he rolls a sum of 7. What's the probability of that?

$$P(X_1 + X_2 = 7) =$$

# EXAMPLE: ROLLING A DIE

- A man rolls a pair of dice. For the game, it is important that he rolls a sum of 7. What's the probability of that?

$$P(X_1 + X_2 = 7) = \frac{6}{6 \cdot 6} = \frac{1}{6} \approx 0.167$$

# EXAMPLE: ROLLING A DIE

- A man rolls a pair of dice. For the game, it is important that he rolls a sum of 7. What's the probability of that?

$$P(X_1 + X_2 = 7) = \frac{6}{6 \cdot 6} = \frac{1}{6} \approx 0.167$$

- The other player is not wearing her glasses and is unable to see the outcome, but sees that there is more than one spot on each die. What is the probability of getting 7 now?

# EXAMPLE: ROLLING A DIE

- A man rolls a pair of dice. For the game, it is important that he rolls a sum of 7. What's the probability of that?

$$P(X_1 + X_2 = 7) = \frac{6}{6 \cdot 6} = \frac{1}{6} \approx 0.167$$

- The other player is not wearing her glasses and is unable to see the outcome, but sees that there is more than one spot on each die. What is the probability of getting 7 now?

$$P(X_1 + X_2 = 7 | \text{multiple spots}) =$$

# EXAMPLE: ROLLING A DIE

- A man rolls a pair of dice. For the game, it is important that he rolls a sum of 7. What's the probability of that?

$$P(X_1 + X_2 = 7) = \frac{6}{6 \cdot 6} = \frac{1}{6} \approx 0.167$$

- The other player is not wearing her glasses and is unable to see the outcome, but sees that there is more than one spot on each die. What is the probability of getting 7 now?

$$P(X_1 + X_2 = 7 | \text{multiple spots}) = \frac{4}{5 \cdot 5} =$$

# EXAMPLE: ROLLING A DIE

- A man rolls a pair of dice. For the game, it is important that he rolls a sum of 7. What's the probability of that?

$$P(X_1 + X_2 = 7) = \frac{6}{6 \cdot 6} = \frac{1}{6} \approx 0.167$$

- The other player is not wearing her glasses and is unable to see the outcome, but sees that there is more than one spot on each die. What is the probability of getting 7 now?

$$P(X_1 + X_2 = 7 | \text{multiple spots}) = \frac{4}{5 \cdot 5} = \frac{4}{25} = 0.16$$



# CONDITIONAL PROBABILITY

- Let  $A$  and  $B$  be events in a sample space  $S$  with  $P(B) > 0$ .

# CONDITIONAL PROBABILITY

- Let  $A$  and  $B$  be events in a sample space  $S$  with  $P(B) > 0$ .
- Then the conditional probability of the event  $A$  given that event  $B$  has occurred is

$$P(A|B) = \frac{P(A \cap B)}{P(B)}$$

# CONDITIONAL PROBABILITY

- Let  $A$  and  $B$  be events in a sample space  $S$  with  $P(B) > 0$ .
- Then the conditional probability of the event  $A$  given that event  $B$  has occurred is

$$P(A|B) = \frac{P(A \cap B)}{P(B)} = \frac{P(AB)}{P(B)} = \frac{P(A \& B)}{P(B)}$$

# FLIPPING A COIN 1

- A coin is flipped 4 times. What is the probability that heads comes up exactly twice given that heads comes up on the first flip?

# FLIPPING A COIN 1

- A coin is flipped 4 times. What is the probability that heads comes up exactly twice given that heads comes up on the first flip?

$A$  — H comes up twice

$B$  — H comes up on the 1<sup>st</sup> flip

# FLIPPING A COIN 1

- A coin is flipped 4 times. What is the probability that heads comes up exactly twice given that heads comes up on the first flip?

$A$  – H comes up twice

$B$  – H comes up on the 1<sup>st</sup> flip

$AB$  – H comes up twice incl. on the 1<sup>st</sup> flip

# FLIPPING A COIN 1

- A coin is flipped 4 times. What is the probability that heads comes up exactly twice given that heads comes up on the first flip?

$A$  – H comes up twice

$P(A) =$

$B$  – H comes up on the 1<sup>st</sup> flip

$AB$  – H comes up twice incl. on the 1<sup>st</sup> flip

# FLIPPING A COIN 1

- A coin is flipped 4 times. What is the probability that heads comes up exactly twice given that heads comes up on the first flip?

$A$  – H comes up twice

$$P(A) = \frac{C(4,2)}{2^4} = \frac{3}{8}$$

$B$  – H comes up on the 1<sup>st</sup> flip

$AB$  – H comes up twice incl. on the 1<sup>st</sup> flip



# FLIPPING A COIN 1

- A coin is flipped 4 times. What is the probability that heads comes up exactly twice given that heads comes up on the first flip?

$A$  – H comes up twice

$$P(A) = \frac{C(4,2)}{2^4} = \frac{3}{8}$$

$B$  – H comes up on the 1<sup>st</sup> flip

$$P(B) =$$

$AB$  – H comes up twice incl. on the 1<sup>st</sup> flip

# FLIPPING A COIN 1

- A coin is flipped 4 times. What is the probability that heads comes up exactly twice given that heads comes up on the first flip?

$A$  – H comes up twice

$$P(A) = \frac{C(4,2)}{2^4} = \frac{3}{8}$$

$B$  – H comes up on the 1<sup>st</sup> flip

$$P(B) = \frac{2^3}{2^4} = \frac{1}{2}$$

$AB$  – H comes up twice incl. on the 1<sup>st</sup> flip

# FLIPPING A COIN 1

- A coin is flipped 4 times. What is the probability that heads comes up exactly twice given that heads comes up on the first flip?

$A$  – H comes up twice

$$P(A) = \frac{C(4,2)}{2^4} = \frac{3}{8}$$

$B$  – H comes up on the 1<sup>st</sup> flip

$$P(B) = \frac{2^3}{2^4} = \frac{1}{2}$$

$AB$  – H comes up twice incl. on the 1<sup>st</sup> flip

$$P(AB) =$$

# FLIPPING A COIN 1

- A coin is flipped 4 times. What is the probability that heads comes up exactly twice given that heads comes up on the first flip?

$A$  – H comes up twice

$$P(A) = \frac{C(4,2)}{2^4} = \frac{3}{8}$$

$B$  – H comes up on the 1<sup>st</sup> flip

$$P(B) = \frac{2^3}{2^4} = \frac{1}{2}$$

$AB$  – H comes up twice incl. on the 1<sup>st</sup> flip

$$P(AB) = \frac{C(3,1)}{2^4} = \frac{3}{16}$$

# FLIPPING A COIN 1

- A coin is flipped 4 times. What is the probability that heads comes up exactly twice given that heads comes up on the first flip?

$A$  – H comes up twice

$$P(A) = \frac{C(4,2)}{2^4} = \frac{3}{8}$$

$B$  – H comes up on the 1<sup>st</sup> flip

$$P(B) = \frac{2^3}{2^4} = \frac{1}{2}$$

$AB$  – H comes up twice incl. on the 1<sup>st</sup> flip

$$P(AB) = \frac{C(3,1)}{2^4} = \frac{3}{16}$$

$$P(A|B) = \frac{P(AB)}{P(B)} =$$

# FLIPPING A COIN 1

- A coin is flipped 4 times. What is the probability that heads comes up exactly twice given that heads comes up on the first flip?

$A$  – H comes up twice

$$P(A) = \frac{C(4,2)}{2^4} = \frac{3}{8}$$

$B$  – H comes up on the 1<sup>st</sup> flip

$$P(B) = \frac{2^3}{2^4} = \frac{1}{2}$$

$AB$  – H comes up twice incl. on the 1<sup>st</sup> flip

$$P(AB) = \frac{C(3,1)}{2^4} = \frac{3}{16}$$

$$P(A|B) = \frac{P(AB)}{P(B)} = \frac{3 \cdot 2}{16 \cdot 1} = \frac{3}{8}$$

# BOWLS AND BALLS

- There are two bowls: bowl A contains 2 blue and 3 red balls, and bowl B contains 1 blue and 4 red balls. You randomly chose a bowl and then pick one ball from it.
- What is the probability of picking a blue ball?
- What is the probability of picking a blue ball if we chose bowl A?
- What is the probability of picking a blue ball if we chose bowl B?

# BOWLS AND BALLS

- There are two bowls: bowl A contains 2 blue and 3 red balls, and bowl B contains 1 blue and 4 red balls. You randomly chose a bowl and then pick one ball from it.
- What is the probability of picking a blue ball?

- What is the probability of picking a blue ball if we chose bowl A?

$$P(\text{Blue}|A) =$$

- What is the probability of picking a blue ball if we chose bowl B?



# BOWLS AND BALLS

- There are two bowls: bowl A contains 2 blue and 3 red balls, and bowl B contains 1 blue and 4 red balls. You randomly chose a bowl and then pick one ball from it.
- What is the probability of picking a blue ball?
- What is the probability of picking a blue ball if we chose bowl A?
$$P(\text{Blue}|A) = \frac{2}{5}$$
- What is the probability of picking a blue ball if we chose bowl B?

# BOWLS AND BALLS

- There are two bowls: bowl A contains 2 blue and 3 red balls, and bowl B contains 1 blue and 4 red balls. You randomly chose a bowl and then pick one ball from it.
- What is the probability of picking a blue ball?

- What is the probability of picking a blue ball if we chose bowl A?

$$P(\text{Blue}|A) = \frac{2}{5}$$

- What is the probability of picking a blue ball if we chose bowl B?

$$P(\text{Blue}|B) =$$

# BOWLS AND BALLS

- There are two bowls: bowl A contains 2 blue and 3 red balls, and bowl B contains 1 blue and 4 red balls. You randomly chose a bowl and then pick one ball from it.
- What is the probability of picking a blue ball?

- What is the probability of picking a blue ball if we chose bowl A?

$$P(\text{Blue}|A) = \frac{2}{5}$$



- What is the probability of picking a blue ball if we chose bowl B?

$$P(\text{Blue}|B) = \frac{1}{5}$$



# The law of total probability

[https://youtu.be/U3\\_783xznQI](https://youtu.be/U3_783xznQI)

# FRUITS





# FRUITS



- What is the probability to pick an apple given that you've chosen box A?



# FRUITS



- What is the probability to pick an apple given that you've chosen box A?

$$P(\text{apple}|A) =$$

# FRUITS





- What is the probability to pick an apple given that you've chosen box A?

$$P(\text{apple}|A) = \frac{5}{10} = 0.5$$





# FRUITS



- What is the probability to pick an apple given that you've chosen box B?



# FRUITS



- What is the probability to pick an apple given that you've chosen box B?

$$P(\text{apple}|B) = \frac{3}{12} = 0.25$$

# FRUITS




- What is the probability to pick an apple?

# THE LAW OF TOTAL PROBABILITY

Suppose that the sample space  $S$  is split into  $n$  disjoint events:

$$S = E_1 \cup E_2 \cup \cdots \cup E_n,$$

$$E_1 \cap E_2 \cap \cdots \cap E_n = \emptyset$$



# THE LAW OF TOTAL PROBABILITY


Suppose that the sample space  $S$  is split into  $n$  disjoint events:

$$S = E_1 \cup E_2 \cup \cdots \cup E_n,$$



$$E_1 \cap E_2 \cap \cdots \cap E_n = \emptyset$$

Then  $P(A)$  can be computed as follows:

$$\begin{aligned} P(A) &= P(A, E_1) + P(A, E_2) + \cdots + P(A, E_n) = \\ &= P(A|E_1) \cdot P(E_1) + P(A|E_2) \cdot P(E_2) + \\ &\quad + \cdots + P(A|E_n) \cdot P(E_n) \end{aligned}$$





# FRUITS



- What is the probability to pick an apple?

$$P(\text{apple}) =$$



# FRUITS



- What is the probability to pick an apple?

$$P(\text{apple}) = P(\text{apple}|A) \cdot P(A) + P(\text{apple}|B) \cdot P(B) =$$

# FRUITS





- What is the probability to pick an apple?

$$\begin{aligned} P(\text{apple}) &= P(\text{apple}|A) \cdot P(A) + P(\text{apple}|B) \cdot P(B) = \\ &= 0.5 \cdot 0.5 + 0.25 \cdot 0.5 = 0.375 \end{aligned}$$





# FRUITS



- What is the probability to pick an orange?



# FRUITS



- What is the probability to pick an orange?

$$P(\text{orange}) =$$



# FRUITS



- What is the probability to pick an orange?

$$P(\text{orange}) = P(\text{orange}|A) \cdot P(A) + P(\text{orange}|B) \cdot P(B) =$$



# FRUITS



- What is the probability to pick an orange?



$$\begin{aligned} P(\text{orange}) &= P(\text{orange}|A) \cdot P(A) + P(\text{orange}|B) \cdot P(B) = \\ &= \frac{1}{2} \cdot \frac{1}{2} + \frac{7}{12} \cdot \frac{1}{2} = \frac{1}{4} + \frac{7}{24} = \frac{13}{24} \end{aligned}$$

# FRUITS



- What is the probability to pick a banana?



# FRUITS



- What is the probability to pick a banana?

$$P(\text{banana}) =$$



# FRUITS



- What is the probability to pick a banana?

$$P(banana) = P(banana|A) \cdot P(A) + P(banana|B) \cdot P(B) =$$

# FRUITS



- What is the probability to pick a banana?

$$\begin{aligned} P(\text{banana}) &= P(\text{banana}|A) \cdot P(A) + P(\text{banana}|B) \cdot P(B) = \\ &= 0 \cdot \frac{1}{2} + \frac{2}{12} \cdot \frac{1}{2} = \frac{1}{12} \end{aligned}$$



# HEIGHTS

- In a class, 40% of students are male and 60% are female. It's known that 60% of the males and 10% of the females are taller than 6 feet. What % of the class is not taller than 6 feet?

# HEIGHTS

- In a class, 40% of students are male and 60% are female. It's known that 60% of the males and 10% of the females are taller than 6 feet. What % of the class is not taller than 6 feet?

$$P(\leq 6) =$$

# HEIGHTS

- In a class, 40% of students are male and 60% are female. It's known that 60% of the males and 10% of the females are taller than 6 feet. What % of the class is not taller than 6 feet?

$$P(\leq 6) = P(\leq 6 \text{ and Male}) + P(\leq 6 \text{ and Female}) =$$

# HEIGHTS

- In a class, 40% of students are male and 60% are female. It's known that 60% of the males and 10% of the females are taller than 6 feet. What % of the class is not taller than 6 feet?

$$\begin{aligned} P(\leq 6) &= P(\leq 6 \text{ and Male}) + P(\leq 6 \text{ and Female}) = \\ &= P(\leq 6|M) \cdot P(M) + P(\leq 6|F) \cdot P(F) = \end{aligned}$$

# HEIGHTS

- In a class, 40% of students are male and 60% are female. It's known that 60% of the males and 10% of the females are taller than 6 feet. What % of the class is not taller than 6 feet?

$$\begin{aligned} P(\leq 6) &= P(\leq 6 \text{ and Male}) + P(\leq 6 \text{ and Female}) = \\ &= P(\leq 6|M) \cdot P(M) + P(\leq 6|F) \cdot P(F) = \\ &= (1 - 0.6) \cdot 0.4 + (1 - 0.1) \cdot 0.6 = 0.7 \end{aligned}$$

# COINS

- A box contains 3 coins with a head on each side, 4 coins with a tail on each side, and 2 fair coins. One of these nine coins is selected at random and tossed once. What is the probability that a head will be obtained?

# COINS

- A box contains 3 coins with a head on each side, 4 coins with a tail on each side, and 2 fair coins. One of these nine coins is selected at random and tossed once. What is the probability that a head will be obtained?

$E_H$  — a coin with 2 H is selected

$E_T$  — a coin with 2 T is selected

$E_F$  — a fair coin is selected

# COINS

- A box contains 3 coins with a head on each side, 4 coins with a tail on each side, and 2 fair coins. One of these nine coins is selected at random and tossed once. What is the probability that a head will be obtained?

$E_H$  — a coin with 2 H is selected

$E_T$  — a coin with 2 T is selected

$E_F$  — a fair coin is selected

$$P(H) = P(H|E_H) \cdot P(E_H) + P(H|E_T) \cdot P(E_T) + P(H|E_F) \cdot P(E_F) =$$



# COINS

- A box contains 3 coins with a head on each side, 4 coins with a tail on each side, and 2 fair coins. One of these nine coins is selected at random and tossed once. What is the probability that a head will be obtained?

$E_H$  — a coin with 2 H is selected

$E_T$  — a coin with 2 T is selected

$E_F$  — a fair coin is selected

$$P(H) = P(H|E_H) \cdot P(E_H) + P(H|E_T) \cdot P(E_T) + P(H|E_F) \cdot P(E_F) =$$

# COINS

- A box contains 3 coins with a head on each side, 4 coins with a tail on each side, and 2 fair coins. One of these nine coins is selected at random and tossed once. What is the probability that a head will be obtained?

$E_H$  — a coin with 2 H is selected       $P(E_H) =$

$E_T$  — a coin with 2 T is selected

$E_F$  — a fair coin is selected

$$P(H) = P(H|E_H) \cdot P(E_H) + P(H|E_T) \cdot P(E_T) + P(H|E_F) \cdot P(E_F) =$$

# COINS

- A box contains 3 coins with a head on each side, 4 coins with a tail on each side, and 2 fair coins. One of these nine coins is selected at random and tossed once. What is the probability that a head will be obtained?

$$E_H \text{ — a coin with 2 H is selected} \qquad P(E_H) = \frac{3}{3+4+2} = \frac{1}{3}$$

$$E_T \text{ — a coin with 2 T is selected}$$

$$E_F \text{ — a fair coin is selected}$$

$$P(H) = P(H|E_H) \cdot P(E_H) + P(H|E_T) \cdot P(E_T) + P(H|E_F) \cdot P(E_F) =$$

# COINS

- A box contains 3 coins with a head on each side, 4 coins with a tail on each side, and 2 fair coins. One of these nine coins is selected at random and tossed once. What is the probability that a head will be obtained?

$$E_H \text{ — a coin with 2 H is selected} \quad P(E_H) = \frac{3}{3+4+2} = \frac{1}{3}$$

$$E_T \text{ — a coin with 2 T is selected} \quad P(E_T) =$$

$$E_F \text{ — a fair coin is selected}$$

$$P(H) = P(H|E_H) \cdot P(E_H) + P(H|E_T) \cdot P(E_T) + P(H|E_F) \cdot P(E_F) =$$

# COINS

- A box contains 3 coins with a head on each side, 4 coins with a tail on each side, and 2 fair coins. One of these nine coins is selected at random and tossed once. What is the probability that a head will be obtained?

$$E_H \text{ — a coin with 2 H is selected} \quad P(E_H) = \frac{3}{3+4+2} = \frac{1}{3}$$

$$E_T \text{ — a coin with 2 T is selected} \quad P(E_T) = \frac{4}{3+4+2} = \frac{4}{9}$$

$$E_F \text{ — a fair coin is selected}$$

$$P(H) = P(H|E_H) \cdot P(E_H) + P(H|E_T) \cdot P(E_T) + P(H|E_F) \cdot P(E_F) =$$

# COINS

- A box contains 3 coins with a head on each side, 4 coins with a tail on each side, and 2 fair coins. One of these nine coins is selected at random and tossed once. What is the probability that a head will be obtained?

$$E_H \text{ — a coin with 2 H is selected} \quad P(E_H) = \frac{3}{3+4+2} = \frac{1}{3}$$

$$E_T \text{ — a coin with 2 T is selected} \quad P(E_T) = \frac{4}{3+4+2} = \frac{4}{9}$$

$$E_F \text{ — a fair coin is selected} \quad P(E_H) =$$

$$P(H) = P(H|E_H) \cdot P(E_H) + P(H|E_T) \cdot P(E_T) + P(H|E_F) \cdot P(E_F) =$$

# COINS

- A box contains 3 coins with a head on each side, 4 coins with a tail on each side, and 2 fair coins. One of these nine coins is selected at random and tossed once. What is the probability that a head will be obtained?

$$E_H \text{ — a coin with 2 H is selected} \quad P(E_H) = \frac{3}{3+4+2} = \frac{1}{3}$$

$$E_T \text{ — a coin with 2 T is selected} \quad P(E_T) = \frac{4}{3+4+2} = \frac{4}{9}$$

$$E_F \text{ — a fair coin is selected} \quad P(E_H) = \frac{2}{3+4+2} = \frac{2}{9}$$

$$P(H) = P(H|E_H) \cdot P(E_H) + P(H|E_T) \cdot P(E_T) + P(H|E_F) \cdot P(E_F) =$$

# COINS

- A box contains 3 coins with a head on each side, 4 coins with a tail on each side, and 2 fair coins. One of these nine coins is selected at random and tossed once. What is the probability that a head will be obtained?

$$E_H \text{ — a coin with 2 H is selected} \quad P(E_H) = \frac{3}{3+4+2} = \frac{1}{3}$$

$$E_T \text{ — a coin with 2 T is selected} \quad P(E_T) = \frac{4}{3+4+2} = \frac{4}{9}$$

$$E_F \text{ — a fair coin is selected} \quad P(E_H) = \frac{2}{3+4+2} = \frac{2}{9}$$

$$P(H) = P(H|E_H) \cdot P(E_H) + P(H|E_T) \cdot P(E_T) + P(H|E_F) \cdot P(E_F) =$$

$$= 1 \cdot \frac{1}{3} + 0 + \frac{1}{2} \cdot \frac{2}{9} =$$



# COINS

- A box contains 3 coins with a head on each side, 4 coins with a tail on each side, and 2 fair coins. One of these nine coins is selected at random and tossed once. What is the probability that a head will be obtained?

$$E_H \text{ — a coin with 2 H is selected} \quad P(E_H) = \frac{3}{3+4+2} = \frac{1}{3}$$

$$E_T \text{ — a coin with 2 T is selected} \quad P(E_T) = \frac{4}{3+4+2} = \frac{4}{9}$$



$$E_F \text{ — a fair coin is selected} \quad P(E_F) = \frac{2}{3+4+2} = \frac{2}{9}$$

$$P(H) = P(H|E_H) \cdot P(E_H) + P(H|E_T) \cdot P(E_T) + P(H|E_F) \cdot P(E_F) =$$

$$= 1 \cdot \frac{1}{3} + 0 + \frac{1}{2} \cdot \frac{2}{9} = \frac{1}{3} + \frac{1}{9} = \frac{4}{9}$$



# Bayes' rule

# FRUITS



- What is the probability that you chose box A given that you picked an apple?

# FRUITS





- What is the probability that you chose box A given that you picked an apple?

$$P(A|apple) =$$

# BAYES' RULE

$$P(A|B) = \frac{P(AB)}{P(B)} = \frac{P(B|A) \cdot P(A)}{P(B)}$$



# FRUITS



- What is the probability that you chose box A given that you picked an apple?

$$P(A|apple) =$$



# FRUITS



- What is the probability that you chose box A given that you picked an apple?

$$P(A|apple) = \frac{P(apple|A) \cdot P(A)}{P(apple)} =$$

# FRUITS





- What is the probability that you chose box A given that you picked an apple?

$$\begin{aligned} P(A|\text{apple}) &= \frac{P(\text{apple}|A) \cdot P(A)}{P(\text{apple})} = \\ &= \frac{P(\text{apple}|A) \cdot P(A)}{P(\text{apple}|A) \cdot P(A) + P(\text{apple}|B) \cdot P(B)} = \end{aligned}$$





# FRUITS



- What is the probability that you chose box A given that you picked an apple?

$$\begin{aligned} P(A|\text{apple}) &= \frac{P(\text{apple}|A) \cdot P(A)}{P(\text{apple})} = \\ &= \frac{P(\text{apple}|A) \cdot P(A)}{P(\text{apple}|A) \cdot P(A) + P(\text{apple}|B) \cdot P(B)} = \frac{0.5 \cdot 0.5}{0.5 \cdot 0.5 + 0.25 \cdot 0.5} = \frac{2}{3} \end{aligned}$$



# FRUITS



- What is the probability that you chose box B given that you picked a banana?

$$P(B|banana) =$$



# FRUITS



- What is the probability that you chose box B given that you picked a banana?

$$P(B|banana) = \frac{P(banana|B) \cdot P(B)}{P(banana)} =$$



# FRUITS



- What is the probability that you chose box B given that you picked a banana?

$$\begin{aligned} P(B|banana) &= \frac{P(banana|B) \cdot P(B)}{P(banana)} = \\ &= \frac{P(banana|B) \cdot P(B)}{P(banana|A) \cdot P(A) + P(banana|B) \cdot P(B)} = \end{aligned}$$



# FRUITS



- What is the probability that you chose box B given that you picked a banana?

$$\begin{aligned} P(B|banana) &= \frac{P(banana|B) \cdot P(B)}{P(banana)} = \\ &= \frac{P(banana|B) \cdot P(B)}{P(banana|A) \cdot P(A) + P(banana|B) \cdot P(B)} = 1 \end{aligned}$$



# FRUITS



- What is the probability that you chose box A given that you picked an orange?

$$P(A|\text{orange}) =$$

# FRUITS





- What is the probability that you chose box A given that you picked an orange?

$$P(A|orange) = \frac{P(orange|A) \cdot P(A)}{P(orange)} =$$

=

# FRUITS



- What is the probability that you chose box A given that you picked an orange?

$$\begin{aligned} P(A|\text{orange}) &= \frac{P(\text{orange}|A) \cdot P(A)}{P(\text{orange})} = \\ &= \frac{P(\text{orange}|A) \cdot P(A)}{P(\text{orange}|A) \cdot P(A) + P(\text{orange}|B) \cdot P(B)} = \frac{0.5 \cdot 0.5}{0.5 \cdot 0.5 + 7/12 \cdot 0.5} \end{aligned}$$



# DRUG TEST

- 0.5% of the population are using drugs. If a person taking the test has used drugs, the test will show a positive result with a 98% chance, and if they haven't used drugs, the test will be negative with a 98% probability. A random person tests positive. What is the probability that this person has actually used drugs?

# DRUGS

- 0.5% of the population are using drugs.

# DRUGS

- 0.5% of the population are using drugs.
- $P(D) = 0.005$

# DRUGS

- 0.5% of the population are using drugs.
  - If a person uses drugs, the test will be positive with a 98% chance.
- $P(D) = 0.005$

# DRUGS

- 0.5% of the population are using drugs.
  - If a person uses drugs, the test will be positive with a 98% chance.
- $P(D) = 0.005$
  - $P(+|drugs) = 0.98$

# DRUGS

- 0.5% of the population are using drugs.
  - If a person uses drugs, the test will be positive with a 98% chance.
  - if they don't use drugs, the test will be negative with a 98% probability.
- $P(D) = 0.005$
  - $P(+|drugs) = 0.98$

# DRUGS

- 0.5% of the population are using drugs.
  - If a person uses drugs, the test will be positive with a 98% chance.
  - if they don't use drugs, the test will be negative with a 98% probability.
- $P(D) = 0.005$
  - $P(+|drugs) = 0.98$
  - $P(-|no\ drugs) = 0.98$

# DRUGS

- 0.5% of the population are using drugs.
  - If a person uses drugs, the test will be positive with a 98% chance.
  - if they don't use drugs, the test will be negative with a 98% probability.
  - What is the probability that this person has actually used drugs?
- $P(D) = 0.005$
  - $P(+|drugs) = 0.98$
  - $P(-|no\ drugs) = 0.98$



# DRUGS

- 0.5% of the population are using drugs.
  - If a person uses drugs, the test will be positive with a 98% chance.
  - if they don't use drugs, the test will be negative with a 98% probability.
  - What is the probability that this person has actually used drugs?
- $P(D) = 0.005$
  - $P(+|drugs) = 0.98$
  - $P(-|no\ drugs) = 0.98$
  - $P(drugs|+) = ?$

# DRUG TEST

- 0.5% of the population are using drugs. If a person taking the test has used drugs, the test will show a positive result with a 98% chance, and if they haven't used drugs, the test will be negative with a 98% probability. A random person tests positive. What is the probability that this person has actually used drugs?

$$P(drugs|+) =$$

# DRUG TEST

- 0.5% of the population are using drugs. If a person taking the test has used drugs, the test will show a positive result with a 98% chance, and if they haven't used drugs, the test will be negative with a 98% probability. A random person tests positive. What is the probability that this person has actually used drugs?

$$P(\text{drugs} | +) = \frac{P(\text{drugs and } +)}{P(+)} =$$

# DRUG TEST

- 0.5% of the population are using drugs. If a person taking the test has used drugs, the test will show a positive result with a 98% chance, and if they haven't used drugs, the test will be negative with a 98% probability. A random person tests positive. What is the probability that this person has actually used drugs?

$$P(\text{drugs} | +) = \frac{P(\text{drugs and } +)}{P(+)} = \frac{P(+ | \text{drugs}) \cdot P(\text{drugs})}{P(+)} =$$

# DRUG TEST

- 0.5% of the population are using drugs. If a person taking the test has used drugs, the test will show a positive result with a 98% chance, and if they haven't used drugs, the test will be negative with a 98% probability. A random person tests positive. What is the probability that this person has actually used drugs?

$$\begin{aligned} P(\text{drugs} | +) &= \frac{P(\text{drugs and } +)}{P(+)} = \frac{P(+ | \text{drugs}) \cdot P(\text{drugs})}{P(+)} = \\ &= \frac{P(+ | \text{drugs}) \cdot P(\text{drugs})}{P(+ | \text{drugs}) \cdot P(\text{drugs}) + P(+ | \text{no drugs}) \cdot P(\text{no drugs})} \end{aligned}$$

# DRUG TEST

- 0.5% of the population are using drugs. If a person taking the test has used drugs, the test will show a positive result with a 98% chance, and if they haven't used drugs, the test will be negative with a 98% probability. A random person tests positive. What is the probability that this person has actually used drugs?

$$P(\text{drugs} | +) = \frac{P(+ | \text{drugs}) \cdot P(\text{drugs})}{P(+ | \text{drugs}) \cdot P(\text{drugs}) + P(+ | \text{no drugs}) \cdot P(\text{no drugs})} =$$

# DRUG TEST

- 0.5% of the population are using drugs. If a person taking the test has used drugs, the test will show a positive result with a 98% chance, and if they haven't used drugs, the test will be negative with a 98% probability. A random person tests positive. What is the probability that this person has actually used drugs?

$$\begin{aligned} P(\text{drugs} | +) &= \frac{P(+ | \text{drugs}) \cdot P(\text{drugs})}{P(+ | \text{drugs}) \cdot P(\text{drugs}) + P(+ | \text{no drugs}) \cdot P(\text{no drugs})} = \\ &= \frac{0.98 \cdot 0.005}{0.98 \cdot 0.005 + (1 - 0.98) \cdot (1 - 0.005)} \approx 0.2 \end{aligned}$$

# Practice Problems



# FLIPPING A COIN 2

- A coin is flipped 4 times. What is the probability of getting an equal number of heads and tails given that the outcomes on the first and the last flips are not the same?

# FLIPPING A COIN 2

- A coin is flipped 4 times. What is the probability of getting an equal number of heads and tails given that the outcomes on the first and the last flips are not the same?

$A$  — equal number of H and T

$B$  — different results on the first and last flips

# FLIPPING A COIN 2

- A coin is flipped 4 times. What is the probability of getting an equal number of heads and tails given that the outcomes on the first and the last flips are not the same?

$A$  — equal number of H and T

$B$  — different results on the first and last flips

$AB$  — different results on the first and last flips  
equal number of H and T

# FLIPPING A COIN 2

- A coin is flipped 4 times. What is the probability of getting an equal number of heads and tails given that the outcomes on the first and the last flips are not the same?

$A$  — equal number of H and T

$B$  — different results on the first and last flips       $P(B) =$

$AB$  — different results on the first and last flips  
equal number of H and T

# FLIPPING A COIN 2

- A coin is flipped 4 times. What is the probability of getting an equal number of heads and tails given that the outcomes on the first and the last flips are not the same?

$A$  — equal number of H and T

$B$  — different results on the first and last flips

$$P(B) = \frac{2^3}{2^4} = \frac{1}{2}$$

$AB$  — different results on the first and last flips  
equal number of H and T

# FLIPPING A COIN 2

- A coin is flipped 4 times. What is the probability of getting an equal number of heads and tails given that the outcomes on the first and the last flips are not the same?

$A$  — equal number of H and T

$B$  — different results on the first and last flips  $P(B) = \frac{2^3}{2^4} = \frac{1}{2}$

$AB$  — different results on the first and last flips  
equal number of H and T  $P(AB) =$

# FLIPPING A COIN 2

- A coin is flipped 4 times. What is the probability of getting an equal number of heads and tails given that the outcomes on the first and the last flips are not the same?

$A$  — equal number of H and T

$B$  — different results on the first and last flips  $P(B) = \frac{2^3}{2^4} = \frac{1}{2}$

$AB$  — different results on the first and last flips  
equal number of H and T  $P(AB) = \frac{2^2}{2^4} = \frac{1}{4}$

# FLIPPING A COIN 2

- A coin is flipped 4 times. What is the probability of getting an equal number of heads and tails given that the outcomes on the first and the last flips are not the same?

$A$  – equal number of H and T

$B$  – different results on the first and last flips  $P(B) = \frac{2^3}{2^4} = \frac{1}{2}$

$AB$  – different results on the first and last flips  
equal number of H and T  $P(AB) = \frac{2^2}{2^4} = \frac{1}{4}$

$$P(A|B) = \frac{P(AB)}{P(B)} =$$



# FLIPPING A COIN 2

- A coin is flipped 4 times. What is the probability of getting an equal number of heads and tails given that the outcomes on the first and the last flips are not the same?

$A$  – equal number of H and T

$B$  – different results on the first and last flips  $P(B) = \frac{2^3}{2^4} = \frac{1}{2}$

$AB$  – different results on the first and last flips  
equal number of H and T  $P(AB) = \frac{2}{2^4} = \frac{1}{8}$

$$P(A|B) = \frac{P(AB)}{P(B)} = \frac{1}{2}$$

# ROLLING A DIE

- We roll two fair dice. What is the probability to have 2 on the first die given that the sum of two numbers is 5?

# ROLLING A DIE

- We roll two fair dice. What is the probability to have 2 on the first die given that the sum of two numbers is 5?

$$P(X_1 = 2 | X_1 + X_2 = 5) =$$

# ROLLING A DIE

- We roll two fair dice. What is the probability to have 2 on the first die given that the sum of two numbers is 5?

$$P(X_1 = 2 | X_1 + X_2 = 5) = \frac{P(X_1 = 2, X_1 + X_2 = 5)}{P(X_1 + X_2 = 5)} =$$

# ROLLING A DIE

- We roll two fair dice. What is the probability to have 2 on the first die given that the sum of two numbers is 5?

$$P(X_1 = 2 | X_1 + X_2 = 5) = \frac{P(X_1 = 2, X_1 + X_2 = 5)}{P(X_1 + X_2 = 5)} =$$

$$P(X_1 = 2 \text{ and } X_1 + X_2 = 5) =$$

# ROLLING A DIE

- We roll two fair dice. What is the probability to have 2 on the first die given that the sum of two numbers is 5?

$$P(X_1 = 2 | X_1 + X_2 = 5) = \frac{P(X_1 = 2, X_1 + X_2 = 5)}{P(X_1 + X_2 = 5)} =$$

$$P(X_1 = 2 \text{ and } X_1 + X_2 = 5) = P(X_1 = 2 \text{ and } X_2 = 3) =$$

# ROLLING A DIE

- We roll two fair dice. What is the probability to have 2 on the first die given that the sum of two numbers is 5?

$$P(X_1 = 2 | X_1 + X_2 = 5) = \frac{P(X_1 = 2, X_1 + X_2 = 5)}{P(X_1 + X_2 = 5)} =$$

$$P(X_1 = 2 \text{ and } X_1 + X_2 = 5) = P(X_1 = 2 \text{ and } X_2 = 3) = \frac{1}{6 \cdot 6}$$

# ROLLING A DIE

- We roll two fair dice. What is the probability to have 2 on the first die given that the sum of two numbers is 5?

$$P(X_1 = 2 | X_1 + X_2 = 5) = \frac{P(X_1 = 2, X_1 + X_2 = 5)}{P(X_1 + X_2 = 5)} =$$

$$P(X_1 = 2 \text{ and } X_1 + X_2 = 5) = P(X_1 = 2 \text{ and } X_2 = 3) = \frac{1}{6 \cdot 6}$$

$$P(X_1 + X_2 = 5) =$$



# ROLLING A DIE

- We roll two fair dice. What is the probability to have 2 on the first die given that the sum of two numbers is 5?

$$P(X_1 = 2 | X_1 + X_2 = 5) = \frac{P(X_1 = 2, X_1 + X_2 = 5)}{P(X_1 + X_2 = 5)} =$$

$$P(X_1 = 2 \text{ and } X_1 + X_2 = 5) = P(X_1 = 2 \text{ and } X_2 = 3) = \frac{1}{6 \cdot 6}$$

$$P(X_1 + X_2 = 5) = \frac{4}{36} = \frac{1}{9}$$

# ROLLING A DIE

- We roll two fair dice. What is the probability to have 2 on the first die given that the sum of two numbers is 5?

$$P(X_1 = 2 | X_1 + X_2 = 5) = \frac{P(X_1 = 2, X_1 + X_2 = 5)}{P(X_1 + X_2 = 5)} = \frac{1}{4}$$

$$P(X_1 = 2 \text{ and } X_1 + X_2 = 5) = P(X_1 = 2 \text{ and } X_2 = 3) = \frac{1}{6 \cdot 6}$$

$$P(X_1 + X_2 = 5) = \frac{4}{36} = \frac{1}{9}$$

# MARKETING

- 1 in 50 potential buyers sees a magazine ad ( $M$ ), 1 in 5 sees a TV ad ( $T$ ) and 1 in 100 sees both. One in 3 purchases the product after seeing the ad, 1 in 10 without seeing it. What is the probability that a randomly selected potential customer will purchase the product ( $P$ )?

$$P(P) =$$

# MARKETING

- 1 in 50 potential buyers sees a magazine ad ( $M$ ), 1 in 5 sees a TV ad ( $T$ ) and 1 in 100 sees both. One in 3 purchases the product after seeing the ad, 1 in 10 without seeing it. What is the probability that a randomly selected potential customer will purchase the product ( $P$ )?

$$P(P) = P(P|A) \cdot P(A) + P(P|\bar{A}) \cdot P(\bar{A}) =$$

# MARKETING

- 1 in 50 potential buyers sees a magazine ad ( $M$ ), 1 in 5 sees a TV ad ( $T$ ) and 1 in 100 sees both. One in 3 purchases the product after seeing the ad, 1 in 10 without seeing it. What is the probability that a randomly selected potential customer will purchase the product ( $P$ )?

$$\begin{aligned} P(P) &= P(P|A) \cdot P(A) + P(P|\bar{A}) \cdot P(\bar{A}) = \\ &= P(P|M \cup T) \cdot P(M \cup T) + P(P|\overline{M \cup T}) \cdot P(\overline{M \cup T}) = \end{aligned}$$

# MARKETING

- 1 in 50 potential buyers sees a magazine ad ( $M$ ), 1 in 5 sees a TV ad ( $T$ ) and 1 in 100 sees both. One in 3 purchases the product after seeing the ad, 1 in 10 without seeing it. What is the probability that a randomly selected potential customer will purchase the product ( $P$ )?

$$\begin{aligned} P(P) &= P(P|A) \cdot P(A) + P(P|\bar{A}) \cdot P(\bar{A}) = \\ &= P(P|M \cup T) \cdot P(M \cup T) + P(P|\overline{M \cup T}) \cdot P(\overline{M \cup T}) = \end{aligned}$$

$$P(M \cup T) =$$

# MARKETING

- 1 in 50 potential buyers sees a magazine ad ( $M$ ), 1 in 5 sees a TV ad ( $T$ ) and 1 in 100 sees both. One in 3 purchases the product after seeing the ad, 1 in 10 without seeing it. What is the probability that a randomly selected potential customer will purchase the product ( $P$ )?

$$\begin{aligned} P(P) &= P(P|A) \cdot P(A) + P(P|\bar{A}) \cdot P(\bar{A}) = \\ &= P(P|M \cup T) \cdot P(M \cup T) + P(P|\overline{M \cup T}) \cdot P(\overline{M \cup T}) = \end{aligned}$$

$$P(M \cup T) = P(M) + P(T) - P(M \cap T) =$$

# MARKETING

- 1 in 50 potential buyers sees a magazine ad ( $M$ ), 1 in 5 sees a TV ad ( $T$ ) and 1 in 100 sees both. One in 3 purchases the product after seeing the ad, 1 in 10 without seeing it. What is the probability that a randomly selected potential customer will purchase the product ( $P$ )?

$$\begin{aligned} P(P) &= P(P|A) \cdot P(A) + P(P|\bar{A}) \cdot P(\bar{A}) = \\ &= P(P|M \cup T) \cdot P(M \cup T) + P(P|\overline{M \cup T}) \cdot P(\overline{M \cup T}) = \end{aligned}$$

$$P(M \cup T) = P(M) + P(T) - P(M \cap T) = 0.02 + 0.2 - 0.01 = 0.21$$



# MARKETING

- 1 in 50 potential buyers sees a magazine ad ( $M$ ), 1 in 5 sees a TV ad ( $T$ ) and 1 in 100 sees both. One in 3 purchases the product after seeing the ad, 1 in 10 without seeing it. What is the probability that a randomly selected potential customer will purchase the product ( $P$ )?

$$\begin{aligned} P(P) &= P(P|A) \cdot P(A) + P(P|\bar{A}) \cdot P(\bar{A}) = \\ &= P(P|M \cup T) \cdot P(M \cup T) + P(P|\overline{M \cup T}) \cdot P(\overline{M \cup T}) = \\ &= \frac{1}{3} \cdot 0.21 + 0.1 \cdot (1 - 0.21) = \end{aligned}$$

$$P(M \cup T) = P(M) + P(T) - P(M \cap T) = 0.02 + 0.2 - 0.01 = 0.21$$

# MARKETING

- 1 in 50 potential buyers sees a magazine ad ( $M$ ), 1 in 5 sees a TV ad ( $T$ ) and 1 in 100 sees both. One in 3 purchases the product after seeing the ad, 1 in 10 without seeing it. What is the probability that a randomly selected potential customer will purchase the product ( $P$ )?

$$\begin{aligned} P(P) &= P(P|A) \cdot P(A) + P(P|\bar{A}) \cdot P(\bar{A}) = \\ &= P(P|M \cup T) \cdot P(M \cup T) + P(P|\overline{M \cup T}) \cdot P(\overline{M \cup T}) = \\ &= \frac{1}{3} \cdot 0.21 + 0.1 \cdot (1 - 0.21) = 0.07 + 0.079 = 0.149 \end{aligned}$$

$$P(M \cup T) = P(M) + P(T) - P(M \cap T) = 0.02 + 0.2 - 0.01 = 0.21$$

# BALLS IN TWO BOWLS

- There are two bowls: bowl A contains 2 blue and 3 red balls, and bowl B contains 1 blue and 4 red balls. A blue ball was randomly picked from one of the bowls. What's the probability that it was bowl A?

# BALLS IN TWO BOWLS

- There are two bowls: bowl A contains 2 blue and 3 red balls, and bowl B contains 1 blue and 4 red balls. A blue ball was randomly picked from one of the bowls. What's the probability that it was bowl A?

$$P(A|blue) =$$

# BALLS IN TWO BOWLS

- There are two bowls: bowl A contains 2 blue and 3 red balls, and bowl B contains 1 blue and 4 red balls. A blue ball was randomly picked from one of the bowls. What's the probability that it was bowl A?

$$P(A|blue) = \frac{P(blue|A) \cdot P(A)}{P(blue)} =$$

# BALLS IN TWO BOWLS

- There are two bowls: bowl A contains 2 blue and 3 red balls, and bowl B contains 1 blue and 4 red balls. A blue ball was randomly picked from one of the bowls. What's the probability that it was bowl A?

$$\begin{aligned} P(A|blue) &= \frac{P(blue|A) \cdot P(A)}{P(blue)} = \\ &= \frac{P(blue|A) \cdot P(A)}{P(blue|A) \cdot P(A) + P(blue|B) \cdot P(B)} = \end{aligned}$$

# BALLS IN TWO BOWLS

- There are two bowls: bowl A contains 2 blue and 3 red balls, and bowl B contains 1 blue and 4 red balls. A blue ball was randomly picked from one of the bowls. What's the probability that it was bowl A?

$$\begin{aligned} P(A|blue) &= \frac{P(blue|A) \cdot P(A)}{P(blue)} = \\ &= \frac{P(blue|A) \cdot P(A)}{P(blue|A) \cdot P(A) + P(blue|B) \cdot P(B)} = \frac{0.4 \cdot 0.5}{0.4 \cdot 0.5 + 0.2 \cdot 0.5} = \\ &= \frac{0.2}{0.3} = \frac{2}{3} \end{aligned}$$

# COINS

- A box contains 3 coins with a head on each side, 4 coins with a tail on each side, and 2 fair coins. One of these nine coins is selected at random and tossed once. Suppose that the toss does result in a head. What is the probability that the coin used for tossing was fair?



# COINS

- A box contains 3 coins with a head on each side, 4 coins with a tail on each side, and 2 fair coins. One of these nine coins is selected at random and tossed once. Suppose that the toss does result in a head. What is the probability that the coin used for tossing was fair?

$E_H$ — a coin with 2 H is selected	$P(E_H) = \frac{3}{3+4+2} = \frac{1}{3}$
$E_T$ — a coin with 2 T is selected	$P(E_T) = \frac{4}{3+4+2} = \frac{4}{9}$
$E_F$ — a fair coin is selected	$P(E_F) = \frac{2}{3+4+2} = \frac{2}{9}$

# COINS

- A box contains 3 coins with a head on each side, 4 coins with a tail on each side, and 2 fair coins. One of these nine coins is selected at random and tossed once. Suppose that the toss does result in a head. What is the probability that the coin used for tossing was fair?

$E_H$ — a coin with 2 H is selected	$P(E_H) = \frac{3}{3+4+2} = \frac{1}{3}$
$E_T$ — a coin with 2 T is selected	$P(E_T) = \frac{4}{3+4+2} = \frac{4}{9}$
$E_F$ — a fair coin is selected	$P(E_F) = \frac{2}{3+4+2} = \frac{2}{9}$

$$P(H) = \frac{4}{9},$$

# COINS

- A box contains 3 coins with a head on each side, 4 coins with a tail on each side, and 2 fair coins. One of these nine coins is selected at random and tossed once. Suppose that the toss does result in a head. What is the probability that the coin used for tossing was fair?

$E_H$ — a coin with 2 H is selected	$P(E_H) = \frac{3}{3+4+2} = \frac{1}{3}$
$E_T$ — a coin with 2 T is selected	$P(E_T) = \frac{4}{3+4+2} = \frac{4}{9}$
$E_F$ — a fair coin is selected	$P(E_F) = \frac{2}{3+4+2} = \frac{2}{9}$

$$P(H) = \frac{4}{9}, \quad P(E_F|H) =$$

# COINS

- A box contains 3 coins with a head on each side, 4 coins with a tail on each side, and 2 fair coins. One of these nine coins is selected at random and tossed once. Suppose that the toss does result in a head. What is the probability that the coin used for tossing was fair?

$E_H$ — a coin with 2 H is selected	$P(E_H) = \frac{3}{3+4+2} = \frac{1}{3}$
$E_T$ — a coin with 2 T is selected	$P(E_T) = \frac{4}{3+4+2} = \frac{4}{9}$
$E_F$ — a fair coin is selected	$P(E_F) = \frac{2}{3+4+2} = \frac{2}{9}$

$$P(H) = \frac{4}{9}, \quad P(E_F|H) = \frac{P(H|E_F) \cdot P(E_F)}{P(H)} =$$

# COINS

- A box contains 3 coins with a head on each side, 4 coins with a tail on each side, and 2 fair coins. One of these nine coins is selected at random and tossed once. Suppose that the toss does result in a head. What is the probability that the coin used for tossing was fair?

$$\begin{array}{ll} E_H - \text{a coin with 2 H is selected} & P(E_H) = \frac{3}{3+4+2} = \frac{1}{3} \\ E_T - \text{a coin with 2 T is selected} & P(E_T) = \frac{4}{3+4+2} = \frac{4}{9} \\ E_F - \text{a fair coin is selected} & P(E_F) = \frac{2}{3+4+2} = \frac{2}{9} \end{array}$$

$$P(H) = \frac{4}{9}, \quad P(E_F|H) = \frac{P(H|E_F) \cdot P(E_F)}{P(H)} = \frac{\frac{1}{2} \cdot \frac{2}{9}}{\frac{4}{9}} = \frac{1}{4}$$