



THE UNIVERSITY *of* EDINBURGH

Edinburgh Research Explorer

The ARIA score of allergic rhinitis using mobile technology correlates with quality-of-life

Citation for published version:

MASK study group, Bousquet, J, Arnavielhe, S, Bedbrook, A, Fonseca, J, Morais Almeida, M, Todo Bom, A, Annesi-Maesano, I, Caimmi, D, Demoly, P, Devillier, P, Siroux, V, Menditto, E, Passalacqua, G, Stellato, C, Ventura, MT, Cruz, AA, Serpa, FS, da Silva, J, Larenas-Linnemann, D, Rodriguez Gonzalez, M, Cabañas, MTB, Bergmann, KC, Keil, T, Klimek, L, Mösges, R, Shamaï, S, Zuberbier, T, Bewick, M, Price, D, Ryan, D, Sheikh, A, Anto, JM, Mullol, J, Valero, A, Haahtela, T, Valovirta, E, Fokkens, WJ, Kuna, P, Samolinski, B, Bindslev-Jensen, C, Eller, E, Bosnic-Anticevich, S, O'Hehir, RE, Tomazic, PV, Yorgancioglu, A, Gemiciloglu, B, Bachert, C, Hellings, PW, Kull, I & Melén, E 2017, 'The ARIA score of allergic rhinitis using mobile technology correlates with quality-of-life: The MASK study', *Allergy*. <https://doi.org/10.1111/all.13307>

Digital Object Identifier (DOI):

[10.1111/all.13307](https://doi.org/10.1111/all.13307)

Link:

[Link to publication record in Edinburgh Research Explorer](#)

Document Version:

Peer reviewed version

Published In:

Allergy

General rights

Copyright for the publications made accessible via the Edinburgh Research Explorer is retained by the author(s) and / or other copyright owners and it is a condition of accessing these publications that users recognise and abide by the legal requirements associated with these rights.

Take down policy

The University of Edinburgh has made every reasonable effort to ensure that Edinburgh Research Explorer content complies with UK legislation. If you believe that the public display of this file breaches copyright please contact openaccess@ed.ac.uk providing details, and we will remove access to the work immediately and investigate your claim.



PROF. PHILIPPE DEVILLIER (Orcid ID : 0000-0003-4107-8317)

PROF. CRISTIANA STELLATO (Orcid ID : 0000-0002-1294-8355)

Article type : Brief Communication

Editor : Thomas Bieber

The ARIA score of allergic rhinitis using mobile technology correlates with quality-of-life: The MASK study

J Bousquet, MD (1, 2), S Arnavielhe, PhD (3), A Bedbrook, BSc (1), J Fonseca, MD (4), M Morais Almeida, MD (5), A Todo Bom, MD (6), I Annesi-Maesano, PhD (7), D Caimmi, MD (8), P Demoly, MD (8), P Devillier, MD (9), V Siroux, PhD (10), E Menditto, PhD, (11), G Passalacqua, MD (12), C Stellato, MD (13), MT Ventura, MD (14), AA Cruz, MD (15), FS Serpa, MD, (16) J da Silva, MD, (17) D Larenas-Linnemann, MD (18) M Rodriguez Gonzalez, MD, (19) MT Burguete Cabañas, MD, (20), KC Bergmann, MD (21), T Keil, MD (22), L Klimek, MD (23), R Mösges, MD (24), S Shamai, MD, (24) T Zuberbier, MD (21), M Bewick, MD, (25), D Price, MD (26), D Ryan, MD (27), A Sheikh, MD (28), JM Anto, MD (29), J Mullol, MD (5), A Valero, MD (30), T Haahtela, MD (31), E Valovirta, MD (30), WJ Fokkens MD, (32), P Kuna, MD (33), B Samolinski, MD, C Bindslev-Jensen, MD (34), E Eller, MD (34), S Bosnic-Anticevich, PhD (35), RE O'Hehir MD, (36), PV Tomazic, MD (37), A Yorgancioglu, MD, (38) B Gemicioglu, MD, (39), C Bachert, MD (40), PW Hellings, MD (41), I Kull, PhD, (42), E Melén, MD (42), M Wickman, MD (43), M van Eerd (44), G De Vries, PhD (44), and the MASK study group

1. MACVIA-France, Contre les MALadies Chroniques pour un Vieillissement Actif en France European Innovation Partnership on Active and Healthy Ageing Reference Site, Montpellier, France.
2. INSERM U 1168, VIMA : Ageing and chronic diseases Epidemiological and public health approaches, Villejuif, Université Versailles St-Quentin-en-Yvelines, UMR-S 1168, Montigny le Bretonneux, France
3. Kyomed , Montpellier France.
4. Center for Health Technology and Services Research- CINTESIS, Faculdade de Medicina, Universidade do Porto; and Allergy Unit, CUF Porto Instituto & Hospital, Porto, Portugal.
5. Allergy Center, CUF-Descobertas Hospital, Lisboa, Portugal.

This article has been accepted for publication and undergone full peer review but has not been through the copyediting, typesetting, pagination and proofreading process, which may lead to differences between this version and the Version of Record. Please cite this article as doi: 10.1111/all.13307

This article is protected by copyright. All rights reserved.

6. Imunoalergologia, Centro Hospitalar Universitário de Coimbra and Faculty of Medicine, University of Coimbra, Portugal.
7. EPAR U707 INSERM, Paris and EPAR UMR-S UPMC, Paris VI, Paris, France.
8. CHRU de Montpellier, Sorbonne Universités, UPMC Paris 06, UMR-S 1136, IPLESP, Equipe EPAR, F-75013 Paris, France.
9. Laboratoire de Pharmacologie Respiratoire UPRES EA220, Pôle des Maladies Respiratoires, Hôpital Foch, Suresnes Université Versailles Saint-Quentin, France.
10. INSERM, Université Grenoble Alpes, IAB, U 1209, Team of Environmental Epidemiology applied to Reproduction and Respiratory Health, Université Joseph Fourier, Grenoble, France.
11. CIRFF, Center of Pharmacoeconomics, University of Naples Federico II, Naples, Italy.
12. Personalized Medicine Clinic Asthma & Allergy, Humanitas University, Humanitas Research Hospital, Rozzano, Milan, Italy
13. Department of Medicine, Surgery and Dentistry "Scuola Medica Salernitana", University of Salerno, Salerno, Italy.
14. University of Bari Medical School, Unit of Geriatric Immunoallergy, Bari, Italy.
15. ProAR – Nucleo de Excelencia em Asma, Federal University of Bahia, Brasil and GARD Executive Committee, Brazil.
16. Asthma Reference Center, Escola Superior de Ciencias da Santa Casa de Misericórdia de Vitória, Brazil.
17. Allergy Service, University Hospital of Federal University of Santa Catarina (HU-UFSC), Florianópolis, Brazil.
18. Center of Excellence in Asthma and Allergy, Hospital Médica Sur, México, Mexico.
19. Mexico City, Mexico.
20. Centro Médico Zambrano Hellion, Monterrey, Mexico.
21. Comprehensive Allergy-Centre-Charité, Department of Dermatology and Allergy, Charité - Universitätsmedizin Berlin; Global Allergy and Asthma European Network (GA²LEN), Berlin, Germany.
22. Institute of Social Medicine, Epidemiology and Health Economics, Charité - Universitätsmedizin Berlin, Berlin, and Institute for Clinical Epidemiology and Biometry, University of Würzburg, Germany
23. Center for Rhinology and Allergology, Wiesbaden, Germany.
24. Institute of Medical Statistics, Informatics and Epidemiology, Medical Faculty, University of Cologne, Germany.
25. iQ4U Consultants Ltd, London, UK.
26. Observational and Pragmatic Research Institute, Singapore, Optimum Patient Care, Cambridge, UK, and Academic Centre of Primary Care, University of Aberdeen, Aberdeen, UK.
27. Allergy and Respiratory Research Group, Usher Institute of Population Health Sciences and Informatics, University of Edinburgh, UK
28. Director, Asthma UK Centre for Applied Research, Centre of Medical Informatics, Usher Institute of Population Health Sciences and Informatics, The University of Edinburgh, Edinburgh, UK.
29. ISGLoBAL, Centre for Research in Environmental Epidemiology (CREAL), Barcelona; IMIM (Hospital del Mar Research Institute); CIBER Epidemiología y Salud Pública (CIBERESP), & Universitat Pompeu Fabra (UPF), Barcelona, Spain.

- Accepted Article
30. Pneumology and Allergy Department Hospital Clínic, Clinical & Experimental Respiratory Immunoallergy, IDIBAPS, CIBERES, University of Barcelona, Spain.
 31. Skin and Allergy Hospital, Helsinki University Hospital, Helsinki, Finland.
 32. Department of Otorhinolaryngology, Academic Medical Centre, Amsterdam, the Netherlands.
 33. Division of Internal Medicine, Asthma and Allergy, Barlicki University Hospital, Medical University of Lodz, Poland.
 34. Department of Dermatology and Allergy Centre, Odense University Hospital, Odense Research Center for Anaphylaxis (ORCA), Odense, Denmark.
 35. Woolcock Institute of Medical Research, University of Sydney and Sydney Local Health District, Glebe, NSW, Australia.
 36. Department of Allergy, Immunology and Respiratory Medicine, Alfred Hospital and Central Clinical School, Monash University, Melbourne, Victoria, Australia; Department of Immunology, Monash University, Melbourne, Victoria, Australia.
 37. Department of ENT, Medical University of Graz, Austria
 38. Celal Bayar University Department of Pulmonology, Manisa, Turkey and GARD Executive Committee, Turkey
 39. Department of Pulmonary Diseases, Istanbul University, Cerrahpasa Faculty of Medicine, Turkey.
 40. Upper Airways Research Laboratory, ENT Dept, Ghent University Hospital, Ghent, Belgium.
 41. Laboratory of Clinical Immunology, Department of Microbiology and Immunology, KU Leuven, Leuven, Belgium
 42. Department of Clinical Science and Education, Södersjukhuset, Karolinska Institutet, Stockholm, Sweden
 43. Sachs' Children and Youth Hospital, Södersjukhuset, Stockholm and Institute of Environmental Medicine, Karolinska Institutet, Stockholm, Sweden.
 44. Peercode DV, Gerdermalsem, The Netherlands.

Short title: Quality of life in rhinitis using a mobile application

Funding source

Unrestricted educational grant from Meda, European Union Development and Structural funds (Région Languedoc Roussillon), MACVIA-LR

Corresponding author

Professor Jean Bousquet

CHU Montpellier, 371 Avenue du Doyen Gaston Giraud, 34295 Montpellier Cedex 5, France Tel
+33 611 42 88 47 jean.bousquet@orange.fr

Abstract

Mobile technology has been used to appraise allergic rhinitis control but more data are needed. In order to better assess the importance of mobile technologies in rhinitis control, the ARIA (Allergic Rhinitis and its Impact on Asthma) score ranging from 0 to 4 of the *Allergy Diary* was compared with EQ-5D (EuroQuol) and WPAI-AS (Work Productivity and Activity Impairment in allergy) in 1,288 users in 18 countries. This study showed that quality-of-life data (EQ-5D visual analogue scale and WPA-IS Question 9) are similar in users without rhinitis and in those with mild rhinitis (scores 0-2). Users with a score of 3 or 4 had a significant impairment in quality-of-life questionnaires.

Key words: rhinitis, EQ-5D, ARIA, MASK, WPAI-AS

Abbreviations

AHA: Active and Healthy Aging

AR: allergic rhinitis

ARIA: Allergic Rhinitis and its Impact on Asthma

EIP on AHA: European Innovation Partnership on Active and Healthy Ageing (DG CONNECT, DG Santé)

EQ-5D: EuroQuol

ICT: information and communications technology

MACVIA: Contre les MALadies Chroniques pour un Vieillessement Actif

MASK: MACVIA-ARIA Sentinel Network

MAFEIP: Monitoring and assessment framework for the EIP on AHA

QOL: Quality of life

Q9: Question 9 of WPAI-AS

SF-36: Short Form 36 questions

This article is protected by copyright. All rights reserved.

VAS: visual analogue scale

WPAI-AS: Work Productivity and Activity Impairment in allergy

Introduction

Measures of allergic rhinitis (AR) control include symptom scores, patients' self-administered visual analogue scales (VAS), objective measures of nasal obstruction, a recent modification of the ARIA severity classification, or patients' reported outcomes such as QOL or scores with several items (1, 2). Mobile technology has been used to appraise AR control (3, 4). More information is however needed to fully understand the importance of these novel approaches.

MASK-rhinitis (MACVIA-ARIA Sentinel Network for allergic rhinitis), an ICT system centred around the patient (5), is one of the implementation tools of the B3 Action Plan of the European Innovation Partnership on Active and Healthy Ageing (EIP on AHA) (6). A mobile phone app (*Allergy Diary*) central to MASK-rhinitis belongs to the Fondation Partenariale FMC VIA-LR (Ministry of Education and Research, France). App users are asked to complete a short demographic questionnaire, EQ-5D (7-10) and WPAI-AS (11, 12), thus providing baseline characteristics of their disease. The *Allergy Diary* has been launched in 21 countries (3, 4). It was found to be an easy and effective method of assessing symptoms of AR and work productivity. The ARIA score is also available in the *Allergy Diary* adding the four components of the impact of AR (sleep, work and school performance, daily activities and bothersome symptoms).

The ARIA score of the *Allergy Diary* was compared with QOL scores of EQ-5D (7-10) and WPAI-AS (11, 12).

Methods

Design of the study

A cross-sectional study on a self-selected population was carried out from June 1, 2016 to June 1, 2017. EQ-5D (7-10) and/or WPAI-AS (11, 12) questionnaires are only available in some countries and not all of the *Allergy Diary* users filled in these questionnaires as they are optional. All users filled in the ARIA score. The ARIA score was compared with the EQ-5D visual analogue scale (VAS) and Question 9 (degree allergy affected regular activities) of the WPAI-AS.

The study is reported according to STROBE.

Setting and users

All consecutive users from June 1, 2016 to June 1, 2017 who answered the questions of the EQ-5D (7-10) and/or WPAI-AS (11, 12) were included in the study in 18 countries. Some demographic characteristics (age, sex, country and language) were recorded. The App was used by people who found it on Internet, Apple App store, Google Play store or any other way. A few users were clinic patients that were asked by their physicians to use the app. However, due to anonymisation of data, no specific information was gathered.

Allergy Diary

The app collects information on AR symptoms experienced (nasal and ocular), disease type (intermittent/persistent), how symptoms impact users' lives, and type(s) of AR treatment used (Table 1 online). The system has been deployed in 21 countries and in 16 languages (translated and back-translated, culturally adapted and legally compliant).

Ethics

The Allergy Diary is CE1 registered but not considered by the Ethical Committee of the Cologne hospital of the MHRA (Medicines and Healthcare products Regulatory Agency - GOV.UK) as a medical device as it does not give any recommendations concerning treatment or diagnosis. The terms of use, translated into all languages and customized according to each country's legislation, allow the use of the results for research purposes. The data are anonymized except for geolocalized data that are never totally anonymous. An Independent Review Board approval was not needed.

Outcomes

The ARIA score was calculated using the four Q4 questions of the *Allergy Diary* which include impact on daily activities, work and sleep and troublesome symptoms (Table 1 online). Each of the 4 items was ascribed a score of 1 ("Yes") or 0 ("No"). The total ARIA score ranged from 0 (no impairment) to 4 (severe impairment).

The electronic form of the EQ-5D-5L questionnaire (<https://euroqol.org>) was applied in the 10 available languages (Danish, Dutch, English, Finnish, French, German, Italian, Polish, Portuguese, Spanish) (Online supplement 1) and we assessed the global VAS level and mobility impairment as this was an absent domain in the assessment of AR impairment.

The electronic form of the WPAI-AS questionnaire was applied in the 10 available languages (same as above for EQ-5D) (11, 12) according to the package obtained from Reilly and associates (www.reillyassociates.net/WPAI_General.html). The percentage of impairment due to allergy for daily activities (Q9) was the outcome used. (Online supplement 2).

Classification of users

Users with any positive answer to Q4 (Table 1 online) were classified as "rhinitis" (score 0-4). Those with a score of zero were classified as "no rhinitis" if they had no symptom (Q3, Table 1 online). Those with a positive answer were classified as "rhinitis" (score 0).

Statistical methods and analyses

Some users filled in EQ-5D or WPAI-AS more than once for a single day. The first data were then used. A non-Gaussian distribution was found for some of the data (Shapiro-Wilk test). However, EQ-5D data are usually reported in means and SD. Since the number of observations was large, we used parametric analyses.

Results

Users

Of the 12,179 registered users, 1,287 filled in the EQ-5D questionnaire and 1,028 the WPAI-AS questionnaire (Table 2 online). Among the 843 users who filled in both questionnaires, there were 507 women (60%) and 336 men (40%), with a mean (\pm SD) age of 35 ± 14 years.

Main results

Similar levels of EQ-5D VAS and WPAI-AS Q9 were found for users with no rhinitis and for those with an ARIA score of 0 to 2. There was a significant reduction of EQ-5D VAS levels and a significant increase of WPAI-AS Q9 levels in users with an ARIA score of 3 or 4 (Table 1).

The repartition of users for both EQ-5D and WPAI-AS (Figure 1) shows that impairment occurred significantly more commonly for ARIA scores of 3 and 4 than for ARIA scores of 0-2. There were from 12 to 16% of users with an EQ-5D VAS level ≥ 60 in ARIA scores 0-2 whereas the level increased to 26 and 27% in users with an ARIA score of 3 or 4. There were from 19 to 31% of users with a Q9 ≥ 50 in ARIA scores of 0-2 whereas the level increased to 51 and 53% in users with an ARIA score of 3 or 4.

Discussion

This pilot study using mobile technology showed that QOL data (EQ-5D VAS and WPAI-AS Q9) are similar in users without rhinitis as in those with mild rhinitis (scores 0-2). Users with a score of 3 or 4 had a significant impairment in QOL.

Strengths and limitations

The strengths and limitations of this study are those of mobile technology lengthily discussed previously (3, 4). In particular, there is a lack of patient characterization that is impossible using an App. However, every observational study we have performed using the *Allergy Diary* has confirmed its interest and was able to identify users with a severe disease. It is likely that mobile technology will become a very important tool for the understanding and management of AR.

One specific problem of the study is that there are more countries with EQ-5D or WPAI-AS reporting than translations in the App. It is not known which translations were employed by users.

In this study, we did not perform sub-analyses assessing the importance of symptoms or other factors. We did not investigate the treatments received. As this is a pilot study, these analyses will be carried out once the number of users will have increased.

Generalizability

The EQ-5D scores observed in the study accord with those of previous studies (8, 10). Users with an ARIA score of 3 to 4 have a level similar to asthmatic patients with uncontrolled asthma (13, 14). Because of the equal weighing score of ARIA, it is difficult to know whether these differences may be due to specific symptoms (e.g. sleep). EQ-5D is a MAFEIP (Monitoring and assessment framework for the EIP on AHA) tool (15) and the present study is in line with the EIP on AHA. This is another important finding since the Transfer of Innovation of the Allergy Diary is an EIP on AHA scaling up project (16).

One of the major findings of the study is the very similar results with both tools supporting the use of the ARIA score to assess AR control using mobile technology. The WPAI-AS scores observed in the study are lower than those reported in patients selected by physicians (11, 17-19). This is because many users have mild rhinitis whereas in clinical trials or in patients selected by physicians AR is usually more severe.


This study also suggests that, in real life, there is a phenotype of severe AR that needs to be considered in terms of public health and cost savings since the severe form causes disability. This phenotype is in focus in the Finnish Allergy Program (20).

Table 1: Mean levels of EQ-5D and WPAI-AS depending on the ARIA score

		EQ-5D VAS		WPAI-AS Q9	
		N	m ± sd	N	m ± sd
No rhinitis		48	80.0±19.0	49	27.5±25.5
ARIA score	0	83	77.0±21.3 ^a	71	29.9±27.3 ^d
	1	403	79.5±19.1	308	20.4±22.1
	2	368	76.2±20.2	268	30.1±23.9
	3	199	72.6±18.7 ^b	164	41.4±27.3
	4	186	67.7±23.0 ^c	168	45.8±27.7 ^e

$p_{a/b} < 0.0001$, $p_{a/c} < 0.0001$, $p_{d/e} < 0.0001$, Student's t test

Figure 1: Repartition of users depending on EQ-5D visual analogue scale (A) and WPAI-AS Q9 (B)



References

1. Schatz M, Meltzer EO, Nathan R, Derebery MJ, Mintz M, Stanford RH, et al. Psychometric validation of the rhinitis control assessment test: a brief patient-completed instrument for evaluating rhinitis symptom control. *Ann Allergy Asthma Immunol.* 2010;104(2):118-24.
2. Bousquet J, Schunemann HJ, Hellings PW, Arnavielhe S, Bachert C, Bedbrook A, et al. MACVIA clinical decision algorithm in adolescents and adults with allergic rhinitis. *J Allergy Clin Immunol.* 2016;138(2):367-74 e2.
3. Bousquet J, Bewick M, Arnavielhe S, Mathieu-Dupas E, Murray R, Bedbrook A, et al. Work productivity in rhinitis using cell phones: The MASK pilot study. *Allergy.* 2017.
4. Bousquet J, Caimmi DP, Bedbrook A, Bewick M, Hellings PW, Devillier P, et al. Pilot study of mobile phone technology in allergic rhinitis in European countries: the MASK-rhinitis study. *Allergy.* 2017;72(6):857-65.
5. Bousquet J, Hellings PW, Agache I, Bedbrook A, Bachert C, Bergmann KC, et al. ARIA 2016: Care pathways implementing emerging technologies for predictive medicine in rhinitis and asthma across the life cycle. *Clin Transl Allergy.* 2016;6:47.
6. Bousquet J, Addis A, Adcock I, Agache I, Agusti A, Alonso A, et al. Integrated care pathways for airway diseases (AIRWAYS-ICPs). *Eur Respir J.* 2014;44(2):304-23.
7. Hoehle LP, Speth MM, Phillips KM, Gaudin RA, Caradonna DS, Gray ST, et al. Association between symptoms of allergic rhinitis with decreased general health-related quality of life. *Am J Rhinol Allergy.* 2017;31(4):235-9.

- Accepted Article
8. Ock M, Jo MW, Gong YH, Lee HJ, Lee J, Sim CS. Estimating the severity distribution of disease in South Korea using EQ-5D-3L: a cross-sectional study. *BMC Public Health*. 2016;16:234.
 9. Petersen KD, Kronborg C, Larsen JN, Dahl R, Gyrd-Hansen D. Patient related outcomes in a real life prospective follow up study: Allergen immunotherapy increase quality of life and reduce sick days. *World Allergy Organ J*. 2013;6(1):15.
 10. Chen H, Cisternas MG, Katz PP, Omachi TA, Trupin L, Yelin EH, et al. Evaluating quality of life in patients with asthma and rhinitis: English adaptation of the rhinasthma questionnaire. *Ann Allergy Asthma Immunol*. 2011;106(2):110-8 e1.
 11. Bousquet J, Neukirch F, Bousquet PJ, Gehano P, Klossek JM, Le Gal M, et al. Severity and impairment of allergic rhinitis in patients consulting in primary care. *J Allergy Clin Immunol*. 2006;117(1):158-62.
 12. Virchow JC, Kay S, Demoly P, Mullol J, Canonica W, Higgins V. Impact of ocular symptoms on quality of life (QoL), work productivity and resource utilisation in allergic rhinitis patients--an observational, cross sectional study in four countries in Europe. *J Med Econ*. 2011;14(3):305-14.
 13. Sullivan PW, Ghushchyan VH, Campbell JD, Globe G, Bender B, Magid DJ. Measurement of utility in asthma: evidence indicating that generic instruments may miss clinically important changes. *Qual Life Res*. 2016;25(12):3017-26.
 14. Terzano C, Cremonesi G, Girbino G, Ingrassia E, Marsico S, Nicolini G, et al. 1-year prospective real life monitoring of asthma control and quality of life in Italy. *Respir Res*. 2012;13:112.
 15. Boehler CE, de Graaf G, Steuten L, Yang Y, Abadie F. Development of a web-based tool for the assessment of health and economic outcomes of the European Innovation Partnership on Active and Healthy Ageing (EIP on AHA). *BMC Med Inform Decis Mak*. 2015;15 Suppl 3:S4.
 16. Bousquet J, Agache I, Aliberti MR, Angles R, Annesi-Maesano I, Anto JM, et al. Transfer of innovation on allergic rhinitis and asthma multimorbidity in the elderly (MACVIA-ARIA) - Reference Site Twinning (EIP on AHA). *Allergy*. 2017.
 17. Stull DE, Roberts L, Frank L, Heithoff K. Relationship of nasal congestion with sleep, mood, and productivity. *Curr Med Res Opin*. 2007;23(4):811-9.
 18. Meltzer EO, Munafo DA, Chung W, Gopalan G, Varghese ST. Intranasal mometasone furoate therapy for allergic rhinitis symptoms and rhinitis-disturbed sleep. *Ann Allergy Asthma Immunol*. 2010;105(1):65-74.
 19. Colas C, Brosa M, Anton E, Montoro J, Navarro A, Dordal MT, et al. Estimate of the total costs of allergic rhinitis in specialized care based on real-world data: the FERIN Study. *Allergy*. 2017;72(6):959-66.
 20. Haahtela T, Valovirta E, Bousquet J, Makela M, and the Allergy Programme Steering G. The Finnish Allergy Programme 2008-2018 works. *Eur Respir J*. 2017;49(6).