

Decidability of the Admissible Rules in Intuitionistic Propositional Logic

Jeroen P. Goudsmit

Utrecht University

Workshop on Admissibility and Unification 2

January 31st 2015

A / Δ admissible



σA is derivable



A / Δ admissible



σC is derivable for some $C \in \Delta$



σA is derivable



$A \rightsquigarrow \Delta$ admissible




σC is derivable for some $C \in \Delta$



Given a rule A/Δ ,
is it admissible?








1975 Friedman





1975 Friedman

1979 Citkin



1975 Friedman

1979 Citkin

1984 Rybakov



1975 Friedman

1979 Citkin

1984 Rybakov

1992 Rozière





1975 Friedman

1979 Citkin


1984 Rybakov

1992 Rozière


1999 Ghilardi









Semantics for Admissible Rules





Semantics for Admissible Rules

Adequate Semantics



Semantics for Admissible Rules

Adequate Semantics

Effective Description



Semantics for Admissible Rules



What is a good notion of
semantics for admissibility?



Definition

Say that A/Δ is **valid** on v , denoted $v \Vdash A/\Delta$, if:


$v \Vdash A$ implies $v \Vdash C$ for some $C \in \Delta$.



Theorem (Rieger, 1949; Nishimura, 1960; Esakia and Grigolia, 1977; Shehtman, 1978; Rybakov, 1984)

For each finite set of variables X , there exists a model $u : U(X) \rightarrow P(X)$ such that:


$$u \Vdash A \text{ iff } \vdash A \text{ for all } A \in \mathcal{L}(X).$$





$v \Vdash A/\Delta$ for
all $v \in \mathcal{K}$



$A \Vdash \Delta$



































Definition

A map $f: u \rightarrow v$ is **definable** if there is a substitution σ such that:

$$u, p \Vdash \sigma A \text{ iff } v, f(p) \Vdash A \text{ for all } A.$$



Definition

A map $f: u \rightarrow v$ is **definable** if there is a substitution σ such that:

$$u, p \Vdash \sigma A \text{ iff } v, f(p) \Vdash A \text{ for all } A.$$

Definition (de Jongh, 1982)

A model v called **exact** if there exists a definable map $u \rightarrow v$.



Definition


A map $f: u \rightarrow v$ is **definable** if there is a substitution σ such that:

$$u, p \Vdash \sigma A \text{ iff } v, f(p) \Vdash A \text{ for all } A.$$

Definition (de Jongh, 1982)

A model v called **exact** if there exists a definable map $u \rightarrow v$, where u is a universal model.





Theorem

The admissible rules of IPC are sound and complete with respect to exact models.



What is a good notion of semantics for admissibility?



What is a good notion of
semantics for admissibility?

Exact models!



Theorem (Fedorishin and Ivanov, 2003; Goudsmit, 2014b)

*The admissible rules of IPC are **not** sound and complete with respect to **finite** exact models.*








Adequate Semantics





What is a *fair* notion of semantics for admissibility?





























Definition

A map $f: u \rightarrow v$ is **adequate** if there exists a substitution such that:

$$u, p \Vdash \sigma A \text{ iff } v, f(p) \Vdash A \text{ for all } A.$$

Definition

A map $f: u \rightarrow v$ is Σ -adequate if there exists a substitution such that:

$$u, p \Vdash \sigma A \text{ iff } v, f(p) \Vdash A \text{ for all } A \in \Sigma.$$

Definition

A map $f: u \rightarrow v$ is Σ -adequate if there exists a substitution such that:

$$u, p \Vdash \sigma A \text{ iff } v, f(p) \Vdash A \text{ for all } A \in \Sigma.$$

Definition

A model v is Σ -adequately exact if there exists a Σ -adequate map $u \rightarrow v$.


Definition

A map $f: u \rightarrow v$ is Σ -adequate if there exists a substitution such that:

$$u, p \Vdash \sigma A \text{ iff } v, f(p) \Vdash A \text{ for all } A \in \Sigma.$$

Definition

A model v is Σ -adequately exact if there exists a Σ -adequate map $u \rightarrow v$, where u is a universal model.



Theorem

*The admissible rules of IPC
are sound and complete with respect to
exact models.*

Theorem

The admissible rules of IPC from an adequate set Σ are sound and complete with respect to Σ -adequately exact models.

Theorem

Let $A \in \Sigma$ and $\Delta \subseteq \Sigma$. The following are equivalent:

- 1. $A \vdash \Delta$;*
- 2. $v \models A/\Delta$ for all Σ -adequately exact models v of size at most $2^{|\Sigma|}$.*

What is a *fair* notion of semantics for admissibility?

What is a *fair* notion of
semantics for admissibility?
Adequately exact models!



IV

Effective Description

When is a model
 Σ -adequately exact?

Theorem (Citkin, 1977)


A finite submodel of a universal model is exact iff it is extendible.

Theorem (Ghilardi, 1999)


A definable submodel of a universal model is exact iff it is extendible.

Extendible


Extendible



Extendible



Extendible



Theorem

Let $U \subseteq \mathcal{U}(X)$ be an upset. Now, U is extendible iff for each finite $W \subseteq U$ there exists a $p \in U$ such that $W \subseteq \uparrow p$ and for all $A \rightarrow B$:

$$p \Vdash A \rightarrow B \text{ iff } (W \Vdash A \rightarrow B \text{ and } p \Vdash A \text{ implies } p \Vdash B).$$

Definition

Let $U \subseteq \mathcal{U}(X)$ be an upset. Now, U is Σ -extendible iff for each finite $W \subseteq U$ there exists a $p \in U$ such that $W \subseteq \uparrow p$ and for all $A \rightarrow B \in \Sigma$:

$$p \Vdash A \rightarrow B \text{ iff } (W \Vdash A \rightarrow B \text{ and } p \Vdash A \text{ implies } p \Vdash B).$$

Theorem

A finite submodel of a universal model is Σ -adequately exact iff it is Σ -extendible.

When is a model
 Σ -adequately exact?
If it's Σ -extendible.

Given a rule A/Δ ,
is it admissible?


Given a rule A/Δ ,
is it admissible?

Take Σ as all subformulae of A and Δ . Compute whether $v \Vdash A/\Delta$ for all Σ -extendible models of size at most $2^{|\Sigma|}$. If it is, then **yes**. Otherwise, **no**.

The admissible rules of IPC
are decidable.

The admissible rules of IPC
are decidable.

Rybakov (1984)
see Goudsmit (2014a) for more details.





Intuitionistic Rules

Admissible Rules of Intermediate Logics

Jeroen P. Goudsmit





References I



Citkin, A. (1977). “On Admissible Rules of Intuitionistic Propositional Logic”. In: *Mathematics of the USSR-Sbornik* 31.2, pp. 279–288. doi: [10.1070/SM1977v031n02ABEH002303](https://doi.org/10.1070/SM1977v031n02ABEH002303). Zbl: [0386.03011](https://zbmath.org/?q=0386.03011) (see p. 67).



– **(1979).** “О Проверке Допустимости Нёкоторых Правил Интуиционистской Логике”. Russian. In: *V-th All-Union Conference in Mathematical Logic*. English translation of title: On verification of admissibility of some rules of intuitionistic logic. Novosibirsk, p. 162 (see pp. 8–11).



Esakia, L. and Grigolia, R. (1977). “The criterion of Brouwerian and closure algebras to be finitely generated.” In: *Bulletin of the Section of Logic* 6, pp. 46–52. ISSN: 0138-0680. MR: [0476400](https://www.ams.org/mathscinet-getitem?mr=0476400). Zbl: [0407.03048](https://zbmath.org/?q=0407.03048) (see p. 20).



Fedorishin, B. and Ivanov, V. (2003). “The finite model property with respect to admissibility for superintuitionistic logics”. In: *Siberian Advances in Mathematics* 13.2, pp. 56–65. MR: [2029995](https://www.ams.org/mathscinet-getitem?mr=2029995). Zbl: [1047.03020](https://zbmath.org/?q=1047.03020) (see p. 42).

References II



Friedman, H. (1975). “One Hundred and Two Problems in Mathematical Logic”. In: *The Journal of Symbolic Logic* 40.2, pp. 113–129. ISSN: 00224812. DOI: [10.2307/2271891](https://doi.org/10.2307/2271891) (see pp. 7–11).



Ghilardi, S. (1999). “Unification in Intuitionistic Logic”. In: *The Journal of Symbolic Logic* 64.2, pp. 859–880. ISSN: 00224812. DOI: [10.2307/2586506](https://doi.org/10.2307/2586506) (see pp. 11, 68).



Goudsmit, J. P. (2014a). “Decidability of Admissibility: On a Problem of Friedman and its Solution by Rybakov”. In: *Logic Group Preprint Series* 322 (see pp. 79, 80).



– **(2014b).** “Finite Frames Fail”. In: *Logic Group Preprint Series* 321. URL: <http://www.phil.uu.nl/preprints/lgps/number/321> (see p. 42).

References III



de Jongh, D. H. J. (1982). “Formulas of One Propositional Variable in Intuitionistic Arithmetic”. In: *The L. E. J. Brouwer Centenary Symposium, Proceedings of the Conference held in Noordwijkerhout*. Ed. by A. S. Troelstra and D. van Dalen. Vol. 110. Studies in Logic and the Foundations of Mathematics. Elsevier, pp. 51–64. DOI: [10.1016/S0049-237X\(09\)70122-3](https://doi.org/10.1016/S0049-237X(09)70122-3) (see pp. 35–37).



Nishimura, I. (1960). “On Formulas of One Variable in Intuitionistic Propositional Calculus”. In: *The Journal of Symbolic Logic* 25.4, pp. 327–331. ISSN: 00224812. DOI: [10.2307/2963526](https://doi.org/10.2307/2963526) (see p. 20).



Rieger, L. (1949). “On the lattice theory of Brouwerian propositional logic”. In: *Acta Facultatis Rerum Naturalium Universitatis Carolinae* 189, pp. 1–40. MR: [0040245](https://www.ams.org/mathscinet/show-statement?mr=0040245) (see p. 20).



Rozière, P. (1992). “Règles admissibles en calcul propositionnel intuitionniste”. PhD thesis. Université de Paris VII (see pp. 10, 11).

References IV



Rybakov, V. V. (1984). “A criterion for admissibility of rules in the model system S4 and the intuitionistic logic”. In: *Algebra and Logic* 23.5, pp. 369–384. ISSN: 0002-5232. DOI: [10.1007/BF01982031](https://doi.org/10.1007/BF01982031) (see pp. [9–11](#), [20](#), [79](#), [80](#)).



Shehtman, V. B. (1978). “Rieger-Nishimura lattices”. In: *Soviet Mathematics Doklady* 19.4, pp. 1014–1018. ISSN: 0197-6788. Zbl: [0412.03010](https://zbmath.org/?q=rieger-nishimura) (see p. [20](#)). Trans. of **V. B. Šehtman**. “Reiger-Nishimura ladders”. In: *Doklady Akademii Nauk SSSR* 241.6, pp. 1288–1291. ISSN: 0002-3264. MR: [504235](#) .