



Utrecht University

Admissible rules of propositional dependence logic

Fan Yang
Utrecht University

Les Diablerets
Jan 30 - Feb 2, 2015

Joint work with Rosalie Iemhoff

Outline

- 1 propositional dependence logic
- 2 flat formulas and projective formulas
- 3 structural completeness

Dependence between first-order variables

First Order Quantifiers:

$$\forall x_1 \exists y_1 \forall x_2 \exists y_2 \phi$$

Dependence between first-order variables

First Order Quantifiers:

$$\forall x_1 \exists y_1 \forall x_2 \exists y_2 \phi$$

Dependence between first-order variables

First Order Quantifiers:

$$\forall x_1 \exists y_1 \forall x_2 \exists y_2 \phi$$

Dependence between first-order variables

First Order Quantifiers:

$$\forall x_1 \exists y_1 \forall x_2 \exists y_2 \phi$$

Henkin Quantifiers (Henkin, 1961):

$$\left(\begin{array}{cc} \forall x_1 & \exists y_1 \\ \forall x_2 & \exists y_2 \end{array} \right) \phi$$

Dependence between first-order variables

First Order Quantifiers:

$$\forall x_1 \exists y_1 \forall x_2 \exists y_2 \phi$$

Henkin Quantifiers (Henkin, 1961):

$$\left(\begin{array}{cc} \forall x_1 & \exists y_1 \\ \forall x_2 & \exists y_2 \end{array} \right) \phi$$

Independence Friendly Logic (Hintikka, Sandu, 1989):

$$\forall x_1 \exists y_1 \forall x_2 \exists y_2 / \{x_1\} \phi$$

Dependence between first-order variables

First Order Quantifiers:

$$\forall x_1 \exists y_1 \forall x_2 \exists y_2 \phi$$

Henkin Quantifiers (Henkin, 1961):

$$\left(\begin{array}{cc} \forall x_1 & \exists y_1 \\ \forall x_2 & \exists y_2 \end{array} \right) \phi$$

Independence Friendly Logic (Hintikka, Sandu, 1989):

$$\forall x_1 \exists y_1 \forall x_2 \exists y_2 / \{x_1\} \phi$$

First-order dependence Logic (Väänänen 2007):

$$\forall x_1 \exists y_1 \forall x_2 \exists y_2 (= (x_2, y_2) \wedge \phi)$$

Dependence between first-order variables

First Order Quantifiers:

$$\forall x_1 \exists y_1 \forall x_2 \exists y_2 \phi$$

Henkin Quantifiers (Henkin, 1961):

$$\left(\begin{array}{cc} \forall x_1 & \exists y_1 \\ \forall x_2 & \exists y_2 \end{array} \right) \phi$$

Independence Friendly Logic (Hintikka, Sandu, 1989):

$$\forall x_1 \exists y_1 \forall x_2 \exists y_2 / \{x_1\} \phi$$

First-order dependence Logic (Väänänen 2007):

$$\forall x_1 \exists y_1 \forall x_2 \exists y_2 (= (x_2, y_2) \wedge \phi)$$

Theorem (Enderton, Walkoe). Over sentences, all of the above extensions of first-order logic are equivalent to Σ_1^1 .

Propositional dependence logic = propositional logic + $=(\vec{p}, q)$

- I will be **absent** depending on whether **he shows up** or not.
- Whether **it rains** depends completely on whether it is **summer** or not.

	summer	rainy	windy
v_1	0	1	0

Propositional dependence logic = propositional logic + $=(\vec{p}, q)$

- I will be **absent** depending on whether **he shows up** or not.
- Whether **it rains** depends completely on whether it is **summer** or not.

	summer	rainy	windy
v_1	0	1	0

$$=(p, q)$$

Propositional dependence logic = propositional logic + $=(\vec{p}, q)$

- I will be **absent** depending on whether **he shows up** or not.
- Whether **it rains** depends completely on whether it is **summer** or not.

	summer	rainy	windy
v_1	0	1	0

$$=(p, q)$$

Propositional dependence logic = propositional logic + $=(\vec{p}, q)$

- I will be **absent** depending on whether **he shows up** or not.
- Whether **it rains** depends completely on whether it is **summer** or not.

	summer	rainy	windy
v_1	0	1	0

$=(\vec{p}, q)$

Propositional dependence logic = propositional logic + $=(\vec{p}, q)$

- I will be **absent** depending on whether **he shows up** or not.
- Whether **it rains** depends completely on whether it is **summer** or not.

Team semantics (Hodges 1997)

	summer	rainy	windy
v_1	0	1	0

$=(\vec{p}, q)$

Propositional dependence logic = propositional logic + $=(\vec{p}, q)$

- I will be **absent** depending on whether **he shows up** or not.
- Whether **it rains** depends completely on whether it is **summer** or not.

Team semantics (Hodges 1997)

	summer	rainy	windy
v_1	0	1	0

$$v_1 \models =(\vec{p}, q)?$$

Propositional dependence logic = propositional logic + $=(\vec{p}, q)$

- I will be **absent** depending on whether **he shows up** or not.
- Whether **it rains** depends completely on whether it is **summer** or not.

Team semantics (Hodges 1997)

	summer	rainy	windy
v_1	0	1	0
v_2	0	1	1
v_3	1	0	1
v_4	1	0	0

$$v_1 \models =(\textcolor{blue}{p}, \textcolor{red}{q})?$$

Propositional dependence logic = propositional logic + $=(\vec{p}, q)$

- I will be **absent** depending on whether **he shows up** or not.
- Whether **it rains** depends completely on whether it is **summer** or not.

Team semantics (Hodges 1997)

	summer	rainy	windy
X { v_1	0	1	0
v_2	0	1	1
v_3	1	0	1
v_4	1	0	0

$=(\textcolor{blue}{p}, \textcolor{red}{q})$

Propositional dependence logic = propositional logic + $=(\vec{p}, q)$

- I will be **absent** depending on whether **he shows up** or not.
- Whether **it rains** depends completely on whether it is **summer** or not.

Team semantics (Hodges 1997)

A *team* X {

	summer	rainy	windy
v_1	0	1	0
v_2	0	1	1
v_3	1	0	1
v_4	1	0	0

$$X \models =(\vec{p}, q)$$

Propositional dependence logic = propositional logic + $=(\vec{p}, q)$

- I will be **absent** depending on whether **he shows up** or not.
- Whether **it rains** depends completely on whether it is **summer** or not.

Team semantics (Hodges 1997)

A *team* X {

	summer	rainy	windy
v_1	0	1	0
v_2	0	1	1
v_3	1	0	1
v_4	1	0	0

$$X \models =(\vec{p}, q)$$

Propositional dependence logic = propositional logic + $=(\vec{p}, q)$

- I will be **absent** depending on whether **he shows up** or not.
- Whether **it rains** depends completely on whether it is **summer** or not.

Team semantics (Hodges 1997)

A *team* X {

	summer	rainy	windy
v_1	0	1	0
v_2	0	1	1
v_3	1	0	1
v_4	1	0	0

$$X \models =(\vec{p}, q)$$

Propositional dependence logic = propositional logic + $=(\vec{p}, q)$

- I will be **absent** depending on whether **he shows up** or not.
- Whether **it rains** depends completely on whether it is **summer** or not.

Team semantics (Hodges 1997)

A *team* X {

	summer	rainy	windy
v_1	0	1	0
v_2	0	1	1
v_3	1	0	1
v_4	1	0	0

$$X \models =(\vec{p}, q)$$

Propositional dependence logic = propositional logic + $=(\vec{p}, q)$

- I will be **absent** depending on whether **he shows up** or not.
- Whether **it rains** depends completely on whether it is **summer** or not.

Team semantics (Hodges 1997)

A *team* X {

	summer	rainy	windy
v_1	0	1	0
v_2	0	1	1
v_3	1	0	1
v_4	1	0	0

$$X \models =(\vec{p}, q)$$

Propositional dependence logic = propositional logic + $=(\vec{p}, q)$

- I will be **absent** depending on whether **he shows up** or not.
- Whether **it rains** depends completely on whether it is **summer** or not.

Team semantics (Hodges 1997)

A *team* X {

	summer	rainy	windy
v_1	0	1	0
v_2	0	1	1
v_3	1	0	1
v_4	1	0	0
v_5	1	1	1

} Y

$$X \models =(\textcolor{blue}{p}, \textcolor{red}{q}), \quad Y \not\models =(\textcolor{blue}{p}, \textcolor{red}{q})$$

Propositional dependence logic and its variants

- Well-formed formulas of *propositional dependence logic* (**PD**) are given by the following grammar

$$\phi ::= p_i \mid \neg p_i \mid =(p_{i_1}, \dots, p_{i_{n-1}}, p_{i_n}) \mid \phi \wedge \phi \mid \phi \otimes \phi$$

- Well-formed formulas of *propositional intuitionistic dependence logic* (**PID**) are given by the following grammar:

$$\phi ::= p_i \mid \perp \mid =(p_{i_1}, \dots, p_{i_{n-1}}, p_{i_n}) \mid \phi \wedge \phi \mid \phi \vee \phi \mid \phi \rightarrow \phi$$

- PD[∨]** is the logic extended from **PD** by adding the connective \vee .

	p_0	p_1	p_2	...
v_1	1	0	0	...

Propositional dependence logic and its variants

- Well-formed formulas of *propositional dependence logic* (**PD**) are given by the following grammar

$$\phi ::= p_i \mid \neg p_i \mid =(p_{i_1}, \dots, p_{i_{n-1}}, p_{i_n}) \mid \phi \wedge \phi \mid \phi \otimes \phi$$

- Well-formed formulas of *propositional intuitionistic dependence logic* (**PID**) are given by the following grammar:

$$\phi ::= p_i \mid \perp \mid =(p_{i_1}, \dots, p_{i_{n-1}}, p_{i_n}) \mid \phi \wedge \phi \mid \phi \vee \phi \mid \phi \rightarrow \phi$$

$(\neg \phi := \phi \rightarrow \perp)$

- PD**[∨] is the logic extended from **PD** by adding the connective ∨.

	p_0	p_1	p_2	...
v_1	1	0	0	...

Propositional dependence logic and its variants

- Well-formed formulas of *propositional dependence logic* (**PD**) are given by the following grammar

$$\phi ::= p_i \mid \neg p_i \mid (p_{i_1}, \dots, p_{i_{n-1}}, p_{i_n}) \mid \phi \wedge \phi \mid \phi \otimes \phi$$

- Well-formed formulas of *propositional intuitionistic dependence logic* (**PID**) are given by the following grammar:

$$\phi ::= p_i \mid \perp \mid (p_{i_1}, \dots, p_{i_{n-1}}, p_{i_n}) \mid \phi \wedge \phi \mid \phi \vee \phi \mid \phi \rightarrow \phi$$

$(\neg \phi := \phi \rightarrow \perp)$

- PD**[∨] is the logic extended from **PD** by adding the connective \vee .

	p_0	p_1	p_2	...
v_1	1	0	0	...

Propositional dependence logic and its variants

- Well-formed formulas of *propositional dependence logic* (**PD**) are given by the following grammar

$$\phi ::= p_i \mid \neg p_i \mid =(p_{i_1}, \dots, p_{i_{n-1}}, p_{i_n}) \mid \phi \wedge \phi \mid \phi \otimes \phi$$

- Well-formed formulas of *propositional intuitionistic dependence logic* (**PID**) are given by the following grammar:

$$\phi ::= p_i \mid \perp \mid =(p_{i_1}, \dots, p_{i_{n-1}}, p_{i_n}) \mid \phi \wedge \phi \mid \phi \vee \phi \mid \phi \rightarrow \phi$$

$(\neg \phi := \phi \rightarrow \perp)$

- PD**[∨] is the logic extended from **PD** by adding the connective \vee .

	p_0	p_1	p_2	\dots
v_1	1	0	0	\dots

Propositional dependence logic and its variants

- Well-formed formulas of *propositional dependence logic* (**PD**) are given by the following grammar

$$\phi ::= p_i \mid \neg p_i \mid (p_{i_1}, \dots, p_{i_{n-1}}, p_{i_n}) \mid \phi \wedge \phi \mid \phi \otimes \phi$$

- Well-formed formulas of *propositional intuitionistic dependence logic* (**PID**) are given by the following grammar:

$$\phi ::= p_i \mid \perp \mid (p_{i_1}, \dots, p_{i_{n-1}}, p_{i_n}) \mid \phi \wedge \phi \mid \phi \vee \phi \mid \phi \rightarrow \phi$$
$$(\neg \phi := \phi \rightarrow \perp)$$

- PD**[∨] is the logic extended from **PD** by adding the connective \vee .

Team semantics: A valuation is a function $v : \mathbb{N} \rightarrow \{0, 1\}$.

	p_0	p_1	p_2	\dots
v_1	1	0	0	\dots

Propositional dependence logic and its variants

- Well-formed formulas of *propositional dependence logic* (**PD**) are given by the following grammar

$$\phi ::= p_i \mid \neg p_i \mid (p_{i_1}, \dots, p_{i_{n-1}}, p_{i_n}) \mid \phi \wedge \phi \mid \phi \otimes \phi$$

- Well-formed formulas of *propositional intuitionistic dependence logic* (**PID**) are given by the following grammar:

$$\phi ::= p_i \mid \perp \mid (p_{i_1}, \dots, p_{i_{n-1}}, p_{i_n}) \mid \phi \wedge \phi \mid \phi \vee \phi \mid \phi \rightarrow \phi$$
$$(\neg \phi := \phi \rightarrow \perp)$$

- PD**[∨] is the logic extended from **PD** by adding the connective \vee .

Team semantics: A valuation is a function $v : \mathbb{N} \rightarrow \{0, 1\}$. A *team* is a set of valuations.

	p_0	p_1	p_2	\dots
v_1	1	0	0	\dots
v_2	1	1	0	\dots
v_4	0	1	0	\dots
\vdots	\vdots	\vdots	\vdots	\vdots

Team Semantics

Let X be a team.

- $X \models p_i$ iff for all $v \in X$, $v(i) = 1$;
- $X \models \neg p_i$ iff for all $v \in X$, $v(i) = 0$;
- $X \models \perp$ iff $X = \emptyset$;
- $X \models \phi$ iff for all $v, v' \in X$
 $[v(i_1) = v'(i_1), \dots, v(i_{n-1}) = v'(i_{n-1})] \implies v(i_n) = v'(i_n)$;
- $X \models \phi \wedge \psi$ iff $X \models \phi$ and $X \models \psi$;
- $X \models \phi \otimes \psi$ iff there exist teams $Y, Z \subseteq X$ with $X = Y \cup Z$ such that
 $Y \models \phi$ and $Z \models \psi$;

	p_0	p_1	p_2
v_1	1	0	0
v_2	1	0	1
v_3	0	1	0
v_4	0	1	1

- $X \models \phi \vee \psi$ iff $X \models \phi$ or $X \models \psi$;

Team Semantics

Let X be a team.

- $X \models p_i$ iff for all $v \in X$, $v(i) = 1$;
- $X \models \neg p_i$ iff for all $v \in X$, $v(i) = 0$;
- $X \models \perp$ iff $X = \emptyset$;
- $X \models \phi$ iff for all $v, v' \in X$
 $[v(i_1) = v'(i_1), \dots, v(i_{n-1}) = v'(i_{n-1})] \implies v(i_n) = v'(i_n)$;
- $X \models \phi \wedge \psi$ iff $X \models \phi$ and $X \models \psi$;
- $X \models \phi \otimes \psi$ iff there exist teams $Y, Z \subseteq X$ with $X = Y \cup Z$ such that
 $Y \models \phi$ and $Z \models \psi$;

		p_0	p_1	p_2	
$X \left\{ \right.$	v_1	1	0	0	$X \models p_0$
	v_2	1	0	1	
	v_3	0	1	0	
	v_4	0	1	1	

- $X \models \phi \vee \psi$ iff $X \models \phi$ or $X \models \psi$;

Team Semantics

Let X be a team.

- $X \models p_i$ iff for all $v \in X$, $v(i) = 1$;
- $X \models \neg p_i$ iff for all $v \in X$, $v(i) = 0$;
- $X \models \perp$ iff $X = \emptyset$;
- $X \models \phi$ iff for all $v, v' \in X$
 $[v(i_1) = v'(i_1), \dots, v(i_{n-1}) = v'(i_{n-1})] \implies v(i_n) = v'(i_n)$;
- $X \models \phi \wedge \psi$ iff $X \models \phi$ and $X \models \psi$;
- $X \models \phi \otimes \psi$ iff there exist teams $Y, Z \subseteq X$ with $X = Y \cup Z$ such that
 $Y \models \phi$ and $Z \models \psi$;

		p_0	p_1	p_2	
X	v_1	1	0	0	$X \models p_0$
	v_2	1	0	1	
Y	v_3	0	1	0	$Y \models \neg p_0$
	v_4	0	1	1	

- $X \models \phi \vee \psi$ iff $X \models \phi$ or $X \models \psi$;

Team Semantics

Let X be a team.

- $X \models p_i$ iff for all $v \in X$, $v(i) = 1$;
- $X \models \neg p_i$ iff for all $v \in X$, $v(i) = 0$;
- $X \models \perp$ iff $X = \emptyset$;
- $X \models \phi$ iff for all $v, v' \in X$
 $[v(i_1) = v'(i_1), \dots, v(i_{n-1}) = v'(i_{n-1})] \implies v(i_n) = v'(i_n)$;
- $X \models \phi \wedge \psi$ iff $X \models \phi$ and $X \models \psi$;
- $X \models \phi \otimes \psi$ iff there exist teams $Y, Z \subseteq X$ with $X = Y \cup Z$ such that
 $Y \models \phi$ and $Z \models \psi$;

		p_0	p_1	p_2
X	v_1	1	0	0
	v_2	1	0	1
Y	v_3	0	1	0
	v_4	0	1	1

$$X \models p_0$$

$$X \cup Y \not\models p_0$$

$$Y \models \neg p_0$$

$$X \cup Y \not\models \neg p_0$$

- $X \models \phi \vee \psi$ iff $X \models \phi$ or $X \models \psi$;

Team Semantics

Let X be a team.

- $X \models p_i$ iff for all $v \in X$, $v(i) = 1$;
- $X \models \neg p_i$ iff for all $v \in X$, $v(i) = 0$;
- $X \models \perp$ iff $X = \emptyset$;
- $X \models \phi(p_{i_1}, \dots, p_{i_n})$ iff for all $v, v' \in X$
 $[v(i_1) = v'(i_1), \dots, v(i_{n-1}) = v'(i_{n-1})] \implies v(i_n) = v'(i_n)$;
- $X \models \phi \wedge \psi$ iff $X \models \phi$ and $X \models \psi$;
- $X \models \phi \otimes \psi$ iff there exist teams $Y, Z \subseteq X$ with $X = Y \cup Z$ such that
 $Y \models \phi$ and $Z \models \psi$;

	p_0	p_1	p_2
v_1	1	0	0
v_2	1	0	1
v_3	0	1	0
v_4	0	1	1

- $X \models \phi \vee \psi$ iff $X \models \phi$ or $X \models \psi$;

Team Semantics

Let X be a team.

- $X \models p_i$ iff for all $v \in X$, $v(i) = 1$;
- $X \models \neg p_i$ iff for all $v \in X$, $v(i) = 0$;
- $X \models \perp$ iff $X = \emptyset$;
- $X \models =(p_{i_1}, \dots, p_{i_n})$ iff for all $v, v' \in X$
 $[v(i_1) = v'(i_1), \dots, v(i_{n-1}) = v'(i_{n-1})] \implies v(i_n) = v'(i_n)$;
- $X \models \phi \wedge \psi$ iff $X \models \phi$ and $X \models \psi$;
- $X \models \phi \otimes \psi$ iff there exist teams $Y, Z \subseteq X$ with $X = Y \cup Z$ such that
 $Y \models \phi$ and $Z \models \psi$;

	p_0	p_1	p_2
v_1	1	0	0
v_2	1	0	1
v_3	0	1	0
v_4	0	1	1

$X \models =(p_0, p_1)$

- $X \models \phi \vee \psi$ iff $X \models \phi$ or $X \models \psi$;

Team Semantics

Let X be a team.

- $X \models p_i$ iff for all $v \in X$, $v(i) = 1$;
- $X \models \neg p_i$ iff for all $v \in X$, $v(i) = 0$;
- $X \models \perp$ iff $X = \emptyset$;
- $X \models =(p_{i_1}, \dots, p_{i_n})$ iff for all $v, v' \in X$
 $[v(i_1) = v'(i_1), \dots, v(i_{n-1}) = v'(i_{n-1})] \implies v(i_n) = v'(i_n)$;
- $X \models \phi \wedge \psi$ iff $X \models \phi$ and $X \models \psi$;
- $X \models \phi \otimes \psi$ iff there exist teams $Y, Z \subseteq X$ with $X = Y \cup Z$ such that
 $Y \models \phi$ and $Z \models \psi$;

	p_0	p_1	p_2
v_1	1	0	0
v_2	1	0	1
v_3	0	1	0
v_4	0	1	1

$X \models =(p_0, p_1)$

- $X \models \phi \vee \psi$ iff $X \models \phi$ or $X \models \psi$;

Team Semantics

Let X be a team.

- $X \models p_i$ iff for all $v \in X$, $v(i) = 1$;
- $X \models \neg p_i$ iff for all $v \in X$, $v(i) = 0$;
- $X \models \perp$ iff $X = \emptyset$;
- $X \models \text{=(}p_{i_1}, \dots, p_{i_n}\text{)}$ iff for all $v, v' \in X$
 $[v(i_1) = v'(i_1), \dots, v(i_{n-1}) = v'(i_{n-1})] \implies v(i_n) = v'(i_n)$;
- $X \models \phi \wedge \psi$ iff $X \models \phi$ and $X \models \psi$;
- $X \models \phi \otimes \psi$ iff there exist teams $Y, Z \subseteq X$ with $X = Y \cup Z$ such that
 $Y \models \phi$ and $Z \models \psi$;
- $X \models \phi \vee \psi$ iff $X \models \phi$ or $X \models \psi$;
- $X \models \phi \rightarrow \psi$ iff for any team $Y \subseteq X$,
 $Y \models \phi \implies Y \models \psi$.

Team Semantics

Let X be a team.

- $X \models p_i$ iff for all $v \in X$, $v(i) = 1$;
- $X \models \neg p_i$ iff for all $v \in X$, $v(i) = 0$;
- $X \models \perp$ iff $X = \emptyset$;
- $X \models \text{=(}p_{i_1}, \dots, p_{i_n}\text{)}$ iff for all $v, v' \in X$
 $[v(i_1) = v'(i_1), \dots, v(i_{n-1}) = v'(i_{n-1})] \implies v(i_n) = v'(i_n)$;
- $X \models \phi \wedge \psi$ iff $X \models \phi$ and $X \models \psi$;
- $X \models \phi \otimes \psi$ iff there exist teams $Y, Z \subseteq X$ with $X = Y \cup Z$ such that
 $Y \models \phi$ and $Z \models \psi$;
- $X \models \phi \vee \psi$ iff $X \models \phi$ or $X \models \psi$;
- $X \models \phi \rightarrow \psi$ iff for any team $Y \subseteq X$,
 $Y \models \phi \implies Y \models \psi$.

Team Semantics

Let X be a team.

- $X \models p_i$ iff for all $v \in X$, $v(i) = 1$;
- $X \models \neg p_i$ iff for all $v \in X$, $v(i) = 0$;
- $X \models \perp$ iff $X = \emptyset$;
- $X \models \phi(p_{i_1}, \dots, p_{i_n})$ iff for all $v, v' \in X$
 $[v(i_1) = v'(i_1), \dots, v(i_{n-1}) = v'(i_{n-1})] \implies v(i_n) = v'(i_n)$;
- $X \models \phi \wedge \psi$ iff $X \models \phi$ and $X \models \psi$;
- $X \models \phi \otimes \psi$ iff there exist teams $Y, Z \subseteq X$ with $X = Y \cup Z$ such that
 $Y \models \phi$ and $Z \models \psi$;
- $X \models \phi \vee \psi$ iff $X \models \phi$ or $X \models \psi$;
- $X \models \phi \rightarrow \psi$ iff for any team $Y \subseteq X$,
 $Y \models \phi \implies Y \models \psi$.

Team Semantics

Let X be a team.

- $X \models p_i$ iff for all $v \in X$, $v(i) = 1$;
- $X \models \neg p_i$ iff for all $v \in X$, $v(i) = 0$;
- $X \models \perp$ iff $X = \emptyset$;
- $X \models =(p_{i_1}, \dots, p_{i_n})$ iff for all $v, v' \in X$
 $[v(i_1) = v'(i_1), \dots, v(i_{n-1}) = v'(i_{n-1})] \implies v(i_n) = v'(i_n)$;
- $X \models \phi \wedge \psi$ iff $X \models \phi$ and $X \models \psi$;
- $X \models \phi \otimes \psi$ iff there exist teams $Y, Z \subseteq X$ with $X = Y \cup Z$ such that
 $Y \models \phi$ and $Z \models \psi$;
- $X \models \phi \vee \psi$ iff $X \models \phi$ or $X \models \psi$;
- $X \models \phi \rightarrow \psi$ iff for any team $Y \subseteq X$,
 $Y \models \phi \implies Y \models \psi$.

The logics **PD**, **PD[∨]**, **PID**

- are **downwards closed**, that is, $X \models \phi$ and $Y \subseteq X \implies Y \models \phi$;
- and have the **empty team property**, that is, $\emptyset \models \phi$, for all ϕ .

Team Semantics

Let X be a team.

- $X \models p_i$ iff for all $v \in X$, $v(i) = 1$;
- $X \models \neg p_i$ iff for all $v \in X$, $v(i) = 0$;
- $X \models \perp$ iff $X = \emptyset$;
- $X \models (p_{i_1}, \dots, p_{i_n})$ iff for all $v, v' \in X$
 $[v(i_1) = v'(i_1), \dots, v(i_{n-1}) = v'(i_{n-1})] \implies v(i_n) = v'(i_n)$;
- $X \models \phi \wedge \psi$ iff $X \models \phi$ and $X \models \psi$;
- $X \models \phi \otimes \psi$ iff there exist teams $Y, Z \subseteq X$ with $X = Y \cup Z$ such that
 $Y \models \phi$ and $Z \models \psi$;
- $X \models \phi \vee \psi$ iff $X \models \phi$ or $X \models \psi$;
- $X \models \phi \rightarrow \psi$ iff for any team $Y \subseteq X$,
 $Y \models \phi \implies Y \models \psi$.

The logics **PD**, **PD[∨]**, **PID**

- are **downwards closed**, that is, $X \models \phi$ and $Y \subseteq X \implies Y \models \phi$;
- and have the **empty team property**, that is, $\emptyset \models \phi$, for all ϕ .

Team Semantics

Let X be a team.

- $X \models p_i$ iff for all $v \in X$, $v(i) = 1$;
- $X \models \neg p_i$ iff for all $v \in X$, $v(i) = 0$;
- $X \models \perp$ iff $X = \emptyset$;
- $X \models =(p_{i_1}, \dots, p_{i_n})$ iff for all $v, v' \in X$
 $[v(i_1) = v'(i_1), \dots, v(i_{n-1}) = v'(i_{n-1})] \implies v(i_n) = v'(i_n)$;
- $X \models \phi \wedge \psi$ iff $X \models \phi$ and $X \models \psi$;
- $X \models \phi \otimes \psi$ iff there exist teams $Y, Z \subseteq X$ with $X = Y \cup Z$ such that
 $Y \models \phi$ and $Z \models \psi$;
- $X \models \phi \vee \psi$ iff $X \models \phi$ or $X \models \psi$;
- $X \models \phi \rightarrow \psi$ iff for any team $Y \subseteq X$,
 $Y \models \phi \implies Y \models \psi$.

Intuitionistic disjunction \vee has the **disjunction property**:

$$\models \phi \vee \psi \implies \models \phi \text{ or } \models \psi.$$

Dependence atoms are definable in the fragment of **PID** and **PD**[∨] without dependence atoms:

$$\begin{aligned} \equiv(p_0, p_1) &\equiv \left(p_0 \wedge (p_1 \vee \neg p_1) \right) \otimes \left(\neg p_0 \wedge (p_1 \vee \neg p_1) \right) \\ &\equiv (p_0 \vee \neg p_0) \rightarrow (p_1 \vee \neg p_1) \end{aligned}$$

	p_0	p_1	p_2
v_1	1	0	0
v_2	1	0	1
v_3	0	1	0
v_4	0	1	1

Dependence atoms are definable in the fragment of **PID** and **PD**[∨] without dependence atoms:

$$\begin{aligned} \equiv(p_0, p_1) &\equiv \left(p_0 \wedge (p_1 \vee \neg p_1) \right) \otimes \left(\neg p_0 \wedge (p_1 \vee \neg p_1) \right) \\ &\equiv (p_0 \vee \neg p_0) \rightarrow (p_1 \vee \neg p_1) \end{aligned}$$

	p_0	p_1	p_2
v_1	1	0	0
v_2	1	0	1
v_3	0	1	0
v_4	0	1	1

Observation (de Jongh, Litak)

PID[−] is equivalent to inquisitive logic (Ciardelli, Roelofsen 2011).

Fix $N = \{i_1, \dots, i_n\} \subseteq \mathbb{N}$.

- An *n -valuation* on N is a function $s : N \rightarrow \{0, 1\}$.
- An *n -team* on N is a set of n -valuations on N .

There are in total 2^n distinct n -valuations, and 2^{2^n} n -teams, among which there exists a biggest team (denoted by 2^n) consisting of all n -valuations on N .

Fix $N = \{i_1, \dots, i_n\} \subseteq \mathbb{N}$.

- An ***n -valuation*** on N is a function $s : N \rightarrow \{0, 1\}$.
- An ***n -team*** on N is a set of n -valuations on N .

10

There are in total 2^n distinct n -valuations, and 2^{2^n} n -teams, among which there exists a biggest team (denoted by **2^n**) consisting of all n -valuations on N .

Fix $N = \{i_1, \dots, i_n\} \subseteq \mathbb{N}$.


- An ***n-valuation*** on N is a function $s : N \rightarrow \{0, 1\}$.
- An ***n-team*** on N is a set of n -valuations on N .



There are in total 2^n distinct n -valuations, and 2^{2^n} n -teams, among which there exists a biggest team (denoted by **2^n**) consisting of all n -valuations on N .

Fix $N = \{i_1, \dots, i_n\} \subseteq \mathbb{N}$.


- An *n -valuation* on N is a function $s : N \rightarrow \{0, 1\}$.
- An *n -team* on N is a set of n -valuations on N .



There are in total 2^n distinct n -valuations, and 2^{2^n} n -teams, among which there exists a biggest team (denoted by 2^n) consisting of all n -valuations on N .

Fix $N = \{i_1, \dots, i_n\} \subseteq \mathbb{N}$.


- An *n -valuation* on N is a function $s : N \rightarrow \{0, 1\}$.
- An *n -team* on N is a set of n -valuations on N .



There are in total 2^n distinct n -valuations, and 2^{2^n} n -teams, among which there exists a biggest team (denoted by 2^n) consisting of all n -valuations on N .

Fix $N = \{i_1, \dots, i_n\} \subseteq \mathbb{N}$.


- An ***n-valuation*** on N is a function $s : N \rightarrow \{0, 1\}$.
- An ***n-team*** on N is a set of n -valuations on N .



There are in total 2^n distinct n -valuations, and 2^{2^n} n -teams, among which there exists a biggest team (denoted by 2^n) consisting of all n -valuations on N .

Fix $N = \{i_1, \dots, i_n\} \subseteq \mathbb{N}$.


- An *n -valuation* on N is a function $s : N \rightarrow \{0, 1\}$.
- An *n -team* on N is a set of n -valuations on N .



There are in total 2^n distinct n -valuations, and 2^{2^n} n -teams, among which there exists a biggest team (denoted by 2^n) consisting of all n -valuations on N .

Fix $N = \{i_1, \dots, i_n\} \subseteq \mathbb{N}$.


- An *n -valuation* on N is a function $s : N \rightarrow \{0, 1\}$.
- An *n -team* on N is a set of n -valuations on N .



There are in total 2^n distinct n -valuations, and 2^{2^n} n -teams, among which there exists a biggest team (denoted by 2^n) consisting of all n -valuations on N .

Fix $N = \{i_1, \dots, i_n\} \subseteq \mathbb{N}$.


- An *n -valuation* on N is a function $s : N \rightarrow \{0, 1\}$.
- An *n -team* on N is a set of n -valuations on N .



There are in total 2^n distinct n -valuations, and 2^{2^n} n -teams, among which there exists a biggest team (denoted by 2^n) consisting of all n -valuations on N .

Fix $N = \{i_1, \dots, i_n\} \subseteq \mathbb{N}$.


- An *n -valuation* on N is a function $s : N \rightarrow \{0, 1\}$.
- An *n -team* on N is a set of n -valuations on N .



There are in total 2^n distinct n -valuations, and 2^{2^n} n -teams, among which there exists a biggest team (denoted by 2^n) consisting of all n -valuations on N .

Fix $N = \{i_1, \dots, i_n\} \subseteq \mathbb{N}$.


- An *n -valuation* on N is a function $s : N \rightarrow \{0, 1\}$.
- An *n -team* on N is a set of n -valuations on N .



There are in total 2^n distinct n -valuations, and 2^{2^n} n -teams, among which there exists a biggest team (denoted by 2^n) consisting of all n -valuations on N .






Fix $N = \{i_1, \dots, i_n\} \subseteq \mathbb{N}$.

- An *n -valuation* on N is a function $s : N \rightarrow \{0, 1\}$.
- An *n -team* on N is a set of n -valuations on N .




There are in total 2^n distinct n -valuations, and 2^{2^n} n -teams, among which there exists a biggest team (denoted by 2^n) consisting of all n -valuations on N .


$$\wp(2^n)$$




$$(\wp(2^n), \supseteq)$$




$$(\wp(2^n) \setminus \{\emptyset\}, \supseteq)$$




A Medvedev frame: $(\wp(2^n) \setminus \{\emptyset\}, \supseteq)$




A Medvedev frame: $(\wp(2^n) \setminus \{\emptyset\}, \supseteq)$




A Medvedev frame: $(\wp(2^n) \setminus \{\emptyset\}, \supseteq)$



A Medvedev frame: $(\wp(2^n) \setminus \{\emptyset\}, \supseteq)$




A Medvedev frame: $(\wp(2^n) \setminus \{\emptyset\}, \supseteq)$



[Ciardelli, Roelofsen 2011]:

$$\mathbf{PID}^- = \mathbf{ML}^\neg = \{\phi \mid \tau(\phi) \in \mathbf{ML}, \text{ where } \tau(p) = \neg p\}$$

A Medvedev frame: $(\wp(2^n) \setminus \{\emptyset\}, \supseteq)$



[Ciardelli, Roelofsen 2011]:

$$\begin{aligned}
 \mathbf{PID}^- &= \mathbf{ML}^\neg = \{\phi \mid \tau(\phi) \in \mathbf{ML}, \text{ where } \tau(p) = \neg p\} \\
 &= \mathbf{KP}^\neg = \mathbf{KP} \oplus \neg\neg p \rightarrow p
 \end{aligned}$$

Fix $N = \{i_1, \dots, i_n\}$. Put

- $\llbracket \phi(p_{i_1}, \dots, p_{i_n}) \rrbracket := \{X \subseteq 2^n \mid X \models \phi\},$
- $\nabla_N := \{\mathcal{K} \subseteq 2^{2^n} \mid \emptyset \in \mathcal{K}, (X \in \mathcal{K}, Y \subseteq X \implies Y \in \mathcal{K})\}.$

For each $\mathcal{K} \in \nabla_N$, consider $\bigvee_{X \in \mathcal{K}} \Theta_X$. For any n -team Y on N ,

$$Y \models \bigvee \Theta_X \iff \exists X \in \mathcal{K} (Y \subseteq X) \iff Y \in \mathcal{K}.$$

Fix $N = \{i_1, \dots, i_n\}$. Put

- $\llbracket \phi(p_{i_1}, \dots, p_{i_n}) \rrbracket := \{X \subseteq 2^n \mid X \models \phi\},$
- $\nabla_N := \{\mathcal{K} \subseteq 2^{2^n} \mid \emptyset \in \mathcal{K}, (X \in \mathcal{K}, Y \subseteq X \implies Y \in \mathcal{K})\}.$

For each $\mathcal{K} \in \nabla_N$, consider $\bigvee_{X \in \mathcal{K}} \Theta_X$. For any n -team Y on N ,

$$Y \models \bigvee \Theta_X \iff \exists X \in \mathcal{K} (Y \subseteq X) \iff Y \in \mathcal{K}.$$

Fix $N = \{i_1, \dots, i_n\}$. Put

- $\llbracket \phi(p_{i_1}, \dots, p_{i_n}) \rrbracket := \{X \subseteq 2^n \mid X \models \phi\},$
- $\nabla_N := \{\mathcal{K} \subseteq 2^{2^n} \mid \emptyset \in \mathcal{K}, (X \in \mathcal{K}, Y \subseteq X \implies Y \in \mathcal{K})\}.$

Theorem (Ciardelli, Huuskonen, Y.)

PID, **PD**[∨], **PD** are maximal downwards closed logics, i.e., for $L \in \{\mathbf{PID}, \mathbf{PD}^\vee, \mathbf{PD}\},$

$$\nabla_N = \{\llbracket \phi \rrbracket \mid \phi(p_{i_1}, \dots, p_{i_n}) \text{ is an } n\text{-formula of } L\}.$$

In particular, **PID** \equiv **PD**[∨] \equiv **PD**.

For each $\mathcal{K} \in \nabla_N$, consider $\bigvee_{X \in \mathcal{K}} \Theta_X$. For any n -team Y on N ,

$$Y \models \bigvee \Theta_X \iff \exists X \in \mathcal{K} (Y \subseteq X) \iff Y \in \mathcal{K}.$$

Fix $N = \{i_1, \dots, i_n\}$. Put

- $\llbracket \phi(p_{i_1}, \dots, p_{i_n}) \rrbracket := \{X \subseteq 2^n \mid X \models \phi\},$
- $\nabla_N := \{\mathcal{K} \subseteq 2^{2^n} \mid \emptyset \in \mathcal{K}, (X \in \mathcal{K}, Y \subseteq X \implies Y \in \mathcal{K})\}.$

Theorem (Ciardelli, Huuskonen, Y.)

PID, **PD**[∨], **PD** are maximal downwards closed logics, i.e., for $L \in \{\mathbf{PID}, \mathbf{PD}^\vee, \mathbf{PD}\},$

$$\nabla_N = \{\llbracket \phi \rrbracket \mid \phi(p_{i_1}, \dots, p_{i_n}) \text{ is an } n\text{-formula of } L\}.$$

In particular, **PID** \equiv **PD**[∨] \equiv **PD**.

Theorem (Y.)

Every instance of \vee and \rightarrow is definable in **PD**, but \vee and \rightarrow are not uniformly definable in **PD**.

Fix $N = \{i_1, \dots, i_n\}$. Put

- $\llbracket \phi(p_{i_1}, \dots, p_{i_n}) \rrbracket := \{X \subseteq 2^n \mid X \models \phi\},$
- $\nabla_N := \{\mathcal{K} \subseteq 2^{2^n} \mid \emptyset \in \mathcal{K}, (X \in \mathcal{K}, Y \subseteq X \implies Y \in \mathcal{K})\}.$

Theorem (Ciardelli, Huuskonen, Y.)

PID, **PD**[∨], **PD** are maximal downwards closed logics, i.e., for $L \in \{\mathbf{PID}, \mathbf{PD}^\vee, \mathbf{PD}\},$

$$\nabla_N = \{\llbracket \phi \rrbracket \mid \phi(p_{i_1}, \dots, p_{i_n}) \text{ is an } n\text{-formula of } L\}.$$

In particular, **PID** \equiv **PD**[∨] \equiv **PD**.

Proof. We only treat **PID** and **PD**[∨].

For each $\mathcal{K} \in \nabla_N$, consider $\bigvee_{X \in \mathcal{K}} \Theta_X$. For any n -team Y on N ,

$$Y \models \bigvee \Theta_X \iff \exists X \in \mathcal{K} (Y \subseteq X) \iff Y \in \mathcal{K}.$$

Fix $N = \{i_1, \dots, i_n\}$. Put

- $\llbracket \phi(p_{i_1}, \dots, p_{i_n}) \rrbracket := \{X \subseteq 2^n \mid X \models \phi\},$
- $\nabla_N := \{\mathcal{K} \subseteq 2^{2^n} \mid \emptyset \in \mathcal{K}, (X \in \mathcal{K}, Y \subseteq X \implies Y \in \mathcal{K})\}.$

Theorem (Ciardelli, Huuskonen, Y.)

PID, **PD**[∨], **PD** are maximal downwards closed logics, i.e., for $L \in \{\mathbf{PID}, \mathbf{PD}^\vee, \mathbf{PD}\},$

$$\nabla_N = \{\llbracket \phi \rrbracket \mid \phi(p_{i_1}, \dots, p_{i_n}) \text{ is an } n\text{-formula of } L\}.$$

In particular, **PID** \equiv **PD**[∨] \equiv **PD**.

Proof. We only treat **PID** and **PD**[∨]. First, consider an n -team:

$$X \left\{ \begin{array}{cc|cc} & p_1 & p_2 & & \\ \hline v_1 & 1 & 1 & & \\ v_2 & 1 & 0 & & \\ v_3 & 0 & 1 & & \end{array} \right.$$

For each $\mathcal{K} \in \nabla_N$, consider $\bigvee_{X \in \mathcal{K}} \Theta_X$. For any n -team Y on N ,

$$Y \models \bigvee \Theta_X \iff \exists X \in \mathcal{K} (Y \subseteq X) \iff Y \in \mathcal{K}.$$

Fix $N = \{i_1, \dots, i_n\}$. Put

- $\llbracket \phi(p_{i_1}, \dots, p_{i_n}) \rrbracket := \{X \subseteq 2^n \mid X \models \phi\},$
- $\nabla_N := \{\mathcal{K} \subseteq 2^{2^n} \mid \emptyset \in \mathcal{K}, (X \in \mathcal{K}, Y \subseteq X \implies Y \in \mathcal{K})\}.$

Theorem (Ciardelli, Huuskonen, Y.)

PID, **PD**[∨], **PD** are maximal downwards closed logics, i.e., for $L \in \{\mathbf{PID}, \mathbf{PD}^\vee, \mathbf{PD}\},$

$$\nabla_N = \{\llbracket \phi \rrbracket \mid \phi(p_{i_1}, \dots, p_{i_n}) \text{ is an } n\text{-formula of } L\}.$$

In particular, **PID** \equiv **PD**[∨] \equiv **PD**.

Proof. We only treat **PID** and **PD**[∨]. First, consider an n -team:

	p_1	p_2
v_1	1	1
v_2	1	0
v_3	0	1

Let

$$\Theta_X := \left\{ \begin{array}{l} \text{for } \mathbf{PD}; \\ \text{for } \mathbf{PID}. \end{array} \right.$$

Then $Y \models \Theta_X \iff Y \subseteq X$, for any n -team Y .

For each $\mathcal{K} \in \nabla_N$, consider $\bigvee_{X \in \mathcal{K}} \Theta_X$. For any n -team Y on N ,

$$Y \models \bigvee \Theta_X \iff \exists X \in \mathcal{K} (Y \subseteq X) \iff Y \in \mathcal{K}.$$

Fix $N = \{i_1, \dots, i_n\}$. Put

- $\llbracket \phi(p_{i_1}, \dots, p_{i_n}) \rrbracket := \{X \subseteq 2^n \mid X \models \phi\},$
- $\nabla_N := \{\mathcal{K} \subseteq 2^{2^n} \mid \emptyset \in \mathcal{K}, (X \in \mathcal{K}, Y \subseteq X \implies Y \in \mathcal{K})\}.$

Theorem (Ciardelli, Huuskonen, Y.)

PID, **PD**[∨], **PD** are maximal downwards closed logics, i.e., for $L \in \{\mathbf{PID}, \mathbf{PD}^\vee, \mathbf{PD}\},$

$$\nabla_N = \{\llbracket \phi \rrbracket \mid \phi(p_{i_1}, \dots, p_{i_n}) \text{ is an } n\text{-formula of } L\}.$$

In particular, **PID** \equiv **PD**[∨] \equiv **PD**.

Proof. We only treat **PID** and **PD**[∨]. First, consider an n -team:

$$X \left\{ \begin{array}{c|cc} & p_1 & p_2 \\ \hline v_1 & 1 & 1 \\ \hline v_2 & 1 & 0 \\ \hline v_3 & 0 & 1 \end{array} \right.$$

Let

$$\Theta_X := \begin{cases} \bigotimes_{v \in X} (p_{i_1}^{v(i_1)} \wedge \dots \wedge p_{i_n}^{v(i_n)}), & \text{for } \mathbf{PD}; \\ \bigvee_{v \in X} (p_{i_1}^{v(i_1)} \wedge \dots \wedge p_{i_n}^{v(i_n)}), & \text{for } \mathbf{PID}. \end{cases}$$

Then $Y \models \Theta_X \iff Y \subseteq X$, for any n -team Y .

For each $\mathcal{K} \in \nabla_N$, consider $\bigvee_{X \in \mathcal{K}} \Theta_X$. For any n -team Y on N ,

$$Y \models \bigvee_{X \in \mathcal{K}} \Theta_X \iff \exists X \in \mathcal{K} (Y \subseteq X) \iff Y \in \mathcal{K}.$$

Fix $N = \{i_1, \dots, i_n\}$. Put

- $\llbracket \phi(p_{i_1}, \dots, p_{i_n}) \rrbracket := \{X \subseteq 2^n \mid X \models \phi\},$
- $\nabla_N := \{\mathcal{K} \subseteq 2^{2^n} \mid \emptyset \in \mathcal{K}, (X \in \mathcal{K}, Y \subseteq X \implies Y \in \mathcal{K})\}.$

Theorem (Ciardelli, Huuskonen, Y.)

PID, **PD**[∨], **PD** are maximal downwards closed logics, i.e., for $L \in \{\mathbf{PID}, \mathbf{PD}^\vee, \mathbf{PD}\},$

$$\nabla_N = \{\llbracket \phi \rrbracket \mid \phi(p_{i_1}, \dots, p_{i_n}) \text{ is an } n\text{-formula of } L\}.$$

In particular, **PID** \equiv **PD**[∨] \equiv **PD**.

Proof. We only treat **PID** and **PD**[∨]. First, consider an n -team:

$$X \left\{ \begin{array}{c|cc} & p_1 & p_2 \\ \hline v_1 & 1 & 1 \\ \hline v_2 & 1 & 0 \\ \hline v_3 & 0 & 1 \end{array} \right.$$

Let

$$\Theta_X := \begin{cases} \bigotimes_{v \in X} (p_{i_1}^{v(i_1)} \wedge \dots \wedge p_{i_n}^{v(i_n)}), & \text{for } \mathbf{PD}; \\ \neg \neg \bigvee_{v \in X} (p_{i_1}^{v(i_1)} \wedge \dots \wedge p_{i_n}^{v(i_n)}), & \text{for } \mathbf{PID}. \end{cases}$$

Then $Y \models \Theta_X \iff Y \subseteq X$, for any n -team Y .

For each $\mathcal{K} \in \nabla_N$, consider $\bigvee_{X \in \mathcal{K}} \Theta_X$. For any n -team Y on N ,

$$Y \models \bigvee_{X \in \mathcal{K}} \Theta_X \iff \exists X \in \mathcal{K} (Y \subseteq X) \iff Y \in \mathcal{K}.$$

Fix $N = \{i_1, \dots, i_n\}$. Put

- $\llbracket \phi(p_{i_1}, \dots, p_{i_n}) \rrbracket := \{X \subseteq 2^n \mid X \models \phi\},$
- $\nabla_N := \{\mathcal{K} \subseteq 2^{2^n} \mid \emptyset \in \mathcal{K}, (X \in \mathcal{K}, Y \subseteq X \implies Y \in \mathcal{K})\}.$

Theorem (Ciardelli, Huuskonen, Y.)

PID, **PD**[∨], **PD** are maximal downwards closed logics, i.e., for $L \in \{\mathbf{PID}, \mathbf{PD}^\vee, \mathbf{PD}\},$

$$\nabla_N = \{\llbracket \phi \rrbracket \mid \phi(p_{i_1}, \dots, p_{i_n}) \text{ is an } n\text{-formula of } L\}.$$

In particular, **PID** \equiv **PD**[∨] \equiv **PD**.

Proof. We only treat **PID** and **PD**[∨]. First, consider an n -team:

$$X \left\{ \begin{array}{c|cc} & p_1 & p_2 \\ \hline v_1 & 1 & 1 \\ \hline v_2 & 1 & 0 \\ \hline v_3 & 0 & 1 \end{array} \right.$$

Let

$$\Theta_X := \begin{cases} \bigotimes_{v \in X} (p_{i_1}^{v(i_1)} \wedge \dots \wedge p_{i_n}^{v(i_n)}), & \text{for } \mathbf{PD}; \\ \neg \neg \bigvee_{v \in X} (p_{i_1}^{v(i_1)} \wedge \dots \wedge p_{i_n}^{v(i_n)}), & \text{for } \mathbf{PID}. \end{cases}$$

Then $Y \models \Theta_X \iff Y \subseteq X$, for any n -team Y .

For each $\mathcal{K} \in \nabla_N$, consider $\bigvee_{X \in \mathcal{K}} \Theta_X$. For any n -team Y on N ,

$$Y \models \bigvee \Theta_X \iff \exists X \in \mathcal{K} (Y \subseteq X) \iff Y \in \mathcal{K}.$$

Theorem (Ciardelli, Huuskonen, Y.)

PID, **PD**[∨], **PD** are maximal downwards closed logics, i.e., for $L \in \{\mathbf{PID}, \mathbf{PD}^\vee, \mathbf{PD}\}$,

$$\nabla_N = \{ \llbracket \phi \rrbracket \mid \phi(p_{i_1}, \dots, p_{i_n}) \text{ is an } n\text{-formula of } L \}.$$

In particular, **PID** \equiv **PD**[∨] \equiv **PD**.

Proof. We only treat **PID** and **PD**[∨]. First, consider an n -team:

$$X \left\{ \begin{array}{c|cc} & p_1 & p_2 \\ \hline v_1 & 1 & 1 \\ v_2 & 1 & 0 \\ \hline v_3 & 0 & 1 \end{array} \right.$$

Let

$$\Theta_X := \begin{cases} \bigotimes_{v \in X} (p_{i_1}^{v(i_1)} \wedge \dots \wedge p_{i_n}^{v(i_n)}), & \text{for } \mathbf{PD}; \\ \neg \neg \bigvee_{v \in X} (p_{i_1}^{v(i_1)} \wedge \dots \wedge p_{i_n}^{v(i_n)}), & \text{for } \mathbf{PID}. \end{cases}$$

Then $Y \models \Theta_X \iff Y \subseteq X$, for any n -team Y .

For each $\mathcal{K} \in \nabla_N$, consider $\bigvee_{X \in \mathcal{K}} \Theta_X$. For any n -team Y on N ,

$$Y \models \bigvee_{X \in \mathcal{K}} \Theta_X \iff \exists X \in \mathcal{K} (Y \subseteq X) \iff Y \in \mathcal{K}.$$

Hence $\llbracket \bigvee_{X \in \mathcal{K}} \Theta_X \rrbracket = \mathcal{K}$.

□

Theorem. **PID**, **PD[∨]**, **PD** are sound and complete w.r.t. their deductive systems.

Corrolary. For every formula ϕ of **PID** and **PD[∨]**, $\phi \dashv\vdash \bigvee_{i \in I} \Theta_{X_i}$.

A Hilbert Style deductive system for **PID⁻** (Ciardelli, Roelofsen 2011)

Axioms:

- all substitution instances of **IPC** axioms
- all substitution instances of

$$(KP) \quad (\neg p_i \rightarrow (p_j \vee p_k)) \rightarrow ((\neg p_i \rightarrow p_j) \vee (\neg p_i \rightarrow p_k)).$$

- $\neg\neg p_i \rightarrow p_i$ for all propositional variables p_i

Rules:

- Modus Ponens

Natural deduction systems for **PD** and **PD[∨]** (Väänänen, Y.)

For $L \in \{\mathbf{PD}, \mathbf{PD}^\vee\}$, if ϕ does not contain any \vee or dependence atoms, then $\vdash_{\mathbf{CPC}} \phi \iff \vdash_L \phi$.

Theorem. **PID**, **PD**[∨], **PD** are sound and complete w.r.t. their deductive systems.

Corrolary. For every formula ϕ of **PID** and **PD**[∨], $\phi \dashv\vdash \bigvee_{i \in I} \Theta_{X_i}$.

A Hilbert Style deductive system for **PID**[−] (Ciardelli, Roelofsen 2011)

Axioms:

- all substitution instances of **IPC** axioms
- all substitution instances of

$$(KP) \quad (\neg p_i \rightarrow (p_j \vee p_k)) \rightarrow ((\neg p_i \rightarrow p_j) \vee (\neg p_i \rightarrow p_k)).$$

- $\neg\neg p_i \rightarrow p_i$ for all propositional variables p_i

Rules:

- Modus Ponens

Natural deduction systems for **PD** and **PD**[∨] (Väänänen, Y.)

For $L \in \{\mathbf{PD}, \mathbf{PD}^\vee\}$, if ϕ does not contain any \vee or dependence atoms, then $\vdash_{\mathbf{CPC}} \phi \iff \vdash_L \phi$.

Theorem. **PID**, **PD**[∨], **PD** are sound and complete w.r.t. their deductive systems.

Corrolary. For every formula ϕ of **PID** and **PD**[∨], $\phi \dashv\vdash \bigvee_{i \in I} \Theta_{X_i}$.

A Hilbert Style deductive system for **PID**[−] (Ciardelli, Roelofsen 2011)

Axioms:

- all substitution instances of **IPC** axioms
- all substitution instances of

$$(KP) \quad (\neg p_i \rightarrow (p_j \vee p_k)) \rightarrow ((\neg p_i \rightarrow p_j) \vee (\neg p_i \rightarrow p_k)).$$

- $\neg\neg p_i \rightarrow p_i$ for all propositional variables p_i

Rules:

- Modus Ponens

Natural deduction systems for **PD** and **PD**[∨] (Väänänen, Y.)

For $L \in \{\mathbf{PD}, \mathbf{PD}^\vee\}$, if ϕ does not contain any \vee or dependence atoms, then $\vdash_{\mathbf{CPC}} \phi \iff \vdash_L \phi$.

Definition


A formula ϕ is said to be **flat** iff for all teams X ,

$$X \models \phi \iff \forall v \in X, \{v\} \models \phi.$$

Definition

A formula ϕ is said to be **flat** iff for all teams X ,


$$X \models \phi \iff \forall v \in X, \{v\} \models \phi.$$



Definition

A formula ϕ is said to be **flat** iff for all teams X ,


$$X \models \phi \iff \forall v \in X, \{v\} \models \phi.$$



Definition

A formula ϕ is said to be **flat** iff for all teams X ,


$$X \models \phi \iff \forall v \in X, \{v\} \models \phi.$$



Definition

A formula ϕ is said to be **flat** iff for all teams X ,

$$X \models \phi \iff \forall v \in X, \{v\} \models \phi.$$




Fact: Formulas with no occurrences of dependence atoms or intuitionistic disjunction \vee are flat.

Definition

A formula ϕ is said to be **flat** iff for all teams X ,

$$X \models \phi \iff \forall v \in X, \{v\} \models \phi.$$




Fact: Formulas with no occurrences of dependence atoms or intuitionistic disjunction \vee are flat.

$$\text{E.g. } \Theta_X = \begin{cases} \bigotimes_{v \in X} (p_{i_1}^{v(i_1)} \wedge \dots \wedge p_{i_n}^{v(i_n)}), & \text{for } \mathbf{PD}; \\ \neg \neg \bigvee_{v \in X} (p_{i_1}^{v(i_1)} \wedge \dots \wedge p_{i_n}^{v(i_n)}), & \text{for } \mathbf{PID}. \end{cases} \quad \text{is flat}$$

Definition

A formula ϕ is said to be **flat** iff for all teams X ,

$$X \models \phi \iff \forall v \in X, \{v\} \models \phi.$$



Fact: Formulas with no occurrences of dependence atoms or intuitionistic disjunction \vee are flat.

Lemma. In **PID**, a formula ϕ is flat if $\vdash \phi \leftrightarrow \neg\neg\phi$.

$$\text{E.g. } \Theta_X = \begin{cases} \bigotimes_{v \in X} (p_{i_1}^{v(i_1)} \wedge \dots \wedge p_{i_n}^{v(i_n)}), & \text{for } \mathbf{PD}; \\ \neg\neg \bigvee_{v \in X} (p_{i_1}^{v(i_1)} \wedge \dots \wedge p_{i_n}^{v(i_n)}), & \text{for } \mathbf{PID}. \end{cases} \quad \text{is flat}$$

Definition

A formula ϕ is said to be **flat** iff for all teams X ,

$$X \models \phi \iff \forall v \in X, \{v\} \models \phi.$$

Lemma

A formula ϕ is flat iff $\phi \dashv\vdash \Theta_X$ for some X .

Definition

A formula ϕ is said to be **flat** iff for all teams X ,

$$X \models \phi \iff \forall v \in X, \{v\} \models \phi.$$

Lemma

A formula ϕ is flat iff $\phi \dashv\vdash \Theta_X$ for some X .

Proof. “ \implies ”: Suppose ϕ is flat. Note that $\phi \dashv\vdash \Theta_{X_1} \vee \dots \vee \Theta_{X_k}$,

Definition

A formula ϕ is said to be **flat** iff for all teams X ,


$$X \models \phi \iff \forall v \in X, \{v\} \models \phi.$$

Fact: $Y \models \Theta_X \iff Y \subseteq X$, for any n -team Y .

Lemma

A formula ϕ is flat iff $\phi \dashv\vdash \Theta_X$ for some X .

Proof. “ \implies ”: Suppose ϕ is flat. Note that $\phi \Vdash \Theta_{X_1} \vee \dots \vee \Theta_{X_k}$, where w.l.o.g. we assume that X_i ’s are maximal. If $k > 1$, then pick



Definition

A formula ϕ is said to be **flat** iff for all teams X ,


$$X \models \phi \iff \forall v \in X, \{v\} \models \phi.$$

Fact: $Y \models \Theta_X \iff Y \subseteq X$, for any n -team Y .

Lemma

A formula ϕ is flat iff $\phi \dashv\vdash \Theta_X$ for some X .

Proof. “ \implies ”: Suppose ϕ is flat. Note that $\phi \dashv\vdash \Theta_{X_1} \vee \dots \vee \Theta_{X_k}$, where w.l.o.g. we assume that X_i 's are maximal. If $k > 1$, then pick



Definition

A formula ϕ is said to be **flat** iff for all teams X ,


$$X \models \phi \iff \forall v \in X, \{v\} \models \phi.$$

Fact: $Y \models \Theta_X \iff Y \subseteq X$, for any n -team Y .

Lemma

A formula ϕ is flat iff $\phi \dashv\vdash \Theta_X$ for some X .

Proof. “ \implies ”: Suppose ϕ is flat. Note that $\phi \dashv\vdash \Theta_{X_1} \vee \dots \vee \Theta_{X_k}$, where w.l.o.g. we assume that X_i 's are maximal. If $k > 1$, then pick



Definition

A formula ϕ is said to be **flat** iff for all teams X ,


$$X \models \phi \iff \forall v \in X, \{v\} \models \phi.$$

Fact: $Y \models \Theta_X \iff Y \subseteq X$, for any n -team Y .

Lemma

A formula ϕ is flat iff $\phi \dashv\vdash \Theta_X$ for some X .

Proof. “ \implies ”: Suppose ϕ is flat. Note that $\phi \dashv\vdash \Theta_{X_1} \vee \dots \vee \Theta_{X_k}$, where w.l.o.g. we assume that X_i 's are maximal. If $k > 1$, then pick



$\{v_i\} \subseteq X_i$ and $\{v_1, \dots, v_k\} \not\subseteq X_i$ for all $1 \leq i \leq k$

$\implies \{v_i\} \models \Theta_{X_i}$ and $\{v_1, \dots, v_k\} \not\models \Theta_{X_i}$ for all $1 \leq i \leq k$

$\implies \{v_i\} \models \phi$ for all $1 \leq i \leq k$ whereas $\{v_1, \dots, v_k\} \not\models \phi$

Definition

A formula ϕ is said to be **flat** iff for all teams X ,


$$X \models \phi \iff \forall v \in X, \{v\} \models \phi.$$

Fact: $Y \models \Theta_X \iff Y \subseteq X$, for any n -team Y .

Lemma

A formula ϕ is flat iff $\phi \dashv\vdash \Theta_X$ for some X .

Proof. “ \implies ”: Suppose ϕ is flat. Note that $\phi \dashv\vdash \Theta_{X_1} \vee \dots \vee \Theta_{X_k}$, where w.l.o.g. we assume that X_i 's are maximal. If $k > 1$, then pick



$\{v_i\} \subseteq X_i$ and $\{v_1, \dots, v_k\} \not\subseteq X_i$ for all $1 \leq i \leq k$

$\implies \{v_i\} \models \Theta_{X_i}$ and $\{v_1, \dots, v_k\} \not\models \Theta_{X_i}$ for all $1 \leq i \leq k$

$\implies \{v_i\} \models \phi$ for all $1 \leq i \leq k$ whereas $\{v_1, \dots, v_k\} \not\models \phi$

$\implies k = 1$ and $\phi \dashv\vdash \Theta_{X_1}$.



We say that a substitution σ of a logic L is **well-behaved**, if

- σ is well-defined: $\sigma(\phi)$ is a well-formed formula of L
- \vdash_L is closed under σ : $\phi \vdash_L \psi$ implies $\sigma(\phi) \vdash_L \sigma(\psi)$

We say that a substitution σ of a logic L is **well-behaved**, if

- σ is well-defined: $\sigma(\phi)$ is a well-formed formula of L
- \vdash_L is closed under σ : $\phi \vdash_L \psi$ implies $\sigma(\phi) \vdash_L \sigma(\psi)$

Recall: Well-formed formulas of **PD** are built from the following grammar:

$$\phi ::= p_i \mid \neg p_i \mid (p_{i_1}, \dots, p_{i_k}) \mid \phi \wedge \phi \mid \phi \otimes \phi$$

where $p_i, p_{i_1}, \dots, p_{i_k}$ are propositional variables.

We say that a substitution σ of a logic L is **well-behaved**, if

- σ is well-defined: $\sigma(\phi)$ is a well-formed formula of L
- \vdash_L is closed under σ : $\phi \vdash_L \psi$ implies $\sigma(\phi) \vdash_L \sigma(\psi)$

For **PID**: $\vdash \neg\neg p \rightarrow p$, but $\nvdash \neg\neg(p \vee \neg p) \rightarrow (p \vee \neg p)$

We say that a substitution σ of a logic L is **well-behaved**, if

- σ is well-defined: $\sigma(\phi)$ is a well-formed formula of L
- \vdash_L is closed under σ : $\phi \vdash_L \psi$ implies $\sigma(\phi) \vdash_L \sigma(\psi)$

For **PID**: $\vdash \neg\neg p \rightarrow p$, but $\not\vdash \neg\neg(p \vee \neg p) \rightarrow (p \vee \neg p)$

Definition

A substitution σ is called a **flat substitution** if $\sigma(p)$ is flat for all p .

Lemma

*Flat substitutions are well-behaved in **PID** and **PD**.*

Proof. For **PID**, it follows from (Ciardelli). For **PD**, nontrivial. □

Definition (Projective formula)

Let S be a set of **well-behaved** substitutions of a logic L . An L -formula ϕ is said to be **S -projective** if there exists $\sigma \in S$ such that

(1) $\vdash_L \sigma(\phi)$

(2) $\phi, \sigma(\psi) \vdash_L \psi$ and $\phi, \psi \vdash_L \sigma(\psi)$ for all L -formulas ψ .

Definition (Projective formula)

Let S be a set of **well-behaved** substitutions of a logic L . An L -formula ϕ is said to be **S -projective** if there exists $\sigma \in S$ such that

$$(1) \vdash_L \sigma(\phi)$$

$$(2) \phi, \sigma(\psi) \vdash_L \psi \text{ and } \phi, \psi \vdash_L \sigma(\psi) \text{ for all } L\text{-formulas } \psi.$$

Lemma

Let $L \in \{\mathbf{PD}, \mathbf{PID}\}$ and F the set of all flat substitutions. An n -formula ϕ of L is F -projective iff $\phi \dashv\vdash \Theta_X$ for some n -team X .

Definition (Projective formula)

Let S be a set of **well-behaved** substitutions of a logic L . An L -formula ϕ is said to be **S -projective** if there exists $\sigma \in S$ such that

$$(1) \vdash_L \sigma(\phi)$$

$$(2) \phi, \sigma(\psi) \vdash_L \psi \text{ and } \phi, \psi \vdash_L \sigma(\psi) \text{ for all } L\text{-formulas } \psi.$$

Lemma

Let $L \in \{\mathbf{PD}, \mathbf{PID}\}$ and F the set of all flat substitutions. An n -formula ϕ of L is F -projective iff $\phi \dashv\vdash \Theta_X$ for some n -team X .

Proof. “ \implies ”: Note that $\phi \equiv \bigvee_{i \in I} \Theta_{x_i}$.

Definition (Projective formula)

Let S be a set of **well-behaved** substitutions of a logic L . An L -formula ϕ is said to be **S -projective** if there exists $\sigma \in S$ such that

$$(1) \vdash_L \sigma(\phi)$$

$$(2) \phi, \sigma(\psi) \vdash_L \psi \text{ and } \phi, \psi \vdash_L \sigma(\psi) \text{ for all } L\text{-formulas } \psi.$$

Lemma

Let $L \in \{\mathbf{PD}, \mathbf{PID}\}$ and F the set of all flat substitutions. An n -formula ϕ of L is F -projective iff $\phi \dashv\vdash \Theta_X$ for some n -team X .

Proof. “ \implies ”: Note that $\phi \equiv \bigvee_{i \in I} \Theta_{X_i}$. Let σ be the F -projective unifier of ϕ . By (1), $\vdash \sigma(\phi) \implies \vdash \sigma(\Theta_{X_i})$ for some $i \in I$ (by DP)
 $\implies \phi \models \Theta_{X_i}$ (by (2)).

Definition (Projective formula)

Let S be a set of **well-behaved** substitutions of a logic L . An L -formula ϕ is said to be **S-projective** if there exists $\sigma \in S$ such that

$$(1) \vdash_L \sigma(\phi)$$

$$(2) \phi, \sigma(\psi) \vdash_L \psi \text{ and } \phi, \psi \vdash_L \sigma(\psi) \text{ for all } L\text{-formulas } \psi.$$

Lemma

Let $L \in \{\mathbf{PD}, \mathbf{PID}\}$ and F the set of all flat substitutions. An n -formula ϕ of L is F -projective iff $\phi \dashv\vdash \Theta_X$ for some n -team X .

Proof. “ \implies ”: Note that $\phi \equiv \bigvee_{i \in I} \Theta_{X_i}$. Let σ be the F -projective unifier of ϕ . By (1), $\vdash \sigma(\phi) \implies \vdash \sigma(\Theta_{X_i})$ for some $i \in I$ (by DP)
 $\implies \phi \models \Theta_{X_i}$ (by (2)).

“ \impliedby ”: Recall:
$$\Theta_X = \begin{cases} \bigotimes_{v \in X} (p_{i_1}^{v(i_1)} \wedge \dots \wedge p_{i_n}^{v(i_n)}), & \text{for } \mathbf{PD}; \\ \neg \neg \bigvee_{v \in X} (p_{i_1}^{v(i_1)} \wedge \dots \wedge p_{i_n}^{v(i_n)}), & \text{for } \mathbf{PID}. \end{cases}$$

Definition (Projective formula)

Let S be a set of **well-behaved** substitutions of a logic L . An L -formula ϕ is said to be **S-projective** if there exists $\sigma \in S$ such that

$$(1) \vdash_L \sigma(\phi)$$

$$(2) \phi, \sigma(\psi) \vdash_L \psi \text{ and } \phi, \psi \vdash_L \sigma(\psi) \text{ for all } L\text{-formulas } \psi.$$

Lemma

Let $L \in \{\mathbf{PD}, \mathbf{PID}\}$ and F the set of all flat substitutions. An n -formula ϕ of L is F -projective iff $\phi \Vdash \Theta_X$ for some n -team X .

Proof. “ \implies ”: Note that $\phi \equiv \bigvee_{i \in I} \Theta_{X_i}$. Let σ be the F -projective unifier of ϕ . By (1), $\models \sigma(\phi) \implies \models \sigma(\Theta_{X_i})$ for some $i \in I$ (by DP)
 $\implies \phi \models \Theta_{X_i}$ (by (2)).

$$\text{“}\Leftarrow\text{”}: \text{Recall: } \Theta_X = \begin{cases} \bigotimes_{v \in X} (p_{i_1}^{v(i_1)} \wedge \dots \wedge p_{i_n}^{v(i_n)}), & \text{for } \mathbf{PD}; \\ \neg \neg \bigvee_{v \in X} (p_{i_1}^{v(i_1)} \wedge \dots \wedge p_{i_n}^{v(i_n)}), & \text{for } \mathbf{PID}. \end{cases}$$

We only give the proof for **PD**.

Definition (Projective formula)

Let S be a set of **well-behaved** substitutions of a logic L . An L -formula ϕ is said to be **S-projective** if there exists $\sigma \in S$ such that

- (1) $\vdash_L \sigma(\phi)$
- (2) $\phi, \sigma(\psi) \vdash_L \psi$ and $\phi, \psi \vdash_L \sigma(\psi)$ for all L -formulas ψ .

Lemma

Let $L \in \{\mathbf{PD}, \mathbf{PID}\}$ and F the set of all flat substitutions. An n -formula ϕ of L is F -projective iff $\phi \dashv\vdash \Theta_X$ for some n -team X .

Proof. (ctd.) View $\phi = \Theta_X = \bigotimes_{v \in X} (p_{i_1}^{v(i_1)} \wedge \dots \wedge p_{i_n}^{v(i_n)})$ as a formula of **CPC**,

Definition (Projective formula)

Let S be a set of **well-behaved** substitutions of a logic L . An L -formula ϕ is said to be **S -projective** if there exists $\sigma \in S$ such that

- (1) $\vdash_L \sigma(\phi)$
- (2) $\phi, \sigma(\psi) \vdash_L \psi$ and $\phi, \psi \vdash_L \sigma(\psi)$ for all L -formulas ψ .

Lemma

Let $L \in \{\mathbf{PD}, \mathbf{PID}\}$ and F the set of all flat substitutions. An n -formula ϕ of L is F -projective iff $\phi \dashv\vdash \Theta_X$ for some n -team X .

Proof. (ctd.) View $\phi = \Theta_X = \bigotimes_{v \in X} (p_{i_1}^{v(i_1)} \wedge \dots \wedge p_{i_n}^{v(i_n)})$ as a formula of **CPC**, then $v(\phi_i) = 1$ for any $v \in X$.

Definition (Projective formula)

Let S be a set of **well-behaved** substitutions of a logic L . An L -formula ϕ is said to be **S -projective** if there exists $\sigma \in S$ such that

$$(1) \vdash_L \sigma(\phi)$$

$$(2) \phi, \sigma(\psi) \vdash_L \psi \text{ and } \phi, \psi \vdash_L \sigma(\psi) \text{ for all } L\text{-formulas } \psi.$$

Lemma

Let $L \in \{\mathbf{PD}, \mathbf{PID}\}$ and F the set of all flat substitutions. An n -formula ϕ of L is F -projective iff $\phi \dashv\vdash \Theta_X$ for some n -team X .

Proof. (ctd.) View $\phi = \Theta_X = \bigotimes_{v \in X} (p_{i_1}^{v(i_1)} \wedge \dots \wedge p_{i_n}^{v(i_n)})$ as a formula of **CPC**, then $v(\phi_i) = 1$ for any $v \in X$. Define σ_v^ϕ as follows:

$$\sigma_v^\phi(p) = \begin{cases} \phi \wedge p, & \text{if } v(p) = 0; \\ \phi \rightarrow p, & \text{if } v(p) = 1. \end{cases} \quad (\text{Prucnal's trick})$$

Definition (Projective formula)

Let S be a set of **well-behaved** substitutions of a logic L . An L -formula ϕ is said to be **S -projective** if there exists $\sigma \in S$ such that

$$(1) \vdash_L \sigma(\phi)$$

$$(2) \phi, \sigma(\psi) \vdash_L \psi \text{ and } \phi, \psi \vdash_L \sigma(\psi) \text{ for all } L\text{-formulas } \psi.$$

Lemma

Let $L \in \{\mathbf{PD}, \mathbf{PID}\}$ and F the set of all flat substitutions. An n -formula ϕ of L is F -projective iff $\phi \dashv\vdash \Theta_X$ for some n -team X .

Proof. (ctd.) View $\phi = \Theta_X = \bigotimes_{v \in X} (p_{i_1}^{v(i_1)} \wedge \dots \wedge p_{i_n}^{v(i_n)})$ as a formula of **CPC**, then $v(\phi_i) = 1$ for any $v \in X$. Define σ_V^ϕ as follows:

$$\sigma_V^\phi(p) = \begin{cases} \phi \wedge p, & \text{if } v(p) = 0; \\ \phi \rightarrow p, & \text{if } v(p) = 1. \end{cases} \quad (\text{Prucnal's trick})$$

Definition (Projective formula)

Let S be a set of **well-behaved** substitutions of a logic L . An L -formula ϕ is said to be **S -projective** if there exists $\sigma \in S$ such that

$$(1) \vdash_L \sigma(\phi)$$

$$(2) \phi, \sigma(\psi) \vdash_L \psi \text{ and } \phi, \psi \vdash_L \sigma(\psi) \text{ for all } L\text{-formulas } \psi.$$

Lemma

Let $L \in \{\mathbf{PD}, \mathbf{PID}\}$ and F the set of all flat substitutions. An n -formula ϕ of L is F -projective iff $\phi \dashv\vdash \Theta_X$ for some n -team X .

Proof. (ctd.) View $\phi = \Theta_X = \bigotimes_{v \in X} (p_{i_1}^{v(i_1)} \wedge \dots \wedge p_{i_n}^{v(i_n)})$ as a formula of **CPC**, then $v(\phi_i) = 1$ for any $v \in X$. Define σ_v^ϕ as follows:

$$\sigma_v^\phi(p) = \begin{cases} \phi \wedge p, & \text{if } v(p) = 0; \\ \phi^\sim \otimes p, & \text{if } v(p) = 1. \end{cases} \quad (\text{Prucnal's trick})$$

Definition (Projective formula)

Let S be a set of **well-behaved** substitutions of a logic L . An L -formula ϕ is said to be **S -projective** if there exists $\sigma \in S$ such that

$$(1) \vdash_L \sigma(\phi)$$

$$(2) \phi, \sigma(\psi) \vdash_L \psi \text{ and } \phi, \psi \vdash_L \sigma(\psi) \text{ for all } L\text{-formulas } \psi.$$

Lemma

Let $L \in \{\mathbf{PD}, \mathbf{PID}\}$ and F the set of all flat substitutions. An n -formula ϕ of L is F -projective iff $\phi \dashv\vdash \Theta_X$ for some n -team X .

Proof. (ctd.) View $\phi = \Theta_X = \bigotimes_{v \in X} (p_{i_1}^{v(i_1)} \wedge \dots \wedge p_{i_n}^{v(i_n)})$ as a formula of **CPC**, then $v(\phi_i) = 1$ for any $v \in X$. Define σ_v^ϕ as follows:

$$\sigma_v^\phi(p) = \begin{cases} \phi \wedge p, & \text{if } v(p) = 0; \\ \phi \sim \otimes p, & \text{if } v(p) = 1. \end{cases} \quad (\text{Prucnal's trick})$$

Put $\sigma = \sigma_v^\phi$. Then $\vdash_{\mathbf{CPC}} \sigma(\phi)$ as $v(\phi) = 1$.

Definition (Projective formula)

Let S be a set of **well-behaved** substitutions of a logic L . An L -formula ϕ is said to be **S -projective** if there exists $\sigma \in S$ such that

$$(1) \vdash_L \sigma(\phi)$$

$$(2) \phi, \sigma(\psi) \vdash_L \psi \text{ and } \phi, \psi \vdash_L \sigma(\psi) \text{ for all } L\text{-formulas } \psi.$$

Lemma

Let $L \in \{\mathbf{PD}, \mathbf{PID}\}$ and F the set of all flat substitutions. An n -formula ϕ of L is F -projective iff $\phi \dashv\vdash \Theta_X$ for some n -team X .

Proof. (ctd.) View $\phi = \Theta_X = \bigotimes_{v \in X} (p_{i_1}^{v(i_1)} \wedge \dots \wedge p_{i_n}^{v(i_n)})$ as a formula of **CPC**, then $v(\phi_i) = 1$ for any $v \in X$. Define σ_v^ϕ as follows:

$$\sigma_v^\phi(p) = \begin{cases} \phi \wedge p, & \text{if } v(p) = 0; \\ \phi \sim \otimes p, & \text{if } v(p) = 1. \end{cases} \quad (\text{Prucnal's trick})$$

Put $\sigma = \sigma_v^\phi$. Then $\vdash_{\mathbf{CPC}} \sigma(\phi)$ as $v(\phi) = 1$. Clearly, $\sigma \in F$

Definition (Projective formula)

Let S be a set of **well-behaved** substitutions of a logic L . An L -formula ϕ is said to be **S -projective** if there exists $\sigma \in S$ such that

$$(1) \vdash_L \sigma(\phi)$$

$$(2) \phi, \sigma(\psi) \vdash_L \psi \text{ and } \phi, \psi \vdash_L \sigma(\psi) \text{ for all } L\text{-formulas } \psi.$$

Lemma

Let $L \in \{\mathbf{PD}, \mathbf{PID}\}$ and F the set of all flat substitutions. An n -formula ϕ of L is F -projective iff $\phi \dashv\vdash \Theta_X$ for some n -team X .

Proof. (ctd.) View $\phi = \Theta_X = \bigotimes_{v \in X} (p_{i_1}^{v(i_1)} \wedge \dots \wedge p_{i_n}^{v(i_n)})$ as a formula of **CPC**, then $v(\phi_i) = 1$ for any $v \in X$. Define σ_v^ϕ as follows:

$$\sigma_v^\phi(p) = \begin{cases} \phi \wedge p, & \text{if } v(p) = 0; \\ \phi \sim \otimes p, & \text{if } v(p) = 1. \end{cases} \quad (\text{Prucnal's trick})$$

Put $\sigma = \sigma_v^\phi$. Then $\vdash_{\mathbf{CPC}} \sigma(\phi)$ as $v(\phi) = 1$. Clearly, $\sigma \in F$ and $\sigma(\phi)$ is classical, thus $\vdash_{\mathbf{PD}} \sigma(\phi)$.

Definition (Projective formula)

Let S be a set of **well-behaved** substitutions of a logic L . An L -formula ϕ is said to be **S -projective** if there exists $\sigma \in S$ such that

$$(1) \vdash_L \sigma(\phi)$$

$$(2) \phi, \sigma(\psi) \vdash_L \psi \text{ and } \phi, \psi \vdash_L \sigma(\psi) \text{ for all } L\text{-formulas } \psi.$$

Lemma

Let $L \in \{\mathbf{PD}, \mathbf{PID}\}$ and F the set of all flat substitutions. An n -formula ϕ of L is F -projective iff $\phi \dashv\vdash \Theta_X$ for some n -team X .

Proof. (ctd.) View $\phi = \Theta_X = \bigotimes_{v \in X} (p_{i_1}^{v(i_1)} \wedge \dots \wedge p_{i_n}^{v(i_n)})$ as a formula of **CPC**, then $v(\phi_i) = 1$ for any $v \in X$. Define σ_v^ϕ as follows:

$$\sigma_v^\phi(p) = \begin{cases} \phi \wedge p, & \text{if } v(p) = 0; \\ \phi^\sim \otimes p, & \text{if } v(p) = 1. \end{cases} \quad (\text{Prucnal's trick})$$

Put $\sigma = \sigma_v^\phi$. Then $\vdash_{\mathbf{CPC}} \sigma(\phi)$ as $v(\phi) = 1$. Clearly, $\sigma \in F$ and $\sigma(\phi)$ is classical, thus $\vdash_{\mathbf{PD}} \sigma(\phi)$. Moreover, (2) is satisfied by the choice of σ . □

Definition (Exact formula)

Let S be a set of **well-behaved** substitutions of a logic L . An L -formula ϕ is said to be **S-exact** if there exists $\sigma \in S$ such that

$$(1) \vdash_L \sigma(\phi)$$

$$(2) \vdash_L \sigma(\psi) \implies \phi \vdash_L \psi \text{ for all } L\text{-formulas } \psi$$

Lemma

Let $L \in \{\mathbf{PD}, \mathbf{PID}\}$ and F the set of all flat substitutions. An n -formula ϕ of L is F -exact iff $\phi \dashv\vdash \Theta_X$ for some n -team X .

Corollary

For formulas ϕ of the logics \mathbf{PD} and \mathbf{PID} , TFAE:

- ϕ is flat;
- ϕ is F -projective;
- ϕ is F -exact;
- $\phi \dashv\vdash \Theta_X$ for some X

Definition (Exact formula)

Let S be a set of **well-behaved** substitutions of a logic L . An L -formula ϕ is said to be **S-exact** if there exists $\sigma \in S$ such that

$$(1) \vdash_L \sigma(\phi)$$

$$(2) \vdash_L \sigma(\psi) \implies \phi \vdash_L \psi \text{ for all } L\text{-formulas } \psi$$

Lemma

Let $L \in \{\mathbf{PD}, \mathbf{PID}\}$ and F the set of all flat substitutions. An n -formula ϕ of L is F -exact iff $\phi \dashv\vdash \Theta_X$ for some n -team X .

Corollary

For formulas ϕ of the logics **PD** and **PID**, TFAE:

- ϕ is flat;
- ϕ is F -projective;
- ϕ is F -exact;
- $\phi \dashv\vdash \Theta_X$ for some X

Definition

Let S be a set of **well-behaved** substitutions of a logic L .

- A rule ϕ/ψ of L is said to be a ***S-admissible rule***, in symbols $\phi \vdash_L^S \psi$ or simply $\phi \vdash^S \psi$, if for all $\sigma \in S$,

$$\vdash_L \sigma(\phi) \implies \vdash_L \sigma(\psi).$$

- L is said to be ***S-structurally complete*** if every S -admissible rule of L is derivable, i.e., $\phi \vdash_L^S \psi \implies \phi \vdash_L \psi$.

Definition

Let S be a set of **well-behaved** substitutions of a logic L .

- A rule ϕ/ψ of L is said to be a ***S-admissible rule***, in symbols $\phi \vdash_L^S \psi$ or simply $\phi \vdash^S \psi$, if for all $\sigma \in S$,

$$\vdash_L \sigma(\phi) \implies \vdash_L \sigma(\psi).$$

- L is said to be ***S-structurally complete*** if every S -admissible rule of L is derivable, i.e., $\phi \vdash_L^S \psi \implies \phi \vdash_L \psi$.

Definition

Let S be a set of **well-behaved** substitutions of a logic L .

- A rule ϕ/ψ of L is said to be a ***S-admissible rule***, in symbols $\phi \vdash_L^S \psi$ or simply $\phi \vdash^S \psi$, if for all $\sigma \in S$,

$$\vdash_L \sigma(\phi) \implies \vdash_L \sigma(\psi).$$

- L is said to be ***S-structurally complete*** if every S -admissible rule of L is derivable, i.e., $\phi \vdash_L^S \psi \implies \phi \vdash_L \psi$.

Clearly, if L is S -structurally complete, then $\phi \vdash_L^S \psi \iff \phi \vdash_L \psi$.

Definition

Let S be a set of **well-behaved** substitutions of a logic L .

- A rule ϕ/ψ of L is said to be a **S -admissible rule**, in symbols $\phi \vdash_L^S \psi$ or simply $\phi \vdash^S \psi$, if for all $\sigma \in S$,

$$\vdash_L \sigma(\phi) \implies \vdash_L \sigma(\psi).$$

- L is said to be **S -structurally complete** if every S -admissible rule of L is derivable, i.e., $\phi \vdash_L^S \psi \implies \phi \vdash_L \psi$.

Clearly, if L is S -structurally complete, then $\phi \vdash_L^S \psi \iff \phi \vdash_L \psi$.

Theorem

PID and **PD** are F -structurally complete, where F is the set of all flat substitutions.

Proof. Since $\phi \equiv \bigvee_{i \in I} \Theta_{X_i}$, where each Θ_{X_i} is S -projective, for every formula ϕ of the logics. □