

Admissible Rules of (Fragments of) R-Mingle

Laura Janina Schnüriger

joint work with George Metcalfe
Universität Bern

Les Diablerets
31 January 2015

Table of contents

1. The logic R-Mingle RM^t
 - 1.1 Notations
 - 1.2 Corresponding algebraic semantics
 - 1.3 Sugihara Monoids
 - 1.4 This talk
2. Finding the bases
 - 2.1 Idea of how to find the bases
 - 2.2 Finding the bases
 - 2.3 The bases
3. Further Work
 - 3.1 Our Conjecture
4. References

R-Mingle

RM

Relevance logic R with Mingle

Mingle

$$p \rightarrow (p \rightarrow p)$$

R-Mingle

RM Relevance logic R with Mingle

Mingle $p \rightarrow (p \rightarrow p)$

RM^t RM with additional constant t

Language $\mathcal{L}_t = \{\wedge, \vee, \rightarrow, \cdot, \neg, t\}$

Definition

- rules are denoted by Γ/φ for finite $\Gamma \cup \{\varphi\} \subset \text{Fm}_{\mathcal{L}}$

Definition

- rules are denoted by Γ/φ for finite $\Gamma \cup \{\varphi\} \subset \text{Fm}_{\mathcal{L}}$
- Γ/φ is derivable in a logic L if $\Gamma \vdash_L \varphi$

Definition

- rules are denoted by Γ/φ for finite $\Gamma \cup \{\varphi\} \subset \text{Fm}_{\mathcal{L}}$
- Γ/φ is derivable in a logic L if $\Gamma \vdash_L \varphi$
- Γ/φ is admissible in a logic L if for all substitutions (homomorphisms) $\sigma: \text{Fm}_{\mathcal{L}} \rightarrow \text{Fm}_{\mathcal{L}}$:

$$\vdash_L \sigma(\psi) \text{ for all } \psi \in \Gamma \quad \Rightarrow \quad \vdash_L \sigma(\varphi)$$

Definition

- rules are denoted by Γ/φ for finite $\Gamma \cup \{\varphi\} \subset \text{Fm}_{\mathcal{L}}$
- Γ/φ is derivable in a logic L if $\Gamma \vdash_L \varphi$
- Γ/φ is admissible in a logic L if for all substitutions (homomorphisms) $\sigma: \text{Fm}_{\mathcal{L}} \rightarrow \text{Fm}_{\mathcal{L}}$:

$$\vdash_L \sigma(\psi) \text{ for all } \psi \in \Gamma \Rightarrow \vdash_L \sigma(\varphi)$$

- $\{\Gamma/\varphi \mid \Gamma/\varphi \text{ is admissible in } L\} =: \sim_L$

Definition

- rules are denoted by Γ/φ for finite $\Gamma \cup \{\varphi\} \subset \text{Fm}_{\mathcal{L}}$
- Γ/φ is derivable in a logic L if $\Gamma \vdash_L \varphi$
- Γ/φ is admissible in a logic L if for all substitutions (homomorphisms) $\sigma: \text{Fm}_{\mathcal{L}} \rightarrow \text{Fm}_{\mathcal{L}}$:

$$\vdash_L \sigma(\psi) \text{ for all } \psi \in \Gamma \Rightarrow \vdash_L \sigma(\varphi)$$

- $\{\Gamma/\varphi \mid \Gamma/\varphi \text{ is admissible in } L\} =: \sim_L$
- Let \mathcal{R} be a set of rules.

$$L + \mathcal{R} = \text{smallest logic containing } L \cup \mathcal{R}$$

Definition

- rules are denoted by Γ/φ for finite $\Gamma \cup \{\varphi\} \subset \text{Fm}_{\mathcal{L}}$
- Γ/φ is derivable in a logic L if $\Gamma \vdash_L \varphi$
- Γ/φ is admissible in a logic L if for all substitutions (homomorphisms) $\sigma: \text{Fm}_{\mathcal{L}} \rightarrow \text{Fm}_{\mathcal{L}}$:

$$\vdash_L \sigma(\psi) \text{ for all } \psi \in \Gamma \Rightarrow \vdash_L \sigma(\varphi)$$

- $\{\Gamma/\varphi \mid \Gamma/\varphi \text{ is admissible in } L\} =: \vdash_L$
- Let \mathcal{R} be a set of rules.

$$L + \mathcal{R} = \text{smallest logic containing } L \cup \mathcal{R}$$

- \mathcal{R} is a basis for the admissible rules of L if $L + \mathcal{R} = \vdash_L$

Corresponding algebraic semantics

$$\mathbf{Z}^\circ = \langle \mathbb{Z} \setminus \{0\}, \min, \max, \rightarrow, \cdot, -, 1 \rangle$$

$$\rightarrow \quad x \rightarrow y := \begin{cases} \max\{-x, y\} & \text{if } x \leq y \\ \min\{-x, y\} & \text{if } x > y \end{cases}$$

$$\cdot \quad x \cdot y := \begin{cases} \min\{x, y\} & \text{if } |x| = |y| \\ y & \text{if } |x| < |y| \\ x & \text{if } |x| > |y| \end{cases}$$

Corresponding algebraic semantics

$$\mathbf{Z}^\circ = \langle \mathbb{Z} \setminus \{0\}, \min, \max, \rightarrow, \cdot, -, 1 \rangle$$

$$\rightarrow \quad x \rightarrow y := \begin{cases} \max\{-x, y\} & \text{if } x \leq y \\ \min\{-x, y\} & \text{if } x > y \end{cases}$$

$$\cdot \quad x \cdot y := \begin{cases} \min\{x, y\} & \text{if } |x| = |y| \\ y & \text{if } |x| < |y| \\ x & \text{if } |x| > |y| \end{cases}$$

$$\mathbf{Z}_{2n} = \langle \{-n, \dots, -1, 1, \dots, n\}, \min, \max, \rightarrow, \cdot, -, 1 \rangle$$

$$\mathbf{Z}_{2n+1} = \langle \{-n, \dots, -1, 0, 1, \dots, n\}, \min, \max, \rightarrow, \cdot, -, 1 \rangle$$

Sugihara Monoids

$\mathcal{SM} = \mathbb{V}(\mathbf{Z}^\circ)$ the variety of Sugihara Monoids generated by \mathbf{Z}° .
 \mathcal{SM} provides an equivalent algebraic semantics for RM^t

$$\begin{aligned} \{\psi \approx |\psi| \mid \psi \in \Gamma\} \models_{\mathcal{SM}} \varphi \approx |\varphi| &\Leftrightarrow: \Gamma \models_{\mathcal{SM}} \varphi \\ &\Leftrightarrow \Gamma \vdash_{\text{RM}^t} \varphi \end{aligned}$$

for any rule Γ/φ .

This talk

Bases for admissible rules of the fragments of RM^t with the following languages

$$\mathcal{L}_1 = \{\rightarrow, t\}$$

$$\mathcal{L}_2 = \{\rightarrow, \cdot, t\}$$

$$\mathcal{L}_m = \{\rightarrow, \neg, t\} = \{\rightarrow, \cdot, \neg, t\} \text{ multiplicative fragment.}$$

This talk

Bases for admissible rules of the fragments of RM^t with the following languages

$$\mathcal{L}_1 = \{\rightarrow, t\}$$

$$\mathcal{L}_2 = \{\rightarrow, \cdot, t\}$$

$$\mathcal{L}_m = \{\rightarrow, \neg, t\} = \{\rightarrow, \cdot, \neg, t\} \text{ multiplicative fragment.}$$

$\mathcal{SM} \upharpoonright \mathcal{L}_i$ algebraic semantics corresponding to the \mathcal{L}_i -fragment of RM^t , $i \in \{1, 2, m\}$

This talk

Bases for admissible rules of the fragments of RM^t with the following languages

$$\mathcal{L}_1 = \{\rightarrow, t\}$$

$$\mathcal{L}_2 = \{\rightarrow, \cdot, t\}$$

$$\mathcal{L}_m = \{\rightarrow, \neg, t\} = \{\rightarrow, \cdot, \neg, t\} \text{ multiplicative fragment.}$$

$\mathcal{SM} \upharpoonright \mathcal{L}_i$ algebraic semantics corresponding to the \mathcal{L}_i -fragment of RM^t , $i \in \{1, 2, m\}$

Remark
Raftery, Olson $\text{RM}^t \upharpoonright \{\wedge, \rightarrow, t\}$ has empty basis (= it is structurally complete).

Idea of how to find the bases

Recall $\mathcal{SM} = \mathbb{V}(\mathcal{SM}) = \mathbb{V}(\mathbf{Z}^\circ)$

Idea of how to find the bases

Recall $\mathcal{SM} = \mathbb{V}(\mathcal{SM}) = \mathbb{V}(\mathbf{Z}^\circ)$

Lemma *S*. $\mathbb{V}(\mathcal{SM} \upharpoonright \mathcal{L}_i) = \mathbb{V}(\mathbf{Z}_4 \upharpoonright \mathcal{L}_i), \quad i \in \{1, 2, m\}$

Idea of how to find the bases

- Recall that if for two varieties \mathbb{V}_1 and \mathbb{V}_2 we have:
 $\mathbb{V}_1 = \mathbb{V}_2$ iff $(\vdash_{\mathbb{V}_1} \varphi \Leftrightarrow \vdash_{\mathbb{V}_2} \varphi \text{ for all formulas } \varphi)$.

Idea of how to find the bases

- Recall that if for two varieties \mathbb{V}_1 and \mathbb{V}_2 we have:
 $\mathbb{V}_1 = \mathbb{V}_2$ iff $(\vdash_{\mathbb{V}_1} \varphi \Leftrightarrow \vdash_{\mathbb{V}_2} \varphi \text{ for all formulas } \varphi)$.
- A rule is admissible in $\text{RM}^t \upharpoonright \mathcal{L}_i$
 \Leftrightarrow it is admissible in $\mathcal{SM} \upharpoonright \mathcal{L}_i$
 \Leftrightarrow it is admissible in $\mathbf{Z}_4 \upharpoonright \mathcal{L}_i$

Idea of how to find the bases

- Recall that if for two varieties \mathbb{V}_1 and \mathbb{V}_2 we have:
 $\mathbb{V}_1 = \mathbb{V}_2$ iff $(\vdash_{\mathbb{V}_1} \varphi \Leftrightarrow \vdash_{\mathbb{V}_2} \varphi \text{ for all formulas } \varphi)$.
- A rule is admissible in $\text{RM}^t \upharpoonright \mathcal{L}_i$
 \Leftrightarrow it is admissible in $\mathcal{SM} \upharpoonright \mathcal{L}_i$
 \Leftrightarrow it is admissible in $\mathbf{Z}_4 \upharpoonright \mathcal{L}_i$
- Interested in algebras s.t. admissibility in $\mathbf{Z}_4 \upharpoonright \mathcal{L}_i$ corresponds to validity in these algebras.

Idea of how to find the bases

- Recall that if for two varieties \mathbb{V}_1 and \mathbb{V}_2 we have:
 $\mathbb{V}_1 = \mathbb{V}_2$ iff $(\vdash_{\mathbb{V}_1} \varphi \Leftrightarrow \vdash_{\mathbb{V}_2} \varphi \text{ for all formulas } \varphi)$.
- A rule is admissible in $\text{RM}^t \upharpoonright \mathcal{L}_i$
 \Leftrightarrow it is admissible in $\mathcal{SM} \upharpoonright \mathcal{L}_i$
 \Leftrightarrow it is admissible in $\mathbf{Z}_4 \upharpoonright \mathcal{L}_i$
- Interested in algebras s.t. admissibility in $\mathbf{Z}_4 \upharpoonright \mathcal{L}_i$ corresponds to validity in these algebras.
- Then: Axiomatize the quasivarieties generated by these algebras to get an axiomatization of the admissible rules of our fragments.

Finding the bases

Theorem Let \mathbf{B} be an algebra and $\mathbf{F}_{\mathbf{B}}(\omega)$ its free algebra on countably infinite many generators. Then

$$\Gamma/\varphi \text{ is } \mathbf{B}\text{-admissible} \quad \Leftrightarrow \quad \Gamma \models_{\mathbf{F}_{\mathbf{B}}(\omega)} \varphi.$$

Finding the bases

Lemma The following are equivalent:

- (i) Γ/φ is **B**-admissible $\Leftrightarrow \Gamma \vDash_{\mathbf{A}} \varphi$
- (ii) $\mathbb{Q}(\mathbf{A}) = \mathbb{Q}(\mathbf{F}_{\mathbf{B}}(\omega))$

Finding the bases

Lemma The following are equivalent:

- (i) Γ/φ is **B**-admissible $\Leftrightarrow \Gamma \vDash_{\mathbf{A}} \varphi$
- (ii) $\mathbb{Q}(\mathbf{A}) = \mathbb{Q}(\mathbf{F}_{\mathbf{B}}(\omega))$

So we want to find **A** which is “easy” to axiomatize - but how?

Finding the bases

Lemma The following are equivalent:

- (i) Γ/φ is **B**-admissible $\Leftrightarrow \Gamma \vDash_{\mathbf{A}} \varphi$
- (ii) $\mathbb{Q}(\mathbf{A}) = \mathbb{Q}(\mathbf{F}_{\mathbf{B}}(\omega))$

So we want to find **A** which is “easy” to axiomatize - but how?

Lemma $\mathbf{A} \subseteq \mathbf{F}_{\mathbf{B}}(\omega), \mathbf{B} \in \mathbb{H}(\mathbf{A}) \Rightarrow \mathbb{Q}(\mathbf{A}) = \mathbb{Q}(\mathbf{F}_{\mathbf{B}}(\omega))$

Lemma S.

The algebras in our case

Let $\mathbf{Z}'_4 \subset (\mathbf{Z}_2 \times \mathbf{Z}_3) \upharpoonright \mathcal{L}_1$,

$\mathbf{Z}''_4 \subset (\mathbf{Z}_2 \times \mathbf{Z}_3) \upharpoonright \mathcal{L}_2$,

$(\mathbf{Z}_2 \times \mathbf{Z}_3) \upharpoonright \mathcal{L}_m$ be the algebras pictured. Then

- (i) $\mathbb{Q}(\mathbf{F}_{\mathbf{Z}_4 \upharpoonright \mathcal{L}_1}(\omega)) = \mathbb{Q}(\mathbf{Z}'_4)$
- (ii) $\mathbb{Q}(\mathbf{F}_{\mathbf{Z}_4 \upharpoonright \mathcal{L}_2}(\omega)) = \mathbb{Q}(\mathbf{Z}''_4)$
- (iii) $\mathbb{Q}(\mathbf{F}_{\mathbf{Z}_4 \upharpoonright \mathcal{L}_m}(\omega)) = \mathbb{Q}((\mathbf{Z}_2 \times \mathbf{Z}_3) \upharpoonright \mathcal{L}_m)$




Figure: \mathbf{Z}'_4 and \mathbf{Z}''_4

Definition

- $|\psi| := \psi \rightarrow \psi$

Definition

- $|\psi| := \psi \rightarrow \psi$
- $\varphi \Rightarrow \psi := (\varphi \rightarrow |\psi|) \rightarrow (\varphi \rightarrow \psi)$

Definition

- $|\psi| := \psi \rightarrow \psi$
- $\varphi \Rightarrow \psi := (\varphi \rightarrow |\psi|) \rightarrow (\varphi \rightarrow \psi)$
- $\{p, p \Rightarrow q\} / q \quad (A)$

Definition

- $|\psi| := \psi \rightarrow \psi$
- $\varphi \Rightarrow \psi := (\varphi \rightarrow |\psi|) \rightarrow (\varphi \rightarrow \psi)$
- $\{p, p \Rightarrow q\} / q$ (A)
- $\varphi \leftrightarrow \psi := (\varphi \rightarrow \psi) \cdot (\psi \rightarrow \varphi)$

Definition

- $|\psi| := \psi \rightarrow \psi$
- $\varphi \Rightarrow \psi := (\varphi \rightarrow |\psi|) \rightarrow (\varphi \rightarrow \psi)$
- $\{p, p \Rightarrow q\} / q \quad (A)$
- $\varphi \leftrightarrow \psi := (\varphi \rightarrow \psi) \cdot (\psi \rightarrow \varphi)$
- $\{\neg(|p_1| \leftrightarrow \dots \leftrightarrow |p_n|)\} / q \quad (R_n), \quad n \in \mathbb{N}.$

The Bases

Lemma 5. We have the following axiomatizations:

- (i) $\text{RM}^t \upharpoonright \mathcal{L}_1 + (A)$ has equivalent q.v. $\mathbb{Q}(\mathbf{Z}'_4)$
- (ii) $\text{RM}^t \upharpoonright \mathcal{L}_2 + (A)$ has equivalent q.v. $\mathbb{Q}(\mathbf{Z}''_4)$
- (iii) $\text{RM}^t \upharpoonright \mathcal{L}_m + (A) + \{(R_n)\}_{n \in \mathbb{N}}$ has eq. q.v.
 $\mathbb{Q}((\mathbf{Z}_2 \times \mathbf{Z}_3) \upharpoonright \mathcal{L}_m)$

The Bases

Lemma S. We have the following axiomatizations:

- (i) $\text{RM}^t \upharpoonright \mathcal{L}_1 + (A)$ has equivalent q.v. $\mathbb{Q}(\mathbf{Z}'_4)$
- (ii) $\text{RM}^t \upharpoonright \mathcal{L}_2 + (A)$ has equivalent q.v. $\mathbb{Q}(\mathbf{Z}''_4)$
- (iii) $\text{RM}^t \upharpoonright \mathcal{L}_m + (A) + \{(R_n)\}_{n \in \mathbb{N}}$ has eq. q.v.
 $\mathbb{Q}((\mathbf{Z}_2 \times \mathbf{Z}_3) \upharpoonright \mathcal{L}_m)$

Theorem S. Then as a Corollary of this lemma

- (i) $\{(A)\}$ is a basis for the $\{\rightarrow, t\}$ - and
 $\{\rightarrow, \cdot, t\}$ -fragment of RM^t .
- (ii) $\{(A)\} \cup \{(R_n)\}_{n \in \mathbb{N}}$ is a basis for $\text{RM}^t \upharpoonright \{\rightarrow, \neg, t\}$.

Our Conjecture

Look again at RM without constant t .

Our Conjecture

Look again at RM without constant t .

$$(B) \quad \{\neg|p| \vee q\}/q$$

Our Conjecture

Look again at RM without constant t .

$$(B) \quad \{\neg|p| \vee q\}/q$$

Conjecture

We hope to prove the following:

(i) $\text{RM} + (B)$ is almost structurally complete, i.e.,

$$\Gamma \sim_{\text{RM}} \varphi \Rightarrow \Gamma \vdash_{\text{RM}} \varphi$$

whenever there is a substitution

$$\sigma: \text{Fm}_{\mathcal{L}} \rightarrow \text{Fm}_{\mathcal{L}} \text{ s.t. for all } \psi \in \Gamma, \\ \vdash_{\text{RM}} \sigma(\psi).$$

(ii) The admissible rules of RM have basis

$$\{(B)\} \cup \{(R_n)\}_{n \in \mathbb{N}}.$$

References

- G. Metcalfe and C. Röthlisberger. *Admissibility in finitely generated quasivarieties*. Logical Methods in Computer Science, vol. 9 (2013), no. 2, pp. 119.
- C. Röthlisberger. *Admissibility in finitely generated quasivarieties*. PhD thesis, 2013.
- G. Metcalfe. *An Avron rule for fragments of R-mingle*. Journal of Logic and Computation, to appear.
- J. Olson and J. Raftery. *Positive Sugihara monoids*. Algebra Universalis, vol. 57 (2005), pp. 7599.