

# Uniform Interpolation and the Congruence Lattice

Sam van Gool

Joint work with George Metcalfe and Constantine Tsinakis

Workshop on Admissible Rules and Unification 2

31 January – 2 February 2015, Les Diablerets

# Uniform interpolation in IPC

## Theorem (Pitts, 1992)

For any formula  $\phi(\bar{x}, \bar{y})$  of intuitionistic propositional logic IPC, there exist **left** and **right uniform interpolants**,

$$\phi^L(\bar{y}) \quad \text{and} \quad \phi^R(\bar{y}),$$

# Uniform interpolation in IPC

## Theorem (Pitts, 1992)

For any formula  $\phi(\bar{x}, \bar{y})$  of intuitionistic propositional logic IPC, there exist **left** and **right uniform interpolants**,

$$\phi^L(\bar{y}) \quad \text{and} \quad \phi^R(\bar{y}),$$

such that for any formula  $\psi(\bar{y}, \bar{z})$ ,

$$\phi(\bar{x}, \bar{y}) \vdash_{\text{IPC}} \psi(\bar{y}, \bar{z}) \quad \Leftrightarrow \quad \phi^R(\bar{y}) \vdash_{\text{IPC}} \psi(\bar{y}, \bar{z})$$

# Uniform interpolation in IPC

## Theorem (Pitts, 1992)

For any formula  $\phi(\bar{x}, \bar{y})$  of intuitionistic propositional logic IPC, there exist **left** and **right uniform interpolants**,

$$\phi^L(\bar{y}) \quad \text{and} \quad \phi^R(\bar{y}),$$

such that for any formula  $\psi(\bar{y}, \bar{z})$ ,

$$\phi(\bar{x}, \bar{y}) \vdash_{\text{IPC}} \psi(\bar{y}, \bar{z}) \quad \Leftrightarrow \quad \phi^R(\bar{y}) \vdash_{\text{IPC}} \psi(\bar{y}, \bar{z})$$

$$\psi(\bar{y}, \bar{z}) \vdash_{\text{IPC}} \phi(\bar{x}, \bar{y}) \quad \Leftrightarrow \quad \psi(\bar{y}, \bar{z}) \vdash_{\text{IPC}} \phi^L(\bar{y}).$$

## Uniform interpolation and restriction

A formula  $\phi^R(\bar{y})$  is a **right uniform interpolant** for  $\phi(\bar{x}, \bar{y})$



# Uniform interpolation and restriction

A formula  $\phi^R(\overline{y})$  is a **right uniform interpolant** for  $\phi(\overline{x}, \overline{y})$



for any formula  $\psi(\overline{y}, \overline{z})$ , if  $\phi(\overline{x}, \overline{y}) \vdash_{\text{IPC}} \psi(\overline{y}, \overline{z})$ ,

then  $\phi(\overline{x}, \overline{y}) \vdash_{\text{IPC}} \phi^R(\overline{y}) \vdash_{\text{IPC}} \psi(\overline{y}, \overline{z})$ .

# Uniform interpolation and restriction

A formula  $\phi^R(\overline{y})$  is a **right uniform interpolant** for  $\phi(\overline{x}, \overline{y})$



for any formula  $\psi(\overline{y}, \overline{z})$ , if  $\phi(\overline{x}, \overline{y}) \vdash_{\text{IPC}} \psi(\overline{y}, \overline{z})$ ,

then  $\phi(\overline{x}, \overline{y}) \vdash_{\text{IPC}} \phi^R(\overline{y}) \vdash_{\text{IPC}} \psi(\overline{y}, \overline{z})$ .



for any formula  $\chi(\overline{y})$ , if  $\phi(\overline{x}, \overline{y}) \vdash_{\text{IPC}} \chi(\overline{y})$ ,

## Uniform interpolation and restriction

A formula  $\phi^R(\overline{y})$  is a **right uniform interpolant** for  $\phi(\overline{x}, \overline{y})$



for any formula  $\psi(\overline{y}, \overline{z})$ , if  $\phi(\overline{x}, \overline{y}) \vdash_{\text{IPC}} \psi(\overline{y}, \overline{z})$ ,

then  $\phi(\overline{x}, \overline{y}) \vdash_{\text{IPC}} \phi^R(\overline{y}) \vdash_{\text{IPC}} \psi(\overline{y}, \overline{z})$ .



for any formula  $\chi(\overline{y})$ , if  $\phi(\overline{x}, \overline{y}) \vdash_{\text{IPC}} \chi(\overline{y})$ ,

then  $\phi(\overline{x}, \overline{y}) \vdash_{\text{IPC}} \phi^R(\overline{y}) \vdash_{\text{IPC}} \chi(\overline{y})$ .



# Uniform interpolation and restriction

A formula  $\phi^R(\overline{y})$  is a **right uniform interpolant** for  $\phi(\overline{x}, \overline{y})$



for any formula  $\psi(\overline{y}, \overline{z})$ , if  $\phi(\overline{x}, \overline{y}) \vdash_{\text{IPC}} \psi(\overline{y}, \overline{z})$ ,

then  $\phi(\overline{x}, \overline{y}) \vdash_{\text{IPC}} \phi^R(\overline{y}) \vdash_{\text{IPC}} \psi(\overline{y}, \overline{z})$ .




for any formula  $\chi(\overline{y})$ , if  $\phi(\overline{x}, \overline{y}) \vdash_{\text{IPC}} \chi(\overline{y})$ ,

then  $\phi(\overline{x}, \overline{y}) \vdash_{\text{IPC}} \phi^R(\overline{y}) \vdash_{\text{IPC}} \chi(\overline{y})$ .


(\*) because IPC has interpolation

## Right uniform restriction, pictorially


**Fm**( $\bar{x}$ ,  $\bar{y}$ )




## Right uniform restriction, pictorially




## Right uniform restriction, pictorially




## Right uniform restriction, pictorially



## Right uniform restriction, pictorially



## Right uniform restriction, pictorially



## This Talk

Which **varieties of algebras** admit uniform interpolation?



# Equational Consequence

The **equational consequence relation** for a variety  $\mathcal{V}$  is defined by

$$\Sigma \models_{\mathcal{V}} \alpha \approx \beta \quad \Leftrightarrow \quad \begin{array}{l} \text{for all } \mathbf{A} \in \mathcal{V} \text{ and } e : \mathbf{F}_{\mathcal{V}}(\omega) \rightarrow \mathbf{A}, \\ \text{if } \Sigma \subseteq \ker(e) \text{ then } \alpha \approx \beta \in \ker(e). \end{array}$$

We will write  $\Sigma \models_{\mathcal{V}} \Delta$  to denote that  $\Sigma \models_{\mathcal{V}} \varepsilon$  **for all**  $\varepsilon \in \Delta$ .

# Right uniform restriction

## Definition

A variety  $\mathcal{V}$  has **right uniform restriction** iff

# Right uniform restriction

## Definition

A variety  $\mathcal{V}$  has **right uniform restriction** iff for any finite set of equations  $\Sigma(\overline{x}, \overline{y})$ ,

## Right uniform restriction

### Definition

A variety  $\mathcal{V}$  has **right uniform restriction** iff for any finite set of equations  $\Sigma(\overline{x}, \overline{y})$ , there exists a finite set of equations  $\Sigma^R(\overline{y})$

## Right uniform restriction

### Definition

A variety  $\mathcal{V}$  has **right uniform restriction** iff for any finite set of equations  $\Sigma(\overline{x}, \overline{y})$ , there exists a finite set of equations  $\Sigma^R(\overline{y})$  such that for any equation  $\epsilon(\overline{y})$ :

## Right uniform restriction

### Definition

A variety  $\mathcal{V}$  has **right uniform restriction** iff for any finite set of equations  $\Sigma(\bar{x}, \bar{y})$ , there exists a finite set of equations  $\Sigma^R(\bar{y})$  such that for any equation  $\epsilon(\bar{y})$ :

$$\Sigma(\bar{x}, \bar{y}) \models_{\mathcal{V}} \epsilon(\bar{y}) \iff \Sigma^R(\bar{y}) \models_{\mathcal{V}} \epsilon(\bar{y}).$$

## Equations and congruences

A set of equations  $\Sigma(\bar{x})$  generates a **congruence**,  $\Theta(\Sigma)$ , on  $\mathbf{F}_{\mathcal{V}}(\bar{x})$ , the free  $\mathcal{V}$ -algebra generated by  $\bar{x}$ .

## Equations and congruences

A set of equations  $\Sigma(\bar{x})$  generates a **congruence**,  $\Theta(\Sigma)$ , on  $\mathbf{F}_{\mathcal{V}}(\bar{x})$ , the free  $\mathcal{V}$ -algebra generated by  $\bar{x}$ .

A congruence  $\theta$  on  $\mathbf{F}_{\mathcal{V}}(\bar{x})$  is **finitely generated** if  $\theta = \Theta(\Sigma)$  for some finite set  $\Sigma$ .



## The congruence lattice

For any algebra **A**, the **lattice of congruences** on **A**,  $\text{Con}(\mathbf{A})$ , is a complete lattice.

## The congruence lattice

For any algebra  $\mathbf{A}$ , the **lattice of congruences** on  $\mathbf{A}$ ,  $\text{Con}(\mathbf{A})$ , is a complete lattice.

An element  $u$  in a complete lattice  $L$  is called **compact** if,

## The congruence lattice

For any algebra **A**, the **lattice of congruences** on **A**,  $\text{Con}(\mathbf{A})$ , is a complete lattice.

An element  $u$  in a complete lattice  $L$  is called **compact** if, for any subset  $S \subseteq L$  such that  $u \leq \bigvee S$ ,

## The congruence lattice

For any algebra **A**, the **lattice of congruences** on **A**,  $\text{Con}(\mathbf{A})$ , is a complete lattice.

An element  $u$  in a complete lattice  $L$  is called **compact** if, for any subset  $S \subseteq L$  such that  $u \leq \bigvee S$ , there exists a finite subset  $F \subseteq S$  such that  $u \leq \bigvee F$ .

## The congruence lattice

For any algebra **A**, the **lattice of congruences** on **A**,  $\text{Con}(\mathbf{A})$ , is a complete lattice.

An element  $u$  in a complete lattice  $L$  is called **compact** if, for any subset  $S \subseteq L$  such that  $u \leq \bigvee S$ , there exists a finite subset  $F \subseteq S$  such that  $u \leq \bigvee F$ .

### Fact

*In a lattice of congruences, the compact elements are exactly the finitely generated congruences.*

## The congruence lattice

For any algebra **A**, the **lattice of congruences** on **A**,  $\text{Con}(\mathbf{A})$ , is a complete lattice.

An element  $u$  in a complete lattice  $L$  is called **compact** if, for any subset  $S \subseteq L$  such that  $u \leq \bigvee S$ , there exists a finite subset  $F \subseteq S$  such that  $u \leq \bigvee F$ .

### Fact

*In a lattice of congruences, the compact elements are exactly the finitely generated congruences.*

We denote by  $\text{KCon}(\mathbf{A})$  the join-semilattice of **compact congruences**.

## Lifting homomorphisms

Any homomorphism  $f : \mathbf{A} \rightarrow \mathbf{B}$  can be **lifted** to the congruence lattices:

## Lifting homomorphisms

Any homomorphism  $f : \mathbf{A} \rightarrow \mathbf{B}$  can be **lifted** to the congruence lattices:

$$f_* : \text{Con}(\mathbf{A}) \rightleftharpoons \text{Con}(\mathbf{B}) : f^{-1}$$



## Lifting homomorphisms

Any homomorphism  $f : \mathbf{A} \rightarrow \mathbf{B}$  can be **lifted** to the congruence lattices:

$$f_* : \text{Con}(\mathbf{A}) \rightleftharpoons \text{Con}(\mathbf{B}) : f^{-1}$$

$$\theta_A \mapsto f_*(\theta_A) \stackrel{\text{def}}{=} \Theta(\{(f(a), f(a')) : (a, a') \in \theta\}),$$

## Lifting homomorphisms

Any homomorphism  $f : \mathbf{A} \rightarrow \mathbf{B}$  can be **lifted** to the congruence lattices:

$$f_* : \text{Con}(\mathbf{A}) \rightleftharpoons \text{Con}(\mathbf{B}) : f^{-1}$$

$$\theta_A \mapsto f_*(\theta_A) \stackrel{\text{def}}{=} \Theta(\{(f(a), f(a')) : (a, a') \in \theta\}),$$

$$f^{-1}(\theta_B) \leftarrow \theta_B.$$

## Lifting homomorphisms

Any homomorphism  $f : \mathbf{A} \rightarrow \mathbf{B}$  can be **lifted** to the congruence lattices:

$$f_* : \text{Con}(\mathbf{A}) \rightleftharpoons \text{Con}(\mathbf{B}) : f^{-1}$$

$$\theta_A \mapsto f_*(\theta_A) \stackrel{\text{def}}{=} \Theta(\{(f(a), f(a')) : (a, a') \in \theta\}),$$

$$f^{-1}(\theta_B) \leftarrow \theta_B.$$

The pair  $(f_*, f^{-1})$  is an **adjunction**

## Lifting homomorphisms

Any homomorphism  $f : \mathbf{A} \rightarrow \mathbf{B}$  can be **lifted** to the congruence lattices:

$$f_* : \text{Con}(\mathbf{A}) \rightleftharpoons \text{Con}(\mathbf{B}) : f^{-1}$$

$$\theta_A \mapsto f_*(\theta_A) \stackrel{\text{def}}{=} \Theta(\{(f(a), f(a')) : (a, a') \in \theta\}),$$

$$f^{-1}(\theta_B) \leftarrow \theta_B.$$

The pair  $(f_*, f^{-1})$  is an **adjunction**, i.e., we have

$$f_*(\theta_A) \subseteq \theta_B \iff \theta_A \subseteq f^{-1}(\theta_B),$$

for any  $\theta_A \in \text{Con}(\mathbf{A})$  and  $\theta_B \in \text{Con}(\mathbf{B})$ .

## Lifting homomorphisms

Any homomorphism  $f : \mathbf{A} \rightarrow \mathbf{B}$  can be **lifted** to the congruence lattices:

$$\begin{aligned} f_* : \text{Con}(\mathbf{A}) &\rightleftarrows \text{Con}(\mathbf{B}) : f^{-1} \\ \theta_A &\mapsto f_*(\theta_A) \stackrel{\text{def}}{=} \Theta(\{(f(a), f(a')) : (a, a') \in \theta\}), \\ f^{-1}(\theta_B) &\leftarrow \theta_B. \end{aligned}$$

The pair  $(f_*, f^{-1})$  is an **adjunction**, i.e., we have

$$f_*(\theta_A) \subseteq \theta_B \iff \theta_A \subseteq f^{-1}(\theta_B),$$

for any  $\theta_A \in \text{Con}(\mathbf{A})$  and  $\theta_B \in \text{Con}(\mathbf{B})$ .


This adjunction **restricts to compact congruences** iff  $f^{-1}$  preserves compact elements.

## Lifting homomorphisms, pictorially

For any  $\overline{x}$  and  $\overline{y}$  finite, let  $i : \mathbf{F}_{\mathcal{V}}(\overline{y}) \hookrightarrow \mathbf{F}_{\mathcal{V}}(\overline{x}, \overline{y})$  denote the natural inclusion map.


## Lifting homomorphisms, pictorially

For any  $\overline{x}$  and  $\overline{y}$  finite, let  $i : \mathbf{F}_V(\overline{y}) \hookrightarrow \mathbf{F}_V(\overline{x}, \overline{y})$  denote the natural inclusion map.



## Lifting homomorphisms, pictorially


For any  $\bar{x}$  and  $\bar{y}$  finite, let  $i : \mathbf{F}_V(\bar{y}) \hookrightarrow \mathbf{F}_V(\bar{x}, \bar{y})$  denote the natural inclusion map.






## Lifting homomorphisms, pictorially

For any  $\bar{x}$  and  $\bar{y}$  finite, let  $i : \mathbf{F}_V(\bar{y}) \hookrightarrow \mathbf{F}_V(\bar{x}, \bar{y})$  denote the natural inclusion map.




## Lifting homomorphisms, pictorially

For any  $\overline{x}$  and  $\overline{y}$  finite, let  $i : \mathbf{F}_V(\overline{y}) \hookrightarrow \mathbf{F}_V(\overline{x}, \overline{y})$  denote the natural inclusion map.



## Lifting homomorphisms, pictorially

For any  $\overline{x}$  and  $\overline{y}$  finite, let  $i : \mathbf{F}_V(\overline{y}) \hookrightarrow \mathbf{F}_V(\overline{x}, \overline{y})$  denote the natural inclusion map.



## Right uniform restriction and existence of adjoints

### Theorem

*For any variety  $\mathcal{V}$ , the following are equivalent:*

- (1)  $\mathcal{V}$  has **right uniform restriction**.

## Right uniform restriction and existence of adjoints

### Theorem

*For any variety  $\mathcal{V}$ , the following are equivalent:*

- (1)  $\mathcal{V}$  has **right uniform restriction**.
- (2) *For finite  $\bar{x}$ ,  $\bar{y}$ , the map  $i_* : \text{KCon}(\mathbf{F}_{\mathcal{V}}(\bar{y})) \rightarrow \text{KCon}(\mathbf{F}_{\mathcal{V}}(\bar{x}, \bar{y}))$  has a right adjoint;*

## Right uniform restriction and existence of adjoints

### Theorem

*For any variety  $\mathcal{V}$ , the following are equivalent:*

- (1)  $\mathcal{V}$  has **right uniform restriction**.
- (2) *For finite  $\bar{x}, \bar{y}$ , the map  $i_* : \text{KCon}(\mathbf{F}_{\mathcal{V}}(\bar{y})) \rightarrow \text{KCon}(\mathbf{F}_{\mathcal{V}}(\bar{x}, \bar{y}))$  has a right adjoint;*
- (3) *For finitely presented  $\mathbf{A}, \mathbf{B} \in \mathcal{V}$  and any homomorphism  $f : \mathbf{A} \rightarrow \mathbf{B}$ , the map  $f_* : \text{KCon}(\mathbf{A}) \rightarrow \text{KCon}(\mathbf{B})$  has a right adjoint.*

## Examples

The following varieties have right uniform restriction:

- Heyting algebras and modal algebras,

## Examples

The following varieties have right uniform restriction:

- Heyting algebras and modal algebras,
- abelian groups, abelian  $\ell$ -groups and MV-algebras,



## Examples

The following varieties have right uniform restriction:

- Heyting algebras and modal algebras,
- abelian groups, abelian  $\ell$ -groups and MV-algebras,
- any locally finite variety.

## Examples

The following varieties have right uniform restriction:

- Heyting algebras and modal algebras,
- abelian groups, abelian  $\ell$ -groups and MV-algebras,
- any locally finite variety.

However, the varieties of **groups** and of **S4-algebras** do not have right uniform restriction.

## Left uniform restriction

### Definition

$\mathcal{V}$  has **left uniform restriction** iff for any finite set of equations  $\Sigma(\bar{x}, \bar{y})$ , there exists a finite set of equations  $\Sigma^L(\bar{y})$  such that for any set of equations  $\Pi(\bar{y})$ :

$$\Pi(\bar{y}) \models_{\mathcal{V}} \Sigma(\bar{x}, \bar{y}) \iff \Pi(\bar{y}) \models_{\mathcal{V}} \Sigma^L(\bar{y}).$$

## Left uniform restriction and compact congruences

### Theorem

*For any variety  $\mathcal{V}$ , the following are equivalent:*

- (1) For finitely presented  $\mathbf{A}, \mathbf{B} \in \mathcal{V}$  and any homomorphism  $f: \mathbf{A} \rightarrow \mathbf{B}$ , the map  $f_*: \text{KCon}(\mathbf{A}) \rightarrow \text{KCon}(\mathbf{B})$  has a left adjoint.*

## Left uniform restriction and compact congruences

### Theorem

*For any variety  $\mathcal{V}$ , the following are equivalent:*

- (1) *For finitely presented  $\mathbf{A}, \mathbf{B} \in \mathcal{V}$  and any homomorphism  $f: \mathbf{A} \rightarrow \mathbf{B}$ , the map  $f_*: \text{KCon}(\mathbf{A}) \rightarrow \text{KCon}(\mathbf{B})$  has a left adjoint.*
- (2)  *$\mathcal{V}$  has left uniform restriction*

## Left uniform restriction and compact congruences

### Theorem

*For any variety  $\mathcal{V}$ , the following are equivalent:*

- (1) *For finitely presented  $\mathbf{A}, \mathbf{B} \in \mathcal{V}$  and any homomorphism  $f: \mathbf{A} \rightarrow \mathbf{B}$ , the map  $f_*: \text{KCon}(\mathbf{A}) \rightarrow \text{KCon}(\mathbf{B})$  has a left adjoint.*
- (2)  *$\mathcal{V}$  has left uniform restriction **and** for finite  $\bar{x}$ , the join-semilattice  $\text{KCon}(\mathbf{F}_{\mathcal{V}}(\bar{x}))$  is residuated.*

## Examples

Heyting algebras have left uniform restriction,

## Examples

Heyting algebras have left uniform restriction, but not the variety  $\mathcal{ISL}$  of implicative semilattices, corresponding to the  $(\wedge, \rightarrow)$ -fragment of IPC.



## Examples

Heyting algebras have left uniform restriction, but not the variety  $\mathcal{ISL}$  of implicative semilattices, corresponding to the  $(\wedge, \rightarrow)$ -fragment of IPC.

For example,

$$\Sigma := \{\top \approx ((x \rightarrow z) \wedge (y \rightarrow z)) \rightarrow z\}$$

is a consequence of both  $\top \approx x$  and  $\top \approx y$ , i.e.,

$$\{\top \approx x\} \models_{\mathcal{ISL}} \Sigma \quad \text{and} \quad \{\top \approx y\} \models_{\mathcal{ISL}} \Sigma,$$

## Examples

Heyting algebras have left uniform restriction, but not the variety  $\mathcal{ISL}$  of implicative semilattices, corresponding to the  $(\wedge, \rightarrow)$ -fragment of IPC.

For example,

$$\Sigma := \{\top \approx ((x \rightarrow z) \wedge (y \rightarrow z)) \rightarrow z\}$$

is a consequence of both  $\top \approx x$  and  $\top \approx y$ , i.e.,

$$\{\top \approx x\} \models_{\mathcal{ISL}} \Sigma \quad \text{and} \quad \{\top \approx y\} \models_{\mathcal{ISL}} \Sigma,$$

but there is no  $\Delta(x, y)$  satisfying

$$\Delta \models_{\mathcal{ISL}} \Sigma, \quad \{\top \approx x\} \models_{\mathcal{ISL}} \Delta, \quad \text{and} \quad \{\top \approx y\} \models_{\mathcal{ISL}} \Delta.$$

## Locally finite case

### Theorem

*Let  $\mathcal{V}$  be a locally finite variety. Then:*

## Locally finite case

### Theorem

*Let  $\mathcal{V}$  be a locally finite variety. Then:*

- 1  $\mathcal{V}$  has right uniform restriction.

## Locally finite case

### Theorem

Let  $\mathcal{V}$  be a locally finite variety. Then:

- 1  $\mathcal{V}$  has right uniform restriction.
- 2  $\mathcal{V}$  has left uniform restriction



$\mathcal{V}$  is congruence-distributive **and** for finite  $\bar{x}, \bar{y}$ ,  
 $i_* : \text{Con}(\mathbf{F}_{\mathcal{V}}(\bar{x})) \rightarrow \text{Con}(\mathbf{F}_{\mathcal{V}}(\bar{x}, \bar{y}))$  preserves intersections.

## Final remarks

We focused on **uniform restriction** and the corresponding properties of **existence of adjoints**.

## Final remarks

We focused on **uniform restriction** and the corresponding properties of **existence of adjoints**.

**Uniform interpolation** is equivalent to uniform restriction + deductive interpolation,

## Final remarks

We focused on **uniform restriction** and the corresponding properties of **existence of adjoints**.

**Uniform interpolation** is equivalent to uniform restriction + deductive interpolation, and also to the existence of adjoints for **countably generated** free algebras:



## Final remarks

We focused on **uniform restriction** and the corresponding properties of **existence of adjoints**.

**Uniform interpolation** is equivalent to uniform restriction + deductive interpolation, and also to the existence of adjoints for **countably generated** free algebras:

### Theorem

*The following are equivalent for any variety  $\mathcal{V}$ :*

- (1)  $\mathcal{V}$  has right uniform interpolation.

## Final remarks

We focused on **uniform restriction** and the corresponding properties of **existence of adjoints**.

**Uniform interpolation** is equivalent to uniform restriction + deductive interpolation, and also to the existence of adjoints for **countably generated** free algebras:

### Theorem

*The following are equivalent for any variety  $\mathcal{V}$ :*

- (1)  $\mathcal{V}$  has right uniform interpolation.*
- (2) For any countable  $X$  and  $Y \subseteq X$ , the natural embedding of  $\mathbf{KCon}(\mathbf{F}_{\mathcal{V}}(Y))$  into  $\mathbf{KCon}(\mathbf{F}_{\mathcal{V}}(X))$  has a right adjoint.*

## Final remarks

Uniform interpolation can be understood as a (weaker) form of **quantifier elimination**.

## Final remarks

Uniform interpolation can be understood as a (weaker) form of **quantifier elimination**.

In particular, under certain conditions (e.g., for varieties of Heyting and modal algebras), uniform interpolation for  $\mathcal{V}$  implies the existence of a **model completion** for the first-order theory of  $\mathcal{V}$ .

S. Ghilardi and M. Zawadowski.  
*Sheaves, Games and Model Completions*,  
Trends in Logic Series, Kluwer (2002).

## Final remarks

Uniform interpolation can be understood as a (weaker) form of **quantifier elimination**.

In particular, under certain conditions (e.g., for varieties of Heyting and modal algebras), uniform interpolation for  $\mathcal{V}$  implies the existence of a **model completion** for the first-order theory of  $\mathcal{V}$ .

S. Ghilardi and M. Zawadowski.  
*Sheaves, Games and Model Completions*,  
Trends in Logic Series, Kluwer (2002).

Can we weaken these conditions to cover other classes of algebras, e.g., quasi-varieties, universal classes?

## Final question

## Final question

What, if any, is the relationship between uniform interpolation and admissibility?

## Final question

What, if any, is the relationship between uniform interpolation and admissibility?



