

# Stable Canonical Rules and Admissibility I

Nick Bezhanishvili

Institute for Logic, Language and Computation

University of Amsterdam

<http://www.phil.uu.nl/~bezhanishvili>

and

Silvio Ghilardi

Department of Mathematics

University of Milan

<http://homes.di.unimi.it/ghilardi>

joint work with G. Bezhanishvili and R. Iemhoff

# History

Axiomatization, the finite model property (fmp) and decidability are some of the most studied properties of non-classical logics.

# History

Axiomatization, the finite model property (fmp) and decidability are some of the most studied properties of non-classical logics.

(Harrop, 1957) If a logic is finitely axiomatizable and has the fmp, then it is decidable.

# History

Axiomatization, the finite model property (fmp) and decidability are some of the most studied properties of non-classical logics.

(Harrop, 1957) If a logic is finitely axiomatizable and has the fmp, then it is decidable.

In the 1960's the research on axiomatization and finite model property was mostly concerned with particular non-classical logics.

# History

Axiomatization, the finite model property (fmp) and decidability are some of the most studied properties of non-classical logics.

(Harrop, 1957) If a logic is finitely axiomatizable and has the fmp, then it is decidable.

In the 1960's the research on axiomatization and finite model property was mostly concerned with particular non-classical logics.

Since the 1970's general methods started to develop for classes of non-classical logics.

# History

- General methods for proving the fmp include **filtration** (Lemmon, Segerberg...) and **selective filtration** (Fine, Zakharyashev...).

# History

- General methods for proving the fmp include **filtration** (Lemmon, Segerberg...) and **selective filtration** (Fine, Zakharyashev...).
- General methods for axiomatizing large classes of logics include **Jankov-de Jongh formulas**, **Fine-Rautenberg formulas**, **subframe formulas**, and **canonical formulas** (Jankov, de Jongh, Fine, Rautenberg, Zakharyashev).

# History

- Jankov (1963, 68) associated with each finite subdirectly irreducible Heyting algebra  $A$  the formula encoding the “behaviour” of  $A$  and showed that uncountably many intermediate logics can be axiomatized by these formulas.



# History

- Jankov (1963, 68) associated with each finite subdirectly irreducible Heyting algebra  $A$  the formula encoding the “behaviour” of  $A$  and showed that uncountably many intermediate logics can be axiomatized by these formulas.

Similar formulas for finite Kripke frames were defined by de Jongh (1968).

# History

- Jankov (1963, 68) associated with each finite subdirectly irreducible Heyting algebra  $A$  the formula encoding the “behaviour” of  $A$  and showed that uncountably many intermediate logics can be axiomatized by these formulas.

Similar formulas for finite Kripke frames were defined by de Jongh (1968).

- Fine (1974) and Rautenberg (1980) introduced modal logic analogues of these formulas.

# History

- [Jankov](#) (1963, 68) associated with each finite subdirectly irreducible Heyting algebra  $A$  the formula encoding the “behaviour” of  $A$  and showed that uncountably many intermediate logics can be axiomatized by these formulas.

Similar formulas for finite Kripke frames were defined by [de Jongh](#) (1968).

- [Fine](#) (1974) and [Rautenberg](#) (1980) introduced modal logic analogues of these formulas.
- [Fine](#) (1985) introduced [subframe formulas](#) and axiomatized large classes of transitive modal logics by these formulas.

# History

- Jankov (1963, 68) associated with each finite subdirectly irreducible Heyting algebra  $A$  the formula encoding the “behaviour” of  $A$  and showed that uncountably many intermediate logics can be axiomatized by these formulas.

Similar formulas for finite Kripke frames were defined by de Jongh (1968).

- Fine (1974) and Rautenberg (1980) introduced modal logic analogues of these formulas.
- Fine (1985) introduced subframe formulas and axiomatized large classes of transitive modal logics by these formulas.
- There exist intermediate and transitive modal logics that are not axiomatizable by Jankov or subframe formulas.

# History

- Zakharyashev (1988-92) refined the Jankov and Fine methods, introduced **canonical formulas** and showed that each intermediate and transitive modal logic is axiomatizable by canonical formulas.

# History

- Zakharyashev (1988-92) refined the Jankov and Fine methods, introduced **canonical formulas** and showed that each intermediate and transitive modal logic is axiomatizable by canonical formulas.
- Jerabek (2009) extended canonical formulas to **canonical rules** and showed that each intermediate and transitive modal rule system is axiomatizable by canonical rules.

# Motivation

Zakharyashev's method of canonical formulas and Jerabek's method of canonical rules do not work in the non-transitive case.

# Motivation

Zakharyashev's method of canonical formulas and Jerabek's method of canonical rules do not work in the non-transitive case.

Whether canonical formulas can be extended to all modal logics (Zakharyashev) and canonical rules to all modal rule systems (Jerabek) was left as an open problem.



# Motivation

Zakharyashev's method of canonical formulas and Jerabek's method of canonical rules do not work in the non-transitive case.

Whether canonical formulas can be extended to all modal logics (Zakharyashev) and canonical rules to all modal rule systems (Jerabek) was left as an open problem.

We will introduce **stable canonical rules** and give a positive solution of Jerabek's problem.

# Motivation

Zakharyashev's method of canonical formulas and Jerabek's method of canonical rules do not work in the non-transitive case.

Whether canonical formulas can be extended to all modal logics (Zakharyashev) and canonical rules to all modal rule systems (Jerabek) was left as an open problem.

We will introduce **stable canonical rules** and give a positive solution of Jerabek's problem.

We also show how to utilise stable canonical rules to axiomatize all modal logics.

# Motivation

Zakharyashev's method of canonical formulas and Jerabek's method of canonical rules do not work in the non-transitive case.

Whether canonical formulas can be extended to all modal logics (Zakharyashev) and canonical rules to all modal rule systems (Jerabek) was left as an open problem.

We will introduce **stable canonical rules** and give a positive solution of Jerabek's problem.

We also show how to utilise stable canonical rules to axiomatize all modal logics.

This gives a positive solution of Zakharyashev's problem. However, the solution is via rules and not formulas.

# Motivation

The key to this is to develop an algebraic approach to canonical formulas and rules for intermediate and modal logics.

# Motivation

The key to this is to develop an algebraic approach to canonical formulas and rules for intermediate and modal logics.

This method relies on locally finite reducts of Heyting and modal algebras.

# Multiple-conclusion rules

A **multiple-conclusion modal rule** is an expression  $\Gamma/\Delta$ , where  $\Gamma, \Delta$  are finite sets of modal formulas.

# Multiple-conclusion rules

A **multiple-conclusion modal rule** is an expression  $\Gamma/\Delta$ , where  $\Gamma, \Delta$  are finite sets of modal formulas.

If  $\Delta = \{\varphi\}$ , then  $\Gamma/\Delta$  is called a **single-conclusion modal rule** and is written  $\Gamma/\varphi$ .

# Multiple-conclusion rules

A **multiple-conclusion modal rule** is an expression  $\Gamma/\Delta$ , where  $\Gamma, \Delta$  are finite sets of modal formulas.

If  $\Delta = \{\varphi\}$ , then  $\Gamma/\Delta$  is called a **single-conclusion modal rule** and is written  $\Gamma/\varphi$ .

If  $\Gamma = \emptyset$ , then  $\Gamma/\Delta$  is called an **assumption-free modal rule** and is written  $/\Delta$ .



# Multiple-conclusion rules

A **multiple-conclusion modal rule** is an expression  $\Gamma/\Delta$ , where  $\Gamma, \Delta$  are finite sets of modal formulas.

If  $\Delta = \{\varphi\}$ , then  $\Gamma/\Delta$  is called a **single-conclusion modal rule** and is written  $\Gamma/\varphi$ .

If  $\Gamma = \emptyset$ , then  $\Gamma/\Delta$  is called an **assumption-free modal rule** and is written  $/\Delta$ .

Assumption-free single-conclusion modal rules  $/\varphi$  can be identified with modal formulas  $\varphi$ .

# Modal rule systems

A **modal rule system** is a set  $\mathcal{S}$  of modal rules such that

# Modal rule systems

A **modal rule system** is a set  $\mathcal{S}$  of modal rules such that

- ①  $\varphi/\varphi \in \mathcal{S}$ .
- ②  $\varphi, \varphi \rightarrow \psi/\psi \in \mathcal{S}$ .
- ③  $\varphi/\Box\varphi \in \mathcal{S}$ .
- ④  $\varphi/\varphi \in \mathcal{S}$  for each theorem  $\varphi$  of **K**.
- ⑤ If  $\Gamma/\Delta \in \mathcal{S}$ , then  $\Gamma, \Gamma'/\Delta, \Delta' \in \mathcal{S}$ .
- ⑥ If  $\Gamma/\Delta, \varphi \in \mathcal{S}$  and  $\Gamma, \varphi/\Delta \in \mathcal{S}$ , then  $\Gamma/\Delta \in \mathcal{S}$ .
- ⑦ If  $\Gamma/\Delta \in \mathcal{S}$  and  $s$  is a substitution, then  $s(\Gamma)/s(\Delta) \in \mathcal{S}$ .

# Modal rule systems

A **modal rule system** is a set  $\mathcal{S}$  of modal rules such that

- ①  $\varphi/\varphi \in \mathcal{S}$ .
- ②  $\varphi, \varphi \rightarrow \psi/\psi \in \mathcal{S}$ .
- ③  $\varphi/\Box\varphi \in \mathcal{S}$ .
- ④  $\varphi/\varphi \in \mathcal{S}$  for each theorem  $\varphi$  of **K**.
- ⑤ If  $\Gamma/\Delta \in \mathcal{S}$ , then  $\Gamma, \Gamma'/\Delta, \Delta' \in \mathcal{S}$ .
- ⑥ If  $\Gamma/\Delta, \varphi \in \mathcal{S}$  and  $\Gamma, \varphi/\Delta \in \mathcal{S}$ , then  $\Gamma/\Delta \in \mathcal{S}$ .
- ⑦ If  $\Gamma/\Delta \in \mathcal{S}$  and  $s$  is a substitution, then  $s(\Gamma)/s(\Delta) \in \mathcal{S}$ .

We denote the least modal rule system by  $\mathbf{S_K}$ , and the complete lattice of modal rule systems by  $\Sigma(\mathbf{S_K})$ .

# Modal rule systems

For a set  $\Xi$  of multiple-conclusion modal rules, let  $\mathbf{S_K} + \Xi$  be the least modal rule system containing  $\Xi$ .

# Modal rule systems

For a set  $\Xi$  of multiple-conclusion modal rules, let  $\mathbf{S_K} + \Xi$  be the least modal rule system containing  $\Xi$ .

If  $\mathcal{S} = \mathbf{S_K} + \Xi$ , then we say that  $\mathcal{S}$  is **axiomatizable** by  $\Xi$ .

# Modal rule systems

For a set  $\Xi$  of multiple-conclusion modal rules, let  $\mathbf{S_K} + \Xi$  be the least modal rule system containing  $\Xi$ .

If  $\mathcal{S} = \mathbf{S_K} + \Xi$ , then we say that  $\mathcal{S}$  is **axiomatizable** by  $\Xi$ .

If  $\rho \in \mathcal{S}$ , then we say that the modal rule system  $\mathcal{S}$  **entails** or **derives** the modal rule  $\rho$ , and write  $\mathcal{S} \vdash \rho$ .

# Modal rule systems and modal logics



# Modal rule systems and modal logics

Let  $\Lambda(\mathbf{K})$  denote the lattice of all modal logics.

# Modal rule systems and modal logics

Let  $\Lambda(\mathbf{K})$  denote the lattice of all modal logics.

Given a modal rule system  $\mathcal{S}$ , let  $\Lambda(\mathcal{S}) = \{\varphi : \vdash \varphi \in \mathcal{S}\}$  be the corresponding modal logic,

# Modal rule systems and modal logics

Let  $\Lambda(\mathbf{K})$  denote the lattice of all modal logics.

Given a modal rule system  $\mathcal{S}$ , let  $\Lambda(\mathcal{S}) = \{\varphi : \text{/\varphi} \in \mathcal{S}\}$  be the corresponding modal logic,

and for a modal logic  $L$ , let  $\Sigma(L) = \mathbf{S_K} + \{\text{/\varphi} : \varphi \in L\}$  be the corresponding modal rule system.

## Modal rule systems and modal logics

Then  $\Lambda : \Sigma(\mathbf{S_K}) \rightarrow \Lambda(\mathbf{K})$  and  $\Sigma : \Lambda(\mathbf{K}) \rightarrow \Sigma(\mathbf{S_K})$  are order-preserving maps such that  $\Lambda(\Sigma(L)) = L$  for each  $L \in \Lambda(\mathbf{K})$  and  $\mathcal{S} \supseteq \Sigma(\Lambda(\mathcal{S}))$  for each  $\mathcal{S} \in \Sigma(\mathbf{S_K})$ .

## Modal rule systems and modal logics

Then  $\Lambda : \Sigma(\mathbf{S_K}) \rightarrow \Lambda(\mathbf{K})$  and  $\Sigma : \Lambda(\mathbf{K}) \rightarrow \Sigma(\mathbf{S_K})$  are order-preserving maps such that  $\Lambda(\Sigma(L)) = L$  for each  $L \in \Lambda(\mathbf{K})$  and  $\mathcal{S} \supseteq \Sigma(\Lambda(\mathcal{S}))$  for each  $\mathcal{S} \in \Sigma(\mathbf{S_K})$ .

Thus,  $\Lambda(\mathbf{K})$  embeds isomorphically into  $\Sigma(\mathbf{S_K})$ . But the embedding is not a lattice embedding.

## Modal rule systems and modal logics

Then  $\Lambda : \Sigma(\mathbf{S_K}) \rightarrow \Lambda(\mathbf{K})$  and  $\Sigma : \Lambda(\mathbf{K}) \rightarrow \Sigma(\mathbf{S_K})$  are order-preserving maps such that  $\Lambda(\Sigma(L)) = L$  for each  $L \in \Lambda(\mathbf{K})$  and  $\mathcal{S} \supseteq \Sigma(\Lambda(\mathcal{S}))$  for each  $\mathcal{S} \in \Sigma(\mathbf{S_K})$ .

Thus,  $\Lambda(\mathbf{K})$  embeds isomorphically into  $\Sigma(\mathbf{S_K})$ . But the embedding is not a lattice embedding.

We say that a modal logic  $L$  is **axiomatized** (over  $\mathbf{K}$ ) by a set  $\Xi$  of multiple-conclusion modal rules if  $L = \Lambda(\mathbf{S_K} + \Xi)$ .

# Modal algebras

A **modal algebra**  $\mathfrak{A} = (A, \Diamond)$  is a Boolean algebra  $A$  endowed with a unary operator  $\Diamond$  satisfying

①  $\Diamond 0 = 0$ ;

②  $\Diamond(a \vee b) = \Diamond a \vee \Diamond b$ .

# Modal algebras and modal rule systems

A modal algebra  $\mathfrak{A} = (A, \Diamond)$  **validates** a multiple-conclusion modal rule  $\Gamma/\Delta$  provided for every valuation  $V$  on  $A$ , if  $V(\gamma) = 1$  for all  $\gamma \in \Gamma$ , then  $V(\delta) = 1$  for some  $\delta \in \Delta$ .



# Modal algebras and modal rule systems

A modal algebra  $\mathfrak{A} = (A, \Diamond)$  **validates** a multiple-conclusion modal rule  $\Gamma/\Delta$  provided for every valuation  $V$  on  $A$ , if  $V(\gamma) = 1$  for all  $\gamma \in \Gamma$ , then  $V(\delta) = 1$  for some  $\delta \in \Delta$ .

Otherwise  $\mathfrak{A}$  **refutes**  $\Gamma/\Delta$ .

# Modal algebras and modal rule systems

A modal algebra  $\mathfrak{A} = (A, \Diamond)$  **validates** a multiple-conclusion modal rule  $\Gamma/\Delta$  provided for every valuation  $V$  on  $A$ , if  $V(\gamma) = 1$  for all  $\gamma \in \Gamma$ , then  $V(\delta) = 1$  for some  $\delta \in \Delta$ .

Otherwise  $\mathfrak{A}$  **refutes**  $\Gamma/\Delta$ .

If  $\mathfrak{A}$  validates  $\Gamma/\Delta$ , we write  $\mathfrak{A} \models \Gamma/\Delta$ , and if  $\mathfrak{A}$  refutes  $\Gamma/\Delta$ , we write  $\mathfrak{A} \not\models \Gamma/\Delta$ .

## Modal rule systems and universal classes

If  $\Gamma = \{\phi_1, \dots, \phi_n\}$ ,  $\Delta = \{\psi_1, \dots, \psi_m\}$ , and  $\phi_i(\underline{x})$  and  $\psi_j(\underline{x})$  are the terms in the first-order language of modal algebras corresponding to the  $\phi_i$  and  $\psi_j$ , then  $\mathfrak{A} \models \Gamma/\Delta$  iff  $\mathfrak{A}$  is a model of the universal sentence  $\forall \underline{x} (\bigwedge_{i=1}^n \phi_i(\underline{x}) = 1 \rightarrow \bigvee_{j=1}^m \psi_j(\underline{x}) = 1)$ .

## Modal rule systems and universal classes

If  $\Gamma = \{\phi_1, \dots, \phi_n\}$ ,  $\Delta = \{\psi_1, \dots, \psi_m\}$ , and  $\phi_i(\underline{x})$  and  $\psi_j(\underline{x})$  are the terms in the first-order language of modal algebras corresponding to the  $\phi_i$  and  $\psi_j$ , then  $\mathfrak{A} \models \Gamma/\Delta$  iff  $\mathfrak{A}$  is a model of the universal sentence  $\forall \underline{x} (\bigwedge_{i=1}^n \phi_i(\underline{x}) = 1 \rightarrow \bigvee_{j=1}^m \psi_j(\underline{x}) = 1)$ .

Modal rule systems correspond to (are complete for) **universal classes** of modal algebras.

# Modal rule systems and universal classes

If  $\Gamma = \{\phi_1, \dots, \phi_n\}$ ,  $\Delta = \{\psi_1, \dots, \psi_m\}$ , and  $\phi_i(\underline{x})$  and  $\psi_j(\underline{x})$  are the terms in the first-order language of modal algebras corresponding to the  $\phi_i$  and  $\psi_j$ , then  $\mathfrak{A} \models \Gamma/\Delta$  iff  $\mathfrak{A}$  is a model of the universal sentence  $\forall \underline{x} (\bigwedge_{i=1}^n \phi_i(\underline{x}) = 1 \rightarrow \bigvee_{j=1}^m \psi_j(\underline{x}) = 1)$ .

Modal rule systems correspond to (are complete for) **universal classes** of modal algebras.

A class of modal algebras is a universal class iff it is closed under isomorphisms, subalgebras, and ultraproducts.

# Modal logics and varieties

Modal logics correspond to (are complete for) equationally definable classes of modal algebras; that is, models of the sentences  $\forall \underline{x} \phi(\underline{x}) = 1$  in the first-order language of modal algebras.

# Modal logics and varieties

Modal logics correspond to (are complete for) equationally definable classes of modal algebras; that is, models of the sentences  $\forall \underline{x} \phi(\underline{x}) = 1$  in the first-order language of modal algebras.

A class of modal algebras is an equational class iff it is a **variety** (closed under homomorphic images, subalgebras, and products).

# Single-conclusion rule systems and quasi-varieties

Modal algebra  $\mathfrak{A}$  validates a single-conclusion modal rule  $\Gamma/\psi$  iff  $\mathfrak{A}$  is a model of the sentence  $\forall \underline{x} (\bigwedge_{i=1}^n \phi_i(\underline{x}) = 1 \rightarrow \psi(\underline{x}) = 1)$ , where  $\Gamma = \{\phi_1, \dots, \phi_n\}$  and  $\phi_i(\underline{x})$  and  $\psi(\underline{x})$  are the terms in the first-order language of modal algebras corresponding to the  $\phi_i$  and  $\psi$ .



# Single-conclusion rule systems and quasi-varieties

Modal algebra  $\mathfrak{A}$  validates a single-conclusion modal rule  $\Gamma/\psi$  iff  $\mathfrak{A}$  is a model of the sentence  $\forall \underline{x} (\bigwedge_{i=1}^n \phi_i(\underline{x}) = 1 \rightarrow \psi(\underline{x}) = 1)$ , where  $\Gamma = \{\phi_1, \dots, \phi_n\}$  and  $\phi_i(\underline{x})$  and  $\psi(\underline{x})$  are the terms in the first-order language of modal algebras corresponding to the  $\phi_i$  and  $\psi$ .

Single-conclusion modal rule systems, which are also known as **modal consequence relations**, correspond to (are complete for) **universal Horn classes** of modal algebras.

# Single-conclusion rule systems and quasi-varieties

Modal algebra  $\mathfrak{A}$  validates a single-conclusion modal rule  $\Gamma/\psi$  iff  $\mathfrak{A}$  is a model of the sentence  $\forall \underline{x} (\bigwedge_{i=1}^n \phi_i(\underline{x}) = 1 \rightarrow \psi(\underline{x}) = 1)$ , where  $\Gamma = \{\phi_1, \dots, \phi_n\}$  and  $\phi_i(\underline{x})$  and  $\psi(\underline{x})$  are the terms in the first-order language of modal algebras corresponding to the  $\phi_i$  and  $\psi$ .

Single-conclusion modal rule systems, which are also known as **modal consequence relations**, correspond to (are complete for) **universal Horn classes** of modal algebras.

A class of modal algebras is a universal Horn class iff it is a **quasi-variety** (closed under isomorphisms, subalgebras, products, and ultraproducts).

# Correspondence

Let  $\mathcal{S}$  be a modal rule system and  $\mathcal{U}$  be the universal class corresponding to  $\mathcal{S}$ .

Then the variety corresponding to the modal logic  $\Lambda(\mathcal{S})$  is the variety generated by  $\mathcal{U}$ .

## Locally finite varieties

A variety is **locally finite** if its every finitely generated algebra is finite.

## Locally finite varieties

A variety is **locally finite** if its every finitely generated algebra is finite.

Boolean algebras **are** locally finite.

## Locally finite varieties

A variety is **locally finite** if its every finitely generated algebra is finite.

Boolean algebras **are** locally finite.

**Theorem.** For each  $n \in \omega$ , the  $n$ -generated free Boolean algebra is isomorphic to the powerset of a  **$2^n$ -element set**.

## Locally finite varieties

A variety is **locally finite** if its every finitely generated algebra is finite.

Boolean algebras **are** locally finite.

**Theorem.** For each  $n \in \omega$ , the  $n$ -generated free Boolean algebra is isomorphic to the powerset of a  **$2^n$ -element set**.

But Heyting algebras **are not**.

# Locally finite varieties

A variety is **locally finite** if its every finitely generated algebra is finite.

Boolean algebras **are** locally finite.

**Theorem.** For each  $n \in \omega$ , the  $n$ -generated free Boolean algebra is isomorphic to the powerset of a  **$2^n$ -element set**.

But Heyting algebras **are not**.

**Theorem** (Rieger, 1949, Nishimura, 1960). The 1-generated free Heyting algebra, also called the **Rieger-Nishimura lattice**, is infinite.



# Locally finite varieties

A variety is **locally finite** if its every finitely generated algebra is finite.

Boolean algebras **are** locally finite.

**Theorem.** For each  $n \in \omega$ , the  $n$ -generated free Boolean algebra is isomorphic to the powerset of a  **$2^n$ -element set**.

But Heyting algebras **are not**.

**Theorem** (Rieger, 1949, Nishimura, 1960). The 1-generated free Heyting algebra, also called the **Rieger-Nishimura lattice**, is infinite.

The varieties of all modal algebras, **K4**-algebras and **S4**-algebras are **not** locally finite.

# Locally finite reducts of Heyting algebras

# Locally finite reducts of Heyting algebras

Heyting algebras  $(A, \wedge, \vee, \rightarrow, 0, 1)$ .

# Locally finite reducts of Heyting algebras

Heyting algebras  $(A, \wedge, \vee, \rightarrow, 0, 1)$ .

$\vee$ -free reducts  $(A, \wedge, \rightarrow, 0, 1)$ : **implicative semilattices**.

# Locally finite reducts of Heyting algebras

Heyting algebras  $(A, \wedge, \vee, \rightarrow, 0, 1)$ .

$\vee$ -free reducts  $(A, \wedge, \rightarrow, 0, 1)$ : **implicative semilattices**.

$\rightarrow$ -free reducts  $(A, \wedge, \vee, 0, 1)$ : **distributive lattices**.

# Locally finite reducts of Heyting algebras

Heyting algebras  $(A, \wedge, \vee, \rightarrow, 0, 1)$ .

$\vee$ -free reducts  $(A, \wedge, \rightarrow, 0, 1)$ : **implicative semilattices**.

$\rightarrow$ -free reducts  $(A, \wedge, \vee, 0, 1)$ : **distributive lattices**.

## Theorem.

- (Diego, 1966). The variety of implicative semilattices **is locally finite**.
- (Folklore). The variety of distributive lattices **is locally finite**.

## Locally finite reducts of Heyting algebras

$(\wedge, \rightarrow, 0)$ -reduct of Heyting algebras leads to  $(\wedge, \rightarrow, 0)$ -canonical formulas that are algebraic analogues of Zakharyashev's canonical formulas for intermediate logics.

# Locally finite reducts of Heyting algebras

$(\wedge, \rightarrow, 0)$ -reduct of Heyting algebras leads to  $(\wedge, \rightarrow, 0)$ -canonical formulas that are algebraic analogues of Zakharyashev's canonical formulas for intermediate logics.

$(\wedge, \vee, 0, 1)$ -reduct of Heyting algebras leads to a new class of  $(\wedge, \vee, 0, 1)$ -canonical formulas.



# Locally finite reducts of Heyting algebras

$(\wedge, \rightarrow, 0)$ -reduct of Heyting algebras leads to  $(\wedge, \rightarrow, 0)$ -canonical formulas that are algebraic analogues of Zakharyashev's canonical formulas for intermediate logics.

$(\wedge, \vee, 0, 1)$ -reduct of Heyting algebras leads to a new class of  $(\wedge, \vee, 0, 1)$ -canonical formulas.

## Theorem.

- (G. B and N. B., 2009). Every intermediate logic is axiomatizable by  $(\wedge, \rightarrow, 0)$ -canonical formulas.
- (G. B and N. B., 2013). Every intermediate logic is axiomatizable by  $(\wedge, \vee, 0, 1)$ -canonical formulas.

## Connection with filtrations

There are two standard methods for proving the finite model property for modal and intermediate logics: **filtration** and **selective filtration**.

## Connection with filtrations

There are two standard methods for proving the finite model property for modal and intermediate logics: **filtration** and **selective filtration**.

Taking  $(\wedge, \rightarrow, 0)$ -free reduct corresponds to **selective filtration**.

## Connection with filtrations

There are two standard methods for proving the finite model property for modal and intermediate logics: **filtration** and **selective filtration**.

Taking  $(\wedge, \rightarrow, 0)$ -free reduct corresponds to **selective filtration**.

Taking  $(\wedge, \vee, 0, 1)$ -free reduct corresponds to **filtration**.

## Connection with filtrations

There are two standard methods for proving the finite model property for modal and intermediate logics: **filtration** and **selective filtration**.

Taking  $(\wedge, \rightarrow, 0)$ -free reduct corresponds to **selective filtration**.

Taking  $(\wedge, \vee, 0, 1)$ -free reduct corresponds to **filtration**.

Modal analogues of  $(\wedge, \rightarrow, 0)$ -canonical formulas for transitive modal logics (extensions of **K4**) and weakly transitive modal logics (extensions of **wK4**) have been developed (G.B. and N.B 2011, 2012)

## Connection with filtrations

There are two standard methods for proving the finite model property for modal and intermediate logics: **filtration** and **selective filtration**.

Taking  $(\wedge, \rightarrow, 0)$ -free reduct corresponds to **selective filtration**.

Taking  $(\wedge, \vee, 0, 1)$ -free reduct corresponds to **filtration**.

Modal analogues of  $(\wedge, \rightarrow, 0)$ -canonical formulas for transitive modal logics (extensions of **K4**) and weakly transitive modal logics (extensions of **wK4**) have been developed (G.B. and N.B 2011, 2012)

These methods are based on an algebraic understanding of selective filtration (G.B., Ghilardi, Jibladze, 2011).

## Connection with filtrations

There are two standard methods for proving the finite model property for modal and intermediate logics: **filtration** and **selective filtration**.

Taking  $(\wedge, \rightarrow, 0)$ -free reduct corresponds to **selective filtration**.

Taking  $(\wedge, \vee, 0, 1)$ -free reduct corresponds to **filtration**.

Modal analogues of  $(\wedge, \rightarrow, 0)$ -canonical formulas for transitive modal logics (extensions of **K4**) and weakly transitive modal logics (extensions of **wK4**) have been developed (G.B. and N.B 2011, 2012)

These methods are based on an algebraic understanding of selective filtration (G.B., Ghilardi, Jibladze, 2011).

These formulas are algebraic analogues of Zakharyashev's canonical formulas for transitive modal logics.

## Connection with filtrations

Selective filtration works well only in the transitive case.



## Connection with filtrations

Selective filtration works well only in the transitive case.

In the non-transitive case one needs to employ standard filtration.

## Connection with filtrations

Selective filtration works well only in the transitive case.

In the non-transitive case one needs to employ standard filtration.

The method of filtration was originally developed by McKinsey and Tarski (1930s and 40s). Their technique was algebraic.

## Connection with filtrations

Selective filtration works well only in the transitive case.

In the non-transitive case one needs to employ standard filtration.

The method of filtration was originally developed by McKinsey and Tarski (1930s and 40s). Their technique was algebraic.

Lemmon (1960s) and Segerberg (1960s and 70s) developed model-theoretic approach to filtrations.

## Connection with filtrations

Selective filtration works well only in the transitive case.

In the non-transitive case one needs to employ standard filtration.

The method of filtration was originally developed by McKinsey and Tarski (1930s and 40s). Their technique was algebraic.

Lemmon (1960s) and Segerberg (1960s and 70s) developed model-theoretic approach to filtrations.

The two are connected through duality.

## Connection with filtrations

Selective filtration works well only in the transitive case.

In the non-transitive case one needs to employ standard filtration.

The method of filtration was originally developed by McKinsey and Tarski (1930s and 40s). Their technique was algebraic.

Lemmon (1960s) and Segerberg (1960s and 70s) developed model-theoretic approach to filtrations.

The two are connected through duality.

The modern account is discussed in Ghilardi (2010) and van Alten et al. (2013).

## Filtrations model theoretically

Let  $\mathfrak{M} = (X, R, V)$  be a Kripke model and let  $\Sigma$  be a set of formulas closed under subformulas.

## Filtrations model theoretically

Let  $\mathfrak{M} = (X, R, V)$  be a Kripke model and let  $\Sigma$  be a set of formulas closed under subformulas.

Define an equivalence relation  $\sim_\Sigma$  on  $X$  by

$$x \sim_\Sigma y \text{ iff } (\forall \varphi \in \Sigma)(x \models \varphi \Leftrightarrow y \models \varphi).$$

Let  $X' = X/\sim_\Sigma$  and let  $V'(p) = \{[x] : x \in V(p)\}$ , where  $[x]$  is the equivalence class of  $x$  with respect to  $\sim_\Sigma$ .

## Filtrations model theoretically

Let  $\mathfrak{M} = (X, R, V)$  be a Kripke model and let  $\Sigma$  be a set of formulas closed under subformulas.

Define an equivalence relation  $\sim_\Sigma$  on  $X$  by

$$x \sim_\Sigma y \text{ iff } (\forall \varphi \in \Sigma)(x \models \varphi \Leftrightarrow y \models \varphi).$$

Let  $X' = X/\sim_\Sigma$  and let  $V'(p) = \{[x] : x \in V(p)\}$ , where  $[x]$  is the equivalence class of  $x$  with respect to  $\sim_\Sigma$ .

**Definition.** For a binary relation  $R'$  on  $X'$ , we say that the triple  $\mathfrak{M}' = (X', R', V')$  is a **filtration of  $\mathfrak{M}$  through  $\Sigma$**  if the following two conditions are satisfied:

(F1)  $xRy \Rightarrow [x]R'[y]$ .

(F2)  $[x]R'[y] \Rightarrow (\forall \Diamond \varphi \in \Sigma)(y \models \varphi \Rightarrow x \models \Diamond \varphi)$ .



# Stable homomorphisms and CDC

The key concepts for developing an algebraic approach to filtrations are stable homomorphisms and the closed domain condition (CDC).

# Stable homomorphisms and CDC

The key concepts for developing an algebraic approach to filtrations are stable homomorphisms and the closed domain condition (CDC).

**Definition.** Let  $\mathfrak{A} = (A, \Diamond)$  and  $\mathfrak{B} = (B, \Diamond)$  be modal algebras and let  $h : A \rightarrow B$  be a Boolean homomorphism. We call  $h$  a **stable homomorphism** provided  $\Diamond h(a) \leq h(\Diamond a)$  for each  $a \in A$ .

# Stable homomorphisms and CDC

The key concepts for developing an algebraic approach to filtrations are stable homomorphisms and the closed domain condition (CDC).

**Definition.** Let  $\mathfrak{A} = (A, \Diamond)$  and  $\mathfrak{B} = (B, \Diamond)$  be modal algebras and let  $h : A \rightarrow B$  be a Boolean homomorphism. We call  $h$  a **stable homomorphism** provided  $\Diamond h(a) \leq h(\Diamond a)$  for each  $a \in A$ .

It is easy to see that  $h : A \rightarrow B$  is stable iff  $h(\Box a) \leq \Box h(a)$  for each  $a \in A$ .

# Stable homomorphisms and CDC

The key concepts for developing an algebraic approach to filtrations are stable homomorphisms and the closed domain condition (CDC).

**Definition.** Let  $\mathfrak{A} = (A, \Diamond)$  and  $\mathfrak{B} = (B, \Diamond)$  be modal algebras and let  $h : A \rightarrow B$  be a Boolean homomorphism. We call  $h$  a **stable homomorphism** provided  $\Diamond h(a) \leq h(\Diamond a)$  for each  $a \in A$ .

It is easy to see that  $h : A \rightarrow B$  is stable iff  $h(\Box a) \leq \Box h(a)$  for each  $a \in A$ .

Stable homomorphisms were studied by G. B., Mines, Morandi (2008), Ghilardi (2010), and Coumans, van Gool (2012).

# Stable homomorphisms and CDC

**Definition.** Let  $\mathfrak{A} = (A, \Diamond)$  and  $\mathfrak{B} = (B, \Diamond)$  be modal algebras and let  $h : A \rightarrow B$  be a stable homomorphism. We say that  $h$  satisfies **the closed domain condition (CDC)** for  $D \subseteq A$  if  $h(\Diamond a) = \Diamond h(a)$  for  $a \in D$ .

## Key idea

Let  $(A, \Diamond)$  and  $(B, \Diamond)$  be modal algebras,  $(X, R)$  and  $(Y, R)$  be their duals,  $h : A \rightarrow B$  be a Boolean homomorphism and  $f : Y \rightarrow X$  be the dual of  $h$ .

## Key idea

Let  $(A, \Diamond)$  and  $(B, \Diamond)$  be modal algebras,  $(X, R)$  and  $(Y, R)$  be their duals,  $h : A \rightarrow B$  be a Boolean homomorphism and  $f : Y \rightarrow X$  be the dual of  $h$ .

Then

- 1  $h$  is one-to-one iff  $f$  is onto.
- 2  $h$  is stable iff  $f$  is stable (that is,  $xRy$  implies  $f(x)Rf(y)$ ).
- 3  $h$  is a modal homomorphism iff  $f$  is a p-morphism.
- 4 If  $h$  is stable but not a modal homomorphism it may still be the case that  $h(\Diamond a) = \Diamond h(a)$  for some  $a \in D \subseteq A$ .

# Key idea

Consequently:

- ① Being a stable homomorphism dually corresponds to satisfying condition (F1) in the definition of filtration.



# Key idea

Consequently:

- 1 Being a stable homomorphism dually corresponds to satisfying condition (F1) in the definition of filtration.
- 2 Satisfying (CDC) dually corresponds to satisfying condition (F2) in the definition of filtration.

# Filtrations and finite refutation patterns

## Refutation Pattern Theorem.

- 1 If  $\mathbf{S}_K \not\models \Gamma/\Delta$ , then there exist  $(\mathfrak{A}_1, D_1), \dots, (\mathfrak{A}_n, D_n)$  such that each  $\mathfrak{A}_i = (A_i, \diamond_i)$  is a finite modal algebra,  $D_i \subseteq A_i$ , and for each modal algebra  $\mathfrak{B} = (B, \diamond)$ , we have  $\mathfrak{B} \not\models \Gamma/\Delta$  iff there is  $i \leq n$  and a stable embedding  $h : A_i \rightarrowtail B$  satisfying (CDC) for  $D_i$ .
- 2 If  $\mathbf{K} \not\models \varphi$ , then there exist  $(\mathfrak{A}_1, D_1), \dots, (\mathfrak{A}_n, D_n)$  such that each  $\mathfrak{A}_i = (A_i, \diamond_i)$  is a finite modal algebra,  $D_i \subseteq A_i$ , and for each modal algebra  $\mathfrak{B} = (B, \diamond)$ , we have  $\mathfrak{B} \not\models \varphi$  iff there is  $i \leq n$  and a stable embedding  $h : A_i \rightarrowtail B$  satisfying (CDC) for  $D_i$ .

## Proof sketch

If  $\mathbf{S_K} \not\models \Gamma/\Delta$ , then there is a modal algebra  $\mathfrak{A} = (A, \Diamond)$  refuting  $\Gamma/\Delta$ .

## Proof sketch

If  $\mathbf{S_K} \not\models \Gamma/\Delta$ , then there is a modal algebra  $\mathfrak{A} = (A, \Diamond)$  refuting  $\Gamma/\Delta$ .

Therefore, there is a valuation  $V$  on  $A$  such that  $V(\gamma) = 1_A$  for each  $\gamma \in \Gamma$  and  $V(\delta) \neq 1_A$  for each  $\delta \in \Delta$ .

## Proof sketch

If  $\mathbf{S_K} \not\models \Gamma/\Delta$ , then there is a modal algebra  $\mathfrak{A} = (A, \Diamond)$  refuting  $\Gamma/\Delta$ .

Therefore, there is a valuation  $V$  on  $A$  such that  $V(\gamma) = 1_A$  for each  $\gamma \in \Gamma$  and  $V(\delta) \neq 1_A$  for each  $\delta \in \Delta$ .

Let  $\Sigma$  be the set of subformulas of  $\Gamma \cup \Delta$ ,  $A'$  be the Boolean subalgebra of  $A$  generated by  $V(\Sigma)$ , and  $\mathfrak{A}' = (A', \Diamond')$  be a filtration of  $\mathfrak{A}$  through  $\Sigma$ .

## Proof sketch

If  $\mathbf{S_K} \not\models \Gamma/\Delta$ , then there is a modal algebra  $\mathfrak{A} = (A, \diamond)$  refuting  $\Gamma/\Delta$ .

Therefore, there is a valuation  $V$  on  $A$  such that  $V(\gamma) = 1_A$  for each  $\gamma \in \Gamma$  and  $V(\delta) \neq 1_A$  for each  $\delta \in \Delta$ .

Let  $\Sigma$  be the set of subformulas of  $\Gamma \cup \Delta$ ,  $A'$  be the Boolean subalgebra of  $A$  generated by  $V(\Sigma)$ , and  $\mathfrak{A}' = (A', \diamond')$  be a filtration of  $\mathfrak{A}$  through  $\Sigma$ .

Then  $\mathfrak{A}'$  is a finite modal algebra refuting  $\Gamma/\Delta$ . In fact,  $|A'| \leq m$ , where  $m = 2^{2^{|\Sigma|}}$  is the size of the free Boolean algebra on  $|\Sigma|$ -generators.

## Proof sketch

Let  $\mathfrak{A}_1, \dots, \mathfrak{A}_n$  be the list of all finite modal algebras  $\mathfrak{A}_i = (A_i, \Diamond_i)$  of size  $\leq m$  refuting  $\Gamma/\Delta$ .

## Proof sketch

Let  $\mathfrak{A}_1, \dots, \mathfrak{A}_n$  be the list of all finite modal algebras  $\mathfrak{A}_i = (A_i, \Diamond_i)$  of size  $\leq m$  refuting  $\Gamma/\Delta$ .

Let  $V_i$  be a valuation on  $A_i$  refuting  $\Gamma/\Delta$ ; that is,  $V_i(\gamma) = 1_{A_i}$  for each  $\gamma \in \Gamma$  and  $V_i(\delta) \neq 1_{A_i}$  for each  $\delta \in \Delta$ . Set  $D_i = \{V_i(\psi) : \Diamond\psi \in \Sigma\}$ .



## Proof sketch

Let  $\mathfrak{A}_1, \dots, \mathfrak{A}_n$  be the list of all finite modal algebras  $\mathfrak{A}_i = (A_i, \Diamond_i)$  of size  $\leq m$  refuting  $\Gamma/\Delta$ .

Let  $V_i$  be a valuation on  $A_i$  refuting  $\Gamma/\Delta$ ; that is,  $V_i(\gamma) = 1_{A_i}$  for each  $\gamma \in \Gamma$  and  $V_i(\delta) \neq 1_{A_i}$  for each  $\delta \in \Delta$ . Set  $D_i = \{V_i(\psi) : \Diamond\psi \in \Sigma\}$ .

**Key step:** Given a modal algebra  $\mathfrak{B} = (B, \Diamond)$ , we show that  $\mathfrak{B} \not\models \Gamma/\Delta$  iff there is  $i \leq n$  and a stable embedding  $h : A_i \rightarrowtail B$  satisfying (CDC) for  $D_i$ .

# Stable canonical rules

**Definition.** Let  $\mathfrak{A} = (A, \Diamond)$  be a finite modal algebra and let  $D$  be a subset of  $A$ . For each  $a \in A$  we introduce a new propositional letter  $p_a$  and define the **stable canonical rule**  $\rho(\mathfrak{A}, D)$  associated with  $\mathfrak{A}$  and  $D$  as  $\Gamma/\Delta$ , where:

$$\begin{aligned}\Gamma = & \{p_{a \vee b} \leftrightarrow p_a \vee p_b : a, b \in A\} \cup \\ & \{p_{\neg a} \leftrightarrow \neg p_a : a \in A\} \cup \\ & \{\Diamond p_a \rightarrow p_{\Diamond a} : a \in A\} \cup \\ & \{p_{\Diamond a} \rightarrow \Diamond p_a : a \in D\},\end{aligned}$$

and

$$\Delta = \{p_a \leftrightarrow p_b : a, b \in A, a \neq b\}.$$

## Stable canonical rules

**Stable Canonical Rule Theorem.** Let  $\mathfrak{A} = (A, \Diamond)$  be a finite modal algebra,  $D \subseteq A$ , and  $\mathfrak{B} = (B, \Diamond)$  be a modal algebra. Then  $\mathfrak{B} \not\models \rho(\mathfrak{A}, D)$  iff there is a stable embedding  $h : A \rightarrowtail B$  satisfying (CDC) for  $D$ .

# Stable canonical rules

## Corollary.

- ① If  $\mathbf{S_K} \not\models \Gamma/\Delta$ , then there exist  $(\mathfrak{A}_1, D_1), \dots, (\mathfrak{A}_n, D_n)$  such that each  $\mathfrak{A}_i = (A_i, \Diamond_i)$  is a finite modal algebra,  $D_i \subseteq A_i$ , and for each modal algebra  $\mathfrak{B} = (B, \Diamond)$ , we have:

$$\mathfrak{B} \models \Gamma/\Delta \text{ iff } \mathfrak{B} \models \rho(\mathfrak{A}_1, D_1), \dots, \rho(\mathfrak{A}_n, D_n).$$

- ② If  $\mathbf{K} \not\models \varphi$ , then there exist  $(\mathfrak{A}_1, D_1), \dots, (\mathfrak{A}_n, D_n)$  such that each  $\mathfrak{A}_i = (A_i, \Diamond_i)$  is a finite modal algebra,  $D_i \subseteq A_i$ , and for each modal algebra  $\mathfrak{B} = (B, \Diamond)$ , we have:

$$\mathfrak{B} \models \varphi \text{ iff } \mathfrak{B} \models \rho(\mathfrak{A}_1, D_1), \dots, \rho(\mathfrak{A}_n, D_n).$$

# Proof

Suppose  $\mathbf{S_K} \not\models \Gamma/\Delta$ .

## Proof

Suppose  $\mathbf{S_K} \not\models \Gamma/\Delta$ .

By the Refutation Pattern Theorem, there exist  $(\mathfrak{A}_1, D_1), \dots, (\mathfrak{A}_n, D_n)$  such that each  $\mathfrak{A}_i = (A_i, \Diamond_i)$  is a finite modal algebra,  $D_i \subseteq A_i$ , and for each modal algebra  $\mathfrak{B} = (B, \Diamond)$ , we have  $\mathfrak{B} \not\models \Gamma/\Delta$  iff there is  $i \leq n$  and a stable embedding  $h : A_i \rightarrowtail B$  satisfying (CDC) for  $D_i$ .

## Proof

Suppose  $\mathbf{S_K} \not\models \Gamma/\Delta$ .

By the Refutation Pattern Theorem, there exist  $(\mathfrak{A}_1, D_1), \dots, (\mathfrak{A}_n, D_n)$  such that each  $\mathfrak{A}_i = (A_i, \Diamond_i)$  is a finite modal algebra,  $D_i \subseteq A_i$ , and for each modal algebra  $\mathfrak{B} = (B, \Diamond)$ , we have  $\mathfrak{B} \not\models \Gamma/\Delta$  iff there is  $i \leq n$  and a stable embedding  $h : A_i \rightarrowtail B$  satisfying (CDC) for  $D_i$ .

By the Stable Canonical Rule Theorem, this is equivalent to the existence of  $i \leq n$  such that  $\mathfrak{B} \not\models \rho(\mathfrak{A}_i, D_i)$ .

# Proof

Suppose  $\mathbf{S_K} \not\models \Gamma/\Delta$ .

By the Refutation Pattern Theorem, there exist  $(\mathfrak{A}_1, D_1), \dots, (\mathfrak{A}_n, D_n)$  such that each  $\mathfrak{A}_i = (A_i, \Diamond_i)$  is a finite modal algebra,  $D_i \subseteq A_i$ , and for each modal algebra  $\mathfrak{B} = (B, \Diamond)$ , we have  $\mathfrak{B} \not\models \Gamma/\Delta$  iff there is  $i \leq n$  and a stable embedding  $h : A_i \rightarrowtail B$  satisfying (CDC) for  $D_i$ .

By the Stable Canonical Rule Theorem, this is equivalent to the existence of  $i \leq n$  such that  $\mathfrak{B} \not\models \rho(\mathfrak{A}_i, D_i)$ .

Thus,  $\mathfrak{B} \models \Gamma/\Delta$  iff  $\mathfrak{B} \models \rho(\mathfrak{A}_1, D_1), \dots, \rho(\mathfrak{A}_n, D_n)$ .



# Main Theorem

- ① Each modal rule system  $S$  over  $\mathbf{S}_K$  is axiomatizable by stable canonical rules.
- ② Each modal logic  $L$  is axiomatizable by stable canonical rules.

# Conclusions

# Conclusions

Part 1 of the Main Theorem yields a solution of an open problem of Jerabek.

# Conclusions

Part 1 of the Main Theorem yields a solution of an open problem of Jerabek.

Part 2 yields a solution of an open problem of Zakharyashev. However, our solution is by means of multiple-conclusion rules rather than formulas.

# Conclusions

Part 1 of the Main Theorem yields a solution of an open problem of Jerabek.

Part 2 yields a solution of an open problem of Zakharyashev. However, our solution is by means of multiple-conclusion rules rather than formulas.

Also our axiomatization requires to work with all finite modal algebras. It is not sufficient to work with only finite s.i. modal algebras.

# Conclusions

Part 1 of the Main Theorem yields a solution of an open problem of Jerabek.

Part 2 yields a solution of an open problem of Zakharyashev. However, our solution is by means of multiple-conclusion rules rather than formulas.

Also our axiomatization requires to work with all finite modal algebras. It is not sufficient to work with only finite s.i. modal algebras.

Various applications of this method will be discussed in Part II of the talk by Silvio Ghilardi.

Thank you!