

Admissible Infinitary Rules in Modal Logic I

Admissible Infinitary Rules in Modal Logic I

Wojciech Dzik

Institute of Mathematics, Silesian University, Katowice, Poland,
wdzik@wdzik.pl

Admissible Infinitary Rules in Modal Logic I

Wojciech Dzik

Institute of Mathematics, Silesian University, Katowice, Poland,
wdzik@wdzik.pl

coauthor

Piotr Wojtylak,

Institute of Mathematics, University of Opole, Opole, Poland,
piotr.wojtylak@gmail.com

Admissible Infinitary Rules in Modal Logic I

Wojciech Dzik

Institute of Mathematics, Silesian University, Katowice, Poland,
wdzik@wdzik.pl

coauthor

Piotr Wojtylak,

Institute of Mathematics, University of Opole, Opole, Poland,
piotr.wojtylak@gmail.com

Workshop on Admissible Rules and Unification II,
Les Diablerets 30 January - 2 February 2015

We admit infinitary rules, i.e. of the form $\{\varphi_i : i < \omega\} / \psi$
and infinitary consequence operations C_n . Why?

We admit infinitary rules, i.e. of the form $\{\varphi_i : i < \omega\} / \psi$
and infinitary consequence operations Cn . Why?

MAIN RESULTS:

- For a consequence operation Cn extending m.l. S4.3 TFAE:
 - (i) Cn is Almost Structurally Complete (ASCpl),
 - (ii) Cn is finitely approximable, i.e. there is a class \mathbb{K} of *finite* algebras such that $Cn = \overrightarrow{\mathbb{K}}$, where $\overrightarrow{\mathbb{K}}$ is a consequence operation determined by \mathbb{K} .

We admit infinitary rules, i.e. of the form $\{\varphi_i : i < \omega\} / \psi$
and infinitary consequence operations Cn . Why?

MAIN RESULTS:

- For a consequence operation Cn extending m.l. S4.3 TFAE:
 - (i) Cn is Almost Structurally Complete (ASCpl),
 - (ii) Cn is finitely approximable, i.e. there is a class \mathbb{K} of *finite* algebras such that $Cn = \overrightarrow{\mathbb{K}}$, where $\overrightarrow{\mathbb{K}}$ is a consequence operation determined by \mathbb{K} .
- An infinite basis for infinitary admissible (non-passive) rules for structurally complete extensions Cn^Σ is provided

We admit infinitary rules, i.e. of the form $\{\varphi_i : i < \omega\} / \psi$ and infinitary consequence operations Cn . Why?

MAIN RESULTS:

- For a consequence operation Cn extending m.l. S4.3 TFAE:
 - (i) Cn is Almost Structurally Complete (ASCpl),
 - (ii) Cn is finitely approximable, i.e. there is a class \mathbb{K} of *finite* algebras such that $Cn = \overrightarrow{\mathbb{K}}$, where $\overrightarrow{\mathbb{K}}$ is a consequence operation determined by \mathbb{K} .
- An infinite basis for infinitary admissible (non-passive) rules for structurally complete extensions Cn^Σ is provided
- a formula provable in Cn^Σ has a syntactic proof of the type $\leq \omega + 1$,

We admit infinitary rules, i.e. of the form $\{\varphi_i : i < \omega\} / \psi$
and infinitary consequence operations Cn . Why?

MAIN RESULTS:

- For a consequence operation Cn extending m.l. S4.3 TFAE:
 - (i) Cn is Almost Structurally Complete (ASCpl),
 - (ii) Cn is finitely approximable, i.e. there is a class \mathbb{K} of *finite* algebras such that $Cn = \overrightarrow{\mathbb{K}}$, where $\overrightarrow{\mathbb{K}}$ is a consequence operation determined by \mathbb{K} .
 - An infinite basis for infinitary admissible (non-passive) rules for structurally complete extensions Cn^Σ is provided
 - a formula provable in Cn^Σ has a syntactic proof of the type $\leq \omega + 1$,
- The key step: all extensions of S4.3 enjoy projective unification (D.-W 2009).

R. Bull (1966) Every normal extension of S4.3 has the *FMP*

R. Bull (1966) Every normal extension of S4.3 has the *FMP*

Kit Fine (1971) Every normal extension of S4.3:

- has the Finite Frame Prop. (for frames)
- is *finitely axiomatizable* and
- is characterized by finite chains of clusters.

Normal Axiomatic Extensions - the lattice NExtS4.3

R. Bull (1966) Every normal extension of S4.3 has the *FMP*

Kit Fine (1971) Every normal extension of S4.3:

- has the Finite Frame Prop. (for frames)
- is *finitely axiomatizable* and
- is characterized by finite chains of clusters.

Normal Axiomatic Extensions - the lattice NExtS4.3

Question: lift the results from theoremhood to derivability

Describe the lattice $\text{EXT}_{fin}\text{S4.3}$

★ Syntactic and semantic description of **finitary** (structural) consequence operations $Cn \in \text{EXT}_{\text{fin}}$ **S4.3**:

- form of *admissible (passive)* rules in consequence oper. Cn ,
- Let \mathcal{K} is a class of subdir. irr. S4.3-algebras characterizing $L \in \text{NExt}$ **S4.3**. Then, for any consequence oper. $Cn \geq Cn_L$:
 - Cn is characterized by a class of algebras of the form of the direct products $\mathcal{A} \times \mathcal{H}_n$, where $\mathcal{A} \in \mathcal{K}$ and \mathcal{H}_n is so called *Henle algebra* with n -atoms, i.e.
 Cn has Strong Finite Model Property (*SFMP*).
 - Cn is *finitely based* (can obtained by adding finitely many rules) and decidable
- The lattice EXT_{fin} **S4.3** of all consequence relations extending S4.3 is countable and distributive (a Heyting algebra).

$Var = \{p_1, p_2, \dots\}$ all propositional variables

Fm all modal formulas built up with \wedge, \neg, \Box, \top ; $Fm_n \{p_i : i \leq n\}$

$\rightarrow, \vee, \leftrightarrow, \Diamond, \perp$ as usual;

$Var = \{p_1, p_2, \dots\}$ all propositional variables

Fm all modal formulas built up with \wedge, \neg, \Box, \top ; $Fm_n \{p_i : i \leq n\}$

$\rightarrow, \vee, \leftrightarrow, \Diamond, \perp$ as usual;

$(Fm, \wedge, \neg, \Box, \top)$ the algebra of modal language, $\varepsilon: Var \rightarrow Fm$

substitution; A *modal logic* - any subset L of Fm containing all classical tautologies, the axiom

$(K) : \Box(\alpha \rightarrow \beta) \rightarrow (\Box\alpha \rightarrow \Box\beta)$ and closed under substit. and

$$MP : \frac{\alpha \rightarrow \beta, \alpha}{\beta} \quad \text{and} \quad RN : \frac{\alpha}{\Box\alpha}.$$

K the least, **S4** = **K** + $(T) : \Box\alpha \rightarrow \alpha$ + $(4) : \Box\Box\alpha \rightarrow \Box\alpha$.

S4.3 = **S4** + $(.3) : \Box(\Box\alpha \rightarrow \Box\beta) \vee \Box(\Box\beta \rightarrow \Box\alpha)$

$Var = \{p_1, p_2, \dots\}$ all propositional variables

Fm all modal formulas built up with \wedge, \neg, \Box, \top ; $Fm_n \{p_i : i \leq n\}$

$\rightarrow, \vee, \leftrightarrow, \Diamond, \perp$ as usual;

$(Fm, \wedge, \neg, \Box, \top)$ the algebra of modal language, $\varepsilon: Var \rightarrow Fm$

substitution; A *modal logic* - any subset L of Fm containing all classical tautologies, the axiom

$(K): \Box(\alpha \rightarrow \beta) \rightarrow (\Box\alpha \rightarrow \Box\beta)$ and closed under substit. and

$$MP: \frac{\alpha \rightarrow \beta, \alpha}{\beta} \quad \text{and} \quad RN: \frac{\alpha}{\Box\alpha}.$$

K the least, **S4** = **K** + $(T): \Box\alpha \rightarrow \alpha$ + $(4): \Box\Box\alpha \rightarrow \Box\alpha$.

S4.3 = **S4** + $(.3): \Box(\Box\alpha \rightarrow \Box\beta) \vee \Box(\Box\beta \rightarrow \Box\alpha)$

$$\text{NExt}\mathbf{S4.3} \ni L \mapsto Cn_L \in \text{EXT}_{\text{fin}}\mathbf{S4.3}$$

its *global consequence relation*; $\alpha \in Cn_L(X)$ means: α can be derived from $X \cup L$ using the rules *MP* and *RN*.

infinitary rules and onsequence operation

infinitary rules, i.e. X/α with X infinite,

infinitary rules and onsequence operation

infinitary rules, i.e. X/α with X infinite, two different notions of (almost) structural completeness

infinitary rules and consequence operation

infinitary rules, i.e. X/α with X infinite, two different notions of (almost) structural completeness

A *consequence operation*, (A.Tarski) - any mapping

$Cn : 2^{Fm} \rightarrow 2^{Fm}$ for each $X, Y \subseteq Fm$

$$X \subseteq Cn(X), \quad Cn(X) \subseteq Cn(X \cup Y), \quad Cn(Cn(X)) \subseteq Cn(X).$$

infinitary rules and consequence operation

infinitary rules, i.e. X/α with X infinite, two different notions of (almost) structural completeness

A *consequence operation*, (A.Tarski) - any mapping $Cn : 2^{Fm} \rightarrow 2^{Fm}$ for each $X, Y \subseteq Fm$

$$X \subseteq Cn(X), \quad Cn(X) \subseteq Cn(X \cup Y), \quad Cn(Cn(X)) \subseteq Cn(X).$$

A consequence operation Cn is *finitary* ($Cn \in Fin$), if

$$Cn(X) = \bigcup \{Cn(Y) : Y \text{ is finite and } Y \subseteq X\}, \quad \text{for each } X \subseteq Fm.$$

For any Cn define its 'finitary fragment' Cn_{fin} putting

$$Cn_{fin}(X) = \bigcup \{Cn(Y) : Y \text{ is finite and } Y \subseteq X\}, \quad \text{for each } X \subseteq Fm.$$

Theorem (Deduction Theorem)

If L is a modal logic and $\mathbf{S4} \subseteq L$, then
 $\alpha \in \mathbf{Cn}_L(X, \beta)$ *iff* $\Box\beta \rightarrow \alpha \in \mathbf{Cn}_L(X)$.

Theorem (Deduction Theorem)

If L is a modal logic and $\mathbf{S4} \subseteq L$, then
 $\alpha \in \mathbf{Cn}_L(X, \beta)$ *iff* $\Box\beta \rightarrow \alpha \in \mathbf{Cn}_L(X)$.

For Cn a modal consequence operation,

$\text{EXT}(Cn)$ (or $\text{EXT}_{fin}(Cn)$) denotes the lattice of all (or all finitary) consequence operations extending Cn .

Theorem (Deduction Theorem)

If L is a modal logic and $\mathbf{S4} \subseteq L$, then
 $\alpha \in Cn_L(X, \beta)$ *iff* $\Box\beta \rightarrow \alpha \in Cn_L(X)$.

For Cn a modal consequence operation,

$\text{EXT}(Cn)$ (or $\text{EXT}_{fin}(Cn)$) denotes the lattice of all (or all finitary) consequence operations extending Cn .

We write $\text{EXT}(L)$ for $\text{EXT}(Cn_L)$.

Theorem (Deduction Theorem)

If L is a modal logic and $\mathbf{S4} \subseteq L$, then
 $\alpha \in Cn_L(X, \beta)$ *iff* $\Box\beta \rightarrow \alpha \in Cn_L(X)$.

For Cn a modal consequence operation,

$\text{EXT}(Cn)$ (or $\text{EXT}_{fin}(Cn)$) denotes the lattice of all (or all finitary) consequence operations extending Cn .

We write $\text{EXT}(L)$ for $\text{EXT}(Cn_L)$.

The lattice: $\text{NExt}(L)$, $\text{EXT}(Cn)$ with \subseteq

A modal algebra $\mathcal{A} = (A, \wedge, \neg, \Box, \top)$, $\Box(a \wedge b) = \Box a \wedge \Box b$,

$\Box \top = \top$; $\text{Log}(\mathcal{A}) = \{\alpha : v(\alpha) = \top, \text{ for all } v : \text{Var} \rightarrow A\}$,

for a class \mathbb{K} , $\text{Log}(\mathbb{K}) = \bigcap \{\text{Log}(\mathcal{A}) : \mathcal{A} \in \mathbb{K}\}$,

Each modal alg. \mathcal{A} generates a modal consequence oper. $\vec{\mathcal{A}}$:

$\alpha \in \vec{\mathcal{A}}(X)$ iff $(v[X] \subseteq \{\top\} \Rightarrow v(\alpha) = \top, \text{ for each } v : \text{Var} \rightarrow A)$.

A modal algebra $\mathcal{A} = (A, \wedge, \neg, \Box, \top)$, $\Box(a \wedge b) = \Box a \wedge \Box b$,
 $\Box \top = \top$; $\text{Log}(\mathcal{A}) = \{\alpha : v(\alpha) = \top, \text{ for all } v : \text{Var} \rightarrow A\}$,
 for a class \mathbb{K} , $\text{Log}(\mathbb{K}) = \bigcap \{\text{Log}(\mathcal{A}) : \mathcal{A} \in \mathbb{K}\}$,

Each modal alg. \mathcal{A} generates a modal consequence oper. $\vec{\mathcal{A}}$:

$\alpha \in \vec{\mathcal{A}}(X)$ iff $(v[X] \subseteq \{\top\} \Rightarrow v(\alpha) = \top, \text{ for each } v : \text{Var} \rightarrow A)$.

$\alpha \in \vec{\mathcal{A}}(\emptyset)$ iff $\alpha \in \text{Log}(\mathcal{A})$. For a class \mathbb{K} of modal algebras, put

$\alpha \in \vec{\mathbb{K}}(X)$ iff $(\alpha \in \vec{\mathcal{A}}(X), \text{ for each } \mathcal{A} \in \mathbb{K})$;

A modal algebra $\mathcal{A} = (A, \wedge, \neg, \Box, \top)$, $\Box(a \wedge b) = \Box a \wedge \Box b$,
 $\Box \top = \top$; $\text{Log}(\mathcal{A}) = \{\alpha : v(\alpha) = \top, \text{ for all } v : \text{Var} \rightarrow A\}$,
 for a class \mathbb{K} , $\text{Log}(\mathbb{K}) = \bigcap \{\text{Log}(\mathcal{A}) : \mathcal{A} \in \mathbb{K}\}$,

Each modal alg. \mathcal{A} generates a modal consequence oper. $\vec{\mathcal{A}}$:

$\alpha \in \vec{\mathcal{A}}(X)$ iff $(v[X] \subseteq \{\top\} \Rightarrow v(\alpha) = \top, \text{ for each } v : \text{Var} \rightarrow A)$.

$\alpha \in \vec{\mathcal{A}}(\emptyset)$ iff $\alpha \in \text{Log}(\mathcal{A})$. For a class \mathbb{K} of modal algebras, put

$\alpha \in \vec{\mathbb{K}}(X)$ iff $(\alpha \in \vec{\mathcal{A}}(X), \text{ for each } \mathcal{A} \in \mathbb{K})$;

A class \mathbb{L} is *strongly adequate* for Cn , if $Cn = \vec{\mathbb{L}}$.

A modal algebra $\mathcal{A} = (A, \wedge, \neg, \Box, \top)$, $\Box(a \wedge b) = \Box a \wedge \Box b$,
 $\Box \top = \top$; $\text{Log}(\mathcal{A}) = \{\alpha : v(\alpha) = \top, \text{ for all } v : \text{Var} \rightarrow A\}$,
 for a class \mathbb{K} , $\text{Log}(\mathbb{K}) = \bigcap \{\text{Log}(\mathcal{A}) : \mathcal{A} \in \mathbb{K}\}$,

Each modal alg. \mathcal{A} generates a modal consequence oper. $\vec{\mathcal{A}}$:

$\alpha \in \vec{\mathcal{A}}(X)$ iff $(v[X] \subseteq \{\top\} \Rightarrow v(\alpha) = \top, \text{ for each } v : \text{Var} \rightarrow A)$.

$\alpha \in \vec{\mathcal{A}}(\emptyset)$ iff $\alpha \in \text{Log}(\mathcal{A})$. For a class \mathbb{K} of modal algebras, put

$\alpha \in \vec{\mathbb{K}}(X)$ iff $(\alpha \in \vec{\mathcal{A}}(X), \text{ for each } \mathcal{A} \in \mathbb{K})$;

A class \mathbb{L} is *strongly adequate* for Cn , if $Cn = \vec{\mathbb{L}}$.

If, additionally, \mathbb{L} is finite, we say that Cn is *strongly finite*, $Cn \in SF$.

A modal algebra $\mathcal{A} = (A, \wedge, \neg, \Box, \top)$, $\Box(a \wedge b) = \Box a \wedge \Box b$,
 $\Box \top = \top$; $\text{Log}(\mathcal{A}) = \{\alpha : v(\alpha) = \top, \text{ for all } v : \text{Var} \rightarrow A\}$,
 for a class \mathbb{K} , $\text{Log}(\mathbb{K}) = \bigcap \{\text{Log}(\mathcal{A}) : \mathcal{A} \in \mathbb{K}\}$,

Each modal alg. \mathcal{A} generates a modal consequence oper. $\vec{\mathcal{A}}$:

$\alpha \in \vec{\mathcal{A}}(X)$ iff $(v[X] \subseteq \{\top\} \Rightarrow v(\alpha) = \top, \text{ for each } v : \text{Var} \rightarrow A)$.

$\alpha \in \vec{\mathcal{A}}(\emptyset)$ iff $\alpha \in \text{Log}(\mathcal{A})$. For a class \mathbb{K} of modal algebras, put

$\alpha \in \vec{\mathbb{K}}(X)$ iff $(\alpha \in \vec{\mathcal{A}}(X), \text{ for each } \mathcal{A} \in \mathbb{K})$;

A class \mathbb{L} is *strongly adequate* for Cn , if $Cn = \vec{\mathbb{L}}$.

If, additionally, \mathbb{L} is finite, we say that Cn is *strongly finite*, $Cn \in SF$.

a consequence operation Cn is *finitely approximable* ($Cn \in FA$)
 if there is a strongly adequate family of finite algebras for Cn ,
 i.e. $Cn = \bigwedge_i \vec{\mathcal{A}}_i$ where \mathcal{A}_i is finite for each i . If

Theorem

If $\mathcal{A} \in S(\mathcal{B})$, then $\vec{\mathcal{B}} \leq \vec{\mathcal{A}}$.

Theorem

If $\mathcal{A} \in S(\mathcal{B})$, then $\vec{\mathcal{B}} \leq \vec{\mathcal{A}}$.

Theorem

- (i) If $X \in \text{Sat}(\mathcal{A}_t)$ for each $t \in T$, then $\overrightarrow{\mathbf{P}_{t \in T} \mathcal{A}_t}(X) = \bigcap_{t \in T} \vec{\mathcal{A}}_t(X)$.
- (ii) If $X \notin \text{Sat}(\mathcal{A}_t)$ for some $t \in T$, then $\overrightarrow{\mathbf{P}_{t \in T} \mathcal{A}_t}(X) = \text{Fm}$.

Theorem

If $\mathcal{A} \in S(\mathcal{B})$, then $\vec{\mathcal{B}} \leq \vec{\mathcal{A}}$.

Theorem

- (i) If $X \in \text{Sat}(\mathcal{A}_t)$ for each $t \in T$, then $\overrightarrow{\mathbf{P}_{t \in T} \mathcal{A}_t}(X) = \bigcap_{t \in T} \vec{\mathcal{A}}_t(X)$.
- (ii) If $X \notin \text{Sat}(\mathcal{A}_t)$ for some $t \in T$, then $\overrightarrow{\mathbf{P}_{t \in T} \mathcal{A}_t}(X) = \text{Fm}$.

it easily follows that $\vec{\mathbb{K}} \leq \vec{\mathcal{A}}$ if $\mathcal{A} \in SP(\mathbb{K})$. We also have

Theorem

Let \mathbb{K} be a class of modal algebras and Cn be a modal consequence operation such that $\vec{\mathbb{K}} \leq Cn$. Then there is a class $\mathbb{L} \subseteq SP(\mathbb{K})$ such that $Cn = \vec{\mathbb{L}}$.

an element a of an algebra is *open* (*closed*) if $\Box a = a$ ($\Diamond a = a$).

Clopen .

an element a of an algebra is *open* (*closed*) if $\Box a = a$ ($\Diamond a = a$).

Clopen .

\mathcal{A} is a finite *TBA*, extend the set *Var* of prop. var. with fresh variables p_a , for each $a \in A$. The *diagram of* \mathcal{A} is

$$\Delta(\mathcal{A}) = \bigwedge \{ (p_a \rightarrow p_b) \leftrightarrow p_{a \rightarrow b} : a, b \in A \} \wedge (p_{\perp} \leftrightarrow \perp) \wedge \bigwedge \{ \Box p_a \leftrightarrow p_{\Box a} :$$

an element a of an algebra is *open* (*closed*) if $\Box a = a$ ($\Diamond a = a$).

Clopen .

\mathcal{A} is a finite *TBA*, extend the set *Var* of prop. var. with fresh variables p_a , for each $a \in A$. The *diagram of* \mathcal{A} is

$$\Delta(\mathcal{A}) = \bigwedge \{ (p_a \rightarrow p_b) \leftrightarrow p_{a \rightarrow b} : a, b \in A \} \wedge (p_{\perp} \leftrightarrow \perp) \wedge \bigwedge \{ \Box p_a \leftrightarrow p_{\Box a} :$$

A finite *TBA* is subdir. irred, iff it contains the greatest non-unit open element. - called *opremum* of \mathcal{A} ; denoted by $\star_{\mathcal{A}}$ or \star .

an element a of an algebra is *open* (*closed*) if $\Box a = a$ ($\Diamond a = a$).

Clopen .

\mathcal{A} is a finite *TBA*, extend the set *Var* of prop. var. with fresh variables p_a , for each $a \in A$. The *diagram* of \mathcal{A} is

$$\Delta(\mathcal{A}) = \bigwedge \{(p_a \rightarrow p_b) \leftrightarrow p_{a \rightarrow b} : a, b \in A\} \wedge (p_{\perp} \leftrightarrow \perp) \wedge \bigwedge \{\Box p_a \leftrightarrow p_{\Box a} :$$

A finite *TBA* is subdir. irred, iff it contains the greatest non-unit open element. - called *opremum* of \mathcal{A} ; denoted by $\star_{\mathcal{A}}$ or \star .

$\chi(\mathcal{A}) = \Box(\Delta(\mathcal{A})) \rightarrow p_{\star}$ - the characteristic formula of \mathcal{A} .

an element a of an algebra is *open* (*closed*) if $\Box a = a$ ($\Diamond a = a$).

Clopen .

\mathcal{A} is a finite *TBA*, extend the set *Var* of prop. var. with fresh variables p_a , for each $a \in A$. The *diagram* of \mathcal{A} is

$$\Delta(\mathcal{A}) = \bigwedge \{ (p_a \rightarrow p_b) \leftrightarrow p_{a \rightarrow b} : a, b \in A \} \wedge (p_{\perp} \leftrightarrow \perp) \wedge \bigwedge \{ \Box p_a \leftrightarrow p_{\Box a} :$$

A finite *TBA* is subdir. irred, iff it contains the greatest non-unit open element. - called *opremum* of \mathcal{A} ; denoted by $\star_{\mathcal{A}}$ or \star .

$\chi(\mathcal{A}) = \Box(\Delta(\mathcal{A})) \rightarrow p_{\star}$ - the characteristic formula of \mathcal{A} .

Theorem

(i) Let \mathcal{A} be a finite *TBA* and $v : Var \rightarrow A$. Then, for $\alpha \in Fm$:

$$\alpha \leftrightarrow p_{v(\alpha)} \in Cn_{S4}(\{ \Delta(\mathcal{A}) \} \cup \{ p \leftrightarrow p_{v(p)} : p \in Var \}).$$

(ii) Let \mathcal{A} be a finite s.i. *TBA* and \mathcal{B} be any *TBA*. Then

$\chi_{\mathcal{A}} \notin \text{Log}(\mathcal{B}) \iff \mathcal{A}$ is embeddable in some homomorphic image of \mathcal{B} .

A *frame* $\mathfrak{F} = (V, R)$: a set V (worlds), $R \subseteq V \times V$

$\text{Log}(\mathfrak{F}) = \{\alpha : (\mathfrak{F}, x) \Vdash \alpha, \text{ for each } x \in V \text{ and each } \Vdash\}$

A *frame* $\mathfrak{F} = (V, R)$: a set V (worlds), $R \subseteq V \times V$

$\text{Log}(\mathfrak{F}) = \{\alpha : (\mathfrak{F}, x) \Vdash \alpha, \text{ for each } x \in V \text{ and each } \Vdash\}$

Complex alg. $\mathfrak{F}^+ = (P(V), \cap, ', \square, V), \square a = \{x \in V : R(x) \subseteq a\},$

A *frame* $\mathfrak{F} = (V, R)$: a set V (worlds), $R \subseteq V \times V$

$\text{Log}(\mathfrak{F}) = \{\alpha : (\mathfrak{F}, x) \Vdash \alpha, \text{ for each } x \in V \text{ and each } \Vdash\}$

Complex alg. $\mathfrak{F}^+ = (P(V), \cap, ', \square, V), \square a = \{x \in V : R(x) \subseteq a\},$

n-element cluster: $\mathfrak{n} = (V_n, R_n), V_n = \{1, \dots, n\}, R_n = V_n \times V_n.$

$1, 2, 3, \dots, n$ denote 1-, 2-, 3-, ... n-element clusters, respectiv.

$1^+, 2^+, 3^+, \dots, n^+$ their complex algebras,

A *frame* $\mathfrak{F} = (V, R)$: a set V (worlds), $R \subseteq V \times V$

$\text{Log}(\mathfrak{F}) = \{\alpha : (\mathfrak{F}, x) \Vdash \alpha, \text{ for each } x \in V \text{ and each } \Vdash\}$

Complex alg. $\mathfrak{F}^+ = (P(V), \cap, ', \Box, V), \Box a = \{x \in V : R(x) \subseteq a\}$,

n-element cluster: $\mathfrak{n} = (V_n, R_n)$, $V_n = \{1, \dots, n\}$, $R_n = V_n \times V_n$.

$1, 2, 3, \dots, n$ denote 1-, 2-, 3-, ... *n*-element clusters, respectiv.

$1^+, 2^+, 3^+, \dots, n^+$ their complex algebras,

A modal algebra \mathcal{A} is a *Henle algebra* if $\Box a = \perp$ for each $a \neq \top$.

Henle algebras are s.i. (simples) for **S5**.

\mathfrak{n}^+ is the Henle algebra with *n* generators.

Note: $1^+ = \mathbf{2} =^{\text{def}} (\{\perp, \top\}, \min, \neg, \Box)$, with $\Box a = a$.

Frames for $L \in \text{NExt}(\mathbf{S4.3})$ - chains of clusters, lists, covering *r*.
(K.Fine)

A *unifier for a formula α in a logic L* - $\varepsilon(\alpha) \in L$.

A *projective unifier for α in L* is a unifier such that $\varepsilon(\beta) \leftrightarrow \beta \in Cn_L(\alpha)$ for each formula β ;

A *unifier for a formula α in a logic L* - $\varepsilon(\alpha) \in L$.

A *projective unifier for α in L* is a unifier such that $\varepsilon(\beta) \leftrightarrow \beta \in Cn_L(\alpha)$ for each formula β ;

A logic L *enjoys projective unification* if each L -unifiable formula has a projective unifier in L .

A *unifier for a formula α in a logic L* - $\varepsilon(\alpha) \in L$.

A *projective unifier for α in L* is a unifier such that $\varepsilon(\beta) \leftrightarrow \beta \in Cn_L(\alpha)$ for each formula β ;

A logic L *enjoys projective unification* if each L -unifiable formula has a projective unifier in L .

Theorem (see [?])

A modal logic $L \supseteq \mathbf{S4}$ *enjoys projective unification* iff $\Box(\Box y \rightarrow \Box z) \vee \Box(\Box z \rightarrow \Box y) \in L$, iff $\mathbf{S4.3} \subseteq L$.

A *unifier for a formula α in a logic L* - $\varepsilon(\alpha) \in L$.

A *projective unifier for α in L* is a unifier such that $\varepsilon(\beta) \leftrightarrow \beta \in Cn_L(\alpha)$ for each formula β ;

A logic L *enjoys projective unification* if each L -unifiable formula has a projective unifier in L .

Theorem (see [?])

A modal logic $L \supseteq \mathbf{S4}$ *enjoys projective unification iff*

$\Box(\Box y \rightarrow \Box z) \vee \Box(\Box z \rightarrow \Box y) \in L$, *iff* $\mathbf{S4.3} \subseteq L$.

The rule X/β is *admissible* for a modal consequence oper. Cn ,
if $\varepsilon[X] \subseteq Cn(\emptyset) \Rightarrow \varepsilon(\beta) \in Cn(\emptyset)$,
for every substitution ε .

The rule is *derivable* for Cn (is Cn -derivable), if $\beta \in Cn(X)$.

(Almost) Structural Completeness

Cn is *structurally complete* ($Cn \in \text{SCpl}$, see Pogorzelski [?])

each admissible rule for Cn is Cn -derivable

$Cn \in \text{SCpl}_{fin}$ means that each finitary rule which is admissible for Cn is Cn -derivable.

(Almost) Structural Completeness

Cn is *structurally complete* ($Cn \in \text{SCpl}$, see Pogorzelski [?])
each admissible rule for Cn is Cn -derivable

$Cn \in \text{SCpl}_{fin}$ means that each finitary rule which is admissible
for Cn is Cn -derivable.

The structurally complete extension of the logic L , Cn_L^{Scpl} , is the
extension of Cn_L with all L -admissible rules; it is the greatest
modal consequence operation Cn such that $Cn(\emptyset) = L$.

(Almost) Structural Completeness

Cn is *structurally complete* ($Cn \in \text{SCpl}$, see Pogorzelski [?])
each admissible rule for Cn is Cn -derivable

$Cn \in \text{SCpl}_{fin}$ means that each finitary rule which is admissible
for Cn is Cn -derivable.

The structurally complete extension of the logic L , Cn_L^{Scpl} , is the
extension of Cn_L with all L -admissible rules; it is the greatest
modal consequence operation Cn such that $Cn(\emptyset) = L$.

X/β is *passive* in L (see Rybakov [?]), if X is not unifiable in L .

(Almost) Structural Completeness

Cn is *structurally complete* ($Cn \in \text{SCpl}$, see Pogorzelski [?])
each admissible rule for Cn is Cn -derivable

$Cn \in \text{SCpl}_{fin}$ means that each finitary rule which is admissible
for Cn is Cn -derivable.

The structurally complete extension of the logic L , Cn_L^{Scpl} , is the
extension of Cn_L with all L -admissible rules; it is the greatest
modal consequence operation Cn such that $Cn(\emptyset) = L$.

X/β is *passive* in L (see Rybakov [?]), if X is not unifiable in L .

S5, and many others **S4.3**,... contrary to **S4**, is not structurally
complete only because the following rule is passive, hence,
admissible, but not derivable:

$$P_2 : \frac{\Diamond \alpha \wedge \Diamond \sim \alpha}{\beta}$$

(Almost) Structural Completeness

Cn is *almost structurally complete*, $Cn \in \text{ASCpl}$; every admissible rule for Cn , which is not passive, is derivable for Cn .
 $Cn \in \text{ASCpl}_{fin}$ if we restrict to finitary rules only.

(Almost) Structural Completeness

Cn is *almost structurally complete*, $Cn \in \text{ASCpl}$; every admissible rule for Cn , which is not passive, is derivable for Cn . $Cn \in \text{ASCpl}_{fin}$ if we restrict to finitary rules only.

Corollary (W.D. P.W. 2011)

$Cn \in \text{ASCpl}_{fin}$, for every modal consequence operation Cn extending $S4.3$.

(Almost) Structural Completeness

Cn is *almost structurally complete*, $Cn \in \text{ASCpl}$; every admissible rule for Cn , which is not passive, is derivable for Cn . $Cn \in \text{ASCpl}_{fin}$ if we restrict to finitary rules only.

Corollary (W.D. P.W. 2011)

$Cn \in \text{ASCpl}_{fin}$, for every modal consequence operation Cn extending $S4.3$.

There are four variants of structural completeness: SCpl , SCpl_{fin} , ASCpl and ASCpl_{fin} . (different, sometimes collapse).

(Almost) Structural Completeness

Cn is *almost structurally complete*, $Cn \in \text{ASCpl}$; every admissible rule for Cn , which is not passive, is derivable for Cn .
 $Cn \in \text{ASCpl}_{fin}$ if we restrict to finitary rules only.

Corollary (W.D. P.W. 2011)

$Cn \in \text{ASCpl}_{fin}$, for every modal consequence operation Cn extending $S4.3$.

There are four variants of structural completeness: SCpl , SCpl_{fin} , ASCpl and ASCpl_{fin} . (different, sometimes collapse).

Theorem

If Cn is modal consequence operation extending $S4.3M$, where $(M) : \Box\Diamond\alpha \rightarrow \Diamond\Box\alpha$ (McKinsey's axiom), then

$$Cn \in \text{ASCpl} \iff Cn \in \text{SCpl}$$

$SCpl \neq SCpl_{fin}$ and $ASCpl \neq ASCpl_{fin}$, more exactly..

$S4.3 \in ASCpl_{fin} \setminus ASCpl$ and $S4.3M \in SCpl_{fin} \setminus SCpl$

$\text{SCpl} \neq \text{SCpl}_{fin}$ and $\text{ASCpl} \neq \text{ASCpl}_{fin}$, more exactly..

$S4.3 \in \text{ASCpl}_{fin} \setminus \text{ASCpl}$ and $S4.3M \in \text{SCpl}_{fin} \setminus \text{SCpl}$ The rules

$$\varrho : \frac{\{\sim \Box(p_i \leftrightarrow p_j) : 0 < i < j\}}{\perp} \quad \varrho' : \frac{\{\Box(p_i \leftrightarrow p_j) \rightarrow p_0 : 0 < i < j\}}{p_0}$$

$\text{SCpl} \neq \text{SCpl}_{fin}$ and $\text{ASCpl} \neq \text{ASCpl}_{fin}$, more exactly:.

$S4.3 \in \text{ASCpl}_{fin} \setminus \text{ASCpl}$ and $S4.3M \in \text{SCpl}_{fin} \setminus \text{SCpl}$ The rules

$$\varrho : \frac{\{\sim \Box(p_i \leftrightarrow p_j) : 0 < i < j\}}{\perp} \quad \varrho' : \frac{\{\Box(p_i \leftrightarrow p_j) \rightarrow p_0 : 0 < i < j\}}{p_0}$$

are valid in every finite TBA \mathcal{A} i.e. derivable for the cons.op. $\vec{\mathcal{A}}$
Hence, by FMP ϱ and ϱ' admissible in $S4.3$ and all extensions.

$\text{SCpl} \neq \text{SCpl}_{fin}$ and $\text{ASCpl} \neq \text{ASCpl}_{fin}$, more exactly:.

$S4.3 \in \text{ASCpl}_{fin} \setminus \text{ASCpl}$ and $S4.3M \in \text{SCpl}_{fin} \setminus \text{SCpl}$ The rules

$$\varrho : \frac{\{\sim \Box(p_i \leftrightarrow p_j) : 0 < i < j\}}{\perp} \quad \varrho' : \frac{\{\Box(p_i \leftrightarrow p_j) \rightarrow p_0 : 0 < i < j\}}{p_0}$$

are valid in every finite TBA \mathcal{A} i.e. derivable for the cons.op. \vec{A}

Hence, by FMP ϱ and ϱ' admissible in $S4.3$ and all extensions.

But,

ϱ' and ϱ are not derivable for $S4.3$ (nor $S4.3M$)

p_0 cannot be deduced, in $S4.3$, from any finite subset of

$\{\Box(p_i \leftrightarrow p_j) \rightarrow p_0 : 0 < i < j\}$.

OBSERVATION: attempts to extend the concept of projective unifier to infinite sets of formulas fails

Theorem

- (i) *Each finitary consequence operation $Cn \geq Cn_{S4.3}$ has a finite basis (of finitary rules) and $Cn = C_{fin}$ for some $C \in FA$.*
- (ii) *The lattice $(EXT_{fin}(\mathbf{S4.3}), \leq)$ is distributive.*

Theorem

- (i) *Each finitary consequence operation $Cn \geq Cn_{S4.3}$ has a finite basis (of finitary rules) and $Cn = C_{fin}$ for some $C \in FA$.*
- (ii) *The lattice $(EXT_{fin}(\mathbf{S4.3}), \leq)$ is distributive.*

By Coroll. each finitary $Cn \geq Cn_{S4.3}$ is an extension of Cn_L , for some $L \in \mathbf{S4.3}$, with passive rules only

Theorem

- (i) *Each finitary consequence operation $Cn \geq Cn_{S4.3}$ has a finite basis (of finitary rules) and $Cn = C_{fin}$ for some $C \in FA$.*
- (ii) *The lattice $(EXT_{fin}(\mathbf{S4.3}), \leq)$ is distributive.*

By Coroll. each finitary $Cn \geq Cn_{S4.3}$ is an extension of Cn_L , for some $L \in \mathbf{S4.3}$, with passive rules only

Fix $n \geq 0$, var.. p_1, \dots, p_n generate a sublanguage of Fm with 2^n (boolean) atoms and they can be written as

Theorem

- (i) Each finitary consequence operation $Cn \geq Cn_{S4.3}$ has a finite basis (of finitary rules) and $Cn = C_{fin}$ for some $C \in FA$.
- (ii) The lattice $(EXT_{fin}(\mathbf{S4.3}), \leq)$ is distributive.

By Coroll. each finitary $Cn \geq Cn_{S4.3}$ is an extension of Cn_L , for some $L \in \mathbf{S4.3}$, with passive rules only

Fix $n \geq 0$, var.. p_1, \dots, p_n generate a sublanguage of Fm with 2^n (boolean) atoms and they can be written as

$$p_1^{\sigma(1)} \wedge \dots \wedge p_n^{\sigma(n)}$$

where $\sigma: \{1, \dots, n\} \rightarrow \{0, 1\}$, and $p^0 = p$, and $p^1 = \sim p$.
Suppose that all these atoms are listed as: $\theta_1, \dots, \theta_{2^n}$.

Corollary

Each $C_n \in \text{EXT}_{fin}(\mathbf{S4.3})$ is an extension of C_{n_L} , for some $L \in \text{NExt}(\mathbf{S4.3})$, $n \geq 0$, with a finite number of passive rules having the form

Corollary

Each $C_n \in \text{EXT}_{\text{fin}}(\mathbf{S4.3})$ is an extension of C_{n_L} , for some $L \in \text{NExt}(\mathbf{S4.3})$, $n \geq 0$, with a finite number of passive rules having the form

$$\frac{\Diamond\theta_1 \wedge \dots \wedge \Diamond\theta_s}{\alpha}$$

where $2 \leq s \leq 2^n$ and $\text{Var}(\alpha) \cap \{p_1, \dots, p_n\} = \emptyset$.

Corollary

Each $C_n \in \text{EXT}_{\text{fin}}(\mathbf{S4.3})$ is an extension of C_{n_L} , for some $L \in \text{NExt}(\mathbf{S4.3})$, $n \geq 0$, with a finite number of passive rules having the form

$$\frac{\Diamond\theta_1 \wedge \cdots \wedge \Diamond\theta_s}{\alpha}$$

where $2 \leq s \leq 2^n$ and $\text{Var}(\alpha) \cap \{p_1, \dots, p_n\} = \emptyset$.

The rule $\Diamond\theta_1 \wedge \cdots \wedge \Diamond\theta_s / \alpha$ is valid in the complex algebra of a cluster of $\leq s - 1$ elements, it is not valid in the cluster s of s elements.

Theorem

If $Cn \in \text{EXT}(\mathbf{S4.3})$ and $\text{Log}(\mathbb{K}) = Cn(\emptyset)$ for some class $\mathbb{K} \subseteq \text{Alg}(\mathbf{S4.3})_{\text{si fin}}$, then one can find $\mathbb{K}_1 \supseteq \cdots \supseteq \mathbb{K}_m$ such that

Theorem

If $Cn \in \text{EXT}(\mathbf{S4.3})$ and $\text{Log}(\mathbb{K}) = Cn(\emptyset)$ for some class $\mathbb{K} \subseteq \text{Alg}(\mathbf{S4.3})_{\text{si fin}}$, then one can find $\mathbb{K}_1 \supseteq \cdots \supseteq \mathbb{K}_m$ such that $S(\mathbb{K}) = \mathbb{K}_1$, and $\mathbb{K}_{i+1} = S(\mathbb{K}_i)$ for $i = 1, \dots, m$, and $Cn =_{\text{fin}} \overrightarrow{\mathbb{L}}$, where

Theorem

If $Cn \in \text{EXT}(\mathbf{S4.3})$ and $\text{Log}(\mathbb{K}) = Cn(\emptyset)$ for some class $\mathbb{K} \subseteq \text{Alg}(\mathbf{S4.3})_{\text{si fin}}$, then one can find $\mathbb{K}_1 \supseteq \dots \supseteq \mathbb{K}_m$ such that $S(\mathbb{K}) = \mathbb{K}_1$, and $\mathbb{K}_{i+1} = S(\mathbb{K}_i)$ for $i = 1, \dots, m$, and $Cn =_{\text{fin}} \overrightarrow{\mathbb{L}}$, where

$$\mathbb{L} = \mathbb{K}_m \cup \left((\mathbb{K}_{m-1} \setminus \mathbb{K}_m) \times (\mathbf{m} - \mathbf{1})^+ \right) \cup \dots \cup \left((\mathbb{K}_1 \setminus \mathbb{K}_2) \times \mathbf{1}^+ \right).$$

Each quasivariety of $S4.3$ -algebras is generated by a class of finite $S4.3$ -algebras of the form:

- s.i. algebra, or
- the direct product of a s.i. algebra and a Henle algebra.

Theorem

If $Cn \in \text{EXT}(\mathbf{S4.3})$ and $\text{Log}(\mathbb{K}) = Cn(\emptyset)$ for some class $\mathbb{K} \subseteq \text{Alg}(\mathbf{S4.3})_{\text{si fin}}$, then one can find $\mathbb{K}_1 \supseteq \dots \supseteq \mathbb{K}_m$ such that $S(\mathbb{K}) = \mathbb{K}_1$, and $\mathbb{K}_{i+1} = S(\mathbb{K}_i)$ for $i = 1, \dots, m$, and $Cn =_{\text{fin}} \overrightarrow{\mathbb{L}}$, where

$$\mathbb{L} = \mathbb{K}_m \cup \left((\mathbb{K}_{m-1} \setminus \mathbb{K}_m) \times (\mathfrak{m} - 1)^+ \right) \cup \dots \cup \left((\mathbb{K}_1 \setminus \mathbb{K}_2) \times 1^+ \right).$$

Each quasivariety of $S4.3$ -algebras is generated by a class of finite $S4.3$ -algebras of the form:

- s.i. algebra, or
- the direct product of a s.i. algebra and a Henle algebra.

With all assumptions/ notations of the above Theorem we have

Theorem

If $\overrightarrow{\mathbb{K}} \leq Cn \leq \overrightarrow{\mathbb{L}}$ and $Cn =_{\text{fin}} \overrightarrow{\mathbb{L}}$, then $Cn = \overrightarrow{\mathbb{L}}$.

Theorem

If a logic $L \in \text{NExt}(\mathbf{S4.3})$ is tabular, then $\text{EXT}(L) \subseteq SF$ and consequently $\text{EXT}(L) = \text{EXT}_{fin}(L)$.

Theorem

If a logic $L \in \text{NExt}(\mathbf{S4.3})$ is tabular, then $\text{EXT}(L) \subseteq \text{SF}$ and consequently $\text{EXT}(L) = \text{EXT}_{\text{fin}}(L)$.

Theorem

A finitary modal consequence operation $C_n \in \text{EXT}(\mathbf{S4.3})$ is finitely approximable, $C_n \in \text{FA}$ iff C_n is strongly finite, $C_n \in \text{SF}$; i.e. $\text{FA} \cap \text{Fin} = \text{SF}$.

Theorem

If a logic $L \in \text{NExt}(\mathbf{S4.3})$ is tabular, then $\text{EXT}(L) \subseteq \text{SF}$ and consequently $\text{EXT}(L) = \text{EXT}_{\text{fin}}(L)$.

Theorem

A finitary modal consequence operation $C_n \in \text{EXT}(\mathbf{S4.3})$ is finitely approximable, $C_n \in \text{FA}$ iff C_n is strongly finite, $C_n \in \text{SF}$; i.e. $\text{FA} \cap \text{Fin} = \text{SF}$.

Theorem

A modal consequence operation $C_n \in \text{EXT}(\mathbf{S4.3})$ is finitely approximable iff C_n is almost structurally complete; i.e. $\text{FA} = \text{ASCpl}$.

Corollary

*If Cn is a finitary consequence operation extending **S4.3**, then Cn is almost structurally complete iff Cn is strongly finite.*

Corollary

*Let $L \in \text{NExt}(\mathbf{S4.3})$ and \mathbb{L} be an adequate class of finite **S4.3**-algebras for L . Then $\overrightarrow{\mathbb{L} \times \mathbf{2}}$ is the structurally complete extension of Cn_L , i.e. $Cn_L^{SCpl} = \overrightarrow{\mathbb{L} \times \mathbf{2}}$.*

Corollary

*If Cn is a finitary consequence operation extending **S4.3**, then Cn is almost structurally complete iff Cn is strongly finite.*

Corollary

*Let $L \in \text{NExt}(\mathbf{S4.3})$ and \mathbb{L} be an adequate class of finite **S4.3**-algebras for L . Then $\overrightarrow{\mathbb{L} \times \mathbf{2}}$ is the structurally complete extension of Cn_L , i.e. $Cn_L^{SCpl} = \overrightarrow{\mathbb{L} \times \mathbf{2}}$.*

The lattices $ASCpl(\mathbf{S4.3})$ and $EXT_{fin}(\mathbf{S4.3})$ are isomorphic, Similarly $ASCpl(\mathbf{L})$ and $EXT_{fin}(\mathbf{L})$.
 $Cn \mapsto Cn_{fin}$ and $Cn \mapsto Cn^\Sigma$

Corollary

*If Cn is a finitary consequence operation extending **S4.3**, then Cn is almost structurally complete iff Cn is strongly finite.*


Corollary


*Let $L \in \text{NExt}(\mathbf{S4.3})$ and \mathbb{L} be an adequate class of finite **S4.3**-algebras for L . Then $\overrightarrow{\mathbb{L} \times \mathbf{2}}$ is the structurally complete extension of Cn_L , i.e. $Cn_L^{SCpl} = \overrightarrow{\mathbb{L} \times \mathbf{2}}$.*

The lattices $ASCpl(\mathbf{S4.3})$ and $EXT_{fin}(\mathbf{S4.3})$ are isomorphic, Similarly $ASCpl(\mathbf{L})$ and $EXT_{fin}(\mathbf{L})$.
 $Cn \mapsto Cn_{fin}$ and $Cn \mapsto Cn^\Sigma$

Σ an infinite basis for infinitary admissible non-passive rules - in Part II.

The lattice $ASCpl(\mathbf{S5})$, Not all $Cn \in EXT_{fin}(\mathbf{S5})$ have a single str. ad. matrix, e.g. $(1^+ \times 3^+) \cap 2^+ \cdot$:





ASCpl(S4.5)

a Logic - a variety $\mathbb{K} = \text{HSP}\mathbb{K}$, a finitary (modal) consequence operation - a quasivariety of (modal) algebras, $\mathbb{K} = \text{ISPP}_U\mathbb{K}$, an arbitrary, (modal) consequence operation - a prevariety of (modal) algebras, $\mathbb{K} = \text{ISP}\mathbb{K}$.

a set of *quasiequations*, that is, by *equational implications* of the form

$$(ei) \quad \left(\bigwedge_{i \in I} p_i(\bar{x}) = q_i(\bar{x}) \right) \Rightarrow p(\bar{x}) = q(\bar{x})$$

a prevariety, $\mathbb{K} = \text{ISP}\mathbb{K}$ is defined by a set of *generalized equational implications*, (ei) , where the set I of indices can be infinite;

Let $F_{\mathbb{K}}$ be the ω -generated free algebra in \mathbb{K} . Admissibility-validity of the (generalized) equational implication (ei) in the free algebra $F_{\mathbb{K}}$,

a generalized equational implication (ei) is *passive* if $\left(\bigwedge_{i \in I} p_i(\bar{x}) = q_i(\bar{x}) \right)$ is not satisfiable in $F_{\mathbb{K}}$.

A prevariety \mathbb{K} is *(almost) structurally complete* iff every generalized equational implication (ei) which is valid in $F_{\mathbb{K}}$ (and is not passive), is valid in the whole prevariety \mathbb{K} .

Theorem

For a prevariety \mathbb{K} of S4.3-algebras TFCE:

- (i) \mathbb{K} is almost structurally complete
- (ii) \mathbb{K} is generated by its finite members.
- (iii) every generalized equational implication (ei) which is valid in $F_{\mathbb{K}}$ and its premises are satisfiable in $F_{\mathbb{K}}$, is valid in the whole prevariety \mathbb{K} .

Let Σ be a set of all increasing functions $f : \mathbb{N} \rightarrow \mathbb{N}$. The following equational implications $(ei)_f$, for $f \in \Sigma$:

$\bigwedge_{n>0} \{ [\bigwedge_{n < j \leq f(n)} \bigvee_{0 < i \leq n} \Box(p_i \leftrightarrow p_j)] \rightarrow p_0 \} = 1 \Rightarrow p_0 = 1$
 form a basis for admissible generalized equational implications in \mathbb{K} .