

# On MV-algebras with convexity operators

Serafina Lapenta,  
University of Salerno

ManyVal'15 - Les Diablerets

11 December 2015

joint work with Tommaso Flaminio

## Driving motivation

- ▶ Convex combinations appear in any area of mathematics as well as in other sciences,

## Driving motivation

- ▶ Convex combinations appear in any area of mathematics as well as in other sciences,
- ▶ MV-algebras are a good candidate to define in abstract way:  
Di Nola and Leuştean - MV-algebras with scalar product,

## Driving motivation

- ▶ Convex combinations appear in any area of mathematics as well as in other sciences,
- ▶ MV-algebras are a good candidate to define in abstract way:  
Di Nola and Leuştean - MV-algebras with scalar product,
- ▶ characterization of states on MV-algebras:  
states are natural generalization of probability on boolean algebras  
and they are convex combinations of homomorphisms.

# MV-algebras

# MV-algebras

- ▶ Algebraic counterpart of Łukasiewicz logic,

# MV-algebras

- ▶ Algebraic counterpart of Łukasiewicz logic,
- ▶ Unit interval of lattice ordered groups with strong unit,

# MV-algebras

- ▶ Algebraic counterpart of Łukasiewicz logic,
- ▶ Unit interval of lattice ordered groups with strong unit,
- ▶ A generalization of boolean algebras.



## The standard model

$$[0, 1]_{MV} = ([0, 1], \oplus, *, 0)$$

►  $x \oplus y = \min(x + y, 1), x^* = 1 - x;$

## The standard model

$$[0, 1]_{MV} = ([0, 1], \oplus, *, 0)$$

- ▶  $x \oplus y = \min(x + y, 1)$ ,  $x^* = 1 - x$ ;
- ▶ is the set of truth values for Łukasiewicz logic;

# The standard model

$$[0, 1]_{MV} = ([0, 1], \oplus, *, 0)$$

- ▶  $x \oplus y = \min(x + y, 1)$ ,  $x^* = 1 - x$ ;
- ▶ is the set of truth values for Łukasiewicz logic;
- ▶ it is closed for product between real numbers!

## MV-algebras

- ▶ MV-algebras are structure  $(A, \oplus, *, 0)$ ,

## MV-algebras

- ▶ MV-algebras are structure  $(A, \oplus, *, 0)$ ,
- ▶ they are the variety generated by  $[0, 1]_{MV}$ ,

## MV-algebras

- ▶ MV-algebras are structure  $(A, \oplus, *, 0)$ ,
- ▶ they are the variety generated by  $[0, 1]_{MV}$ ,
- ▶  $A$  is an MV-algebra iff  $A \in HSP([0, 1]_{MV})$ .

## MV-algebras

- ▶ MV-algebras are structure  $(A, \oplus, *, 0)$ ,
- ▶ they are the variety generated by  $[0, 1]_{MV}$ ,
- ▶  $A$  is an MV-algebra iff  $A \in HSP([0, 1]_{MV})$ .

## Natural lattice order

if  $x \rightarrow y = x^* \oplus y$  then  $x \leq y$  iff  $x \rightarrow y = 1$

## MV-algebras

- ▶ MV-algebras are structure  $(A, \oplus, *, 0)$ ,
- ▶ they are the variety generated by  $[0, 1]_{MV}$ ,
- ▶  $A$  is an MV-algebra iff  $A \in HSP([0, 1]_{MV})$ .

## Natural lattice order

if  $x \rightarrow y = x^* \oplus y$  then  $x \leq y$  iff  $x \rightarrow y = 1$

## Relation with boolean algebras

$B(A) = \{x \mid x \oplus x = x\}$  is a boolean algebra

$B(A)$  is called boolean skeleton of  $A$ .



## Homomorphisms and ideals

- ▶  $A, B$  MV-algebras,  $h : A \rightarrow B$  such that

$$h(0) = 0,$$

$$h(x \oplus y) = h(x) \oplus h(y),$$

$$h(x^*) = h(x)^*.$$

- ▶  $I \subseteq A$ , ideal if

$$0 \in I,$$

$$x, y \in I \text{ implies } x \oplus y \in I,$$

$$x \in I, y \leq x \text{ then } y \in I.$$

# MV-algebras via groups

# MV-algebras via groups


$(G, +, 0, \leq)$  is  $\ell$ -group

- ▶ if  $(G, +, 0)$  group,
- ▶  $(G, \leq)$  lattice,
- ▶  $x \leq y$  implies  $x + z \leq y + z$  for any  $x, y, z \in G$ .


$\ell u$ -groups

$u \in G$  is a **strong unit**:  $u \geq 0$ , for any  $x \in G$  there is  $n \geq 1$  s.t.  $x \leq nu$ .

An **abelian**  $\ell$ -group with strong unit is an  $\ell u$ -group.




For any  $x \geq 0$  in  $G$  there are  $x_1, \dots, x_n \in [0, u]$  s.t.  $x = x_1 + \dots + x_n$ .



For any  $x \geq 0$  in  $G$  there are  $x_1, \dots, x_n \in [0, u]$  s.t.  $x = x_1 + \dots + x_n$ .

$(G, u)$   $\ell u$ -group,  $x, y \in G$

$$x \oplus y = (x +_G y) \wedge u, \quad x^* = u -_G x$$




For any  $x \geq 0$  in  $G$  there are  $x_1, \dots, x_n \in [0, u]$  s.t.  $x = x_1 + \dots + x_n$ .

$(G, u)$   $\ell u$ -group,  $x, y \in G$

$$x \oplus y = (x +_G y) \wedge u, \quad x^* = u -_G x$$

$([0, u]_G, \oplus, *, 0)$  is an MV-algebra



For any  $x \geq 0$  in  $G$  there are  $x_1, \dots, x_n \in [0, u]$  s.t.  $x = x_1 + \dots + x_n$ .

$(G, u)$   $\ell u$ -group,  $x, y \in G$

$$x \oplus y = (x +_G y) \wedge u, \quad x^* = u -_G x$$

$([0, u]_G, \oplus, *, 0)$  is an MV-algebra

Categorical equivalence. Mundici, 1986

For any MV-algebra  $A$  there exists a  $\ell u$ -group  $(G, u)$  such that  $A \simeq [0, u]_G$ .

The category of MV-algebras is equivalent with the category of Abelian lattice-ordered groups with strong unit with unit preserving homomorphism.

# MV-algebras with product



# MV-algebras with product

MV-algebras  $P$  endowed with a  
“ring-like” product.

## MV-algebras with product

MV-algebras  $P$  endowed with a  
“ring-like” product.

PMV-algebras (Montagna,  
2000)

# MV-algebras with product

MV-algebras  $P$  endowed with a  
“ring-like” product.

PMV-algebras (Montagna,  
2000)

- ▶ they form a variety,
- ▶ categorical equivalence with  $\ell U$ -rings,
- ▶  $[0, 1]_{PMV}$  generates a proper sub-variety.

## MV-algebras with product

MV-algebras  $P$  endowed with a “ring-like” product.

PMV-algebras (Montagna, 2000)

- ▶ they form a variety,
- ▶ categorical equivalence with  $\ell U$ -rings,
- ▶  $[0, 1]_{PMV}$  generates a proper sub-variety.

MV-algebras  $R$  endowed with a scalar multiplication with scalars from  $[0, 1]$ .

## MV-algebras with product

MV-algebras  $P$  endowed with a “ring-like” product.

PMV-algebras (Montagna, 2000)

- ▶ they form a variety,
- ▶ categorical equivalence with  $\ell U$ -rings,
- ▶  $[0, 1]_{PMV}$  generates a proper sub-variety.

MV-algebras  $R$  endowed with a scalar multiplication with scalars from  $[0, 1]$ .

Riesz MV-algebras (Di Nola, Leuştean, 2014)

## MV-algebras with product

MV-algebras  $P$  endowed with a “ring-like” product.

PMV-algebras (Montagna, 2000)

- ▶ they form a variety,
- ▶ categorical equivalence with  $\ell U$ -rings,
- ▶  $[0, 1]_{PMV}$  generates a proper sub-variety.

MV-algebras  $R$  endowed with a scalar multiplication with scalars from  $[0, 1]$ .

Riesz MV-algebras (Di Nola, Leuştean, 2014)

- ▶ they form a variety,
- ▶ categorical equivalence with Riesz Spaces (vector lattices) with strong unit,
- ▶  $RMV = HSP([0, 1]_{RMV})$ .

# The partial sum

## The partial sum

If  $x \odot y = 0$ , then  $x + y = x \oplus y$



## The partial sum

If  $x \odot y = 0$ , then  $x + y = x \oplus y$

$+$  is the sum in the corresponding  $\ell u$ -group.

## The partial sum

If  $x \odot y = 0$ , then  $x + y = x \oplus y$

$+$  is the sum in the corresponding  $\ell u$ -group.

For example

In  $[0, 1]$   $x + y$  is defined when  $x +_{\mathbb{R}} y \leq 1$ !

# Probability on MV-algebras: states

## Probability on MV-algebras: states

A an MV-algebra,  $s : A \rightarrow [0, 1]$  such that

- ▶  $s(1) = 1$ ,
- ▶  $s(x \oplus y) = s(x) + s(y)$  when  $x \odot y = 0$ .

## Probability on MV-algebras: states

A an MV-algebra,  $s : A \rightarrow [0, 1]$  such that

- ▶  $s(1) = 1$ ,
- ▶  $s(x \oplus y) = s(x) + s(y)$  when  $x \odot y = 0$ .



Mundici D., *Averaging the Truth-Value in Łukasiewicz Logic*, *Studia Logica* 55 (1995) 113-127.

## Probability on MV-algebras: states

A an MV-algebra,  $s : A \rightarrow [0, 1]$  such that

- ▶  $s(1) = 1$ ,
- ▶  $s(x \oplus y) = s(x) + s(y)$  when  $x \odot y = 0$ .



Mundici D., *Averaging the Truth-Value in Łukasiewicz Logic*, *Studia Logica* 55 (1995) 113-127.

On the boolean skeleton of an MV-algebra we obtain a *finitely additive* probability measure,

## Probability on MV-algebras: states

A an MV-algebra,  $s : A \rightarrow [0, 1]$  such that

- ▶  $s(1) = 1$ ,
- ▶  $s(x \oplus y) = s(x) + s(y)$  when  $x \odot y = 0$ .



Mundici D., *Averaging the Truth-Value in Łukasiewicz Logic*, *Studia Logica* 55 (1995) 113-127.

On the boolean skeleton of an MV-algebra we obtain a *finitely additive* probability measure,

If  $A = [0, 1]^k$ , then  $S(A) = co(H(A))!$

## Probability on MV-algebras: states

$A$  an MV-algebra,  $s : A \rightarrow [0, 1]$  such that

- ▶  $s(1) = 1$ ,
- ▶  $s(x \oplus y) = s(x) + s(y)$  when  $x \odot y = 0$ .



Mundici D., *Averaging the Truth-Value in Łukasiewicz Logic*, *Studia Logica* 55 (1995) 113-127.

On the boolean skeleton of an MV-algebra we obtain a *finitely additive* probability measure,

If  $A = [0, 1]^k$ , then  $S(A) = co(H(A))!$

states are convex combinations of homomorphism



# Convex combinations in MV-algebras

# Convex combinations in MV-algebras

We want to find a set of axioms that captures

for any  $x, y, \alpha \in [0, 1]$

$$\alpha x + (1 - \alpha)y$$

# MV-algebras with convexity operators (CMV-algebras)

## MV-algebras with convexity operators (CMV-algebras)

MV-algebra **A** together with a family of binary operators

$\mathcal{C} = \{cc_\alpha\}_{\alpha \in [0,1]}$  on  $A$  such that

(C1)  $cc_0(x, y) = y$ ;

## MV-algebras with convexity operators (CMV-algebras)

MV-algebra  $\mathbf{A}$  together with a family of binary operators

$\mathcal{C} = \{cc_\alpha\}_{\alpha \in [0,1]}$  on  $A$  such that

(C1)  $cc_0(x, y) = y$ ;

(C2)  $cc_\alpha(x, y) = cc_{1-\alpha}(y, x)$ ;

## MV-algebras with convexity operators (CMV-algebras)

MV-algebra  $\mathbf{A}$  together with a family of binary operators

$\mathcal{C} = \{cc_\alpha\}_{\alpha \in [0,1]}$  on  $A$  such that

(C1)  $cc_0(x, y) = y$ ;

(C2)  $cc_\alpha(x, y) = cc_{1-\alpha}(y, x)$ ;

(C3)  $cc_\alpha(x, x) = x$ ;

## MV-algebras with convexity operators (CMV-algebras)

MV-algebra **A** together with a family of binary operators

$\mathcal{C} = \{cc_\alpha\}_{\alpha \in [0,1]}$  on **A** such that

(C1)  $cc_0(x, y) = y;$

(C2)  $cc_\alpha(x, y) = cc_{1-\alpha}(y, x);$

(C3)  $cc_\alpha(x, x) = x;$

(C4)  $cc_\alpha(cc_\beta(x, y), z) = cc_{\alpha\beta}(x, cc_\gamma(y, z)),$  with  $\gamma$  arbitrary if  $\alpha = \beta = 1$   
and  $\gamma = \frac{\alpha(1-\beta)}{1-\alpha\beta}$  otherwise;

## MV-algebras with convexity operators (CMV-algebras)

MV-algebra  $\mathbf{A}$  together with a family of binary operators

$\mathcal{C} = \{cc_\alpha\}_{\alpha \in [0,1]}$  on  $A$  such that

- (C1)  $cc_0(x, y) = y$ ;
- (C2)  $cc_\alpha(x, y) = cc_{1-\alpha}(y, x)$ ;
- (C3)  $cc_\alpha(x, x) = x$ ;
- (C4)  $cc_\alpha(cc_\beta(x, y), z) = cc_{\alpha\beta}(x, cc_\gamma(y, z))$ , with  $\gamma$  arbitrary if  $\alpha = \beta = 1$  and  $\gamma = \frac{\alpha(1-\beta)}{1-\alpha\beta}$  otherwise;
- (C5) For all  $\alpha, \beta \in [0, 1]$  such that  $\alpha + \beta \leq 1$ ,  $cc_\alpha(x, 0) + cc_\beta(x, 0)$  is defined and it coincides with  $cc_{\alpha+\beta}(x, 0)$ ;



## MV-algebras with convexity operators (CMV-algebras)

MV-algebra  $\mathbf{A}$  together with a family of binary operators

$\mathcal{C} = \{cc_\alpha\}_{\alpha \in [0,1]}$  on  $A$  such that

- (C1)  $cc_0(x, y) = y$ ;
- (C2)  $cc_\alpha(x, y) = cc_{1-\alpha}(y, x)$ ;
- (C3)  $cc_\alpha(x, x) = x$ ;
- (C4)  $cc_\alpha(cc_\beta(x, y), z) = cc_{\alpha\beta}(x, cc_\gamma(y, z))$ , with  $\gamma$  arbitrary if  $\alpha = \beta = 1$  and  $\gamma = \frac{\alpha(1-\beta)}{1-\alpha\beta}$  otherwise;
- (C5) For all  $\alpha, \beta \in [0, 1]$  such that  $\alpha + \beta \leq 1$ ,  $cc_\alpha(x, 0) + cc_\beta(x, 0)$  is defined and it coincides with  $cc_{\alpha+\beta}(x, 0)$ ;
- (C6) If  $x + x'$  and  $y + y'$  are defined, so is  $cc_\alpha(x + y) + cc_\alpha(x', y')$  and it coincides with  $cc_\alpha(x + x', y + y')$ ;

## MV-algebras with convexity operators (CMV-algebras)

MV-algebra  $\mathbf{A}$  together with a family of binary operators

$\mathcal{C} = \{cc_\alpha\}_{\alpha \in [0,1]}$  on  $A$  such that

- (C1)  $cc_0(x, y) = y$ ;
- (C2)  $cc_\alpha(x, y) = cc_{1-\alpha}(y, x)$ ;
- (C3)  $cc_\alpha(x, x) = x$ ;
- (C4)  $cc_\alpha(cc_\beta(x, y), z) = cc_{\alpha\beta}(x, cc_\gamma(y, z))$ , with  $\gamma$  arbitrary if  $\alpha = \beta = 1$  and  $\gamma = \frac{\alpha(1-\beta)}{1-\alpha\beta}$  otherwise;
- (C5) For all  $\alpha, \beta \in [0, 1]$  such that  $\alpha + \beta \leq 1$ ,  $cc_\alpha(x, 0) + cc_\beta(x, 0)$  is defined and it coincides with  $cc_{\alpha+\beta}(x, 0)$ ;
- (C6) If  $x + x'$  and  $y + y'$  are defined, so is  $cc_\alpha(x + y) + cc_\alpha(x', y')$  and it coincides with  $cc_\alpha(x + x', y + y')$ ;
- (C7)  $cc_\alpha(x, y)^* = cc_\alpha(x^*, y^*)$ .

# Examples

- ▶ Let  $[0, 1]_{PMV}$  be the standard PMV-algebra and, for every  $x, y, \alpha \in [0, 1]$ , define

$$cc_{\alpha}(x, y) = \alpha x \oplus (1 - \alpha)y.$$

In this case  $\oplus$  is the usual sum between real numbers!

Then  $([0, 1]_{MV}, \mathcal{C})$  is a CMV-algebra, with  $\mathcal{C} = \{cc_{\alpha}\}_{\alpha \in [0, 1]}$


# Consequences of the axioms

## Consequences of the axioms


$$(i) \quad cc_1(x, y) = x$$

## Consequences of the axioms


- (i)  $cc_1(x, y) = x$
- (ii)  $cc_\alpha(cc_\beta(x, y), y) = cc_{\alpha\beta}(x, y)$ ;



(iii)  $cc_\alpha(cc_\beta(x, z), cc_\gamma(y, z)) = cc_\mu(cc_\nu(x, y), z)$ , with  $\mu = \alpha\beta + (1 - \alpha)\gamma$  and  $\nu = \frac{\alpha\beta}{\mu}$ ;




$$(iv) \quad cc_{\alpha}(cc_{\beta}(a, b), cc_{\beta}(x, y)) = cc_{\beta}(cc_{\alpha}(a, x), cc_{\alpha}(b, y));$$





$$(iv) \quad cc_{\alpha}(cc_{\beta}(a, b), cc_{\beta}(x, y)) = cc_{\beta}(cc_{\alpha}(a, x), cc_{\alpha}(b, y));$$



$$(v) \quad \text{If } \alpha \leq \beta, \text{ then } cc_{\alpha}(x, 0) \leq cc_{\beta}(x, 0).$$

# Consequences of the axioms

## Consequences of the axioms

(vi)  $x \leq y$  implies  $cc_\alpha(x, z) \leq cc_\alpha(y, z)$ , for any  $z$ ,

## Consequences of the axioms

(vi)  $x \leq y$  implies  $cc_\alpha(x, z) \leq cc_\alpha(y, z)$ , for any  $z$ ,

(vii)  $x \odot y \leq x \wedge y \leq cc_\alpha(x, y) \leq x \vee y \leq x \oplus y$ ,

## Consequences of the axioms

(vi)  $x \leq y$  implies  $cc_\alpha(x, z) \leq cc_\alpha(y, z)$ , for any  $z$ ,

(vii)  $x \odot y \leq x \wedge y \leq cc_\alpha(x, y) \leq x \vee y \leq x \oplus y$ ,

(viii)  $cc_\alpha(x, 0) \leq x$  and  $cc_\alpha(0, y) \leq y$ .

# The main result

# The main result

$\mathbf{CMV}$  and  $\mathbf{RMV}$  are term-wise equivalent

## How it works

$$\mathcal{R} : \mathbb{CMV} \rightarrow \mathbb{RMV}$$



## How it works

$$\mathcal{R} : \mathbb{CMV} \rightarrow \mathbb{RMV}$$

$$(\mathbf{A}, \mathcal{C}) \in \mathbb{CMV},$$

## How it works

$$\mathcal{R} : \mathbb{CMV} \rightarrow \mathbb{RMV}$$

$$(\mathbf{A}, \mathcal{C}) \in \mathbb{CMV},$$

$$(\mathbf{A}, \mathcal{C}) \xrightarrow{\mathcal{R}} (\mathbf{A}, \{\alpha^{\mathcal{R}}(\cdot)\}_{\alpha \in [0,1]})$$

## How it works

$$\mathcal{R} : \mathbb{CMV} \rightarrow \mathbb{RMV}$$

$$(\mathbf{A}, \mathcal{C}) \in \mathbb{CMV},$$

$$(\mathbf{A}, \mathcal{C}) \xrightarrow{\mathcal{R}} (\mathbf{A}, \{\alpha^{\mathcal{R}}(\cdot)\}_{\alpha \in [0,1]})$$
$$\alpha^{\mathcal{R}}(x) = cc_{\alpha}(x, 0).$$

## How it works

$$\mathcal{R} : \mathbb{CMV} \rightarrow \mathbb{RMV}$$

$$(\mathbf{A}, \mathcal{C}) \in \mathbb{CMV},$$

$$\begin{aligned} (\mathbf{A}, \mathcal{C}) &\xrightarrow{\mathcal{R}} (\mathbf{A}, \{\alpha^{\mathcal{R}}(\cdot)\}_{\alpha \in [0,1]}) \\ \alpha^{\mathcal{R}}(x) &= cc_{\alpha}(x, 0). \end{aligned}$$

$$\mathcal{R}(\mathbf{A}, \mathcal{C}) \text{ is a Riesz MV-algebra}$$

## How it works

$$\mathcal{C} : \mathbf{RMV} \rightarrow \mathbf{CMV}$$

## How it works

$$\mathcal{C} : \mathbb{RMV} \rightarrow \mathbb{CMV}$$

$$(\mathbf{A}, \{\alpha(\cdot)\}_{\alpha \in [0,1]}) \in \mathbb{RMV},$$

## How it works

$$\mathcal{C} : \mathbb{RMV} \rightarrow \mathbb{CMV}$$

$$(\mathbf{A}, \{\alpha(\cdot)\}_{\alpha \in [0,1]}) \in \mathbb{RMV},$$

$$\begin{aligned} (\mathbf{A}, \{\alpha(\cdot)\}_{\alpha \in [0,1]}) &\xrightarrow{\mathcal{C}} (\mathbf{A}, \mathcal{C}) \\ \mathcal{C} &= \{\overline{\mathcal{C}}_{\alpha}(\cdot, \cdot)\}_{\alpha \in [0,1]}, \end{aligned}$$

## How it works

$$\mathcal{C} : \mathbb{RMV} \rightarrow \mathbb{CMV}$$

$$(\mathbf{A}, \{\alpha(\cdot)\}_{\alpha \in [0,1]}) \in \mathbb{RMV},$$

$$(\mathbf{A}, \{\alpha(\cdot)\}_{\alpha \in [0,1]}) \xrightarrow{\mathcal{C}} (\mathbf{A}, \mathcal{C})$$

$$\mathcal{C} = \{\overline{\mathcal{C}}_{\alpha}(\cdot, \cdot)\}_{\alpha \in [0,1]},$$

$$\overline{\mathcal{C}}_{\alpha}(x, y) = \alpha x \oplus (1 - \alpha)y.$$



## How it works

$$\mathcal{C} : \mathbb{RMV} \rightarrow \mathbb{CMV}$$

$$(\mathbf{A}, \{\alpha(\cdot)\}_{\alpha \in [0,1]}) \in \mathbb{RMV},$$

$$(\mathbf{A}, \{\alpha(\cdot)\}_{\alpha \in [0,1]}) \xrightarrow{\mathcal{C}} (\mathbf{A}, \mathcal{C})$$

$$\mathcal{C} = \{\overline{\mathcal{C}}_{\alpha}(\cdot, \cdot)\}_{\alpha \in [0,1]},$$

$$\overline{\mathcal{C}}_{\alpha}(x, y) = \alpha x \oplus (1 - \alpha)y.$$

$$\mathcal{C}(\mathbf{A}, \{\alpha(\cdot)\}_{\alpha \in [0,1]}) \text{ is a CMV-algebra.}$$

## Straightforward consequences

## Straightforward consequences

- ▶ A **convex ideal** of  $A$  is a subset  $I$  of  $A$  which is a MV-ideal and such that, if  $x, y \in I$ , then  $cc_\alpha(x, y) \in I$  for every  $\alpha \in [0, 1]$ ,

## Straightforward consequences

- ▶ A **convex ideal** of  $A$  is a subset  $I$  of  $A$  which is a MV-ideal and such that, if  $x, y \in I$ , then  $cc_\alpha(x, y) \in I$  for every  $\alpha \in [0, 1]$ ,  
Every CMV-algebra  $\mathbf{A}$  and its MV-reduct  $\mathbf{A}^-$  have the **same congruences**,

## Straightforward consequences

- ▶ A **convex ideal** of  $A$  is a subset  $I$  of  $A$  which is a MV-ideal and such that, if  $x, y \in I$ , then  $cc_\alpha(x, y) \in I$  for every  $\alpha \in [0, 1]$ ,  
Every CMV-algebra  $\mathbf{A}$  and its MV-reduct  $\mathbf{A}^-$  have the **same congruences**,
- ▶ **Homomorphism** of CMV-algebras are homogeneous homomorphisms of MV-algebras,

## Straightforward consequences

- ▶ A **convex ideal** of  $A$  is a subset  $I$  of  $A$  which is a MV-ideal and such that, if  $x, y \in I$ , then  $cc_\alpha(x, y) \in I$  for every  $\alpha \in [0, 1]$ ,  
Every CMV-algebra  $\mathbf{A}$  and its MV-reduct  $\mathbf{A}^-$  have the **same congruences**,
- ▶ **Homomorphism** of CMV-algebras are homogeneous homomorphisms of MV-algebras,
- ▶ CMV-algebras form a **variety**,

## Straightforward consequences

- ▶ A **convex ideal** of  $A$  is a subset  $I$  of  $A$  which is a MV-ideal and such that, if  $x, y \in I$ , then  $cc_\alpha(x, y) \in I$  for every  $\alpha \in [0, 1]$ ,  
Every CMV-algebra  $\mathbf{A}$  and its MV-reduct  $\mathbf{A}^-$  have the **same congruences**,
- ▶ **Homomorphism** of CMV-algebras are homogeneous homomorphisms of MV-algebras,
- ▶ CMV-algebras form a **variety**,
- ▶ Any CMV-algebra is **subdirect product** of totally ordered CMV-algebras,

# Straightforward consequences

- ▶ A **convex ideal** of  $A$  is a subset  $I$  of  $A$  which is a MV-ideal and such that, if  $x, y \in I$ , then  $cc_\alpha(x, y) \in I$  for every  $\alpha \in [0, 1]$ ,  
Every CMV-algebra  $\mathbf{A}$  and its MV-reduct  $\mathbf{A}^-$  have the **same congruences**,
- ▶ **Homomorphism** of CMV-algebras are homogeneous homomorphisms of MV-algebras,
- ▶ CMV-algebras form a **variety**,
- ▶ Any CMV-algebra is **subdirect product** of totally ordered CMV-algebras,
- ▶  $\mathbf{CMV} = \mathbf{HSP}([0, 1])$ .



# Application to states

## Application to states

We consider states on  $[0, 1]^k$ ,  $k \in \mathbb{N}$ ,

## Application to states

We consider states on  $[0, 1]^k$ ,  $k \in \mathbb{N}$ ,

In  $[0, 1]^k$ , a state is a convex combination of the projections  
 $\{\pi_1, \dots, \pi_k\}$ ,

## Application to states

We consider states on  $[0, 1]^k$ ,  $k \in \mathbb{N}$ ,

In  $[0, 1]^k$ , a state is a convex combination of the projections

$$\{\pi_1, \dots, \pi_k\},$$

$\{\pi_1, \dots, \pi_k\}$  generates the free MV-algebra over  $k$  generators

$$F_{MV}(k),$$

## Application to states

We consider states on  $[0, 1]^k$ ,  $k \in \mathbb{N}$ ,

In  $[0, 1]^k$ , a state is a convex combination of the projections  
 $\{\pi_1, \dots, \pi_k\}$ ,

$\{\pi_1, \dots, \pi_k\}$  generates the free MV-algebra over  $k$  generators  
 $F_{MV}(k)$ ,

$F_{MV}(k) \subseteq [0, 1]^{[0, 1]^k}$  is the subalgebra of functions which are  
piecewise linear with integer coefficients,

## Application to states

We consider states on  $[0, 1]^k$ ,  $k \in \mathbb{N}$ ,

In  $[0, 1]^k$ , a state is a convex combination of the projections  $\{\pi_1, \dots, \pi_k\}$ ,

$\{\pi_1, \dots, \pi_k\}$  generates the free MV-algebra over  $k$  generators  $F_{MV}(k)$ ,

$F_{MV}(k) \subseteq [0, 1]^{[0, 1]^k}$  is the subalgebra of functions which are piecewise linear with integer coefficients,

$F_{MV}(k)$  is not closed for convex combinations!

## Application to states

We consider states on  $[0, 1]^k$ ,  $k \in \mathbb{N}$ ,

In  $[0, 1]^k$ , a state is a convex combination of the projections  $\{\pi_1, \dots, \pi_k\}$ ,

$\{\pi_1, \dots, \pi_k\}$  generates the free MV-algebra over  $k$  generators  $F_{MV}(k)$ ,

$F_{MV}(k) \subseteq [0, 1]^{[0, 1]^k}$  is the subalgebra of functions which are piecewise linear with integer coefficients,

$F_{MV}(k)$  is not closed for convex combinations!

We consider the Riesz Hull and we get

$$R(F_{MV}(k)) \cong [0, 1] \otimes F_{MV}(k) \cong F_{RMV}(k),$$

$F_{RMV}(k)$  is the algebra of piecewise linear functions with real coefficients.



$F_{RMV}(k)$  is the algebra of piecewise linear functions with real coefficients.

## Goal

To write a states  $s : [0, 1]^k \rightarrow [0, 1]$  as a term in  $F_{RMV}(k)$ !

## Notation

for every  $\alpha_1, \dots, \alpha_{k-1} \in [0, 1]$  we set  $[\alpha_i] = \alpha_1, \dots, \alpha_{k-1}$  and for every  $f_1, \dots, f_k \in F_{RMV}(k)$ , we write

$$\begin{aligned} & cc_{[\alpha_i]}(f_1, \dots, f_k) \\ &= \\ & cc_{\alpha_1}(f_1, cc_{\alpha_2}(f_2, \dots cc_{\alpha_{k-1}}(f_{k-1}, f_k) \dots)). \end{aligned}$$

And finally...

# And finally...

For every  $\alpha_1, \dots, \alpha_{k-1} \in [0, 1]$ ,

$cc_{[\alpha_i]}(\pi_1, \dots, \pi_k)$  is a state of  $[0, 1]^k$ .

# And finally...

For every  $\alpha_1, \dots, \alpha_{k-1} \in [0, 1]$ ,

$cc_{[\alpha_i]}(\pi_1, \dots, \pi_k)$  is a state of  $[0, 1]^k$ .

(by the term-wise equivalence)

# And finally...

For every  $\alpha_1, \dots, \alpha_{k-1} \in [0, 1]$ ,

$cc_{[\alpha_i]}(\pi_1, \dots, \pi_k)$  is a state of  $[0, 1]^k$ .

(by the term-wise equivalence)

For every state  $s \in S([0, 1]^k)$ , there exists  $f \in (F_{RMV}(k), \mathcal{C})$  such that,  
for every  $x \in A$ ,

$$s(x) = f(x).$$

# And finally...

For every  $\alpha_1, \dots, \alpha_{k-1} \in [0, 1]$ ,

$cc_{[\alpha_i]}(\pi_1, \dots, \pi_k)$  is a state of  $[0, 1]^k$ .

(by the term-wise equivalence)

For every state  $s \in S([0, 1]^k)$ , there exists  $f \in (F_{RMV}(k), \mathcal{C})$  such that,  
for every  $x \in A$ ,

$$s(x) = f(x).$$

$(S([0, 1]^k) = co(H([0, 1]^k)))$  + the fact that projections generate  $F_{RMV}(k)$

THANK YOU!