

On modal expansions of left-continuous t-norm logics

Amanda Vidal

Joint work with
Félix Bou, Francesc Esteva and Lluís Godó



Institute of Computer Science,
Czech Academy of Sciences



Summary of contents

1. Introduction

2. Strong standard completeness

MTL with rational constants and Δ

Modally well-behaved axiomatizations

3. Modal expansions

4. Open problems

Summary of contents

1. Introduction

2. Strong standard completeness

MTL with rational constants and Δ

Modally well-behaved axiomatizations

3. Modal expansions

4. Open problems

Fuzzy & modal logics




Figure: Hierarchy of t-norm based logics and extensions

Fuzzy & modal logics




Figure: Hierarchy of t-norm based logics and extensions

- Reasoning over qualification of sentences (necessity, possibility...)

Fuzzy & modal logics




Figure: Hierarchy of t-norm based logics and extensions

- ▶ Reasoning over qualification of sentences (necessity, possibility...)
- ▶ Usual modalities: \Box, \Diamond . Kripke semantics.

Definition

$A \in \text{MTL-algebra}$. An A -**Kripke model** ($*$ -Kripke model) $M = \langle W, R, e \rangle$
s.t. $[W \neq \emptyset]$ $[R : W \times W \rightarrow A]$ $[e : W \times \text{Var} \rightarrow A]$:

$$e(w, \Box \varphi) := \inf \{ R w v \Rightarrow e(v, \varphi) : v \in W \}$$

$$e(w, \Diamond \varphi) := \sup \{ R w v \odot e(v, \varphi) : v \in W \}$$

Definition

$A \in \text{MTL-algebra}$. An A -**Kripke model** ($*$ -Kripke model) $M = \langle W, R, e \rangle$
s.t. $[W \neq \emptyset]$ $[R : W \times W \rightarrow A]$ $[e : W \times \text{Var} \rightarrow A]$:

$$e(w, \Box \varphi) := \inf \{ R w v \Rightarrow e(v, \varphi) : v \in W \}$$

$$e(w, \Diamond \varphi) := \sup \{ R w v \odot e(v, \varphi) : v \in W \}$$

crisp: $R \subseteq W \times W$.

Definition

$A \in \text{MTL-algebra}$. An **A -Kripke model** ($*$ -Kripke model) $M = \langle W, R, e \rangle$ s.t. $[W \neq \emptyset]$ $[R : W \times W \rightarrow A]$ $[e : W \times \text{Var} \rightarrow A]$:

$$e(w, \Box \varphi) := \inf \{ R w v \Rightarrow e(v, \varphi) : v \in W \}$$

$$e(w, \Diamond \varphi) := \sup \{ R w v \odot e(v, \varphi) : v \in W \}$$

crisp: $R \subseteq W \times W$.

- **Łukasiewicz** [Hansoul-Teheux] : crisp Kripke models over $[0, 1]_{\mathbb{L}}$.
linfinitary rule (propositional strong completeness).

Definition

$A \in \text{MTL-algebra}$. An **A-Kripke model** ($*$ -Kripke model) $M = \langle W, R, e \rangle$ s.t. $[W \neq \emptyset]$ $[R : W \times W \rightarrow A]$ $[e : W \times \text{Var} \rightarrow A]$:

$$e(w, \Box \varphi) := \inf \{ R w v \Rightarrow e(v, \varphi) : v \in W \}$$

$$e(w, \Diamond \varphi) := \sup \{ R w v \odot e(v, \varphi) : v \in W \}$$

crisp: $R \subseteq W \times W$.

- ▶ **Łukasiewicz** [Hansoul-Teheux] : crisp Kripke models over $[0, 1]_{\mathbb{L}}$.
infinitary rule (propositional strong completeness).
- ▶ **Finite residuated lattices** [Bou et. al]: Kripke models over finite residuated lattices, only \Box operator. One constant in the language for each element in the algebra.

Definition

$A \in \text{MTL-algebra}$. An **A-Kripke model** ($*$ -Kripke model) $M = \langle W, R, e \rangle$ s.t. $[W \neq \emptyset]$ $[R : W \times W \rightarrow A]$ $[e : W \times \text{Var} \rightarrow A]$:

$$e(w, \Box \varphi) := \inf \{ R w v \Rightarrow e(v, \varphi) : v \in W \}$$

$$e(w, \Diamond \varphi) := \sup \{ R w v \odot e(v, \varphi) : v \in W \}$$

crisp: $R \subseteq W \times W$.

- ▶ **Łukasiewicz** [Hansoul-Teheux] : crisp Kripke models over $[0, 1]_{\mathbb{L}}$.
infinitary rule (propositional strong completeness).
- ▶ **Finite residuated lattices** [Bou et. al]: Kripke models over finite residuated lattices, only \Box operator. One constant in the language for each element in the algebra.
- ▶ **Gödel** [Caicedo-Rodríguez]: Kripke models over $[0, 1]_G$, both \Box and \Diamond . Strong standard completeness of Gödel logic and of characteristics of $[0, 1]_G$ -endomorphisms. **Finite model property**[+ Metcalfe-Rogger]

Summary of contents

1. Introduction

2. Strong standard completeness

MTL with rational constants and Δ

Modally well-behaved axiomatizations

3. Modal expansions

4. Open problems

Strong completeness

objective: for a left-continuous t-norm $*$ » an **(strong) axiomatization** of $[0, 1]_*^{\mathbb{Q}}$: standard algebra of $*$ with Δ and canonical (rational) constants.

Strong completeness

objective: for a left-continuous t-norm $*$ » an **(strong) axiomatization** of $[0, 1]_*^{\mathbb{Q}}$: standard algebra of $*$ with Δ and canonical (rational) constants.

Proposition [Cintula]

Let A be an expansion of an standard *MTL*-algebra with a non-continuous operation, and L_A a finitary axiomatic system for A . Then no finitary rational expansion of L_A enjoys the Pavelka-style completeness.

$$\blacktriangleright \Pi \text{ logic} + \Delta \text{ op} \quad \frac{\{p \rightarrow \bar{c}\}_{c \in (0,1)_{\mathbb{Q}}}}{\neg p}, \quad \frac{\{\bar{c} \rightarrow p\}_{c \in (0,1)_{\mathbb{Q}}}}{\Delta p}$$

Strong completeness

objective: for a left-continuous t-norm $*$ » an **(strong) axiomatization** of $[0, 1]_*^{\mathbb{Q}}$: standard algebra of $*$ with Δ and canonical (rational) constants.

Proposition [Cintula]

Let A be an expansion of an standard *MTL*-algebra with a non-continuous operation, and L_A a finitary axiomatic system for A . Then no finitary rational expansion of L_A enjoys the Pavelka-style completeness.

- ▶ Π logic + Δ op $\frac{\{p \rightarrow \bar{c}\}_{c \in (0,1)_{\mathbb{Q}}}}{\neg p}, \quad \frac{\{\bar{c} \rightarrow p\}_{c \in (0,1)_{\mathbb{Q}}}}{\Delta p}$
- ▶ General: an axiomatic system of $[0, 1]_*$ validates a certain infinitary rule for each discontinuity point of the operations & is seminilinear \implies it is Pavelka complete.

Strong completeness

objective: for a left-continuous t-norm $*$ » an **(strong) axiomatization** of $[0, 1]_*^{\mathbb{Q}}$: standard algebra of $*$ with Δ and canonical (rational) constants.

Proposition [Cintula]

Let A be an expansion of an standard *MTL*-algebra with a non-continuous operation, and L_A a finitary axiomatic system for A . Then no finitary rational expansion of L_A enjoys the Pavelka-style completeness.

- ▶ Π logic + Δ op $\frac{\{p \rightarrow \bar{c}\}_{c \in (0,1)_{\mathbb{Q}}}}{\neg p}, \quad \frac{\{\bar{c} \rightarrow p\}_{c \in (0,1)_{\mathbb{Q}}}}{\Delta p}$
- ▶ General: an axiomatic system of $[0, 1]_*$ validates a certain infinitary rule for each discontinuity point of the operations & is semilinear \implies it is Pavelka complete.

* many cases: uncountable infinitary rules

* how to know when a logic with infinitary rules is semilinear?

Semilinearity Lemma

Let L be an implicative logic expanding MTL_{Δ} such that

- ▶ There is a countable amount of (infinitary) inference rules (with a finite number of variables each)
- ▶ The derivations of L are closed under \vee

Then L is semilinear.

Semilinearity Lemma

Let L be an implicative logic expanding MTL_{Δ} such that

- ▶ There is a countable amount of (infinitary) inference rules (with a finite number of variables each)
- ▶ The derivations of L are closed under \vee

Then L is semilinear.

Proof based on a prime-theory construction.

Semilinearity Lemma

Let L be an implicative logic expanding MTL_Δ such that

- ▶ There is a countable amount of (infinitary) inference rules (with a finite number of variables each)
- ▶ The derivations of L are closed under \vee

Then L is semilinear.

Proof based on a prime-theory construction.

goal: finding a countable amount of infinitary rules for treating **all the discontinuity points**

A strong SC axiomatization

Takeuti and Titani's density rule can be adapted

$$D^{\infty} : \frac{\{(p \rightarrow \bar{c}) \vee (\bar{c} \rightarrow q)\}_{c \in Q_*}}{(p \rightarrow q)}$$

A strong SC axiomatization

Takeuti and Titani's density rule can be adapted

$$D^{\infty} : \frac{\{(p \rightarrow \bar{c}) \vee (\bar{c} \rightarrow q)\}_{c \in \mathbb{Q}_*}}{(p \rightarrow q)}$$

Definition

$L_*^{\infty} :=$ extension of MTL_{Δ} by the book-keeping axioms for \mathbb{Q}_* and D^{∞} .

A strong SC axiomatization

Takeuti and Titani's density rule can be adapted

$$D^{\infty} : \frac{\{(p \rightarrow \bar{c}) \vee (\bar{c} \rightarrow q)\}_{c \in \mathbb{Q}_*}}{(p \rightarrow q)}$$

Definition

$L_*^{\infty} :=$ extension of MTL_{Δ} by the book-keeping axioms for \mathbb{Q}_* and D^{∞} .

- Deductions on L_*^{∞} are closed under $\vee \gg L_*^{\infty}$ is semilinear.

A strong SC axiomatization


Takeuti and Titani's density rule can be adapted

$$D^{\infty} : \frac{\{(p \rightarrow \bar{c}) \vee (\bar{c} \rightarrow q)\}_{c \in \mathbb{Q}_*}}{(p \rightarrow q)}$$

Definition

$L_*^{\infty} :=$ extension of MTL_{Δ} by the book-keeping axioms for \mathbb{Q}_* and D^{∞} .

- ▶ Deductions on L_*^{∞} are closed under $\vee \gg L_*^{\infty}$ is semilinear.
- ▶ Constants are dense on linearly ordered L_*^{∞} -algebras.



A strong SC axiomatization


Takeuti and Titani's density rule can be adapted

$$D^{\infty} : \frac{\{(p \rightarrow \bar{c}) \vee (\bar{c} \rightarrow q)\}_{c \in \mathbb{Q}_*}}{(p \rightarrow q)}$$

Definition

L_*^{∞} := extension of MTL_{Δ} by the book-keeping axioms for \mathbb{Q}_* and D^{∞} .

- ▶ Deductions on L_*^{∞} are closed under $\vee \gg L_*^{\infty}$ is semilinear.
- ▶ Constants are dense on linearly ordered L_*^{∞} -algebras.



Strong Standard Completeness

$$\Gamma \vdash_{L_*^{\infty}} \varphi$$

iff

$$\Gamma \models_{[0,1]_*^{\mathbb{Q}_*}} \varphi$$

Conjunctive rules

The density rule is not well-behaved when the logic is expanded with modalities: not clear how to prove that it is closed under \Box .

Conjunctive rules

The density rule is not well-behaved when the logic is expanded with modalities: not clear how to prove that it is closed under \Box .

Definition

Conjunctive rules: for $x \in [0, 1]$

$$R_x^\infty : \frac{\{(p \rightarrow \bar{c}) \wedge (\bar{d} \rightarrow q)\}_{d \in [0, x]_{\mathbb{Q}_*}, c \in (x, 1]_{\mathbb{Q}_*}}}{p \rightarrow q}$$

Conjunctive rules

The density rule is not well-behaved when the logic is expanded with modalities: not clear how to prove that it is closed under \Box .

Definition

Conjunctive rules: for $x \in [0, 1]$

$$R_x^\infty : \frac{\{(p \rightarrow \bar{c}) \wedge (\bar{d} \rightarrow q)\}_{d \in [0, x)_{\mathbb{Q}_*}, c \in (x, 1]_{\mathbb{Q}_*}}}{p \rightarrow q}$$

Lemma

Let A be a $MTL_{*}^{\mathbb{Q}_*}$ -chain. Then the following are equivalent:

1. A validates D^∞ ,
2. A validates R_x^∞ for all $x \in [0, 1]$,
3. for all $a < b \in A$, there is $c \in \mathbb{Q}_*$ such that $a < \bar{c}^A < b$.

Definition

A t-norm $*$ **accepts a conjunctive axiomatization** when there exists an axiomatic system L such that:

- ▶ L is strongly complete with respect to $[0, 1]_*^{\mathbb{Q}}$,
- ▶ L extends $MTL_*^{\mathbb{Q}}$,
- ▶ The only inference rules added to $MTL_*^{\mathbb{Q}}$ are conjunctive.

Definition

A t-norm $*$ **accepts a conjunctive axiomatization** when there exists an axiomatic system L such that:

- ▶ L is strongly complete with respect to $[0, 1]_{*}^{\mathbb{Q}}$,
- ▶ L extends $MTL_{*}^{\mathbb{Q}}$,
- ▶ The only inference rules added to $MTL_{*}^{\mathbb{Q}}$ are conjunctive.

✗ Gödel t-norm does not accept a conjunctive axiomatization: there is a Gödel algebra that validates all R_x^{∞} but not D^{∞} .

Definition

A t-norm $*$ **accepts a conjunctive axiomatization** when there exists an axiomatic system L such that:

- ▶ L is strongly complete with respect to $[0, 1]_*^{\mathbb{Q}}$,
- ▶ L extends $MTL_*^{\mathbb{Q}}$,
- ▶ The only inference rules added to $MTL_*^{\mathbb{Q}}$ are conjunctive.

✗ Gödel t-norm does not accept a conjunctive axiomatization: there is a Gödel algebra that validates all R_x^∞ but not D^∞ .

✓ Ordinal sums of Lukasiewicz and Product t-norms.

Definition

A t-norm $*$ **accepts a conjunctive axiomatization** when there exists an axiomatic system L such that:

- ▶ L is strongly complete with respect to $[0, 1]_*^{\mathbb{Q}}$,
- ▶ L extends $MTL_*^{\mathbb{Q}}$,
- ▶ The only inference rules added to $MTL_*^{\mathbb{Q}}$ are conjunctive.

✗ Gödel t-norm does not accept a conjunctive axiomatization: there is a Gödel algebra that validates all R_x^∞ but not D^∞ .

- ✓ Ordinal sums of Lukasiewicz and Product t-norms.
- ✓ Left-continuous t-norms with a countable number of discontinuity points on the diagonal of its residuum.

Summary of contents

1. Introduction

2. Strong standard completeness

MTL with rational constants and Δ

Modally well-behaved axiomatizations

3. Modal expansions

4. Open problems

In what follows, $*$ is a **left-continuous t-norm accepting a conjunctive axiomatization**.

In what follows, $*$ is a **left-continuous t-norm accepting a conjunctive axiomatization**.

- ▶ SM_* : class of all **crisp** $[0, 1]_{*}^{\mathbb{Q}}$ -Kripke models.
- ▶ KM_* : class of all **safe crisp** A -Kripke models for A a L_*^{∞} -algebra.

In what follows, $*$ is a **left-continuous t-norm accepting a conjunctive axiomatization**.

- ▶ SM_* : class of all **crisp** $[0, 1]_{*}^{\mathbb{Q}}$ -Kripke models.
- ▶ KM_* : class of all **safe crisp** A -Kripke models for A a L_*^{∞} -algebra.

Local modal logic: $\Gamma \Vdash_{\mathbb{M}} \varphi$ iff for any $M \in \mathbb{M}$,

for any $w \in W$, if $e(w, \Gamma) = \{1\}$ then $e(w, \varphi) = 1$.

Global modal logic: $\Gamma \Vdash_{\mathbb{M}}^g \varphi$ iff for any $M \in \mathbb{M}$,

if for any $w \in W$ $e(w, \Gamma) = \{1\}$ then for any $w \in W$ $e(w, \varphi) = 1$.

Axiomatic systems: K_* and K_*^g

- **Local modal logic:** K_* is the expansion of L_*^∞ with

$$K : \Box(p \rightarrow q) \rightarrow (\Box p \rightarrow \Box q)$$

$$\Box 1 : \Box(\bar{c} \rightarrow p) \leftrightarrow (\bar{c} \rightarrow \Box p)$$

$$\Box 2 : \Delta \Box p \rightarrow \Box \Delta p$$

$$\Diamond 1 : \Box(p \rightarrow \bar{c}) \leftrightarrow (\Diamond p \rightarrow \bar{c})$$

$$FS1 : (\Diamond p \rightarrow \Box q) \rightarrow \Box(p \rightarrow q)$$

$$N_\Box : \text{From } \emptyset \vdash \varphi \text{ infer } \emptyset \vdash \Box \varphi.$$

Axiomatic systems: K_* and K_*^g

- **Local modal logic:** K_* is the expansion of L_*^∞ with

$$K : \Box(p \rightarrow q) \rightarrow (\Box p \rightarrow \Box q)$$

$$\Box 1 : \Box(\bar{c} \rightarrow p) \leftrightarrow (\bar{c} \rightarrow \Box p)$$

$$\Box 2 : \Delta \Box p \rightarrow \Box \Delta p$$

$$\Diamond 1 : \Box(p \rightarrow \bar{c}) \leftrightarrow (\Diamond p \rightarrow \bar{c})$$

$$FS1 : (\Diamond p \rightarrow \Box q) \rightarrow \Box(p \rightarrow q)$$

$$N_\Box : \text{From } \emptyset \vdash \varphi \text{ infer } \emptyset \vdash \Box \varphi.$$

- **Global modal logic:** K_*^g coincides with K_* except for the N_\Box rule, turned to $N_\Box^g : \varphi \vdash \Box \varphi$

Axiomatic systems: K_* and K_*^g

- **Local modal logic:** K_* is the expansion of L_*^∞ with

$$K : \Box(p \rightarrow q) \rightarrow (\Box p \rightarrow \Box q)$$

$$\Box 1 : \Box(\bar{c} \rightarrow p) \leftrightarrow (\bar{c} \rightarrow \Box p)$$

$$\Box 2 : \Delta \Box p \rightarrow \Box \Delta p$$

$$\Diamond 1 : \Box(p \rightarrow \bar{c}) \leftrightarrow (\Diamond p \rightarrow \bar{c})$$

$$FS1 : (\Diamond p \rightarrow \Box q) \rightarrow \Box(p \rightarrow q)$$

$$N_\Box : \text{From } \emptyset \vdash \varphi \text{ infer } \emptyset \vdash \Box \varphi.$$

- **Global modal logic:** K_*^g coincides with K_* except for the N_\Box rule, turned to $N_\Box^g : \varphi \vdash \Box \varphi$

Some important properties of K_*

1. $\Gamma \vdash_{K_*} \varphi$ if and only if $(\Gamma \cup Th(K_*))^\# \vdash_{L_*^\infty} \varphi^\#$
translates formulas $\Box \varphi, \Diamond \varphi$ to new propositional variables $\varphi_\Box, \varphi_\Diamond$.

Axiomatic systems: K_* and K_*^g

- **Local modal logic:** K_* is the expansion of L_*^∞ with

$$K : \Box(p \rightarrow q) \rightarrow (\Box p \rightarrow \Box q)$$

$$\Box 1 : \Box(\bar{c} \rightarrow p) \leftrightarrow (\bar{c} \rightarrow \Box p)$$

$$\Box 2 : \Delta \Box p \rightarrow \Box \Delta p$$

$$\Diamond 1 : \Box(p \rightarrow \bar{c}) \leftrightarrow (\Diamond p \rightarrow \bar{c})$$

$$FS1 : (\Diamond p \rightarrow \Box q) \rightarrow \Box(p \rightarrow q)$$

$$N_\Box : \text{From } \emptyset \vdash \varphi \text{ infer } \emptyset \vdash \Box \varphi.$$

- **Global modal logic:** K_*^g coincides with K_* except for the N_\Box rule, turned to $N_\Box^g : \varphi \vdash \Box \varphi$

Some important properties of K_*

1. $\Gamma \vdash_{K_*} \varphi$ if and only if $(\Gamma \cup Th(K_*))^\# \vdash_{L_*^\infty} \varphi^\#$
translates formulas $\Box \varphi, \Diamond \varphi$ to new propositional variables $\varphi_\Box, \varphi_\Diamond$.
2. $\Gamma \vdash_{K_*} \varphi$ implies that $\Box \Gamma \vdash_{K_*} \Box \varphi$.

Canonical model for the local logic

The **canonical model of K_*** is the $[0, 1]_{**}^{\mathbb{Q}}$ -Kripke model

$M_c^* = \langle W_c^*, R_c^*, e_c^* \rangle$ with:

- ▶ $W_c^* = \{h \in \text{Hom}(Fm^\#, [0, 1]_{**}^{\mathbb{Q}}) : h(\text{Th}(K_*)^\#) = \{1\}\}$
- ▶ R_c^*vw when for any φ such that $v(\varphi_\square) = 1$ then $w(\varphi^\#) = 1$.
- ▶ $e_c^*(v, p) = v(p)$ for the propositional variables.

Canonical model for the local logic

The **canonical model of K_*** is the $[0, 1]_*^{\mathbb{Q}_*}$ -Kripke model

$M_c^* = \langle W_c^*, R_c^*, e_c^* \rangle$ with:

- ▶ $W_c^* = \{h \in \text{Hom}(Fm^\#, [0, 1]_*^{\mathbb{Q}_*}) : h(\text{Th}(K_*)^\#) = \{1\}\}$
- ▶ R_c^*vw when for any φ such that $v(\varphi_\square) = 1$ then $w(\varphi^\#) = 1$.
- ▶ $e_c^*(v, p) = v(p)$ for the propositional variables.

Some important properties of M_c^*

1. R_c^*vw iff for all φ both $v(\varphi_\square) \leq w(\varphi^\#)$ and $v(\varphi_\diamond) \geq w(\varphi^\#)$.

Canonical model for the local logic

The **canonical model of K_*** is the $[0, 1]_{**}^{\mathbb{Q}}$ -Kripke model

$M_c^* = \langle W_c^*, R_c^*, e_c^* \rangle$ with:

- ▶ $W_c^* = \{h \in \text{Hom}(Fm^\#, [0, 1]_{**}^{\mathbb{Q}}) : h(Th(K_*)^\#) = \{1\}\}$
- ▶ R_c^*vw when for any φ such that $v(\varphi_\square) = 1$ then $w(\varphi^\#) = 1$.
- ▶ $e_c^*(v, p) = v(p)$ for the propositional variables.

Some important properties of M_c^*

1. R_c^*vw iff for all φ both $v(\varphi_\square) \leq w(\varphi^\#)$ and $v(\varphi_\diamond) \geq w(\varphi^\#)$.
2. Let $v \in W_c^*$ and φ be such that $w(\varphi^\#) = 1$ for all $w \in W_c^*$ with R_c^*vw . Then $v(\varphi_\square) = 1$.

Completeness of the local modal logic

Truth Lemma

For any formula φ and any $v \in W_c^*$

$$e_c^*(v, \varphi) = v(\varphi^\#).$$

Completeness of the local modal logic

Truth Lemma

For any formula φ and any $v \in W_c^*$

$$e_c^*(v, \varphi) = v(\varphi^\#).$$

Strong standard completeness of K_*

For any $\Gamma \cup \{\varphi\} \subseteq Fm$, the following are equivalent

1. $\Gamma \vdash_{K_*} \varphi$,
2. $\Gamma \Vdash_{SM_*} \varphi$,
3. $\Gamma \Vdash_{KM_*} \varphi$.

- ▶ soundness is easy to check (wrt 3.)
- ▶ If $\Gamma \not\vdash_{K_*} \varphi$ then $(\Gamma \cup Th(K_*))^\# \not\vdash_{L_*^\infty} \varphi^\#$. Non-modal SSC provides a world w from the canonical model where $e_c^*(w, \Gamma) = w(\Gamma^\#) = \{1\}$ and $e_c^*(w, \varphi) = w(\varphi^\#) < 1$.

The global modal logic

Instead of a unique canonical model, one model for each set of formulas.

For Γ a set of formulas, the Γ -**canonical model of K_*^g is the $[0, 1]_*^{\mathbb{Q}}$ -Kripke model $M_c^*[\Gamma] = \langle W_c^*[\Gamma], R_c^*[\Gamma], e_c^*[\Gamma] \rangle$ with:**

- ▶ $W_c^*[\Gamma] = \{h \in \text{Hom}(Fm^\#, [0, 1]_*^{\mathbb{Q}}) : h(\text{Con}_{K_*^g}(\Gamma)^\#) = \{1\}\}$
- ▶ $R_c[\Gamma]vw$ if and only if for any formula φ such that $v(\varphi_\square) = 1$ it holds that $w(\varphi^\#) = 1$.
- ▶ $e_c^*(v, p)[\Gamma] = v(p)$ for the propositional variables.

The global modal logic

Instead of a unique canonical model, one model for each set of formulas.

For Γ a set of formulas, the Γ -**canonical model** of K_*^g is the $[0, 1]_*^{\mathbb{Q}}$ -**Kripke model** $M_c^*[\Gamma] = \langle W_c^*[\Gamma], R_c^*[\Gamma], e_c^*[\Gamma] \rangle$ with:

- ▶ $W_c^*[\Gamma] = \{ h \in \text{Hom}(Fm^\#, [0, 1]_*^{\mathbb{Q}}) : h(\text{Con}_{K_*^g}(\Gamma)^\#) = \{1\} \}$
- ▶ $R_c[\Gamma]vw$ if and only if for any formula φ such that $v(\varphi_\square) = 1$ it holds that $w(\varphi^\#) = 1$.
- ▶ $e_c^*(v, p)[\Gamma] = v(p)$ for the propositional variables.

Strong standard completeness of K_*^g

For any $\Gamma \cup \{ \varphi \} \subseteq Fm$, the following are equivalent

1. $\Gamma \vdash_{K_*^g} \varphi$,
2. $\Gamma \Vdash_{\text{SM}_*}^g \varphi$,
3. $\Gamma \Vdash_{\text{KM}_*}^g \varphi$.

Summary of contents

1. Introduction

2. Strong standard completeness

MTL with rational constants and Δ

Modally well-behaved axiomatizations

3. Modal expansions

4. Open problems

Open problems

- ▶ Axiomatic system for logics arising from an arbitrary left-continuous t -norm.
- ▶ Modal expansion of MTL -logics without rational constants.
- ▶ Axiomatic system of the finitary modal logics -evaluated over infinite-valued algebras-.
- ▶ Logics arising from Kripke models evaluated over MTL -algebras whose accessibility relation is no longer crisp.
- ▶ Decidability and complexity problems.
- ▶ ...

Thank you!