


A
Definition of next-mutation recall
Future phylogeny
Testing examples
MFVWLVLV
V6I ✗ True
V3A ✓ mutations
antiGen
V3A L4C Predicted
V6I V8L top 10%
V3L F2W
MFVWLILV
F2Y ✓ True mutation
antiGen
F2Y L4C Predicted
I6V V3L top 10%
V8L L5V
Time
Past (training)
Future (testing)
B
Case study: BA.2.86 to KP.3
BA.2.86
JN.1
JN.1.11.1
KP.3
L455S
V483D
R346T
F456L
D483V
T346R
Q493E
still changed in KP.3
reverts along the way
top decile
antiGen rank
antiGen
S×S-MCM interp.
antiGen / PFM gate
MCM
S×M
PFM
S×S
BLOSUM62
S×M-MCM interp.
EVEscape
C
Top decile recall, all RBD cells (per cell)
P(cell in top decile)
Train / test cutoff
D
Top decile recall, all RBD cells (per mutation)
P(obs. mut. in top decile)
Train / test cutoff
Model type
antiGen
Statistical models
Deep Mutational Scans
Pandemic Data in Training Window
Pre-Pandemic Data Only
Designed in this work
External


CoG-UK escape list, 2021-01-01 cutoff
All RBD cells, 2023-01-01 cutoff
E All escape, per cell
F All escape, per mutation
G De novo escape, per cell
H De novo escape, per mutation
I All RBD, per cell
J All RBD, per mutation
K De novo RBD, per cell
L De novo RBD, per mutation
antiGen (CV x10)
S×S
S×S-MCM interp.
PFM
PFM Conservation
MSA Transformer
EVE
Tranception
Codon-B62 mixture
BA.1 DMS (bind)
BA.1 DMS (expression)
BA.2 DMS (bind)
BA.2 DMS (expression)
EVEscape
ESM 2 (35M), untrained
BLOSUM62
Codon table
Recall at q = 0.1