
Recall at top 10%
A
All substitutions
antiGen (multi−tree)
antiGen (single−tree)
Where x which
Frequency (memorization)
Substitution matrix
PSSM (sequence composition)
Where only
Codon accessibility
Uniform
0.88
0.83
0.75
0.73
0.69
0.65
0.45
0.29
0.10
0.00
0.25
0.50
0.75
Recall at top 10%
B
De novo only
antiGen (multi−tree)
antiGen (single−tree)
Where x which
Frequency (memorization)
Substitution matrix
PSSM (sequence composition)
Where only
Codon accessibility
Uniform
0.61
0.44
0.50
0.05
0.49
0.07
0.19
0.36
0.10
0.0
0.2
0.4
0.6
Recall at top 10%
All substitutions
De novo only
95% interval excludes zero
interval includes zero
C
Versus single−tree
SARS−CoV−2 S
H5Nx HA
Measles H
RSV A F
RSV B F
DENV1 E
DENV2 E
DENV3 E
DENV4 E
H3N2 HA
H1N1pdm HA
Pooled
−0.5
0.0
0.5
1.0
Multi−tree minus comparator, recall at top 10%
D
Versus where x which
SARS−CoV−2 S
H5Nx HA
Measles H
RSV A F
RSV B F
DENV1 E
DENV2 E
DENV3 E
DENV4 E
H3N2 HA
H1N1pdm HA
Pooled
−0.5
0.0
0.5
1.0
Multi−tree minus comparator, recall at top 10%
antiGen (multi−tree)
Where x which
E
All substitutions, recall at top 10%
Which changes are flagged
113
16%
22
3%
507
72%
caught by neither: 66 (9%) of 708
F
De novo only, recall at top 10%
35
21%
16
9%
68
40%
caught by neither: 50 (30%) of 169