
State effect on per-set copy number (95% CI; one-sided p for multicellular > unicellular)
R2-P2 motif-binding (ELM, 115 families)
confirmatory
R2-P3 cell death (12 widespread families)
confirmatory
P4 cell-type TFs (11 widespread families)
exploratory
main
BM (lambda=1)
no disorder covariate
grey-zone excluded
all 187 proteomes
all 187 + transcriptome & BUSCO covariates
drop metazoa
drop fungi_hyphae
drop dictyostelia
drop streptophyta
drop rhodophyta
drop px_filaments
drop volvocine_tgv
p=0.722
p=0.722
p=0.677
p=0.839
p=0.850
p=0.424
p=0.934
p=0.552
p=0.743
p=0.739
p=0.629
p=0.752
p=0.464
p=0.005
p=0.019
p=0.003
p=0.009
p=0.005
p=9.7e-04
p=0.010
p=0.004
p=0.004
p=0.078
p=0.012
p=0.004
p=2.0e-04
p=0.298
p=0.298
p=0.264
p=0.377
p=0.315
p=0.102
p=0.356
p=0.436
p=0.286
p=0.444
p=0.242
p=0.311
p=0.077
−0.15 −0.10 −0.05 0.00 0.05 0.10
0.0 0.1 0.2 0.3 0.4
−0.1 0.0 0.1 0.2
multicellular − unicellular coefficient (SD units of the equal-weight mean of standardised log copy number)
primary PGLS (λ), 148 genomes
sensitivity
leave one primary origin out