

# Growth curve analyses in poultry science

D. NARINÇ<sup>1\*</sup>, N. ÖKSÜZ NARINÇ<sup>2</sup> and A. AYGÜN<sup>3</sup>

<sup>1</sup>Faculty of Veterinary Medicine, Department of Genetics, Namık Kemal University, Turkey; <sup>2</sup>Vocational School of Social Sciences, Namık Kemal University, Turkey;

<sup>3</sup>Faculty of Agriculture, Department of Animal Science, Selçuk University, Turkey

\*Corresponding author: [dnarinc@nku.edu.tr](mailto:dnarinc@nku.edu.tr)

---

Growth is a key characteristic of animals and can be defined as any change in body size per time unit, and is influenced by genotype and environment. Mathematical functions called ‘growth models’ have been used to explain the growth patterns of poultry species. These semi-mechanistic growth models have a non-linear structure, sigmoid shape, and certain biologically meaningful parameters. In poultry science, Gompertz, Logistic, Richards and von Bertalanffy functions have been commonly used to model the growth patterns of birds. In this review, the studies concerned have been summarised under the titles ‘determination of the best-fitting growth model’, ‘a comparison of the growth of poultry species or various experimental groups’, and ‘genetic parameter estimates for growth curve parameters’. This review discusses existing and new approaches to growth modelling.

---

**Keywords:** growth; growth functions; poultry species; mechanistic modelling

## Introduction

For about two hundred years, mathematical models have been used to explain and interpret data obtained through observations and measurements and to reveal cause and effect relationships in different fields of science. Mathematical functions enable scientists to reach unbiased results by considering the events in an abstract manner. The evaluation and processing of data collected from samples within the population or the system as a whole, clarification of the findings, their validation on wide scales and estimations obtained for the following cases are more accurate and significant through the use of mathematical functions. In addition, it is possible to intervene in the changes at various times and in different cases considering the estimations obtained as a result of the modelling of a system (Panik, 2014). Since the growth of a living organism is affected by many factors in biology, it is not always possible to identify some of them numerically. Therefore, the methods used in exact sciences such as physics and chemistry cannot be used with the same success for animals. Besides, it is not compulsory to represent all factors in the model when making the mathematical definition of the growth of a system or an organism. Conversely, representing the factors with fewer parameters and a simple function is essential in terms of both

© World's Poultry Science Association 2017

World's Poultry Science Journal, Vol. 73, June 2017

*Received for publication October 18, 2016*

*Accepted for publication November 24, 2016*


comprehensibility and practice. If the basic model is presented, it may be allowed to solve similar problems under different conditions by changing or transforming the parameters of the model (Konarzewski *et al.*, 1998).

Mostly 'asymptotic' functions and 'parabolic' functions - although at a small rate - are used to model data from agriculture. If the dependent variable tends to approach the maximum point according to the levels of the independent variable, it is expressed with an asymptotic function. However, if the dependent variable tends to decrease after it has reached the maximum, it represents a parabolic process. In plants and animals, growth is generally of asymptotic structure. Growth is of parabolic structure in plants such as parsnip, corn, and cotton. The models used in the field of agriculture are divided into three categories as teleonomic, empirical, and mechanistic (Thornley and France, 2007). Teleonomic models act towards the target and are generally multi-stage mathematical models which are clearly formulated in terms of the target. The use of teleonomic modelling is rather limited in explaining biological events (France and Dijkstra, 2006). The mathematical models for growth may be either empirical or mechanistic. A mechanistic model should be preferred to represent the biochemical processes controlling growth as much as possible. However, unless the mechanism which manages the process is known in detail and the data obtained with an experiment are interpreted, the mathematical function of the process may be empirical and model fitness evaluates the ability of fitness to empirical data according to the statistical criteria. Empirical models are expressions which define the action of the dependent variable without attempting to identify the causes or to explain the event. This means neither that empirical functions do not provide biologically realistic estimations nor that they fall short of biologically-based equations. The most widely used form of empirical equations is linear, and although they have non-linear forms as well. In fact, when more information about the system is available, empirical models can be transformed into a more mechanistic structure (Zwitering *et al.*, 1990).

## Growth modelling in the poultry

Many mathematical models have been used to model growth in poultry, and the majority are asymptotic and mechanistic. These models consist of functions which accept, on the basis of the reality of the biological growth of an animal, that the dependent variable has an estimated asymptotic value when the independent variable is at infinity. These growth models use non-linear functions, and the majority of them have a sigmoid structure; however, their courses of growth may vary. A comprehensive review and history of the growth models used in poultry species has been published by Darmani Kuhi *et al.* (2010). Many models have been made more mechanistic by adding biologically significant parameters to an empirical infrastructure. In this way, many models have been derived from each other.

As a result of the survey carried out in four different categories on the 'Web of Science' database by Thomson Reuters (agriculture dairy animal science, veterinary science, agriculture multidisciplinary, and mathematical computational biology) and covering the years from 1970 to 2016, it was determined that Gompertz was the most commonly used growth model in the publications on these subjects (*Figure 1*).



**Figure 1** Numbers of growth curve models according to database of Web of Science.

The non-linear regression models which are most widely used to model growth in the poultry are the three-parameter Gompertz, Logistic, and von Bertalanffy models and the four-parameter Richards functions. On the other hand, other models which are less mentioned in scientific studies are von Bertalanffy, Weibull, Monomolecular, France, Michaelis-Menten, and Morgan-Mercer-Flodin. The expressions of the Gompertz, Logistic, von Bertalanffy and Richards growth models and their coordinates of the point of inflection are presented in *Table 1*.

**Table 1** Expressions and point of inflections of commonly used growth functions.

| Growth Model    | Equation  | IPT                                 | IPW  |
|-----------------|---|-------------------------------------|--|
| Gompertz        | $Y_t = \beta_0 \cdot e^{-\beta_1 e^{-\beta_2 t}}$           | $\ln(\beta_1)/\beta_2$              | $\beta_0/e$                                |
| Logistic        | $Y_t = \beta_0(1 + \beta_1 \cdot e^{-\beta_2 t})^{-1}$      | $-\ln(1/\beta_1)/\beta_2$           | $\beta_0/2$                                |
| Richards        | $Y_t = \beta_0(1 + \beta_1 \cdot e^{-\beta_2 t})^{\beta_3}$ | $\beta_2^{-1} \ln(\beta_3 \beta_1)$ | $\beta_0((\beta_3 - 1)/\beta_3)^{\beta_3}$ |
| Von Bertalanffy | $Y_t = \beta_0(1 - \beta_1 \cdot e^{-\beta_2 t})^3$         | $\ln(3\beta_1)/\beta_2$             | $8\beta_0/27$                              |

In the equations, 't' denotes time, 'y' weight, 'β0' the maximum body weight the animal is assumed to be able to reach, 'β1' the biological constant about the shape of the curve, 'β2' the biological constant about the growth rate and 'β3' the shape parameter. These parameter models are special cases of the more flexible Richards function, which a variable point of inflection is specified by the shape parameter (β3). The Logistic, Gompertz and von Bertalanffy functions have fixed growth forms with the point of inflection at about 50, 37 and 30% of the asymptote, respectively (Porter *et al.*, 2010). It is possible to examine the studies on poultry by using growth models under three categories, namely 'determination of the best-fitting model', 'a comparison of various application results with the growth models', and 'the genetic structure of the growth curve'.

## Determination of the best-fitting growth model

The goodness-of-fit criteria used widely for non-linear regression equations are presented in Table 2 (Nariñ et al., 2010b; Karadas et al., 2017).

Table 2 Some goodness of fit statistics.

| Goodness of Fit Criteria           | Formula  |
|------------------------------------|--|
| Determination Coefficient,         | $R^2 = 1 - \left( \frac{SSE}{SST} \right)$   |
| Adjusted Determination Coefficient | $\bar{R}^2 = R^2 - \left[ \left( \frac{k-1}{n-k} \right) (1 - R^2) \right]$  |
| Mean Square Error                  | $MSE = \frac{SSE}{(n-k)}$  |
| Akaike's Information Criterion     | $AIC = n \cdot \ln \left( \frac{SSE}{n} \right) + 2k$  |
| Bayesian Information Criterion     | $BIC = n \cdot \ln \left( \frac{SSE}{n} \right) + k \cdot \ln(n)$  |
| Standard Deviation Ratio           | $SD_{Rat} = \frac{\sqrt{\frac{1}{n-1} \sum (\varepsilon - \bar{\varepsilon})^2}}{\sqrt{\frac{1}{n-1} \sum (Y - \bar{Y})^2}}$ |

SSE: sum of square errors, SST: total sum of squares, n: the number of observations, k: the number of parameters,  $\varepsilon$ : actual residual value, Y: actual yield value

There are a large number of studies which were performed to determine the best-fitting growth models in different poultry species or genotypic groups in the same species. In a pioneering study, Ricklefs (1967) determined that the best-fitting model was Gompertz for growth data of some birds. Sigmoid, Logistic, and polynomial models have been fitted to growth curves of chickens (Grossman and Bohren, 1982). Tzeng and Becker (1981) analysed the growth data of male chickens from a purebred sire line using Gompertz, Logistic and von Bertalanffy functions. They reported that Gompertz was the best-fitting model. Having carried out a similar study, Anthony *et al.* (1986) reported, as a result of their analyses with the growth data of the quail lines treated with divergent selection, that the best-fitting model for the line with decreased weight was Logistic and that the best-fitting model for the line with increased weight was Gompertz. Having analysed the growth data of turkeys selected for increased body weight and its randombred control line by means of Gompertz, Logistic and von Bertalanffy, Anthony *et al.* (1991a) reported that the best-fitting model was Gompertz. Akbaş and Oğuz (1998), who analysed the growth patterns of Japanese quail lines selected for low and high body weight, reported that the best-fitting model was Gompertz. Maruyama *et al.* (1998) analysed the growth data of male and female turkeys of two genetic lines (a fast-growing line and a randombred control line) by means of the Logistic, Gompertz, von Bertalanffy, Richards, Weibull, and Morgan-Mercer-Flodin functions. In a study by Yakupoglu and Atil (2001), the values of weekly body weight in Cobb and Hubbard commercial broiler flocks were analysed using Gompertz and von Bertalanffy functions, and they reported that the Gompertz model explained outcomes better than von Bertalanffy.

Aggrey (2002) compared three non-linear models (Richards, Logistic, and Gompertz) and a spline linear regression model for describing chicken growth curves. The spline model predicted the hatching weight better than the Gompertz model; however, the spline model had the poorest fit to the data compared with the non-linear models. Narushin and Takma (2003) carried out a study to determine the best predictive model for the accurate description of the average flock growth of laying hens (Shaver white laying hen line) using Logistic, Gompertz, von Bertalanffy, Richards, Weibull and Morgan-Mercer-Flodin functions, polynomial regression and Narushin-Takma models. The researchers found that the Narushin-Takma models they had developed themselves had functions which well-defined both growth data and egg yields. However, these empirical models with no biologically significant parameters were not used much in the studies carried out to model growth in the poultry. Having analysed the growth data of meat and egg strains of chicken by using the Gompertz, Logistic, Lopez, Richards, France, and von Bertalanffy models, Darmani Kuhi *et al.* (2003) reported that the Richards model was better than the others according to various goodness-of-fit criteria. Darmani Kuhi *et al.* (2003) stated that it would be more useful to use flexible models like Richards instead of models with a fixed point of inflection such as Gompertz and Logistic. Growth curve parameters were estimated for large white turkey flock by Sengul and Kiraz (2005) with four different non-linear models (Gompertz, Logistic, Morgan-Mercer-Flodin, and Richards). They reported that Gompertz, Logistic and Richards models were more suitable models for growth data of large white turkey. Flexible non-linear models were evaluated statistically for their suitability in predicting weights of carcass parts by Zuidhof (2005). The analysis employed four sigmoidal (S) models (Gompertz, modified Gompertz, Richards, and Lopez) describing carcass part weight as a function of age, as well as three diminishing returns models (Lopez, Mitscherlich, and log linear), and a log-linear proportional yield model, which described carcass part yield and weight. Zuidhof (2005) reported that the sigmoidal models with a flexible point of inflection were better able to predict carcass part weights than the Gompertz model with a fixed point of inflection and the diminishing returns models. In a study carried out to model two sets of growth data from the literature, male broiler body weight at 168 and 170 days of age were used (Roush and Branton, 2005). Researchers analysed growth data using two forms of Logistic model, two forms of Gompertz model, and the Saturated Kinetic models via the non-linear regression and the genetic algorithm approaches. The genetic algorithm approach was found to successfully determine the coefficients of growth equations (Roush and Branton, 2005). Roush *et al.* (2006), published the findings of the first use of the neural network model in poultry (growth data of male chicks from Ross x Ross 308) in order to estimate growth curve parameters. A comparison was made for the Gompertz function between the non-linear regression and neural network methodologies. Accuracy of the models was determined by mean square error, mean absolute deviation, mean absolute percentage error, and bias. When measured by bias, the Gompertz equation underestimated values, whereas the genetic algorithm produced little or no overestimation of the observed body weight responses (Roush *et al.*, 2006). Nahashon *et al.* (2006a) compared two non-linear mathematical functions (Gompertz and Logistic) using weekly body weight data of the French guinea fowl. The Gompertz model was found to be better than the Logistic model for growth patterns in French guinea fowl. Nahashon *et al.* (2006b) compared Richards, Gompertz, and Logistic functions for describing the growth of the pearl gray Guinea fowl using weekly live weight data from hatch to 22 weeks of age. It was determined that the Logistic model was not compatible with the data of pearl gray guinea fowl compared with the Gompertz and Richards models. In another study by Norris *et al.* (2007), the Gompertz, Logistic and Richards models were compared for the live weight of

indigenous Venda and Naked Neck chickens and analyses was carried out to determine the existence of differences in the growth patterns of these breeds. The Gompertz model was most suitable for explaining the growth of the chickens. Ahmadi and Mottaghtalab (2007) evolved three growth models out of a family of flexible curve ideas, the so called hyperbolic models. The researchers reported that the greater flexibility of a hyperbolic model and Richards function may lead to more accurate prediction and better fit to the broiler growth data than Gompertz. Balcioglu *et al.* (2009) compared the growth models (Gompertz, Bertalanffy, Logistic functions) using data from chukar partridges (*Alectoris chukar*). According to the goodness-of-fit criteria, the Gompertz model was found to be the best to determine the growth pattern of partridges. In another study, three flexible growth functions (Von Bertalanffy, Richards, and Morgan) were compared with the Gompertz function for growth profiles of turkey hens from commercial flocks (Porter *et al.*, 2010). The results showed that the fixed point of inflection of the Gompertz equation can be a limitation and that the relationship between body weight and age in turkeys was best described using flexible growth functions. The Morgan equation provided the best fit to the data set and was used for characterising growth profiles of turkey hens. Four non-linear models (Gompertz, Logistic, Richards, and Weibull) and a spline linear regression model were evaluated with regard to their ability to describe the growth pattern of male mule duck (Vitezica *et al.*, 2010). Furthermore, fixed and mixed effects models were compared to analyse growth, and of the non-linear models, the mixed effects Weibull model had the best overall fit, with the mixed effects spline regression model being the second best model. Nariñç *et al.* (2010b) compared ten growth functions (Gompertz, Richards, Logistic, Bertalanffy, Brody, Negative Exponential, Morgan-Mercer-Flodin and three Hyperbolic models) for growth data of a randombred Japanese quail flock. The Gompertz model was found to be the best for Japanese quail among all the candidate models according to the goodness-of-fit criteria. Tompić *et al.* (2011) compared three non-linear growth functions (Logistic, Gompertz, and Richards) and three polynomial functions (linear, second-order, and third-order) for describing the growth profiles of Ross 308 broiler breeder flocks from hatch to 35 weeks of age.

The third-order polynomial function gave the optimal results for body weight data of Ross 308 parents, when the Richards model was the best fitted function among the non-linear models. Ramos *et al.* (2013) investigated the most appropriate growth model for the body weights of ostriches in a Brazilian population. Two non-linear growth functions (Gompertz and Logistic) and a third-order polynomial function were utilised. The results of this study showed that the non-linear models were more compatible for describing ostrich growth. In another study (Rizzi *et al.*, 2013), the linear and three non-linear models (Gompertz, Logistic and Richards) were compared for growth patterns of five local Italian chicken lines. Researchers reported that Richards model was determined to be the most suitable for explaining the differences of growth patterns of the local lines (Rizzi *et al.*, 2013). Zhao *et al.* (2015) compared three growth functions (Logistics, Gompertz, and von Bertalanffy) and evaluated the relationship between body weight and age in growing three types of indigenous chicken in China (Shaobo, Huaixiang, and Youxi). The authors reported that the Gompertz model was the best due to less bias.



## A comparison of growth in poultry species or in various experimental groups

One of the studies in which growth samples were compared was performed by Anthony *et al.* (1991b), where quail, chicken and turkey species were used, and the parameters of the Gompertz model gave the best model according to some goodness-of-fit criteria. The effects of selection on the point of inflection in the species concerned were compared in the same study. Once the growth samples in chicken, turkey, duck, and goose species were analysed by using the Richards function, Knizetova *et al.* (1995) reported that the ages at the points of inflection were 47.7, 74.0, 25.5, and 21.1 days, respectively and stated that there were differences among the species in terms of other parameters as well. In the studies performed by Knizetova *et al.* (1991a; 1991b; 1994), the growth samples in eight broiler, nine duck, and three goose lines were analysed by using the Richards function, and growth differences among the lines and between the sexes were presented. In addition to these studies, there are others where the growth of different lines or genetic groups in different poultry species were compared with the growth curves. In the studies where the Gompertz model was used Scheuermann *et al.* (2003) used data from a total of eight broiler lines and crosses; Marcato *et al.* (2008) analysed two commercial broiler lines (Cobb and Ross); and Sakomura *et al.* (2011) compared the growth of 56-day major body components of two commercial broiler lines (Cobb and Ross). Pis (2012) analysed the growth patterns of the grey partridge (*Perdix perdix*) and chukar (*Alectoris chukar*) from hatch to 120 days of age. The growth rate and development of body mass, wing, tarsus, and bill length were measured and fitted by Gompertz function. The growth of body weight, breast weight, and leg weight in the male broiler chickens of two commercial lines were analysed using the Richards function by Goliomytis *et al.* (2003). Sezer and Tarhan (2005) presented the growth curves in three meat type quail lines and Ersoy *et al.* (2006) revealed the growth differences by sex in bronze turkeys using the Richards function.

Individual phenotypic selection for high and low body weights at four weeks of age was conducted for a long term in Japanese quail under two selection environments (Marks, 1978).

The growth patterns of four experimental lines have been analysed using the Logistic function. Marks (1978) reported that growth curve parameters were changed by selection. Selection for high body weight increased the relative growth rate and, the inflection point was shifted to a younger age (Marks, 1978). The growth patterns of a line of turkeys selected for increased body weight and its randombred control line were compared via the Gompertz growth model (Anthony *et al.*, 1991b). It was concluded that the selection line had a higher slope at inflection point and an asymptotic body weight than the control line. The age at point of inflection and  $\beta_0$  parameter were achieved later for the control flock. Likewise, there are other studies in which the differences among the growth samples of different lines obtained via selection experiments were presented by using growth functions. Akbaş and Oğuz (1998) compared Gompertz, Logistic and von Bertalanffy for the growth patterns of a long term selected line and its control line in Japanese quail. They reported that the mature weight parameter ( $\beta_0$ ) of the birds increased by 15% as a result of five-generation selection in order to increase body weight at five weeks of age. A similar study was performed by Rezvannejad *et al.* (2013) using divergent lines of Japanese quail and their crosses. They utilised Gompertz, Logistic, and Richards functions and reported that the Logistic model described the growth patterns of Japanese quail and their crosses better than other models. Mignon-Grasteau *et al.* (2000) used the growth data of five meat type chicken lines to investigate the effects of selection on the parameters of the Gompertz model. Having

examined growth in four selected duck lines, Maruyama *et al.* (2001) determined the differences in the traits of adult weight, age at the point of inflection and growth rate among the lines with the Weibull growth function. Aggrey *et al.* (2003) examined the growth samples of females and males in two lines and a control line obtained as a result of long-term divergent selection in Japanese quail by using the Richards function. The parameter of mature weight ( $\beta_0$ ) was reported to have increased by 104% in males of the line with increased weight as a result of 30-generation selection in quail (Aggrey *et al.*, 2003). It was also stated by other researchers that the variations of weight resulting from short-term selection in quail populations affected the parameters of Richards (Hyankova *et al.*, 2001; Beiki *et al.*, 2013) and Gompertz (Anthony *et al.*, 1986; Balcioğlu *et al.*, 2005; Alkan *et al.*, 2009) growth curves.

There are studies in which the effects of various environmental (feeding or management) applications applied in different poultry species have been examined using growth curves. Having raised two different turkey genotypes (Nicholas and British United Turkeys) under two lighting schedules and two feeding schedules, Lilburn *et al.* (1992; 1993) compared the growth samples of the experimental groups by means of the Gompertz model. It was concluded that the slope of the growth curve was significantly less in British United Turkeys, but the weight and age at the point of inflection were increased compared with Nicholas toms. There were no significant diet effects on any growth curve parameters (Lilburn *et al.*, 1993). In research where they used a commercial broiler line and two domestic genotypes (Barred Plymouth Rock and Jærhøn), Ali and Brenoe (2002) applied feed restriction and *ad libitum* feeding schedules and analysed the growth samples of the experimental groups by means of the Gompertz model. In work by Schinckel *et al.* (2005), an analysis was made by means of the Weibull model and non-linear mixed effects by using the weekly body weight data of the ducks fed with six different diets. Having applied two different feed restriction schedules in Ross 308 broilers, Mendes *et al.* (2007) examined the differences between the values of body mass index of the experimental groups and the control group chicks fed *ad libitum* by means of the Gompertz growth model. According to the maximal week increment values estimated with the Gompertz model, they put forward that fat deposition would be higher in the chicks fed *ad libitum*. Alkan *et al.* (2012), who applied a high temperature during the embryonic period to provide the Japanese quail flock with the ability to cope with the heat stress, compared the growth samples of treatment groups by means of the Gompertz model. It was determined that the thermal manipulation treatment created a difference in the growth curves of the quail. It was found that thermal manipulation in the early period caused a lower mature weight and led to an increase in the rate of maturation. On the other hand, it was observed that thermal manipulation in the later period did not lead to any difference in the growth parameters compared to the control group.

There are few studies that have been performed with respect to analysis in slow-growing broilers managed in alternative rearing systems. Santos *et al.* (2005), Nariñç *et al.* (2010c), and Eleroğlu *et al.* (2014) performed analyses for growth data of slow-growing broiler lines reared in indoors, semi-extensive and organic systems, respectively. N'Dri *et al.* (2006) estimated parameters of the Gompertz function for slow-growing broilers reared under the label rouge system. Dottavio *et al.* (2007) and Dourado *et al.* (2009) used the Gompertz model to examine the growth of slow-growing broilers (SASSO and ISA Label) reared in the free range system.



## Genetic parameter estimates for growth curves

The number of studies with estimated genetic parameters for growth models in poultry is rather limited. This is mainly because body weights at fixed ages were used as the selection criteria in almost all the genetic improvement studies on poultry growth. There is no selection study where parameters such as the growth curve parameters or the point of inflection are used in the poultry.

As a result of the analyses made for the growth curves of broilers by Grossman and Bohren (1985) and Barbato (1991), it was revealed that the main parameters were traits with a high heritability. In another study in chickens (Mignon-Grasteau *et al.*, 1999), the heritabilities of the Gompertz growth curve parameters were found to be moderate to high and ranged from 0.31 to 0.54. Mignon-Grasteau *et al.* (2000) carried out Gibbs sampling estimations for the genetic parameter estimates of the Gompertz growth curve parameters on 10,671 male and female chickens originating from five lines. It was reported that the heritabilities were high for all three parameters (ranging from 0.43 to 0.60). The genetic correlation of  $\beta_0$  with  $\beta_1$  was highly positive, whereas the genetic correlations of  $\beta_2$  and  $\beta_0$  or  $\beta_1$  were estimated as moderately negative. N'Dri *et al.* (2006) reported genetic relationships among feed conversion ratio, using Gompertz growth curve parameters and carcass composition traits for 1,061 slow-growing chickens. The growth curve parameters were found moderately heritable (0.25 to 0.34). Genetic correlations between growth curve parameters and either feed conversion ratio or residual feed consumption were estimated to be low to moderate (0.31 to 0.51).

There are some studies for genetic parameter estimates of growth curve parameters in Japanese quail. But the researchers used only Gompertz function in these studies. Heritability estimates for the mature weight parameter ( $\beta_0$ ) were found mostly high (from 0.38 to 0.60) by Gebhardt-Henrich and Marks (1993), Akbaş and Oğuz (1998), and Nariñç *et al.* (2010a; 2014). However, the  $\beta_0$  parameter is a lower heritable trait according to Akbaş and Yaylak (2000), they estimated a low heritability (0.18) for this trait from the Gompertz model. For parameter  $\beta_1$ , the heritability was estimated as 0.21 by Nariñç *et al.* (2010a). In parallel with this, its heritability was low to moderate levels in the other studies (Akbaş and Oğuz, 1998; Akbaş and Yaylak, 2000). The heritability for the parameter of instantaneous growth rate ( $\beta_2$ ) was estimated as 0.38 by Akbaş and Oğuz (1998), and 0.32 by Akbaş and Yaylak (2000), and it has been reported that the heritability estimations of the traits of age and weight at the point of inflection were also at low to moderate levels. Similar results were also reported by Nariñç *et al.* (2010a), who estimated the heritabilities for the age and weight at the point of inflection as 0.08 and 0.36, respectively.

Aslam *et al.* (2011), analysed the growth data of two commercial turkey parent flocks by using the Logistic function, and reported that the heritability estimated for parameter  $\beta_0$  was 0.30, while the other parameters had low levels (0.05–0.11). Genetic correlations between the growth curve parameters and various slaughter-carcass traits and meat quality traits were estimated in the same research.

## Conclusions

For many years researchers have used models to analyse growth in poultry. Most of the models used are of sigmoidal structure, while some of them have fixed or flexible structures in terms of point of inflection. In a recent study (Darmani Kuhi *et al.*, 2010), it was stated that the equations with a fixed point of inflection such as the Gompertz and Logistic may cause limitations, whereas the use of the models with a

flexible inflection point, such as Richards, can give better results. This view is widely held; however, Gompertz and Logistic have been the two most used models in the studies performed so far. In this review, the studies concerned have been summarised under the titles ‘determination of the best-fitting growth model’, ‘a comparison of the growth of poultry species or various experimental groups’, and ‘genetic parameter estimates for growth curve parameters’. When the variation of the studies by year is examined, it is possible to state that the studies to determine the best-fitting model for different poultry species have decreased. Furthermore, there is a need for studies to transform the existing growth models into a more mechanistic form. It has been seen that there was an increase in the studies concerned upon the use of different algorithms (e.g. artificial neural networks, data mining, and the Bayesian framework) in growth modelling particularly for the last decade, and there are only a limited number of published studies where the methods concerned are used for growth in poultry. It is estimated that there will be an increase in the use of non-linear mixed effects models in the studies to be performed to compare the growth of different groups (e.g. sex, genotype, and various experiments). It is necessary to have studies on the utilisation of growth curves in the inheritance of growth curve parameters and genetic-genomic selection studies in different poultry species.

## References

- AGGREY, S.E. (2002) Comparison of three nonlinear and spline regression models for describing chicken growth curves. *Poultry Science* **81**: 1782-1788.
- AGGREY, S.E., ANKRA-BADU, G.A. and MARKS, H.L. (2003) Effect of long-term selection on growth characteristics in Japanese quail. *Poultry Science* **82**: 538-542.
- AHMADI, H. and MOTTAGHITALAB, M. (2007) Hyperbolic models as a new powerful tool to describe broiler growth kinetics. *Poultry Science* **86**: 2461-2465.
- AKBAŞ, Y. and OĞUZ, I. (1998) Growth curve parameters of lines of Japanese quail (*Coturnix coturnix japonica*), unselected and selected for four-week body weight. *Archiv Für Gefügelkunde* **62**: 104-109.
- AKBAŞ, Y. and YAYLAK, E. (2000) Heritability estimates of growth curve parameters and genetic correlations between the growth curve parameters and weights at different age of Japanese quail. *Archiv Für Gefügelkunde* **64**: 141-146.
- ALI, K.O. and BRENOE, U.T. (2002) Comparing genotypes of different body sizes for growth-related traits in chickens. I. Live weight and growth performance under intensive and feed-restricted extensive systems. *Acta Agricultural Scandinavica, Section A. - Animal Science* **52**: 1-10.
- ALKAN, S., MENDES, M., KARABAG, K. and BALCIOĞLU, M.S. (2009) Effects short term divergent selection for 5-week body weight on growth characteristics in Japanese quail. *Archiv Für Gefügelkunde* **73**: 124-131.
- ALKAN, S., NARİNC, D., KARSLI, T., KARABAĞ, K., BALCIOĞLU, M.S. (2012) Effects of thermal manipulations during early and late embryogenesis on growth characteristics in Japanese quails (*Coturnix coturnix japonica*). *Archiv Für Gefügelkunde* **76**: 184-190.
- ANTHONY, N.B., NESTOR, K.E. and BACON, W.L. (1986) Growth curves of Japanese quail as modified by divergent selection for 4-week body weight. *Poultry Science* **65**: 1825-1833.
- ANTHONY, N.B., EMMERSON, D.A. and NESTOR, K.E. (1991a) Research Note: Influence of body weight selection on the growth curve of Turkeys. *Poultry Science* **70**: 192-194.
- ANTHONY, N.B., EMMERSON, D.A., NESTOR, K.E. and BACON, W.L. (1991b) Comparison of growth curves of weight selected populations of turkeys, quail and chickens. *Poultry Science* **70**: 13-19.
- ASLAM, M., BASTIAANSEN, J., CROOIJMANS, R., DUCRO, B., VEREIJKEN, A. and GROENEN, M. (2011) Genetic variances, heritabilities and maternal effects on body weight, breast meat yield, meat quality traits and the shape of the growth curve in turkey birds. *BMC Genetics* **12**: 14-23.
- BALCIOĞLU, M.S., KIZILKAYA, K., YOLCU, H.I. and KARABAĞ, K. (2005) Analysis of growth characteristics in short-term divergently selected Japanese quail. *South African Journal of Animal Science* **35**: 83-89.
- BALCIOĞLU, M.S., KIZILKAYA, K., KARABAĞ, K., ALKAN, S., YOLCU, H.I. and ŞAHİN, E. (2009) Comparison of growth characteristics of chukar partridges (*Alectoris chukar*) raised in captivity. *Journal of Applied Animal Research* **35**: 21-24.

- BARBATO, G.F. (1991) Genetic architecture of growth curve parameters in chickens. *Theoretical and Applied Genetics* **83**: 24-32.
- BEIKI, H., PAKDEL, A., MORADI-SHAHRBABA, M. and MEHRBAN, H. (2013) Evaluation of growth functions on Japanese quail lines. *The Journal of Poultry Science* **50**: 20-27
- DARMANI KUH, H., KEBREAB, E., LOPEZ, S. and FRANCE, J. (2003) An evaluation of different growth functions for describing the profile of live weight with time (age) in meat and egg strains of chicken. *Poultry Science* **82**: 1536-1543.
- DARMANI KUH, H., PORTER, T., LOPEZ, S., KEBREAB, E., STRATHE, A.B., DUMAS, A., DIJKSTRA, J. and FRANCE, J. (2010) A review of mathematical functions for the analysis of growth in poultry. *World's Poultry Science Journal* **66**: 227-239.
- DOTTAIO, A.M., ÁLVAREZ, M., CANET, Z.E., FONT, M.T. and DI MASSO, R.J. (2007) Growth pattern of experimental hybrids for free range broiler production. *Revista Argentina de Producción Animal* **27**: 75-82.
- DOURADO, L.R.B., SAKOMURA, N.K., NASCIMENTO, D.C.N., DORIGAM, J.C., MARCATO, S.M. and FERNANDES, J.B.K. (2009) Growth and performance of naked neck broiler reared in free-range system. *Ciência e Agrotecnologia* **33**: 875-881.
- ELEROĞLU, H., YILDIRIM, A., ŞEKEROĞLU, A., ÇOKSÖYLER, F.N. and DUMAN, M. (2014) Comparison of growth curves by growth models in slow-growing chicken genotypes raised the organic system. *International Journal of Agriculture and Biology* **16**: 529-535.
- ERSOY, I.E., MENDEŞ, M. and AKTAN, S. (2006) Growth Curve Establishment for American Bronze Turkeys. *Archiv Tierzucht* **49**: 293-299.
- FRANCE, J. and DIJKSTRA, J. (2006) Scientific progress and mathematical modelling: Different approaches to modelling animal systems, in: GOUS, R. FISHER, C. & MORRIS, T.R. (Eds) *Mechanistic Modelling in Pig and Poultry Production*, pp. 6-22 (CAB International, New York).
- GEHARDT-HENRICH, S.G. and MARKS, H.L. (1993) Heritability of growth curve parameters and age-specific expression of genetic variation under two different feeding regimes in Japanese quail (*Coturnix coturnix japonica*). *Genetical Research* **62**: 45-55.
- GOLIOMYTIS, M., PANOPULOU, E. and ROGDAKIS, E. (2003) Growth curves for body weight and major component parts, feed consumption, and mortality of male broiler chickens raised to maturity. *Poultry Science* **82**: 1061-1068.
- GROSSMAN, M. and BOHREN, B.B. (1982) Comparison of proposed growth curve functions in chickens. *Growth* **46**: 259-274.
- GROSSMAN, M. and BOHREN, B.B. (1985) Logistic growth curve of chickens: heritability of parameters. *Journal of Heredity* **76**: 459-462.
- HYANKOVA, L., KNIZETOVA, H., DEDKOVA, L. and HORT, J. (2001) Divergent selection shape of growth curve in Japanese quail 1. Responses in Growth parameters and food conversion. *British Poultry Science* **42**: 583-589.
- KARADAS, K., TARIQ, M., TARIQ, M.M. and EYDURAN, E. (2017) Measuring predictive performance of data mining and artificial neural network algorithms for predicting lactation milk yield in indigenous Akkaraman Sheep. *Pakistan Journal of Zoology* **49**: 1-7.
- KNIZETOVA, H., HYANEK, J., KNIZE, B. and ROUBICEK, J. (1991a) Analysis of growth curves in fowl. I. Chickens. *British Poultry Science* **32**: 1027-1038.
- KNIZETOVA, H., HYANEK, J., KNIZE, B. and PROCHAZKOVA, H. (1991b) Analysis of growth curves in fowl. II. Ducks. *British Poultry Science* **32**: 1039-1053.
- KNIZETOVA, H., HYANEK, J. and VESELSKY, A. (1994) Analysis of growth curves in fowl. III. Geese. *British Poultry Science* **35**: 335-344.
- KNIZETOVA, H., HYANEK, J., HYANKOVA, L. and BELICEK, P. (1995) Comparative study of growth curves in poultry. *Genetics Selection Evolution* **27**: 365-375.
- KONARZEWSKI, M., KOIJMAN, S.A.L.M. and RICKLEFS, R.E. (1998) Models for avian growth and development, in: STARCK, J.M. & RICKLEFS, R.E. (Eds) *Avian Growth and Development: Evolution within the Altricial-precocial Spectrum*, pp.340-361 (Oxford University Press, New York).
- LILBURN, M.S., RENNER, P.A. and ANTHONY, N.B. (1992) Interaction between step-up versus step-down lighting from four to six teen weeks on growth and development in turkey hens from two commercial strains. *Poultry Science* **71**: 419-426.
- LILBURN, M.S. and EMMERSON, D. (1993) The influence of differences of dietary amino acids during the early growing period on growth and development of Nicholas and British United Turkey toms. *Poultry Science* **72**: 1722-1730.
- MARCATO, S.M., SAKOMURA, N.K., MUNARI, D.P., FERNANDES, J.B.K., KAWAUCHI, I.M. and BONATO, M.A. (2008) Growth and Body Nutrient Deposition of Two Broiler Commercial Genetic Lines. *Brazilian Journal of Poultry Science* **10**: 117-123.
- MARKS, H.L. (1978) Growth curve changes associated with long-term selection for body weight in Japanese quail. *Growth* **42**: 129-140.

- MARUYAMA, K., POTTS, W.J.E., BACON, W.L. and NESTOR, K.E. (1998) Modeling turkey growth with the relative growth rate. *Growth Development and Aging* **62**: 123-139.
- MARUYAMA, K., VINYARD, B., AKBAR, M.K., SHAFER, D.J. and TURK, C.M. (2001) Growth curve analyses in selected duck lines. *British Poultry Science* **42**: 574-582.
- MENDES, M., DINCER, E. and ARSLAN, E. (2007) Profile Analysis and Growth Curve for Body Mass Index of Broiler Chickens Reared Under Different Feed Restrictions in Early Age. *Archiv für Tierzucht* **50**: 403-411.
- MIGNON-GRASTEAU, S., BEAUMONT, C., LE BIHAN-DUVAL, E., POIVEY, J.P., DE ROCHAMBEAU, H. and RICHARD, F.H. (1999) Genetic parameters of growth curve parameters in male and female chickens. *British Poultry Science* **40**: 44-51.
- MIGNON-GRASTEAU, S., PILES, M., VARONA, L., DE ROCHAMBEAU, H., POIVEY, J.P., BLASCO, A. and BEAUMONT, C. (2000) Genetic analysis of growth curve parameters for male and female chickens resulting from selection on shape of growth curve. *Journal of Animal Science* **78**: 2515-2524.
- NAHASHON, S.N., AGGREY, S.E., ADEFOPE, N.A. and AMENYENU, A. (2006a). Modeling growth characteristics of meat-type guinea fowl. *Poultry Science* **85**: 943-946.
- NAHASHON, S.N., AGGREY, S.E., ADEFOPE, N.A., AMENYENU, A. and WRIGHT, D. (2006b). Growth Characteristics of Pearl Gray Guinea Fowl as Predicted by the Richards, Gompertz, and Logistic Models. *Poultry Science* **85**: 359-363.
- NARINÇ, D., AKSOY, T. and KARAMAN, E. (2010a) Genetic parameters of growth curve parameters and weekly body weights in Japanese quail. *Journal of Animal and Veterinary Advances* **9**: 501-507.
- NARINÇ, D., KARAMAN, E., FIRAT, M.Z. and AKSOY, T. (2010b) Comparison of non-linear growth models to describe the growth in Japanese quail. *Journal of Animal and Veterinary Advances* **9**: 1961-1966.
- NARINÇ, D., KARAMAN, E., AKSOY T. and DENİZ, Ç. (2010c) Analysis of fitting growth models in medium growing chicken raised indoor system. *Trends Animal and Veterinary Science Journal* **1**: 12-18.
- NARINÇ, D., KARAMAN, E., AKSOY, T. and FIRAT, M.Z. (2014) Genetic parameter estimates of growth curve and reproduction traits in Japanese quail. *Poultry Science* **93**: 24-30.
- NARUSHIN, V.G. and TAKMA, C. (2003) Sigmoid model for the evaluation of growth and production curves in laying hens. *Biosystems Engineering* **84**: 343-348.
- N'DRI, A.L., MIGNON-GRASTEAU, S., SELLIER, S., TIXIER-BOICHARD, M. and BEAUMONT, C. (2006) Genetic relationships between feed conversion ratio, growth curve and body composition in slow growing chickens. *British Poultry Science* **47**: 273-280.
- NORRIS, D., NGAMBI, J.W., BENYI, K., MAKGAHLELA, M.L., SHIMELIS, H.A. and NESAMVUNI, E.A. (2007) Analysis of growth curves of indigenous male Venda and Naked Neck chickens. *South African Journal of Animal Science* **37**: 21-26.
- PANIK, M.J. (2014) Growth Curve Modeling: Theory and Applications. John Wiley & Sons, USA.
- PIS, T. (2012) Growth and development of chicks of two species of partridge: the Grey partridge (*Perdix perdix*) and the chukar (*Alectoris chukar*). *British Poultry Science* **53**: 141-144.
- PORTER, T., KEBREAB, E., DARMANI KUHI, H., LOPEZ, S., STRATHE, A.B. and FRANCE, J. (2010) Flexible alternatives to the Gompertz equation for describing growth with age in turkey hens. *Poultry Science* **89**: 371-378.
- RAMOS, S.B., CAETANO, S.L., SAVEGNAGO, R.P., NUNES, B.N., RAMOS, A.A. and MUNARI, D.P. (2013) Growth curves for ostriches (*Struthiocamelus*) in a Brazilian population. *Poultry Science* **92**: 277-282.
- REZVANNEJAD, E., PAKDEL, A., MIRAE ASHTIANEE, S.R., MEHRABANI YEGANEH, H. and YAGHOobi, M.M. (2013) Analysis of growth characteristics in short-term divergently selected Japanese quail lines and their cross. *The Journal of Applied Poultry Research* **22**: 663-670.
- RICKLEFS, R.E. (1967) Relative growth, body constituents, and energy content of nestling barn swallows and red-winged blackbirds. *The Auk* **84**: 560-570.
- RIZZI, C., CONTIERO, B. and CASSANDRO, M. (2013) Growth patterns of Italian local chicken populations. *Poultry Science* **92**: 2226-2235.
- ROUSH, W.B. and BRANTON, S.L. (2005) A comparison of fitting growth models with a genetic algorithm and nonlinear regression. *Poultry Science* **84**: 494-502.
- ROUSH, W.B., DOZIER 3<sup>rd</sup>, W.A. and BRANTON, S.L. (2006) Comparison of Gompertz and Neural Network models of broiler growth. *Poultry Science* **85**: 794-797.
- SAKOMURA, N.K., GOUS, R.M., MARCATO, S.M. and FERNANDES, J.B.K. (2011) A description of the growth of the major body components of 2 broiler chicken strains. *Poultry Science* **90**: 2888-2896.
- SANTOS, A.L., SAKOMURA, N.K., FREITAS, E.R., FORTES, C.M.S. and CARRILHO, E.N.V.M. (2005) Comparison of free range broiler chicken strains raised in confined or semi-confined systems. *Revista Brasileira de CiênciaAvícola* **7**: 85-92.
- SCHEUERMANN, G.N., BILGILI, S.F., HESS, J.B. and MULVANEY, D.R. (2003) Breast muscle development in commercial broiler chickens. *Poultry Science* **82**: 1648-1658.

- SCHINCKEL, A.P., ADEOLA, O. and EINSTEIN, M.E. (2005) Evaluation of alternative nonlinear mixed effects models of duck growth. *Poultry Science* **84**: 256-264.
- SENGUL, T. and KIRAZ, S. (2005) Non-linear models of growth curves in large white turkeys. *Turkish Journal of Veterinary and Animal Sciences* **29**: 331-337.
- SEZER, M. and TARHAN, S. (2005) Model parameters of growth curves of three meat-type lines of Japanese quail. *Czech Journal of Animal Science* **50**: 22-30.
- THORNLEY, J.H.M. and FRANCE, J. (2007) Mathematical models in agriculture: quantitative methods for the plant, animal and ecological sciences. Second Edition. Wallingford, UK: CABI Publishing, 923 pp.
- TOMPIĆ, T., DOBŠA, J., LEGEN, S., TOMPIĆ, N. and MEDIĆ, H. (2011) Modeling the growth pattern of in-season and off-season Ross 308 broiler breeder flocks. *Poultry Science* **90**: 2879-2887.
- TZENG, R.Y. and BECKER, W.A. (1981) Growth patterns of body and abdominal fat weight in male broiler chickens. *Poultry Science* **60**: 1101-1106.
- VITEZICA, Z.G., MARIE-ETANCELIN, C., BERNADET, M.D., FERNANDEZ, X. and ROBERT-GRANIE, C. (2010) Comparison of nonlinear and spline regression models for describing mule duck growth curves. *Poultry Science* **89**: 1778-1784.
- YAKUPOGLU, C. and ATIL, H. (2001) Comparison of growth curve models on broilers growth curve I: Parameters estimation. *Online Journal of Biological Sciences* **1**: 680-681.
- ZHAO, Z., LI, S., HUANG, H., LI, C., WANG, Q. and XUE, L. (2015) Comparative Study on Growth and Developmental Model of Indigenous Chicken Breeds in China. *Open Journal of Animal Sciences* **5**: 219-223.
- ZUIDHOF, M.J. (2005) Mathematical characterisation of broiler carcass yield dynamics. *Poultry Science* **84**: 1108-1122.
- ZWITERING, M.H., JONGENBURGER, I., ROMBOUTS, F.M. and VAN 'T RIET, K. (1990) Modelling of the bacterial growth curve. *Applied and Environmental Microbiology* **56**: 1875-1991.