

1 Fast Fourier Transform (Section 6.12)

Fast Fourier Transform can be used to efficiently evaluate an interpolating trigonometric polynomial,

$$P(x_j) = f(x_j),$$

for $x_j = \frac{2\pi j}{N}$, $E_k(x) = e^{ikx}$, and $i = \sqrt{-1}$, for $0 \leq j \leq N-1$ (recall again that N is the number of nodes). Then, we can define P as

$$P(x) = \sum_{k=0}^{N-1} c_k E_k(x), \quad c_k = \langle f, E_k \rangle_N = \frac{1}{N} \sum_{j=0}^{N-1} f(x_j) \overline{E_k(x_j)}. \quad (1)$$

Computing $P(x)$ (i.e., computing the c_k coefficients) without structure results in $\mathcal{O}(N^2)$ flops operations. How can we improve this runtime?

1.1 Algorithm for Power of 2 Polynomial

The algorithm we'll consider is when $N = 2^m$, with $m \in \mathbb{N}$. This algorithm for evaluating such a polynomial depends on the recursion on intermediate polynomials. In particular, we'll be working with the polynomial,


$$P_k^{(n)}(x), \quad 0 \leq n \leq m, \quad 0 \leq k \leq 2^{m-n} - 1.$$

Keep in mind that N and m are *fixed*, but n and k are *not* fixed. In any case, the formula is given by

$$P_k^{(n+1)}(x) = \frac{1}{2} \left(1 + e^{i2^n x} \right) P_k^{(n)}(x) + \frac{1}{2} \left(1 - e^{i2^n x} \right) P_{k+2^{m-n-1}}^{(n)} \left(x - \frac{\pi}{2^n} \right).$$

Note that the superscript (m) is just an iteration counter.

(Example.) Suppose we wanted to compute $P_0^{(3)}$. This can be obtained by computing $P_0^{(2)}$ and $P_1^{(2)}$. Each of these can be obtained from the four polynomials of lower order. For example, $P_0^{(2)}$ can be obtained by $P_0^{(1)}$ and $P_{0+2^{3-1-1}}^{(1)} = P_2^{(1)}$.



1.1.1 The Idea

Each $P_k^{(n)}$ can be represented by a set of coefficients, $C_{kj}^{(n)}$, for $0 \leq n \leq m$ and $0 \leq k \leq 2^{m-n} - 1$ and $0 \leq j \leq 2^n - 1$. More specifically, the polynomial can be written as

$$P_k^{(n)}(x) = \sum_{j=0}^{2^n-1} C_{kj}^{(n)} E_j(x) = \sum_{j=0}^{2^n-1} C_{kj}^{(m)} e^{ijx}.$$


Note that

$$C_{kj}^{(n+1)} = \frac{1}{2} \left(C_{kj}^{(n)} + e^{-\frac{ij\pi}{2^n}} C_{k+2^{m-n-1},j}^{(m)} \right)$$

and

$$C_{k,j+2^n}^{(n+1)} = \frac{1}{2} \left(C_{kj}^{(n)} - e^{-\frac{ij\pi}{2^n}} C_{k+2^{m-n-1},j}^{(m)} \right)$$

So, for a fixed n where $0 \leq k \leq 2^{m-n} - 1$ and $0 \leq j \leq 2^n - 1$, we have a matrix with 2^{m-n} rows and 2^n columns. Together, there's $2^{m-n}2^n = 2^m = N$ entries. Instead, we'd like to map each matrix row to some elements in an array, something like



The long array has N elements in total. To get each element in this array, we can write

$$C(2^n k + j) = C_{kj}^{(n)} \quad 0 \leq k \leq 2^{m-n} - 1, 0 \leq j \leq 2^n - 1$$

$$D(2^{n+1} k + j) = C_{kj}^{(n+1)} \quad 0 \leq k \leq 2^{m-n-1} - 1, 0 \leq j \leq 2^{n+1} - 1.$$

Next, we want to precompute $Z(j)$; that is,

$$Z(j) = e^{-\frac{2\pi i j}{N}}$$

and write

$$Z(j2^{m-n-1}) = e^{-\frac{ij\pi}{2^n}}.$$

1.2 The Algorithm

With the ideas in mind, we can write the algorithm below. Note that this has inputs m and $f(x)$.

Algorithm 1 Fast Fourier Transform

```

1: function FFT( $m, f$ )
2:    $N \leftarrow 2^m$ 
3:    $w \leftarrow e^{-\frac{2\pi i}{N}}$ 
4:   for  $k \leftarrow 0$  to  $N - 1$  do
5:      $Z(k) \leftarrow w^k$ 
6:      $C(k) \leftarrow f\left(\frac{2\pi k}{N}\right)$ 
7:   end for
8:   for  $n \leftarrow 0$  to  $m - 1$  do
9:     for  $k \leftarrow 0$  to  $2^{m-n-1} - 1$  do
10:      for  $j \leftarrow 0$  to  $2^n - 1$  do
11:         $u \leftarrow C(2^n k + j)$ 
12:         $v \leftarrow Z(j2^{m-n-1})C(2^n k + 2^{m-1} + j)$ 
13:         $D(2^{n+1}k + j) \leftarrow \frac{u+v}{2}$ 
14:         $D(2^{n+1}k + j + 2^n) \leftarrow \frac{u-v}{2}$ 
15:      end for
16:    end for
17:    for  $j \leftarrow 0$  to  $N - 1$  do
18:       $C(j) \leftarrow D(j)$ 
19:    end for
20:  end for
21:  return  $C$ 
22: end function

```

The elements in C can then be used in the interpolating formula (1).