

Computational Complexity of Smooth Differential Equations

Akitoshi Kawamura¹, Hiroyuki Ota¹, Carsten Rösnick², and Martin Ziegler²

¹ University of Tokyo

² TU Darmstadt

Abstract. The computational complexity of the solution h to the ordinary differential equation $h(0) = 0$, $h'(t) = g(t, h(t))$ under various assumptions on the function g has been investigated in hope of understanding the intrinsic hardness of solving the equation numerically. Kawamura showed in 2010 that the solution h can be PSPACE-hard even if g is assumed to be Lipschitz continuous. We place further requirements on the smoothness of g and obtain the following results: the solution h is still PSPACE-hard if g is assumed to be continuously differentiable; for each $k \geq 2$, the solution h is hard for the counting hierarchy if g is assumed to be k -times continuously differentiable.

1 Introduction

Let $g: [0, 1] \times \mathbf{R} \rightarrow \mathbf{R}$ be continuous and consider the following differential equation:

$$h(0) = 0, \quad Dh(t) = g(t, h(t)) \quad t \in [0, 1], \quad (1.1)$$

where Dh denotes the derivative of h . How complex can the solution h be, assuming that g is polynomial-time computable? Here, the polynomial-time computability and other notions of complexity are from the field of *Computable Analysis* [13] and measure how hard it is to approximate real functions with specified precisions (Section 2).

If we put no assumption on g other than being polynomial-time computable, the solution h (which is not unique in general) can be non-computable. Table 1.1 summarizes known results about the complexity of h under various assumptions on g , with the assumptions getting stronger as we go down. In particular, if g is (globally) Lipschitz continuous, then the (unique) solution h is known to be polynomial-space computable but still can be PSPACE-hard [2]. In this paper, we study the complexity of h when we put stronger assumptions about the smoothness of g .

In numerical analysis, knowledge about smoothness of the input function (such as being differentiable enough times) is often beneficial in applying certain algorithms or simplifying their analysis. However, to our knowledge this casual understanding that smoothness is good has not been rigorously substantiated in terms of computational complexity theory. This motivates us to ask whether,

Table 1.1. The complexity of the solution h of (1.1) assuming g is polynomial-time computable.

Assumptions	Upper bounds	Lower bounds
—	—	can be all non-computable [10]
h is the unique solution	computable [1]	can take arbitrarily long time [5, 8]
the Lipschitz condition	polynomial-space [5]	can be PSPACE-hard [2]
g is of class $C^{(\infty,1)}$	polynomial-space	can be PSPACE-hard (Theorem 1)
g is of class $C^{(\infty,k)}$ (for any constant k)	polynomial-space	can be CH-hard (Theorem 2)
g is analytic	polynomial-time [7, 9]	—

for our differential equation (1.1), smoothness helps to reduce the complexity of the solution.

At the extreme is the case where g is analytic: as the last row of the table shows, h can then be shown to be polynomial-time computable by an argument based on Taylor series. Thus our interest is in the cases between Lipschitz and analytic (the fourth and fifth rows in the table). We say that g is of class $C^{(i,j)}$ if the partial derivative $D^{(i,j)}g$ (often also denoted $\partial^{i+j}g(t, y)/\partial t^i \partial y^j$) exists and is continuous³; it is said to be of class $C^{(\infty,j)}$ if it is of class $C^{(i,j)}$ for all $i \in \mathbf{N}$.

Theorem 1. *There is a polynomial-time computable function $g: [0, 1] \times [-1, 1] \rightarrow \mathbf{R}$ of class $C^{(\infty,1)}$ such that the equation (1.1) has a PSPACE-hard solution $h: [0, 1] \rightarrow \mathbf{R}$.*

Theorem 2. *Let k be a positive integer. There is a polynomial-time computable function $g: [0, 1] \times [-1, 1] \rightarrow \mathbf{R}$ of class $C^{(\infty,k)}$ such that the equation (1.1) has a CH-hard solution $h: [0, 1] \rightarrow \mathbf{R}$, where $\text{CH} \subseteq \text{PSPACE}$ is the Counting Hierarchy (see Section 3.2).*

We said $g: [0, 1] \times [-1, 1] \rightarrow \mathbf{R}$ instead of $g: [0, 1] \times \mathbf{R} \rightarrow \mathbf{R}$, because the notion of polynomial-time computability of real functions is defined in this paper only when the domain is bounded closed region. This notational choice makes the validity of the equation (1.1) ill-defined in case h ever takes a value outside $[-1, 1]$; by saying that h is a solution in Theorem 1, we are implicitly also claiming that $h(t) \in [-1, 1]$ for all $t \in [0, 1]$. In any case, since we are putting stronger assumptions on g than Lipschitz continuity, such a solution h , if it exists, is unique.

The questions of whether smoothness of the input function reduces the complexity of the output have been asked for operations other than solving differential equations, and some negative results are known. The integral of a

³ Another common terminology is to say that g is of class C^k if it is of class $C^{(i,j)}$ for all i, j with $i + j \leq k$.

polynomial-time computable real function is $\#P$ -hard, and this does not change even if the input is restricted to C^∞ (infinitely differentiable) functions [6, Theorem 5.33]. Similarly, the function obtained by maximization from a polynomial-time computable real function is NP -hard, and this is still so even if the input function is restricted to C^∞ [6, Theorem 3.7]⁴. (Restricting to analytic inputs renders the output polynomial-time computable, again because of the argument based on Taylor series.) In contrast, although we have shown Theorem 2 for each k , we do not know about the complexity of h when g is assumed to be infinitely differentiable.

Notation Let \mathbf{N} denote the set of natural numbers, \mathbf{Z} denote the set of integer numbers, \mathbf{Q} denote the set of rational numbers and \mathbf{R} denote the set of real numbers.

Let A and B be bounded closed intervals in \mathbf{R} . We denote $|f|$ as $\sup_{x \in A} f(x)$ where $f: A \rightarrow \mathbf{R}$. A function $f: A \rightarrow \mathbf{R}$ is *class* C^i (i -times continuously differentiable) if there exist the derivatives $Df, D^2f, \dots, D^i f$ and all of them are continuous. A function g is of *class* C^k if it is k -times continuously differentiable.

Let g be a differentiable function of two variable, we denote D_1g as the derivative of g with respect to the first variable, and D_2g as the derivative of g with respect to the second variable. A function $g: A \times B \rightarrow \mathbf{R}$ is of *class* $C^{(i,j)}$ ((i,j) -times continuously differentiable) if for each $n \in \{0, \dots, i\}$ and $m \in \{0, \dots, j\}$, there exists the derivative $D_1^n D_2^m g$ and it is continuous. A function g is of *class* $C^{(\infty,j)}$ if g is of class $C^{(i,j)}$ for all $i \in \mathbf{N}$. When a function g is of class $C^{(i,j)}$, we write $D^{(i,j)}g$ for the derivative $D_1^i D_2^j g$.

2 Computational Complexity of Real Functions

2.1 Computation of Real Function

We start by fixing a way to encode real numbers by functions from strings to strings.

Definition 3. A function $\phi: \{0\}^* \rightarrow \{0, 1\}^*$ is a name of a real number x if for all $n \in \mathbf{N}$, $\phi(0^n)$ is the binary representation of $\lfloor x \cdot 2^n \rfloor$ or $\lceil x \cdot 2^n \rceil$, where $\lfloor \cdot \rfloor$ and $\lceil \cdot \rceil$ mean rounding down and up to the nearest integer.

In effect, a name of a real number x receives 0^n and returns an approximation of x with precision 2^{-n} .

We use *oracle Turing machines* (henceforth just machines) to work on names of real numbers (Figure 2.1).

⁴ The proof of this fact in [6, Theorem 3.7] needs to be fixed by redefining

$$f(x) = \begin{cases} u_s & \text{if not } R(s, t), \\ u_s + 2^{-(p(n)+2n+1) \cdot n} \cdot h_1(2^{p(n)+2n+1}(x - y_{s,t})) & \text{if } R(s, t). \end{cases}$$




Fig. 2.1. A machine M computing a real function f .

Let M be a machine and ϕ be a function from strings to strings. We write $M^\phi(0^n)$ for the output string when M is given ϕ as oracle and string 0^n as input. Thus we also regard M^ϕ as a function from strings to strings.

Definition 4. Let A be a bounded closed interval of \mathbf{R} . A machine M computes a real function $f: A \rightarrow \mathbf{R}$ if for any $x \in A$ and any name ϕ_x of it, M^{ϕ_x} is a name of $f(x)$.

When A is a bounded closed interval of \mathbf{R}^2 , we define a machine computing $f: A \rightarrow \mathbf{R}$ in a similar way using machines with two oracles. A real function is (*polynomial-time*) *computable* if there exists some machine that computes it (in polynomial time). Polynomial-time computability of a real function f implies that for any $n \in \mathbf{N}$, an approximation of $f(x)$ with error less than 2^{-n} is computable in time polynomial in n independent of real number x .

Computing a real function in polynomial time, machines only know about x by its approximation within polynomial size (with polynomial precision). This fact yields in (2.2) in the following lemma.

Lemma 5. A real function is (*polynomial-time*) *computable* if and only if there exist a (*polynomial-time*) *computable* function $\phi: (\mathbf{Q} \cap [0, 1]) \times \{0\}^* \rightarrow \mathbf{Q}$ and polynomial $p: \mathbf{N} \rightarrow \mathbf{N}$ such that for all $d \in \mathbf{Q} \cap [0, 1]$ and $n \in \mathbf{N}$,

$$|\phi(d, 0^n) - f(d)| \leq 2^{-n}, \quad (2.1)$$

and for all $x, y \in [0, 1]$, $n \in \mathbf{N}$,

$$|x - y| \leq 2^{-p(n)} \Rightarrow |f(x) - f(y)| \leq 2^{-n}, \quad (2.2)$$

where each rational number in \mathbf{Q} is represented by a pair of integers in binary representation.

2.2 Reduction and Hardness

A language $L \subseteq \{0, 1\}^*$ is identified with the function $L: \{0, 1\}^* \rightarrow \{0, 1\}$ such that $L(u) = 1$ if and only if $u \in L$.




Fig. 2.2. Reduction from a language L to a function $f: [0, 1] \rightarrow \mathbf{R}$

Definition 6 (Reduction). A Language L reduces to a real function $f: [0, 1] \rightarrow \mathbf{R}$ if there exists a polynomial-time function S and a polynomial-time oracle Turing machine M (Figure 2.2) such that for any string u :

- (i) $S(u, \cdot)$ is a name of x_u ;
- (ii) $M^\phi(u)$ accepts if and only if $u \in L$ for any name ϕ of $f(x_u)$.

As a matter of form this definition is different from that by Kawamura but they have same power as reduction. Let C be a complexity class, a function f is C -hard if for all language in C is reducible to f .

3 Proof of the Theorems

The proofs of Theorems 1 and 2 proceed as follows. In Section 3.1, we define *difference equations*, a discrete version of the differential equations. In Section 3.2, we show the PSPACE- and CH-hardness of difference equations with certain restrictions. In Section 3.3, we show that these classes of difference equations are simulable by certain families of differential equations given by $C^{(\infty, 1)}$ and $C^{(\infty, k)}$ functions. In Section 3.4, we put these families of functions together into one real function to obtain the smooth differential equations stated in the theorems.

The idea of simulating difference equations with differential equations is essentially from the proof of the Lipschitz version [2]. In this paper we focus on the structure of difference equations to analyze precisely the effect of smoothness assumptions. Consequently we show differential equations given by $C^{(\infty, k)}$ functions can simulate difference equations of restricted *height*, and this leads to the proof of CH-hardness.

3.1 Difference Equations

In this section, we define difference equations, a discrete version of differential equations, and show the PSPACE- and CH-hardness of families of difference equations with height restrictions.

Let $[n]$ denote $\{0, \dots, n-1\}$. Let $G: [P] \times [Q] \times [R] \rightarrow \{-1, 0, 1\}$ and $H: [P+1] \times [Q+1] \rightarrow [R]$. We say that H is the solution of the *difference equation* given by G if for all $i \in [P]$ and $T \in [Q]$ (Figure 3.1),

⁵ In [2, Lemma 4.7] it is stated that the language class recognized by uniform families with polynomial height coincides **PSPACE**.

The definition of the counting hierarchy CH, its connection to difference equations and the proof of Lemma 8 will be presented in Section 3.2.

3.2 The Counting Hierarchy and Difference Equations of Logarithmic Height

The polynomial hierarchy PH is defined using non-deterministic polynomial-time oracle Turing machines:

$$\Sigma_0^P = P, \quad \Sigma_{n+1}^P = \text{NP}^{\Sigma_n^P}, \quad \text{PH} = \bigcup_n \Sigma_n^P. \quad (3.3)$$

In the same way, the counting hierarchy CH [12] is defined using probabilistic polynomial-time oracle Turing machines⁶:

$$\text{C}_0 P = P, \quad \text{C}_{n+1} P = \text{PP}^{\text{C}_n P}, \quad \text{CH} = \bigcup_n \text{C}_n P. \quad (3.4)$$

It is known that $\text{PH} \subseteq \text{CH} \subseteq \text{PSPACE}$, but we do not know whether $\text{PH} = \text{PSPACE}$.

Each level of the counting hierarchy has a complete problem defined as follows. For every formula $\phi(X)$ with the list X of l free propositional variables, we write

$$\text{C}^m X \phi(X) \longleftrightarrow \sum_{X \in \{0,1\}^l} \phi(X) \geq m, \quad (3.5)$$

where $\phi(X)$ is identified with the function $\phi: \{0,1\}^l \rightarrow \{0,1\}$ such that $\phi(X) = 1$ if and only if $\phi(X)$ is true. This “counting quantifier” C^m generalizes the usual quantifiers \exists and \forall , because $\text{C}^1 = \exists$ and $\text{C}^{2^l} = \forall$. For lists X_1, \dots, X_n of variables and a formula $\phi(X_1, \dots, X_n)$ with all free variables listed, we define

$$\langle \phi(X_1, \dots, X_n), m_1, \dots, m_n \rangle \in \text{C}_n B_{be} \longleftrightarrow \text{C}^{m_1} X_1 \dots \text{C}^{m_n} X_n \phi(X_1, \dots, X_n). \quad (3.6)$$

Lemma 9 ([12, Theorem 7]). *For every $n \geq 1$, the problem $\text{C}_n B_{be}$ is $\text{C}_n P$ -complete.*

We define a problem $\text{C}_{\log} B_{be}$ by

$$\langle 0^{2^n}, u \rangle \in \text{C}_{\log} B_{be} \longleftrightarrow u \in \text{C}_n B_{be}. \quad (3.7)$$

We show that $\text{C}_{\log} B_{be}$ is CH-hard and recognized by a logarithmic-height uniform function family, as required in Lemma 8.

Proof (Lemma 8). First we prove that $\text{C}_{\log} B_{be}$ is CH-hard. For each problem A in CH, there is a constant n such that $A \in \text{C}_n P$. From Lemma 9, for each $u \in \{0,1\}^*$ there is a polynomial-time function f_n such that $u \in A \leftrightarrow f_n(u) \in \text{C}_n B_{be}$. So

$$u \in A \longleftrightarrow \langle 0^{2^n}, f_n(u) \rangle \in \text{C}_{\log} B_{be}. \quad (3.8)$$

⁶ This characterization, introduced by Torán [11], is different from Wagner’s.

Since $\langle 0^{2^n}, f_n(\cdot) \rangle$ is polynomial time computable, A is reducible to $C_{\log B_{be}}$.

Next we construct a logarithmic-height uniform function family $(G_u)_u$ recognizing $C_{\log B_{be}}$. Let $u = \langle 0^{2^n}, \langle \phi(X_1, \dots, X_n), m_1, \dots, m_n \rangle \rangle$, where n, m_1, \dots, m_n are nonnegative integers and ϕ is a formula. (If u is not of this form, then $u \notin C_{\log B_{be}}$.)

We write $l_i = |X_i|$ and $s_i = i + \sum_{j=1}^i l_j$. For each $i \in \{0, \dots, n\}$ and $Y_{i+1} \in \{0, 1\}^{l_{i+1}}, \dots, Y_n \in \{0, 1\}^{l_n}$, we write $\phi_i(Y_{i+1}, \dots, Y_n)$ for the truth value of the subformula $C^{m_i} X_i \dots C^{m_1} X_1 \phi(X_1, \dots, X_i, Y_{i+1}, \dots, Y_n)$, so that $\phi_0 = \phi$ and $\phi_n() = C_{\log B_{be}}(u)$. We regard the quantifier C^m as a function from \mathbf{N} to $\{0, 1\}$:

$$C^m(x) = \begin{cases} 1 & \text{if } x \geq m, \\ 0 & \text{if } x < m. \end{cases} \quad (3.9)$$

Thus,

$$\phi_{i+1}(Y_{i+2}, \dots, Y_n) = C^{m_{i+1}} \left(\sum_{X_{i+1} \in \{0, 1\}^{l_i}} \phi_i(X_{i+1}, Y_{i+2}, \dots, Y_n) \right). \quad (3.10)$$

For $T \in \mathbf{N}$, we write T_i for the i th digit of T written in binary, and $T_{[i,j]}$ for the string $T_{j-1}T_{j-2} \dots T_{i+1}T_i$.

For each $(i, T, Y) \in [n+1] \times [2^{s_n} + 1] \times [2^{|u|}]$, we define $G_u(i, T, Y)$ as follows. The first row is given by

$$G_u(0, T, Y) = (-1)^{T_{s_1}} \phi(T_{[1, s_1]}, T_{[s_1+1, s_2]}, \dots, T_{[s_{n-1}+1, s_n]}), \quad (3.11)$$

and for $i \neq 0$, we define

$$G_u(i, T, Y) = \begin{cases} (-1)^{T_{s_{i+1}}} C^{m_i}(Y) & \text{if } T_{[1, s_{i+1}]} = 10 \dots 0, \\ 0 & \text{otherwise.} \end{cases} \quad (3.12)$$

Define H_u from G_u by (3.1) and (3.2).

We prove by induction on i that $H_u(i, T) \in [2^{l_i}]$ for all T , and that

$$G_u(i, T, H_u(i, V)) = (-1)^{V_{s_{i+1}}} \phi_i(V_{[s_i+1, s_{i+1}]}, \dots, V_{[s_{n-1}+1, s_n]}) \quad (3.13)$$

if $V_{[1, s_{i+1}]} = 10 \dots 0$ (otherwise it is immediate from the definition that $G_u(i, V, H_u(n, V)) = 0$).

For $i = 0$, the claims follows from (3.11). For the induction step, assume (3.13). We have

$$H_u(i+1, T) = \sum_{V=0}^{T-1} G_u(i, V, H_u(i, V)). \quad (3.14)$$

Since the assumption (3.13) implies that flipping the bit $V_{s_{i+1}}$ of any V reverses the sign of $G_u(i, V, H_u(i, V))$, most of the summands in (3.14) cancel out. The terms that is can survive satisfy that $V_{[1, s_{i+1}]} = 10 \dots 0$ and that V is

between $\overline{T_{s_n} \dots T_{s_{i+1}+1} 00 \dots 0}$ and $\overline{T_{s_n} \dots T_{s_{i+1}+1} 01 \dots 1}$, where we write \overline{U} for the number represented by string U in binary. Since these terms are 0 or 1, $H_u(i+1, T) \in [2^{l_i}]$. Then if $T_{[1, s_{i+1}+1]} = 10 \dots 0$,

$$H_u(i+1, T) = \sum_{X \in \{0,1\}^{l_i}} \phi_i(X, T_{[s_{i+1}+1, s_{i+2}]}, \dots, T_{[s_{n-1}+1, s_n]}). \quad (3.15)$$

By this equation and (3.10),

$$\begin{aligned} G_u(i+1, T, H_u(i+1, T)) &= (-1)^{T_{s_{i+2}}} C^{m_{i+1}}(H_u(i+1, T)) \\ &= (-1)^{T_{s_{i+2}}} \phi_{i+1}(T_{[s_{i+1}+1, s_{i+2}]}, \dots, T_{[s_{n-1}+1, s_n]}), \end{aligned} \quad (3.16)$$

completing the induction steps.

By substituting n for i and 2^{s_n} for T in (3.13), we get $G_u(n, 2^{s_n}, H_u(n, 2^{s_n})) = \phi_n() = C_{\log B_{be}}(u)$. Hence $H_u(n+1, 2^{s_n}+1) = C_{\log B_{be}}(u)$.

We show that $(G_u)_u$ is uniform and has logarithmic height. The height $n+1$, the width $2^{s_n}+1$, and the cell size $2^{|u|}$ of G_u are polynomial-time computable from u , and $n+1 \leq \log |0^{2^n}| + 1 \leq \log |u| + 1$. \square

The language class recognized by uniform function families with i rows contains C_iP (the i th level of the counting hierarchy) and is contained in $C_{i+1}P$. While the class C_iP is defined by (3.4) using oracle Turing machines, it is also characterized as those languages Karp-reducible to C_iB_{be} , or as those accepted by a polynomial-time alternating Turing machine extended with “threshold states” and having at most i alternations. Likewise, the language class accepted by uniform function families of logarithmic height coincided with languages Karp-reducible to $C_{\log B_{be}}$ and with those accepted by an extended alternating Turing machine with logarithmic alternations. Since this class contains CH, we only state CH-hard in Lemma 10 and Theorem 2, but it is not known such class how hard the class is between CH and PSPACE.

3.3 Families of Real Functions Simulating Difference Equations

We show that certain families of smooth differential equations can simulate PSPACE-hard or CH-hard difference equations stated in previous section.

Before stating Lemma 10 and Lemma 11, we extend the definition of polynomial-time computability of real function to families of real functions. A machine M computes a family $(f_u)_u$ of functions $f_u: A \rightarrow \mathbf{R}$ indexed by strings u if for any $x \in A$ and any name ϕ_x of x , the function taking v to $M^{\phi_x}(u, v)$ is a name of $f_u(x)$. We say a family of real functions $(f_u)_u$ is polynomial-time if there is a polynomial-time machine computing $(f_u)_u$.

Lemma 10. *There exist a CH-hard language L and a polynomial μ , such that for any $k \geq 1$ and polynomials γ , there are a polynomial ρ and families $(g_u)_u$, $(h_u)_u$ of real functions such that $(g_u)_u$ is polynomial-time computable and for any string u :*

- (i) $g_u: [0, 1] \times [-1, 1] \rightarrow \mathbf{R}$, $h_u: [0, 1] \rightarrow [-1, 1]$;
- (ii) $h_u(0) = 0$ and $Dh_u(t) = g_u(t, h_u(t))$ for all $t \in [0, 1]$;
- (iii) g_u is of class $C^{(\infty, k)}$;
- (iv) $D^{(i, 0)}g_u(0, y) = D^{(i, 0)}g_u(1, y) = 0$ for all $i \in \mathbf{N}$ and $y \in [-1, 1]$;
- (v) $|D^{(i, j)}g_u(t, y)| \leq 2^{\mu(i, |u|) - \gamma(|u|)}$ for all $i \in \mathbf{N}$ and $j \in \{0, \dots, k\}$;
- (vi) $h_u(1) = 2^{-\rho(|u|)}L(u)$.

Lemma 11. *There exist a PSPACE-hard language L and a polynomial μ , such that for any polynomial γ , there are a polynomial ρ and families $(g_u)_u$, $(h_u)_u$ of real functions such that $(g_u)_u$ is polynomial-time computable and for any string u satisfying (i)–(vi) of Lemma 10 with $k = 1$.*

We will prove Lemma 10 using Lemma 8 as follows. Let a function family $(G_u)_u$ be as in Lemma 8, and let $(H_u)_u$ be the solution of the difference equation given by $(G_u)_u$. We construct h_u and g_u from H_u and G_u such that $h_u(T/2^{q(|u|)}) = \sum_{i=0}^{p(|u|)} H_u(i, T)/B^{d_u(i)}$ for each $T = 0, \dots, 2^{q(|u|)}$ and $Dh_u(t) = g_u(t, h_u(t))$. The polynomial-time computability of $(g_u)_u$ follows from that of $(G_u)_u$. We can prove Lemma 11 from Lemma 7 in the same way.

In Lemma 10, we have the new conditions (iii)–(v) about the smoothness and the derivatives of g_u that were not present in [2, Lemma 4.1]. To satisfy these conditions, we construct g_u using the smooth function f in following lemma.

Lemma 12 ([6, Lemma 3.6]). *There exist a polynomial-time function $f: [0, 1] \rightarrow \mathbf{R}$ of class C^∞ and a polynomial s such that*

- (i) $f(0) = 0$ and $f(1) = 1$;
- (ii) $D^n f(0) = D^n f(1) = 0$ for all $n \geq 1$;
- (iii) f is strictly increasing;
- (iv) $D^n f$ is polynomial-time computable for all $n \geq 1$;
- (v) $|D^n f| \leq s(n)$ for all $n \geq 1$.

Although the existence of the polynomial s satisfying the condition (v) is not stated in [6, Lemma 3.6], it can be shown easily.

We only prove Lemma 10 here and omit the analogous and easier proof of Lemma 11.

Proof (Lemma 10). Let L and $(G_u)_u$ be as in Lemma 8, and let a function family $(H_u)_u$ be the solution of the difference equation given by $(G_u)_u$.

By a similar argument to the beginning of the proof of [2, Lemma 4.1], we may assume that there exist polynomial-time functions p , j_u and polynomials q , r satisfying the following properties:

$$G_u: [p(|u|)] \times [2^{q(|u|)}] \times [2^{r(|u|)}] \rightarrow \{-1, 0, 1\}, \quad (3.17)$$

$$H_u(i, 2^{q(|u|)}) = \begin{cases} L(u) & \text{if } i = p(|u|), \\ 0 & \text{if } i < p(|u|), \end{cases} \quad (3.18)$$

$$G_u(i, T, Y) \neq 0 \rightarrow i = j_u(T). \quad (3.19)$$

Since G_u has logarithmic height, there exists a polynomial σ such that $(k+1)^{p(x)} \leq \sigma(x)$

We construct the families of real functions $(g_u)_u$ and $(h_u)_u$ simulating G_u and H_u in the sense that $h_u(T/2^{q(|u|)}) = \sum_{i=0}^{p(|u|)} H_u(i, T)/B^{d_u(i)}$, where the constant B and the function $d_u: [p(|u|)+1] \rightarrow \mathbf{N}$ are defined by

$$B = 2^{\gamma(|u|)+r(|u|)+s(k)+k+3}, \quad d_u(i) = \begin{cases} \sigma(|u|) & \text{if } i = p(|u|), \\ (k+1)^i & \text{if } i < p(|u|). \end{cases} \quad (3.20)$$

For each $(t, y) \in [0, 1] \times [-1, 1]$, there exist unique $N \in \mathbf{N}$, $\theta \in [0, 1)$, $Y \in \mathbf{Z}$ and $\eta \in [-1/4, 3/4]$ such that $t = (T+\theta)2^{-q(|u|)}$ and $y = (Y+\eta)B^{-d_u(j_u(T))}$. Using f and a polynomial s of Lemma 12, we define $\delta_{u,Y}: [0, 1] \rightarrow \mathbf{R}$, $g_u: [0, 1] \times [-1, 1] \rightarrow \mathbf{R}$ and $h_u: [0, 1] \rightarrow [-1, 1]$ by

$$\delta_{u,Y}(t) = \frac{2^{q(|u|)} Df(\theta)}{B^{d_u(j_u(T)+1)}} G_u(j_u(T), T, Y \bmod 2^{r(|u|)}), \quad (3.21)$$

$$g_u(t, y) = \begin{cases} \delta_{u,Y}(t) & \text{if } \eta \leq \frac{1}{4}, \\ (1 - f(\frac{4\eta-1}{2}))\delta_{u,Y}(t) + f(\frac{4\eta-1}{2})\delta_{u,Y+1}(t) & \text{if } \eta > \frac{1}{4}, \end{cases} \quad (3.22)$$

$$h_u(t) = \sum_{i=0}^{p(|u|)} \frac{H_u(i, T)}{B^{d_u(i)}} + \frac{f(\theta)}{B^{d_u(j_u(T)+1)}} G_u(j_u(T), T, H_u(j_u(T), T)). \quad (3.23)$$

We will verify that $(g_u)_u$ and $(h_u)_u$ defined above satisfy all the conditions stated in Lemma 10. Polynomial-time computability of $(g_u)_u$ can be verified using Lemma 5. The condition (i) is immediate from (3.22) and the condition (ii) is verified by the same argument as [2, Lemma 4.1].

We prove that the condition (iii) holds, which states that g_u is of class $C^{(\infty, k)}$. This condition follows from that $D_2^j D_1^i g_u$ exists and is continuous for each $i \in \mathbf{N}$ and $j \in \{0, \dots, k\}$. We can show it by induction on i and j . For $j = i = 0$, it follows immediately from the definition of g_u . For $i \neq 0$ and $j = 0$, assuming that g_u is of $C^{(i-1, 0)}$ by the induction hypothesis, we get

$$D^i \delta_{u,Y}(t) = \frac{2^{(i+1)q(|u|)} D^{i+1} f(\theta)}{B^{d_u(j_u(T)+1)}} G_u(j_u(T), T, Y \bmod 2^{r(|u|)}), N \quad (3.24)$$

$$D_1^i g_u(t, y) = \begin{cases} D^i \delta_{u,Y}(t) & \text{if } \eta \leq \frac{1}{4}, \\ (1 - f(\frac{4\eta-1}{2})) D^i \delta_{u,Y}(t) + f(\frac{4\eta-1}{2}) D^i \delta_{u,Y+1}(t) & \text{if } \frac{1}{4} < \eta. \end{cases} \quad (3.25)$$

And it is continuous by the definition of f (Lemma 12). For $i \neq 0$ and $j \neq 0$, assuming that g_u is $C^{(i, j-1)}$ by the induction hypothesis, we get

$$D_2^j D_1^i g_u(t, y) = \begin{cases} 0 & \text{if } -\frac{1}{4} < \eta < \frac{1}{4}, \\ (2B^{d_u(j_u(T))})^j D^j f(\frac{4\eta-1}{2})(D^i \delta_{u,Y+1}(t) - D^i \delta_{u,Y}(t)) & \text{if } \frac{1}{4} < \eta < \frac{3}{4}. \end{cases} \quad (3.26)$$

and it is continuous. Here we complete the induction step.

Substituting $t = 0, 1$ ($\theta = 0$) into (3.25), we get $D^{(i,0)}g_u(0, y) = D^{(i,0)}g_u(1, y) = 0$, so the condition (iv) holds.

We show that the condition (v) holds with $\mu(x, y) = (x + 1)q(y) + s(x + 1)$. Note that μ is a polynomial and independent of k and γ . Since $|D^i \delta_{u,Y}(t)| \leq 2^{(i+1)q(|u|)+s(i+1)} B^{-d_u(j_u(|u|)+1)}$ by (3.21), for all $i \in \mathbb{N}$ and $j \in \{0, \dots, k\}$, we have

$$|D^{(i,j)}g_u| \leq 2^k B^{k \cdot j_u(T)} 2^{s(k)} \cdot 2 \cdot \frac{2^{(i+1)q(|u|)+s(i+1)}}{B^{d_u(j_u(|u|)+1)}} \leq \frac{2^{\mu(i,|u|)+s(k)+k+1}}{B} \leq 2^{\mu(i,|u|)-\gamma(|u|)} \quad (3.27)$$

by (3.25), (3.26) and our choice of B .

We have (vi) with $\rho(x) = \sigma(x) \cdot (\gamma(x) + r(x) + s(k) + k + 3)$, because

$$h_u(1) = \frac{H_u(p(|u|), 2^{q(|u|)})}{B^{d_u(p(|u|))}} = \frac{L(u)}{2^{\sigma(|u|) \cdot (\gamma(|u|)+r(|u|)+s(k)+k+3)}} = 2^{-\rho(|u|)} L(u). \quad (3.28)$$

□

To prove Lemma 11, let L and $(G_u)_u$ be as Lemma 7, and let $(H_u)_u$ be the solution of the difference equation given by $(G_u)_u$. Define $(g_u)_u$ and $(h_u)_u$ as (3.22) and (3.23) with $d_u(i) = i$. It is shown in the same way as above that they meet all the conditions stated in Lemma 11.

3.4 Proof of the Main Theorems

Using the function families $(g_u)_u$ and $(h_u)_u$ obtained from Lemmas 10 or 11, we construct the functions g and h in Theorems 1 and 2 as follows. Divide $[0, 1]$ into infinitely many subintervals $[l_u^-, l_u^+]$, with midpoints c_u . We construct h by putting a scaled copy of h_u onto $[l_u^-, c_u]$ and putting a horizontally reversed scaled copy of h_u onto $[c_u, l_u^+]$ so that $h(l_u^-) = 0$, $h(c_u) = 2^{-\rho'(|u|)} L(u)$ and $h(l_u^+) = 0$ where ρ' is a polynomial. In the same way, g is constructed from $(g_u)_u$ so that g and h satisfy (1.1). We give the details of the proof of Theorem 2 from Lemma 10, and omit the analogous proof of Theorem 1 from Lemma 11.

Proof (Theorem 2). Let L and μ be as Lemma 11. Define

$$\lambda(x) = 2x + 2, \quad \gamma(x) = x\mu(x, x) + x\lambda(x), \quad (3.29)$$

and for each u let

$$\Lambda_u = 2^{\lambda(|u|)}, \quad c_u = 1 - \frac{1}{2^{|u|}} + \frac{2\bar{u} + 1}{\Lambda_u}, \quad l_u^\mp = c_u \mp \frac{1}{\Lambda_u}, \quad (3.30)$$

where $\bar{u} \in \{0, \dots, 2^{|u|} - 1\}$ is the number represented by u in binary notation. Let ρ , $(g_u)_u$, $(h_u)_u$ be as in Lemma 10 corresponding to the above γ .

We define

$$g\left(l_u^\mp \pm \frac{t}{\Lambda_u}, \frac{y}{\Lambda_u}\right) = \begin{cases} \pm \sum_{l=0}^k \frac{D^{(0,l)} g_u(t, 1)}{l!} (y-1)^l & \text{if } 1 < y, \\ \pm g_u(t, y) & \text{if } -1 \leq y \leq 1, \\ \pm \sum_{l=0}^k \frac{D^{(0,l)} g_u(t, -1)}{l!} (y+1)^l & \text{if } 1 < y, \end{cases} \quad (3.31)$$

$$h\left(l_u^\mp \pm \frac{t}{\Lambda_u}\right) = \frac{h_u(t)}{\Lambda_u} \quad (3.32)$$

for each string u and $t \in [0, 1)$, $y \in [-1, 1]$. Let $D_1^i g(1, y) = 0$ and $h(1) = 0$ for any $y \in [-1, 1]$ and $i \in \mathbf{N}$.

It can be shown similarly to the Lipschitz version [2, Theorem 3.2] that g and h satisfy (1.1) and g is polynomial-time computable. Here we only prove that g is of class $C^{(\infty, k)}$.

We claim that for each $i \in \mathbf{N}$ and $j \in \{0, \dots, k\}$, the derivative $D_1^i D_2^j g$ is given by

$$D_1^i D_2^j g\left(l_u^\mp \pm \frac{t}{\Lambda_u}, \frac{y}{\Lambda_u}\right) = \begin{cases} \pm \Lambda_u^{i+j} \sum_{l=j}^k \frac{D^{(i,l)} g_u(t, 1)}{(l-j)!} (y-1)^l & \text{if } y < -1, \\ \pm \Lambda_u^{i+j} D^{(i,j)} g_u(t, y) & \text{if } -1 \leq y \leq 1, \\ \pm \Lambda_u^{i+j} \sum_{l=j}^k \frac{D^{(i,l)} g_u(t, -1)}{(l-j)!} (y+1)^l & \text{if } 1 < y \end{cases} \quad (3.33)$$

for each $l_u^\mp \pm t/\Lambda_u \in [0, 1)$ and $y/\Lambda_u \in [-1, 1]$, and by $D_1^i D_2^j g(1, y) = 0$. This is verified by induction on $i+j$. The equation (3.33) follows from calculation (note that this means verifying that (3.33) follows from the definition of g when $i=j=0$, from the induction hypothesis about $D_2^{j-1} g$ when $i=0$ and $j>0$, and from the induction hypothesis about $D_1^{i-1} D_2^j g$ when $i>0$). That $D_1^i D_2^j g(1, y) = 0$ also follows easily from induction hypothesis when $i=0$. When $i>0$, we verify the existence of $D_1^i D_2^j g(1, y)$ by the existence of $D_1^i D_2^j g(t, y)$ when t tends to 1, i.e., $|u|$ tends to infinity. From Lemma 10 (v), we get

$$\begin{aligned} \left| D_1^i D_2^j g\left(l_u^\mp \pm \frac{t}{\Lambda_u}, \frac{y}{\Lambda_u}\right) \right| &\leq \Lambda_u^{i+j} \sum_{l=j}^k |D^{(i,l)} g_u|(\Lambda_u + 1)^l \\ &\leq \Lambda_u^{i+j} \cdot k \cdot 2^{\mu(i, |u|) - \gamma(|u|)} \cdot (2\Lambda_u)^k \\ &\leq 2^{(i+j+k)\lambda(|u|) + 2k + \mu(i, |u|) - \gamma(|u|)}. \end{aligned} \quad (3.34)$$

By our choice of γ , it converges to 0 when $|u| \rightarrow \infty$. Hence $D_1^i D_2^j g(1, y)$ exists and

$$D_1^i D_2^j g(1, y) = \lim_{t \rightarrow 1-0} D_1^i D_2^j g(t, y) = 0. \quad (3.35)$$

The continuity of $D_1^i D_2^j g$ on $[0, 1) \times [-1, 1]$ follows from (3.33) and Lemma 10 (iv). Since (3.34) holds for any $i \in \mathbf{N}$ and $j \in \{0, \dots, k\}$, it follows from (3.35)

that $D_1^i D_2^j g$ is continuous when the first variable is 1. Here we complete the induction steps. □

4 Complexity of Operations

Both Theorem 1 and 2 states the complexity of the solution h under the assumption that g is polynomial-time computable. How complex can it be to “solve” differential equations, i.e., how complex is the operator computing h by given g ? To make an answer to this problem, we need to define complexity of operators from real functions to real functions.

To discuss complexity of real functions, we encode real numbers by functions from string to string. We use string functions as names of elements in \mathbf{R} , such relation is called representation of \mathbf{R} . In the same way, we represent real functions as string functions to discuss complexity of real operators. So we need to define representations as string functions of class $C_{[0,1]}$ of continuous real functions $h: [0, 1] \rightarrow \mathbf{R}$ and class $C_{L[0,1] \times [-1,1]}$ of Lipschitz continuous real functions. The choice of representations is important since the computability and complexity is depending on these representations. Following [4], we use δ_\square as the representation of $C_{[0,1]}$ and $\delta_{\square L}$ as the representation of $C_{L[0,1] \times [-1,1]}$. It is known that in some sense δ_\square is a unique natural representation of $C_{[0,1]}$ [3]. And $\delta_{\square L}$ is a representation added information about the Lipschitz constant to δ_\square .

Since size of string functions are not finite, by restricting the amount of resources (time and space) of oracle Turing machine using second order polynomial, we define second-order complexity classes (e.g. **FPSPACE**, polynomial-space computable), reductions (e.g. \leq_W , polynomial-time Weihrauch reduction), and hardness. Combining it with the representations of real functions described above, we can write theorems of this paper in constructive form.

Let ODE be an operator mapping a real function g of class $C_{L[0,1] \times [-1,1]}$ to the solution h of (1.1) of class $C_{[0,1]}$. The operator ODE is a partial projection from $C_{L[0,1] \times [-1,1]}$ to $C_{[0,1]}$. In [4, Theorem 4.9], $(\delta_{\square L}, \delta_\square)$ -**FPSPACE**- \leq_W -completeness of ODE is proven by rewriting the proof of the result of the third row in Table 1.1 in the constructive form. In a similar way, Theorem 1 can be rewritten in the constructive form. That is, let ODE_k as the operator ODE whose input is restricted to class $C^{(\infty, k)}$,

Theorem 13. *The operator ODE_1 is $(\delta_{\square L}, \delta_\square)$ -**FPSPACE**- \leq_W -complete.*

To show this theorem, we need to verify that the information used to construct functions in the proof of Theorem 1 can be acquired easily from inputs. We omit the proof since it does not need any new technique. This constructive theorem is stronger than non-constructive one [4, Lemma 3.7 and 3.8], i.e., Theorem 1 follows immediately from Theorem 13.

The constructive version of Theorem 2 is also true. In fact for each $k \in \mathbf{N}$, ODE_k is $(\delta_{\square L}, \delta_\square)$ -**CH**- \leq_W -hard. We will treat it in another paper since we need more discussion on how to handle hierarchy to define **CH** in the frame work of [4].

References

1. E.A. Coddington and N. Levinson. *Theory of Ordinary Differential Equations*. McGraw-Hill, 1955.
2. A. Kawamura. Lipschitz continuous ordinary differential equations are polynomial-space complete. *Computational Complexity*, 19(2):305–332, 2010.
3. A. Kawamura. On function space representations and polynomial-time computability. Dagstuhl Seminar 11411: Computing with Infinite Data, 2011. <http://www.imai.is.s.u-tokyo.ac.jp/~kawamura/dagstuhl.pdf>.
4. A. Kawamura and S. Cook. Complexity theory for operators in analysis. In *Proceedings of the 42nd ACM Symposium on Theory of Computing*, pages 495–502. ACM, 2010.
5. K.I. Ko. On the computational complexity of ordinary differential equations. *Information and Control*, 58(1-3):157–194, 1983.
6. K.I. Ko. *Complexity Theory of Real Functions*. Birkhäuser Boston, 1991.
7. K.I. Ko and H. Friedman. Computing power series in polynomial time. *Advances in Applied Mathematics*, 9(1):40–50, 1988.
8. W. Miller. Recursive function theory and numerical analysis. *Journal of Computer and System Sciences*, 4(5):465–472, 1970.
9. N.T. Müller. Uniform computational complexity of Taylor series. *Automata, Languages and Programming*, pages 435–444, 1987.
10. M.B. Pour-el and I. Richards. A computable ordinary differential equation which possesses no computable solution. *Annals of Mathematical Logic*, 17(1-2):61–90, 1979.
11. J. Torán. Complexity classes defined by counting quantifiers. *Journal of the ACM (JACM)*, 38(3):752–773, 1991.
12. K.W. Wagner. The complexity of combinatorial problems with succinct input representation. *Acta Informatica*, 23(3):325–356, 1986.
13. Klaus Weihrauch. *Computable Analysis: An Introduction*. Texts in Theoretical Computer Science. Springer, 2000.