

# **Pesticide-contaminated eggs, Are they really dangerous?**

**Jaekwon Kim  
Seokhwan Park**

# Controversy at 2017



# Contents



**1.Fipronil**

**2.Bifenthrin**

**3.DDT**

**4.Harmfulness**

**5.Conclusion**

# **1. Fipronil**

## **Chemical property**


# 1.Fipronil

2.Bifenthrin

3.DDT

4.Harmfulness

5.Conclusion




**Chemical formula**

**C<sub>12</sub>H<sub>4</sub>Cl<sub>2</sub>F<sub>6</sub>N<sub>4</sub>OS**



**Density**

**1.477-1.626 g/cm<sup>3</sup>**



**Melting point**    **200.5 °C**


(RS)-5-Amino-1-[2,6-dichloro-4-(trifluoromethyl)phenyl]-4-(trifluoromethylsulfinyl)pyrazole-3-carbonitrile




# **1. Fipronil**

## **Biological characteristics**

## Glutamate-gated chloride channels(GLuCl): Fipronil targets present in insects



GLuCl in Musca





# 1.Fipronil

## 2.Bifenthrin

## 3.DDT

## 4.Harmfulness

## 5.Conclusion




**GABA**( $\gamma$ -AminoButyricAcid)

Pipronil acts on the nervous system and blocks GABA and glutamate receptors.

Inhibit signal not generated by blocking  
-> signal confused and excitation  
-> fully discharged

High effect for insects?

1. high affinity with GABA
2. GLuCl is not in mammals





### Toxicity?


#### For Human

No Long-term Research on Human body

ingestion: accumulation in the body  
-> damage to the liver and kidneys

the production and use of chlordane  
and DDT are strictly limited by the  
Stockholm convention

classed as a WHO Class II moderately  
hazardous pesticide



## **2. Bifenthrin**

### **Chemical property**


## Chemical formula




## Density

$$1.3 \pm 0.1 \text{ g/cm}^3$$

Melting point 79.6 °C




2-Methyl-3-phenylphenyl)methyl (1*S*,3*S*)-3-[(*Z*)-2-chloro-3,3,3-trifluoroprop-1-enyl]- 2,2-dimethylcyclopropane-1-carboxylate



## **2. Bifenthrin**

### **Biological characteristics**



## Famous for fire ant repellent



based on the affinity to the voltage-gated sodium channels

**Type 1**  
Temporary combination and separation -> keep excitation

**Type 2**  
open the sodium-channel permanently -> keep depolarized






## Low effect for mammals?

mammals' higher body temperature,  
larger body size, and enough  
degrading enzymes

lower bifenthrin-sodium channel  
affinity than insects

only cats among vertebrates do not  
have glucuronidase(degrading enzyme)  
-> High effect for cats



## Toxicity?



### For Human

there are chronic toxicity, endocrine disruption in study, HOWEVER **no human data** were found

skin contact: itchy, prickly, tickling



ingestion: vomiting, diarrhea, headache



### Carcinogen

the U.S. EPA classified bifenthrin as a Group C, possible human carcinogen



Group C - Possibly Carcinogenic to Humans: Agents with limited animal evidence and little or no human data




### **3. DDT** (Dichloro-Diphenyl-Trichloroethane)

## **Chemical property**





Chemical formula





Density

0.99 g/cm<sup>3</sup>

Melting point 108.5 °C

1,1'-(2,2,2-Trichloroethane-1,1-diyl)bis(4-chlorobenzene)



### **3. DDT** (Dichloro-Diphenyl-Trichloroethane)

## **Biological characteristics**


# The most famous insecticide



Published in 1962

Paul Hermann Müller won the Nobel prize in Physiology or Medicine in 1948 for his discovery of the high efficiency of DDT

Rachel Carson introduced DDT's harm to Reader's Digest






## HOW does DDT kill insects?

bind to open sodium channels and stabilize the open state

DDT opens sodium ion channel in neurons, causing them to fire spontaneously, which leads to **spasms** and **eventual death**

used to kill **Pediculus(0|)**, but limited in 1986



# carcinogen?




**Toxic**

DDT is WHO IARC Group 2-A

WHO IARC Group 2-A: **Probably carcinogenic to humans**

Other WHO IARC Group 2-A: Very hot beverages (above 65 degrees Celsius), Red meat,





**Harmfulness?**

Why are carbon compounds dangerous?

In human's body, The action of several enzymes is important

must be digested what we ate

Halogen-free in human's body




**HOW about the eggs?**



## DANGEROUS but..

### 1. very high lethal dose



Fipronil – 97mg/kg  
Bifenthrin – 70mg/kg  
DDT – 113mg/kg

Shows LD50



## DANGEROUS but..

2. small amount in the egg


In Korea..

**Fipronil – 0.272mg/kg**

**Bifenthrin – 0.272mg/kg**

(Maximum detected value)

The biggest egg is 70mg(0.07kg)





**NOT** proven to be harmful to humans





High threshold



JUST eat!



HOWEVER, the unpleasant..??



# Citation

1. Narahashi, T., Zhao, X., Ikeda, T., Salgado, V. L., & Yeh, J. Z. (2010). Glutamate-activated chloride channels: Unique fipronil targets present in insects but not in mammals. *Pesticide biochemistry and physiology*, 97(2), 149–152. <https://doi.org/10.1016/j.pestbp.2009.07.008>
2. Silver, K. S., Du, Y., Nomura, Y., Oliveira, E. E., Salgado, V. L., Zhorov, B. S., & Dong, K. (2014). Voltage-Gated Sodium Channels as Insecticide Targets. *Advances in insect physiology*, 46, 389–433. <https://doi.org/10.1016/B978-0-12-417010-0.00005-7>
3. Yang, L., & Li, L. (2015). Actions of the pyrethroid insecticide bifenthrin on sodium channels expressed in rat cerebral cortical neurons. *Toxicology mechanisms and methods*, 25(1), 63–69. <https://doi.org/10.3109/15376516.2014.985355>
4. Lund Albert E., Narahashi Toshio (1983). "Kinetics of sodium channel modification as the basis for the variation in the nerve membrane effects of pyrethroids and DDT analogs". *Pesticide Biochemistry and Physiology*. **20** (2): 203–216.
5. Johnson, M.; Luukinen, B.; Gervais, J.; Buhl, K.; Stone, D. 2010. *Bifenthrin Technical Fact Sheet*, National Pesticide Information Center, Oregon State University Extension Services. <http://npic.orst.edu/factsheets/archive/biftech.html>
6. Al-Amin (2011). "Evaluation of Bifenthrin 80 SC, as a wall treatment against *Culex quinquefasciatus* Say (Diptera: Culicidae), a vector of *Wuchereria bancrofti* Cobbold, an etiological agent of Human Lymphatic Filariasis". *Terrestrial Arthropod Reviews*. **4** (3): 183–202. [doi:10.1163/187498311X577405](https://doi.org/10.1163/187498311X577405).
7. <https://m.blog.naver.com/PostView.nhn?blogId=almostcurable&logNo=221080275275&proxyReferer=https:%2F%2Fwww.google.com%2F>
8. Ratra, G. S.; Casida, J. E. (2001). "GABA receptor subunit composition relative to insecticide potency and selectivity". *Toxicol. Lett.* 122 (3): 215–222. doi:10.1016/s0378-4274(01)00366-6. PMID 11489356.



Created by Gan Khoon Lay  
from Noun Project

# Thank you for your attention



Created by anbilero adaleru  
from Noun Project