


2020 Fall General Physics 2 Homework #10

1. (5pts) When two events occur at the location A and B. The time interval between two locations is proper time interval. (O/X) **Answer: X Proper time is the time as measured by a clock with observe two events at the same location.**

2. (5pts) If two observers are in relative motion. One observer finds two events to be simultaneous, the other observer will not. (O/X) **Answer: O**

3. (5pts) A light source S moves toward the right direction. Which points the Doppler shift occur? **Answer: (e) In contrast with classical mechanics, Doppler shift occurs in the case that the source moves transverse to the detector.**

- (a) A
- (b) B
- (c) C
- (d) A,C
- (e) All



4. (5pts) There are people A and B which have same age. B boards a spaceship and travels to the other star. B arrives at the stare, B returns to Earth immediately. Then, which one is older? **Answer: (a) By time dilation, time flows slowly in a moving frame.**

- (a) A
- (b) B
- (c) same

5. (15pts) A spaceship moves away from Earth with speed v and fires a shuttle craft in the forward direction at a speed v relative to the spaceship. The pilot of the shuttle craft launches a probe in the forward direction at speed v relative to the shuttle craft. What is the speed of the probe relative to the Earth?

The speed of shuttle craft relative to the Earth is v' .

$$v = \frac{v' - v}{1 - \frac{v'v}{c^2}} \quad (+4)$$

$$\text{Then, } v' = \frac{2v}{1 + \frac{v^2}{c^2}} \quad (+4)$$

Similarly, speed of the probe relative to the Earth is v'' .

$$v = \frac{v'' - v'}{1 - \frac{v''v'}{c^2}} \quad (+4)$$

$$v'' = \frac{v' + v}{1 + \frac{v'v}{c^2}} = \frac{v^2 + 3c^2}{c^2 + 3v^2} v \quad (+3)$$

6. (15pts) When an object's speed increases by 10 %, its momentum increases by a factor of 10. What was its original speed?

The original speed is v and original momentum is $\gamma mv = \frac{mv}{\sqrt{1-\frac{v^2}{c^2}}}$.

If speed increases by 10 %, then the speed is $\frac{11}{10}v$.

Then, the momentum becomes $\gamma' m * \frac{11}{10}v = \frac{11mv}{10\sqrt{1-\frac{121v^2}{100c^2}}} \quad (+5)$

And it is same with $\frac{10mv}{\sqrt{1-\frac{v^2}{c^2}}}$.

$$\frac{11mv}{10\sqrt{1-\frac{121v^2}{100c^2}}} = \frac{10mv}{\sqrt{1-\frac{v^2}{c^2}}} \quad (+5)$$

$$11\sqrt{1-\frac{v^2}{c^2}} = 100\sqrt{1-\frac{121v^2}{100c^2}}$$

$$121\left(1-\frac{v^2}{c^2}\right) = 10000\left(1-\frac{121v^2}{100c^2}\right)$$

$$v=0.91c \quad (+5)$$

7. (15pts) The only invariant time in relativity is proper time which means that a length of time at a rest frame of an object. Show that infinitesimal proper time ($d\tau$) is proportional to dS ($dS^2 = c^2 dt^2 - dx^2 - dy^2 - dz^2$). Calculate ratio between infinitesimal proper time and infinitesimal time in any inertial frame (ct, x, y, z), and explain the meaning of the ratio.

At rest frame ($dx_r = dy_r = dz_r = 0$)

$$c^2 d\tau^2 - dx_r^2 - dy_r^2 - dz_r^2 = c^2 d\tau^2$$

$$c^2 d\tau^2 = c^2 dt^2 - dx^2 - dy^2 - dz^2 = dS^2$$

(for any inertial frame)

$$d\tau \propto dS$$

4pts

$$c^2 \left(\frac{d\tau}{dt} \right)^2 = c^2 - \left(\left(\frac{dx}{dt} \right)^2 + \left(\frac{dy}{dt} \right)^2 + \left(\frac{dz}{dt} \right)^2 \right)$$

$$= c^2 - v^2 \quad \beta = v/c$$

$$\frac{d\tau}{dt} = \sqrt{1 - \beta^2} = \frac{1}{\gamma}$$

4pts

$$\frac{dt}{d\tau} = \gamma \Rightarrow \text{Time Dilation}$$


2pts

8. (15pts) A source emitting light with frequency f moves toward you at speed u . By using the Lorentz transformation and drawing proper spacetime diagram, show that you measure a Doppler-shifted frequency given by,

$$f' = f \sqrt{\frac{c+u}{c-u}}$$

Use the binomial approximation to show that this result can be written in the form,

$$f' = \frac{f}{1-u/c}, (u \ll c)$$



$$T = \frac{1}{f} \quad \begin{pmatrix} c\Delta t \\ \Delta x \end{pmatrix} = \begin{pmatrix} \gamma & \beta\gamma \\ \beta\gamma & \gamma \end{pmatrix} \begin{pmatrix} T \\ 0 \end{pmatrix}$$

$$\beta = \frac{u}{c}$$

$$T' = \gamma T - \beta\gamma T = \gamma T (1 - \beta)$$

$$\frac{1}{f'} = \gamma (1 - \beta) \frac{1}{f}$$

Calculation 3pts

$$f' = f \frac{1}{1-\beta} \sqrt{1-\beta^2}$$

$$= f \sqrt{\frac{1+\beta}{1-\beta}}$$

for $u \ll c$

$$(1+\beta)^{1/2} \approx (1 + \frac{1}{2}\beta)$$

$$(1-\beta)^{1/2} \approx (1 - \frac{1}{2}\beta)$$


$$f' = f (1+\beta)^{1/2} (1-\beta)^{1/2} \approx f (1-\beta)^{-1} = \frac{f}{1-\beta}$$


Calculation 2pts

[Advanced Problem]


(20pts) Consider a line of positive charge with line charge density λ as measured in a frame S at rest with respect to the charges.


- (a) (2pts) Show that the electric field a distance r from this charged line has magnitude $E = \lambda/2\pi\epsilon_0 r$, and that there is no magnetic field (no relativity needed here).
- (b) (5pts) Now consider the situation in a frame S' moving at speed v parallel to the line of charge. Show that the charge density measured in S' is given by $\lambda' = \gamma\lambda$, where $\gamma = 1/\sqrt{1-v^2/c^2}$.
- (c) (4pts) Use the result of (b) to find the electric field in S' . Since the charge is moving with respect to S' , there is a current in S' . Also, find an expression for this current and the magnetic field it produces.
- (d) (9pts) Determine the values of the quantities $\vec{E} \cdot \vec{B}$ and $E^2 - c^2 B^2$ in both reference frames, and show that these quantities are invariant. Explain physical meaning of this invariance.

(a)  $\oint \vec{E} \cdot d\vec{A} = \frac{Q}{\epsilon_0}$ 1pt
 $|\vec{E}|(2\pi r l) = \frac{\lambda l}{\epsilon_0}$
 $|\vec{E}| = \frac{\lambda}{2\pi\epsilon_0 r}$ 1pt

(b) $\lambda' = \frac{Q'}{l'} = \gamma \frac{Q}{l} = \gamma \lambda$ 1pt

 $Q' = Q$ 2pts (charge conservation)
 $l' = \frac{l}{\gamma}$ 2pts (length contraction)

(c) $I = \frac{dQ'}{dt} = \frac{dQ'}{dx} \frac{dx}{dt} = \lambda' v$ 3pts
 $\oint \vec{B} \cdot d\vec{\ell} = \mu_0 I$ $|\vec{B}|(2\pi r) = \mu_0 \lambda v$
 $|\vec{B}| = \frac{\mu_0 \lambda v}{2\pi r}$ 1pt

(d)  $\vec{E} = \frac{\lambda}{2\pi\epsilon_0 r} \hat{r}$
 $\vec{B} = 0$ 3pts

 $\vec{E}' = \frac{\lambda'}{2\pi\epsilon_0 r} \hat{r}$
 $\vec{B}' = \frac{\mu_0 \lambda v}{2\pi r} \hat{\phi}$ 3pts

$\vec{E} \cdot \vec{B} = \vec{E}' \cdot \vec{B}' = 0$ 1pt

$|\vec{E}|^2 - c^2 |\vec{B}|^2 = |\vec{E}'|^2 - c^2 |\vec{B}'|^2$
 $= \frac{\lambda^2}{4\pi^2 \epsilon_0^2 r^2} = \frac{\lambda'^2}{4\pi^2 \epsilon_0^2 r^2} \left(\frac{\gamma^2}{\epsilon_0^2} - c^2 \mu_0^2 \gamma^2 v^2 \right)$
 $= \frac{\lambda'^2}{4\pi^2 \epsilon_0^2 r^2} \left(\frac{\gamma^2}{\epsilon_0^2} - \frac{\gamma^2 v^2}{c^2} \right) \quad \beta = \frac{v}{c}$
 $= \frac{\lambda'^2}{4\pi^2 \epsilon_0^2 r^2}$ 2pts