

Two Functions, Two Variables

$$(X_1, X_2) \sim \underline{f}_X(X_1, X_2)$$

(Y_1, Y_2) : some functions of (X_1, X_2)

$$\begin{cases} Y_1 = u_1(X_1, X_2) \\ Y_2 = u_2(X_1, X_2) \end{cases}$$

$$X_1 = v_1(Y_1, Y_2)$$

$$X_2 = v_2(Y_1, Y_2)$$

pdf of Y_1, Y_2

$$: \underline{f}_Y(y_1, y_2) = f_X(v_1(y_1, y_2), v_2(y_1, y_2)) \cdot |J|$$

$$J = \det \begin{pmatrix} \frac{\partial v_1}{\partial y_1} & \frac{\partial v_1}{\partial y_2} \\ \frac{\partial v_2}{\partial y_1} & \frac{\partial v_2}{\partial y_2} \end{pmatrix}$$

Eg) $X_1, X_2 \sim \text{iid Exp}(1)$

$$f_X(x_1, x_2) = e^{-x_1} e^{-x_2}, \quad x_1 > 0 \quad x_2 > 0$$


$$\begin{cases} Y_1 = \underline{X_1 - X_2} \\ Y_2 = X_1 + X_2 \end{cases}$$

$$\Rightarrow \begin{cases} X_1 = \frac{Y_1 + Y_2}{2} \\ X_2 = \frac{Y_2 - Y_1}{2} \end{cases}$$

$$Y_1 = \frac{y_1 + y_2}{2}$$

$$X_2 = \frac{y_2 - y_1}{2}$$

$$J = \det \begin{pmatrix} \frac{1}{2} & \frac{1}{2} \\ -\frac{1}{2} & \frac{1}{2} \end{pmatrix} = \frac{1}{2}$$



$$f_Y(y_1, y_2) = e^{-\frac{y_1+y_2}{2}} \cdot e^{-\frac{y_2-y_1}{2}} \cdot \frac{1}{2} = \frac{1}{2} e^{-y_2}, \quad y_2 > 0$$

$$x_1 > 0 \Rightarrow 0 < y_1 + y_2 < \infty$$

$$x_2 > 0 \Rightarrow 0 < y_2 - y_1 < \infty$$

\Rightarrow


$$-y_2 < y_1 < y_2$$

$$0 < y_2 < \infty$$

Find the marginals:

$$f(y_2) = \int_{-y_2}^{y_2} \frac{1}{2} e^{-y_2} dy_1 = y_2 e^{-y_2}, \quad y_2 > 0 \quad ; \text{Gamma}(2, 1)$$

$$f(y_1) = \begin{cases} \int_{-y_1}^{\infty} \frac{1}{2} e^{-y_2} dy_2 = \frac{1}{2} e^{-y_1}, & -\infty < y_1 < 0 \\ \int_{y_1}^{\infty} \frac{1}{2} e^{-y_2} dy_2 = \frac{1}{2} e^{-y_1}, & 0 < y_1 < \infty \end{cases}$$



• Double Exponentials

• Laplace

Eg) $Y_1, Y_2 \sim U(0, 1)$ $0 < Y_1 < 1$ $0 < Y_2 < 1$

$$\begin{cases} U = Y_1 Y_2 \\ V = Y_2 \end{cases} \quad \begin{cases} Y_1 = \frac{U}{V} \\ Y_2 = V \end{cases} \quad \begin{cases} y_1 = \frac{u}{v} \\ y_2 = v \end{cases} \quad J = \det \begin{pmatrix} \frac{1}{v} & -\frac{u}{v^2} \\ 0 & 1 \end{pmatrix}$$

$$0 < \frac{u}{v} < 1 \Rightarrow 0 < u < v \quad = \frac{1}{v}$$

$$f_{u,v}(u, v) = \frac{1}{v}, \quad 0 < u < v < 1$$

$$f_u(u) = \int_u^1 \frac{1}{v} dv = -\log u, \quad 0 < u < 1.$$

The Method of Moment-Generating Functions

$m_X(t), m_Y(t)$: mgfs of r.v.'s X and Y . If they exist and $m_X(t) = m_Y(t)$ for all values of t , then X and Y have the same probability distribution.

U : a function of r.v.'s Y_1, Y_2, \dots, Y_n .

1. Find the mgf for U , $m_U(t)$.
2. Compare $m_U(t)$ with other well-known mgfs. If $m_U(t) = m_V(t)$ for all values of t , U and V have identical distribution.

Use this to prove that

• Sum of independent Poisson is Poisson.

$$X_i \sim \text{Poisson}(\lambda_i) \quad i = 1, \dots, n$$

$$X_1 + \dots + X_n \sim \text{Poisson}(\lambda_1 + \dots + \lambda_n)$$

$$E(e^{t(X_1 + \dots + X_n)}) = E(e^{tX_1}) \dots E(e^{tX_n})$$

$$= \exp(e^{\lambda_1(t-1)}) \dots \exp(e^{\lambda_n(t-1)})$$

$$= \exp(e^{(\sum \lambda_i)(t-1)}) : \text{Poisson}(\sum \lambda_i)$$

DIY

- Sum of Indep Binomids with same p . is Binomial
- Sum of Indep. Gamma with same β is Gamma.

$$Z \sim N(0, 1)$$

$$Y = Z^2$$

$$E(e^{tY}) = \int_{-\infty}^{\infty} e^{tZ^2} \cdot \frac{1}{\sqrt{2\pi}} e^{-\frac{1}{2}Z^2} dZ = \int_{-\infty}^{\infty} \frac{1}{\sqrt{2\pi}} e^{(-\frac{1}{2} + t)Z^2} dZ$$

...

need $t < \frac{1}{2}$

DIY

is a part of $N(0, \Sigma)$

$$= (1-2t)^{-\frac{1}{2}}$$

$$\therefore \text{Gamma}\left(\frac{1}{2}, 2\right) = \chi^2(1)$$

Super useful for normal-related problems.

$$Y_1, \dots, Y_n \stackrel{\text{i.i.d.}}{\sim} N(\mu, \sigma^2)$$

$$\star \quad \bar{Y} \sim N\left(\mu, \frac{\sigma^2}{n}\right)$$

$$\therefore E(e^{t\bar{Y}}) = E\left(e^{\frac{1}{n}t(Y_1 + \dots + Y_n)}\right)$$

$$= E\left(e^{\frac{1}{n}tY_1}\right) \dots \dots E\left(e^{\frac{1}{n}tY_n}\right)$$

$$\left(\underbrace{\exp\left(\mu t' + \frac{1}{2}\sigma^2 t'^2\right)}_{\parallel} \right)$$

$$= \left(\exp\left(\frac{1}{n}\mu t + \frac{1}{2}\frac{\sigma^2}{n} t^2\right) \right)^n = \exp\left(\mu t + \frac{1}{2}\frac{\sigma^2}{n} t^2\right)$$

$$\cdot Y_i \sim N(\mu_i, \sigma_i^2) \quad i=1, \dots, n. \quad \text{indep.}$$

Show that using mgf method

$$\star\star \quad \sum a_i Y_i \sim N(\sum a_i \mu_i, \sum a_i^2 \sigma_i^2)$$

e.g.

$$\left[\begin{array}{l} X_1 \dots X_n \stackrel{\text{iid}}{\sim} N(\mu_1, \sigma_1^2) \\ Y_1 \dots Y_m \stackrel{\text{iid}}{\sim} N(\mu_2, \sigma_2^2) \end{array} \right] \text{ indep}$$

Check!!

$$\bar{X} - \bar{Y} \sim N(\mu_1 - \mu_2, \frac{\sigma_1^2}{n} + \frac{\sigma_2^2}{m})$$