

Introduction to Computational Fluid Dynamics using OpenFOAM and Octave


Dr. Lakshman Anumolu (Sr. Research Engineer)
Kumaresh Selvakumar (PhD candidate)
(Session-10)

*Instructions: Mon, Wed, Thu (3:30PM-4:30PM IST)
Query session: Sundays 8:30AM-9:00AM IST*

Quick Recap

What Did We Discuss?

$$\frac{\partial T}{\partial t} = \alpha \frac{\partial^2 T}{\partial x^2}$$




```
1  #include <iostream>
2  #include <vector>
3  #include <cmath>
4
5  std::vector<float> compute_rate_of_convergence(std::vector<float> errors, std::vector<float> dx) {
6      std::vector<float> roc;
7
8      for (int i = 0; i < errors.size()-1; ++i) {
9          float r = log(errors[i]/errors[i+1])/log(dx[i]/dx[i+1]);
10
11          roc.push_back(r);
12      }
13
14      return roc;
15  }
16
17  int main()
18  {
19      std::vector<float> errors({0.16, 0.0775, 0.038125});
20      std::vector<float> dx({0.1, 0.05, 0.025});
21
22      std::vector<float> roc = compute_rate_of_convergence(errors, dx);
23
24      for (int i = 0; i < roc.size(); ++i) {
25          std::cout << roc[i] << std::endl;
26      }
27
28      return 0;
29  }
```

Current Session

Overview

- **OpenFOAM:** Numerical Solution to Diffusion Equation
- Introduction to C++ for OpenFOAM (contd.)

Numerical Solution to Diffusion Equation



$$\frac{\partial T}{\partial t} = \alpha \frac{\partial^2 T}{\partial x^2}$$

Introduction to C++ for OpenFOAM

b10_roc.cpp

Next Session

- Finite difference method to solve convection equation in Octave
- Introduction to C++ for OpenFOAM (Contd.)

Thank you