# Introduction to Computational Fluid Dynamics using OpenFOAM and Octave

Lakshman Anumolu Kumaresh Selvakumar (Session-6)


Instructions: Wed, Fri (4:30-5:30PM IST), Sat (4PM-5PM IST)

Query sessions: Sundays 9:00AM-9:30AM IST

## Quick Recap

#### What Did We Discuss?

#### **Taylor Series**



$$\rho(x^*) = \rho(x_i) + \frac{(x^* - x_i)}{1!} \left(\frac{d\rho}{dx}\right)_i + \frac{(x^* - x_i)^2}{2!} \left(\frac{d^2\rho}{dx^2}\right)_i + \frac{(x^* - x_i)^3}{3!} \left(\frac{d^3\rho}{dx^3}\right)_i + \cdots$$




$$\left(\frac{d\rho}{dx}\right)_{i} \approx \frac{\rho(x_{i+1}) - \rho(x_{i})}{\Delta x_{i}} \qquad \left(\frac{d\rho}{dx}\right)_{i}$$

First order upwind

$$\left(\frac{d\rho}{dx}\right)_{i} \approx \frac{\rho(x_{i+1}) - \rho(x_{i-1})}{2\Delta x_{i}}$$

Second order central difference



## **Current Session**

#### Overview

- Introduction to Octave programming
- Stability Analysis

## Introducing Octave


6\_a\_octave\_introduction.m

Numerical approach should not magnify the error that appears in the solution.

$$\frac{d\rho}{dx} = -c\rho$$

$$\int_{\rho_o}^{\rho} \frac{d\rho}{\rho} = \int_{x_o}^{x} -cdx$$

$$\rho = \rho_o e^{-c(x - x_o)}$$



Numerical discretization

$$\frac{d\rho}{dx} = -c\rho \qquad \qquad \chi_{i-1} \qquad \chi_i$$

$$\frac{\rho_{i+1} - \rho_i}{\Delta x} = -c\rho_i$$

 $x_{i+1}$ 

$$\rho_{i+1} = \rho_i (1 - c\Delta x)$$


Numerical discretization

$$\frac{d\rho}{dx} = -c\rho$$

$$x_{i-1}$$
  $x_i$   $x_{i+1}$   $x_{i+2}$ 

$$\frac{\rho_{i+1} - \rho_i}{\Delta x} = -c\rho_i$$


$$\rho_{i+1} = \rho_i (1 - c\Delta x)$$





$$\frac{d\rho}{dx} = -c\rho$$


$$x_{i-1}$$
  $x_i$   $x_{i+1}$   $x_{i+2}$ 


$$\rho_{i+1} = \rho_i (1 - c\Delta x)$$













$$\frac{d\rho}{dx} = -c\rho$$

$$x_{i-1}$$
  $x_i$   $x_{i+1}$   $x_{i+2}$ 

$$\left|\frac{\rho_{i+1}}{\rho_i}\right| < 1$$

$$\rho_{i+1} = \rho_i (1 - c\Delta x)$$





$$c = 4, x_o = 0, \rho_o = 1, x \in [0,10], \Delta x = 0.1$$
  $c = 4, x_o = 0, \rho_o = 1, x \in [0,10], \Delta x = 0.01$ 

$$\frac{d\rho}{dx} = -c\rho$$

$$\frac{d\rho}{dx} = -c\rho \qquad \qquad x_{i-1} \quad x_i \qquad x_{i+1} \quad x_{i+2}$$

$$\left|\frac{\rho_{i+1}}{\rho_i}\right| < 1$$

$$\rho_{i+1} = \rho_i (1 - c\Delta x)$$

$$\left|\frac{\rho_{i+1}}{\rho_i}\right| = |(1 - c\Delta x)| < 1$$

$$-1 < (1 - c\Delta x) < 1$$


$$0 < \Delta x < 2/c$$


$$\Delta x < 2/c$$


$$\frac{d\rho}{dx} = -c\rho$$


$$x_{i-1}$$
  $x_i$   $x_{i+1}$   $x_{i+2}$ 


$$\rho_{i+1} = \rho_i (1 - c\Delta x)$$











$$\Delta x < \frac{2}{c}$$

$$\frac{2}{c} = \frac{2}{4} = 0.5$$

#### **Next Session**

- Brief look at stability analysis again
- Introduction to C++ for OpenFOAM

# Thank you