

1 Tensor Basics

1.1 Definitions

TODO

1.2 Tensor Manipulations and Operations

TODO

2 Tensor Derivatives

let T be a contravariant tensor, then:

$$\bar{T}^i = T^r \frac{\partial \bar{x}^i}{\partial x^r}, \text{ where } T = T^i(x(t))$$

Take derivative wrt. t :

$$\frac{\partial \bar{T}^i}{\partial t} = \frac{\partial T^r}{\partial t} \frac{\partial \bar{x}^i}{\partial x^r} + T^r \frac{\partial^2 \bar{x}^i}{\partial x^s \partial x^r} \frac{\partial x^s}{\partial t} \quad (\text{not a general tensor})$$

\bar{x}^i being a linear functions of $x^r \implies \frac{\partial \bar{T}^i}{\partial t} = \frac{\partial T^r}{\partial t} \frac{\partial \bar{x}^i}{\partial x^r}$
(it's a general tensor)

Goal: Would like the curvature of a curve to be an intrinsic concept, independent of coordinate systems. Use Christoffel symbols to solve it.

2.1 Christoffel Symbol of the 1st Kind

$$\Gamma_{ijk} = \frac{1}{2} \left(-\frac{\partial(g_{ij})}{\partial x^k} + \frac{\partial(g_{jk})}{\partial x^i} + \frac{\partial(g_{ki})}{\partial x^j} \right)$$

$$\text{Let } g_{ijk} = \frac{\partial(g_{ij})}{\partial(x^k)}$$

$$\Gamma_{ijk} = \frac{1}{2}(-g_{ijk} + g_{jki} + g_{kij})$$

Cyclic permute and add:

$$\Gamma_{ijk} + \Gamma_{jki} = g_{kij}$$

Driving transformation law for Γ_{ijk} :

By using g_{ij} derivative and its cyclic permutatations,

$$\bar{\Gamma}_{ijk} = \frac{1}{2}(-\bar{g}_{ijk} + \bar{g}_{jki} + \bar{g}_{kij})$$

where:

$$\bar{g}_{ijk} = \frac{\partial}{\partial \bar{x}^k} (g_{rs} \frac{\partial x^r}{\partial \bar{x}^i} \frac{\partial x^s}{\partial \bar{x}^j}) \quad (\text{expand this})$$

Rearrange and simplify to:

$$\bar{\Gamma}_{ijk} = \Gamma_{rst} \frac{\partial x^r}{\partial \bar{x}^i} \frac{\partial x^s}{\partial \bar{x}^j} \frac{\partial x^t}{\partial \bar{x}^k} + g_{rs} \frac{\partial^2 x^r}{\partial \bar{x}^i \partial \bar{x}^j} \frac{\partial x^s}{\partial \bar{x}^k} \quad (\text{not a general tensor})$$

2.2 Christoffel Symbol of the 2nd Kind

$\Gamma_{ij}^k g^{kr} \Gamma_{ijk}$ (raising an index of Christoffel Symbol of the 1st kind.

Transformation rule for Γ_{ij}^k ; use Γ_{ijk} in derivation:

$$\bar{\Gamma}_{ij}^k = \bar{g}^{kr} \bar{\Gamma}_{ijr}$$

$$\bar{\Gamma}_{ij}^k = \left(g^{sr} \frac{\partial \bar{x}^k}{\partial x^s} \frac{\partial \bar{x}^r}{\partial x^t} \right) \bar{\Gamma}_{ijr}$$

$$\bar{\Gamma}_{ij}^k = \left(g^{sr} \frac{\partial \bar{x}^k}{\partial x^s} \frac{\partial \bar{x}^r}{\partial x^t} \right) \left(\Gamma_{uvw} \frac{\partial x^u}{\partial \bar{x}^i} \frac{\partial x^v}{\partial \bar{x}^j} \frac{\partial x^w}{\partial \bar{x}^r} + g_{uv} \frac{\partial^2 x^u}{\partial \bar{x}^i \partial \bar{x}^j} \frac{\partial x^v}{\partial \bar{x}^r} \right)$$

$$\bar{\Gamma}_{ij}^k = \Gamma_{uv}^w \frac{\partial \bar{x}^k}{\partial x^w} \frac{\partial x^u}{\partial \bar{x}^i} \frac{\partial x^v}{\partial \bar{x}^j} + \frac{\partial \bar{x}^k}{\partial x^w} \frac{\partial^2 x^w}{\partial \bar{x}^i \partial \bar{x}^j} \quad (\text{not a general tensor})$$

linear coordinate transform $\implies \Gamma_{ij}^k$ becomes a tensor

$$\frac{\partial^2 x^u}{\partial \bar{x}^i \partial \bar{x}^j} = \bar{\Gamma}_{ij}^k \frac{\partial x^u}{\partial \bar{x}^k} - \Gamma_{uv}^w \frac{\partial x^u}{\partial \bar{x}^i} \frac{\partial x^v}{\partial \bar{x}^j}$$

2.3 Covariant Derivative of Covariant Vector

let T be a covariant tensor, then:

$$\bar{T}_i = T_r \frac{x^r}{\partial \bar{x}^i}$$

$$\frac{\bar{T}_i}{\partial \bar{x}^k} = \frac{\partial T_r}{\partial \bar{x}^k} \frac{\partial x^r}{\partial \bar{x}^i} + T_r \frac{\partial^2 x^r}{\partial \bar{x}^k \partial \bar{x}^i}$$

use $\frac{\partial^2 x^r}{\partial \bar{x}^i \partial \bar{x}^j}$ from earlier:

$$\frac{\bar{T}_i}{\partial \bar{x}^k} = \frac{\partial T_r}{\partial x^s} \frac{\partial x^s}{\partial \bar{x}^k} \frac{\partial x^r}{\partial \bar{x}^i} + T_r \left(\bar{\Gamma}_{ik}^s \frac{\partial x^r}{\partial \bar{x}^s} - \Gamma_{st}^r \frac{\partial x^s}{\partial \bar{x}^i} \frac{\partial x^t}{\partial \bar{x}^k} \right)$$

$$\frac{\bar{T}_i}{\partial \bar{x}^k} = \frac{\partial T_r}{\partial x^s} \frac{\partial x^s}{\partial \bar{x}^k} \frac{x^r}{\partial \bar{x}^i} + \bar{\Gamma}_{ik}^s \bar{T}_s - \Gamma_{st}^r T_r \frac{\partial x^s}{\partial \bar{x}^i} \frac{\partial x^t}{\partial \bar{x}^k}$$

rename indices:

$$\frac{\bar{T}_i}{\partial \bar{x}^k} = \frac{\partial T_r}{\partial x^s} \frac{\partial x^s}{\partial \bar{x}^k} \frac{x^r}{\partial \bar{x}^i} + \bar{\Gamma}_{ik}^t \bar{T}_t - \Gamma_{rs}^t T_t \frac{\partial x^r}{\partial \bar{x}^i} \frac{\partial x^s}{\partial \bar{x}^k}$$

$$\frac{\bar{T}_i}{\partial \bar{x}^k} = \bar{\Gamma}_{ik}^t \bar{T}_t + \frac{\partial x^s}{\partial \bar{x}^k} \frac{\partial x^r}{\partial \bar{x}^i} \left(\frac{\partial T_r}{\partial x^s} - \Gamma_{rs}^t T_t \right)$$

$$\frac{\bar{T}_i}{\partial \bar{x}^k} - \bar{\Gamma}_{ik}^t \bar{T}_t = \frac{\partial x^s}{\partial \bar{x}^k} \frac{\partial x^r}{\partial \bar{x}^i} \left(\frac{\partial T_r}{\partial x^s} - \Gamma_{rs}^t T_t \right)$$

$$\frac{\bar{T}_i}{\partial \bar{x}^k} - \bar{\Gamma}_{ik}^t \bar{T}_t \text{ is a } (0,2)\text{-tensor}$$

components of covariant derivative wrt. x^k of a covariant vector $T = (T_i)$:

$$T_{,k} = (T_{i,k}) = \frac{\partial T_i}{\partial x^k} - \Gamma_{ik}^t T_t$$

g_{ij} are constants \implies covariant derivatives and partial derivatives coincide

2.4 Covariant Derivative of Contravariant Vector

Similar to previous section, covariant derivative wrt. x^k of a contravariant vector $T = (T^i)$ is:

$$T_{,k} = (T^i_{,k}) = \frac{\partial T^i}{\partial x^k} + \Gamma_{tk}^i T^t$$

Covariant derivative of any tensor is a tensor that has 1 additional covariant order more than the original tensor.

2.5 Absolute Differentiation Along Curve

TODO