

Algebraic Geometry 5

Lecture notes with examples

July 29, 2026

Contents

1 Problem 2 - Intersections of Affine Open Subschemes

Problem. Let X be a scheme, with open affine subsets

$$U = \operatorname{Spec} A, \quad V = \operatorname{Spec} B.$$

Show that $U \cap V$ can be covered by open affine subschemes $\{U_i\}$ such that there are elements

$$f_i \in A, \quad g_i \in B,$$

with

$$U_i = D(f_i) \subseteq U \quad \text{and} \quad U_i = D(g_i) \subseteq V.$$

Solution.

We use the fact that distinguished open subsets form a basis for the topology of an affine scheme.

Let

$$x \in U \cap V.$$

Since $U \cap V$ is open in $U = \operatorname{Spec} A$, and distinguished opens form a basis for $\operatorname{Spec} A$, there exists $f \in A$ such that

$$x \in D(f) \subseteq U \cap V.$$

Thus $D(f)$ is an affine open subset of U , and hence an affine open subscheme of X .

Now $D(f) \subseteq V = \operatorname{Spec} B$. Since $D(f)$ is open in V , and distinguished opens form a basis for V , there exists $g \in B$ such that

$$x \in D(g) \subseteq D(f) \subseteq V.$$

Here $D(g)$ is a distinguished open subset of V .

It remains to see that, after possibly shrinking, this same open set is also distinguished in U . Since

$$D(f) \cong \operatorname{Spec} A_f,$$

every distinguished open inside $D(f)$ is of the form

$$D\left(\frac{h}{1}\right) \subseteq \operatorname{Spec} A_f$$

for some $h \in A$. Under the identification

$$D(f) \subseteq \operatorname{Spec} A,$$

this distinguished open corresponds to

$$D(fh) \subseteq \operatorname{Spec} A.$$

Therefore, after replacing our open neighborhood by a smaller distinguished open if necessary, we obtain an affine open neighborhood U_x of x such that

$$U_x = D(f_x) \subseteq U$$

for some $f_x \in A$, and also

$$U_x = D(g_x) \subseteq V$$

for some $g_x \in B$.

As x varies over $U \cap V$, the sets U_x cover $U \cap V$. Hence $U \cap V$ can be covered by affine opens $\{U_i\}$ such that

$$U_i = D(f_i) \subseteq U, \quad U_i = D(g_i) \subseteq V.$$

Thus

$$\boxed{U \cap V \text{ can be covered by affine opens distinguished in both } U \text{ and } V.}$$

2 Problem 3 - Integral Schemes, Reduced Schemes, and Irreducible Schemes

Problem. We say a scheme X is *irreducible* if it is irreducible as a topological space, i.e. whenever

$$X = X_1 \cup X_2$$

with X_1, X_2 closed subsets, then either $X_1 = X$ or $X_2 = X$.

We say a scheme X is *reduced* if for every open subset $U \subseteq X$, the ring

$$\mathcal{O}_X(U)$$

has no nilpotents.

We say a scheme X is *integral* if for every open subset $U \subseteq X$, the ring

$$\mathcal{O}_X(U)$$

is an integral domain.

Show that a scheme is integral if and only if it is reduced and irreducible.

Solution.

We prove both implications.

Integral implies reduced and irreducible

Assume that X is integral. By definition, for every open subset $U \subseteq X$, the ring

$$\mathcal{O}_X(U)$$

is an integral domain. Every integral domain is reduced. Therefore $\mathcal{O}_X(U)$ has no nilpotents for every open U , so X is reduced.

It remains to prove that X is irreducible.

Recall that a topological space is irreducible if and only if every two nonempty open subsets intersect. Suppose, for contradiction, that X is not irreducible. Then there exist nonempty open subsets

$$U_1, U_2 \subseteq X$$

such that

$$U_1 \cap U_2 = \emptyset.$$

Consider the open subset

$$W = U_1 \cup U_2.$$

Since U_1 and U_2 are disjoint open subsets, the sheaf property gives

$$\mathcal{O}_X(W) \cong \mathcal{O}_X(U_1) \times \mathcal{O}_X(U_2).$$

But the product of two nonzero rings has zero divisors. Indeed,

$$(1, 0)(0, 1) = (0, 0),$$

while

$$(1, 0) \neq 0, \quad (0, 1) \neq 0.$$

Therefore $\mathcal{O}_X(W)$ is not an integral domain. This contradicts the assumption that X is integral.

Hence every two nonempty open subsets of X intersect, so X is irreducible.

Therefore

$$X \text{ integral} \implies X \text{ reduced and irreducible.}$$

Reduced and irreducible implies integral

Now assume that X is reduced and irreducible. We must show that for every open subset $U \subseteq X$,

$$\mathcal{O}_X(U)$$

is an integral domain.

First observe that every nonempty open subset of an irreducible topological space is irreducible. Hence every nonempty open subset $U \subseteq X$ is irreducible.

Let $U \subseteq X$ be a nonempty open subset. Take

$$s, t \in \mathcal{O}_X(U)$$

such that

$$st = 0.$$

We want to show that either $s = 0$ or $t = 0$.

Suppose, for contradiction, that

$$s \neq 0 \quad \text{and} \quad t \neq 0.$$

Then there exist points $x, y \in U$ such that

$$s_x \neq 0, \quad t_y \neq 0.$$

The nonvanishing locus of a section is open, so the sets

$$U_s = \{p \in U : s_p \neq 0\}, \quad U_t = \{p \in U : t_p \neq 0\}$$

are nonempty open subsets of U .

Since U is irreducible, two nonempty open subsets of U must intersect. Therefore

$$U_s \cap U_t \neq \emptyset.$$

Choose

$$p \in U_s \cap U_t.$$

Then

$$s_p \neq 0, \quad t_p \neq 0.$$

Now choose an affine open neighborhood

$$W = \operatorname{Spec} R$$

of p contained in U . Since X is reduced and irreducible, the open subscheme W is also reduced and irreducible. Therefore R is a reduced ring and $\operatorname{Spec} R$ is irreducible.

For an affine scheme $\operatorname{Spec} R$, irreducibility of $\operatorname{Spec} R$ means that the nilradical of R is a prime ideal. Since R is reduced, its nilradical is zero. Hence

$$(0) \subseteq R$$

is prime, so R is an integral domain.

Thus the local ring

$$\mathcal{O}_{X,p}$$

is a localization of an integral domain, hence is itself an integral domain.

But

$$(st)_p = s_p t_p = 0$$

in $\mathcal{O}_{X,p}$. Since $\mathcal{O}_{X,p}$ is an integral domain, this implies

$$s_p = 0 \quad \text{or} \quad t_p = 0,$$

contradicting the choice of p .

Therefore our assumption was false, and so either

$$s = 0 \quad \text{or} \quad t = 0.$$

Hence $\mathcal{O}_X(U)$ has no zero divisors.

Since X is reduced, $\mathcal{O}_X(U)$ also has no nilpotents. Therefore $\mathcal{O}_X(U)$ is an integral domain.

Thus X is integral.

Conclusion

We have proved both directions, so

$X \text{ is integral} \iff X \text{ is reduced and irreducible.}$
--

3 Problem 4 - Locally Noetherian Schemes

Problem. We say a scheme X is *locally Noetherian* if it can be covered by affine open subsets

$$\operatorname{Spec} A_i$$

with A_i a Noetherian ring.

We say a scheme X is *Noetherian* if it can be covered by a finite number of affine open subsets

$$\operatorname{Spec} A_i$$

with A_i Noetherian.

Show that a scheme X is locally Noetherian if and only if for every open affine subset

$$U = \operatorname{Spec} A,$$

the ring A is Noetherian.

Solution.

We prove both implications.

If every affine open has Noetherian coordinate ring, then X is locally Noetherian

Assume that for every affine open subset

$$U = \operatorname{Spec} A \subseteq X,$$

the ring A is Noetherian.

Since X is a scheme, it has an affine open cover

$$X = \bigcup_i U_i, \quad U_i = \operatorname{Spec} A_i.$$

By assumption, each A_i is Noetherian. Therefore X is covered by affine open subsets whose coordinate rings are Noetherian.

Hence X is locally Noetherian.

This direction is immediate from the definition.

If X is locally Noetherian, then every affine open has Noetherian coordinate ring

Now assume that X is locally Noetherian.

Let

$$U = \operatorname{Spec} A$$

be an arbitrary affine open subset of X . We must show that A is Noetherian.

Since X is locally Noetherian, X has an affine open cover

$$X = \bigcup_{\lambda} V_{\lambda}, \quad V_{\lambda} = \operatorname{Spec} B_{\lambda},$$

where every B_{λ} is Noetherian.

Intersecting with U , we get

$$U = \bigcup_{\lambda} (U \cap V_{\lambda}).$$

By the previous problem, each intersection $U \cap V_{\lambda}$ can be covered by affine open subsets which are distinguished both in U and in V_{λ} . Therefore U can be covered by distinguished open subsets

$$D(f_i) \subseteq U = \operatorname{Spec} A$$

such that each $D(f_i)$ is also distinguished inside some Noetherian affine open $V_{\lambda} = \operatorname{Spec} B_{\lambda}$.

Since $U = \operatorname{Spec} A$ is affine, it is quasi-compact. Therefore we may choose finitely many such distinguished opens:

$$U = D(f_1) \cup \cdots \cup D(f_r), \quad f_i \in A.$$

For each i , the open set $D(f_i) \subseteq U$ is affine with coordinate ring

$$\mathcal{O}_X(D(f_i)) \cong A_{f_i}.$$

But $D(f_i)$ is also distinguished inside some $\operatorname{Spec} B_{\lambda}$, with B_{λ} Noetherian. Hence its coordinate ring is a localization of a Noetherian ring, and therefore is Noetherian.

Thus

$$A_{f_i}$$

is Noetherian for every $i = 1, \dots, r$.

Because

$$\operatorname{Spec} A = D(f_1) \cup \cdots \cup D(f_r),$$

the elements $f_1, \dots, f_r$ generate the unit ideal:

$$(f_1, \dots, f_r) = A.$$

We now use the following algebra lemma.

Lemma

Let A be a ring and let $f_1, \dots, f_r \in A$ generate the unit ideal:

$$(f_1, \dots, f_r) = A.$$

If each localization

$$A_{f_i}$$

is Noetherian, then A is Noetherian.

Proof of the lemma.

Let $I \subseteq A$ be an ideal. We show that I is finitely generated.

Since A_{f_i} is Noetherian, the localized ideal

$$I_{f_i} \subseteq A_{f_i}$$

is finitely generated. Hence, for each i , there exist elements

$$a_{i1}, \dots, a_{im_i} \in I$$

such that

$$I_{f_i} = \left(\frac{a_{i1}}{1}, \dots, \frac{a_{im_i}}{1} \right) \subseteq A_{f_i}.$$

Let $J \subseteq A$ be the ideal generated by all these finitely many elements:

$$J = (a_{ij}).$$

Then

$$J \subseteq I.$$

Moreover, after localizing at each f_i , we get

$$J_{f_i} = I_{f_i}.$$

Now take any element

$$a \in I.$$

Since

$$\frac{a}{1} \in I_{f_i} = J_{f_i},$$

there exists an integer $N_i \geq 0$ such that

$$f_i^{N_i} a \in J.$$

Choose N large enough so that

$$N \geq N_i$$

for every i . Then

$$f_i^N a \in J$$

for all $i = 1, \dots, r$.

Since $f_1, \dots, f_r$ generate the unit ideal, the powers

$$f_1^N, \dots, f_r^N$$

also generate the unit ideal. Hence there exist elements

$$b_1, \dots, b_r \in A$$

such that

$$1 = b_1 f_1^N + \dots + b_r f_r^N.$$

Multiplying by a , we obtain

$$a = b_1 f_1^N a + \dots + b_r f_r^N a.$$

Each term $b_i f_i^N a$ lies in J , so

$$a \in J.$$

Therefore

$$I \subseteq J.$$

Since $J \subseteq I$, we get

$$I = J.$$

Thus I is finitely generated.

Since every ideal $I \subseteq A$ is finitely generated, A is Noetherian.

This proves the lemma.

Finishing the proof

We have shown that $U = \operatorname{Spec} A$ admits a finite distinguished open cover

$$U = D(f_1) \cup \cdots \cup D(f_r)$$

such that each localization

$$A_{f_i}$$

is Noetherian, and such that

$$(f_1, \dots, f_r) = A.$$

By the lemma, A is Noetherian.

Since $U = \operatorname{Spec} A$ was an arbitrary affine open subset of X , every affine open subset of X has Noetherian coordinate ring.

Conclusion

Therefore

$$X \text{ is locally Noetherian} \iff \text{for every affine open } U = \operatorname{Spec} A, A \text{ is Noetherian.}$$

4 Theory Behind Problems 2–4

The three problems are connected by one central theme:

$$\boxed{\text{Many properties of schemes are controlled by affine open pieces.}}$$

A scheme is built by gluing affine schemes. Therefore, to understand a scheme X , one repeatedly studies affine open subsets

$$U = \operatorname{Spec} A$$

and translates geometric questions about U into algebraic questions about the ring A .

The three problems develop this philosophy in different directions:

Problem	Main idea	Purpose
2	Affine overlaps can be refined by common distinguished opens.	This lets us compare two affine charts on the same scheme.
3	Integral means reduced and irreducible.	This identifies integral schemes as one-piece schemes with no nilpotent thickening.
4	Locally Noetherian is independent of the chosen affine cover.	This shows that Noetherianity is an intrinsic local property of a scheme.

4.1 Theory for Problem 2: Intersections of Affine Open Subschemes

Let X be a scheme, and let

$$U = \operatorname{Spec} A, \quad V = \operatorname{Spec} B$$

be affine open subsets of X . The problem asks us to show that the overlap

$$U \cap V$$

can be covered by affine open subschemes which are simultaneously distinguished open subsets in U and in V .

The key fact is:

Distinguished open subsets form a basis for the topology of an affine scheme.

For an affine scheme $\operatorname{Spec} A$, the distinguished open subsets are

$$D(f) = \{\mathfrak{p} \in \operatorname{Spec} A : f \notin \mathfrak{p}\}, \quad f \in A.$$

They are not merely open subsets. They are affine open subschemes, because

$$D(f) \cong \operatorname{Spec} A_f.$$

Thus the passage

$$A \longmapsto A_f$$

has the geometric meaning of restricting from $\operatorname{Spec} A$ to the open subset where f is nonzero.

Distinguished opens are stable under intersection

If $f, g \in A$, then

$$D(f) \cap D(g) = D(fg).$$

Indeed, for a prime ideal $\mathfrak{p} \subseteq A$,

$$\mathfrak{p} \in D(f) \cap D(g)$$

means

$$f \notin \mathfrak{p} \quad \text{and} \quad g \notin \mathfrak{p}.$$

Since $\mathfrak{p}$ is prime, this is equivalent to

$$fg \notin \mathfrak{p}.$$

Therefore

$$D(f) \cap D(g) = D(fg).$$

This is one of the reasons affine schemes are so convenient: their standard open subsets behave algebraically.

Distinguished opens inside distinguished opens

Suppose

$$D(f) \subseteq \operatorname{Spec} A.$$

Then

$$D(f) \cong \operatorname{Spec} A_f.$$

A distinguished open subset inside $D(f)$ has the form

$$D\left(\frac{h}{1}\right) \subseteq \operatorname{Spec} A_f$$

for some $h \in A$.

Under the identification

$$D(f) \subseteq \operatorname{Spec} A,$$

this open subset corresponds to

$$D(fh) \subseteq \operatorname{Spec} A.$$

Hence we have the important rule:

A distinguished open inside a distinguished open is again distinguished.

This allows us to refine affine open covers without leaving the class of distinguished opens.

Why the intersection itself need not be affine

If U and V are affine open subsets of a scheme X , it is not true in general that

$$U \cap V$$

is affine.

However, if X is separated, then intersections of affine opens are affine. More precisely,

$$X \text{ separated} \implies U \cap V \text{ is affine for affine opens } U, V.$$

In Problem 2, no separatedness assumption is made. Therefore we cannot prove that $U \cap V$ itself is affine. Instead, we prove the weaker but extremely useful statement that

$$U \cap V$$

can be covered by smaller affine opens which are distinguished from both affine perspectives.

Thus we obtain affine opens U_i such that

$$U_i = D(f_i) \subseteq U = \operatorname{Spec} A$$

and also

$$U_i = D(g_i) \subseteq V = \operatorname{Spec} B.$$

Geometric meaning

The affine opens

$$U = \operatorname{Spec} A, \quad V = \operatorname{Spec} B$$

are like coordinate charts on X . Their overlap

$$U \cap V$$

is the region where both coordinate systems are valid.

Problem 2 says that this overlap can be broken into smaller affine pieces

$$U_i$$

such that each U_i is visible algebraically from both sides:

$$U_i = D(f_i) \subseteq \operatorname{Spec} A, \quad U_i = D(g_i) \subseteq \operatorname{Spec} B.$$

Therefore, on U_i , we can compute either using the ring A or using the ring B .

This is fundamental for gluing schemes and for proving that properties are local.

4.2 Theory for Problem 3: Integral, Reduced, and Irreducible Schemes

Problem 3 proves the equivalence

$$\boxed{X \text{ integral} \iff X \text{ reduced and irreducible.}}$$

This is one of the basic structural results about schemes.

Reduced schemes

A scheme X is reduced if for every open subset $U \subseteq X$, the ring

$$\mathcal{O}_X(U)$$

has no nonzero nilpotent elements.

A nilpotent element is an element a such that

$$a^n = 0$$

for some integer $n > 0$.

Geometrically, nilpotents represent infinitesimal thickening.

For example,

$$\operatorname{Spec} k[x]/(x^2)$$

is not reduced. Its underlying topological space is the same as the space of

$$\operatorname{Spec} k[x]/(x),$$

but its structure sheaf remembers an infinitesimal thickening.

Thus reduced means:

There are no hidden infinitesimal directions.

Irreducible schemes

A topological space X is irreducible if it cannot be written as a union of two proper closed subsets.

Equivalently,

$$X \text{ is irreducible}$$

if and only if every two nonempty open subsets of X intersect.

So irreducibility means that X is one geometric piece.

For example,

$$\operatorname{Spec} k[x, y]/(xy)$$

is reducible, because the equation

$$xy = 0$$

describes the union of two coordinate axes:

$$x = 0 \quad \text{or} \quad y = 0.$$

Algebraically, this reducibility appears through zero divisors:

$$x \cdot y = 0$$

in the ring $k[x, y]/(xy)$, even though neither x nor y is zero.

Integral schemes

A ring A is an integral domain if

$$ab = 0 \implies a = 0 \text{ or } b = 0.$$

An affine scheme

$$\operatorname{Spec} A$$

is integral precisely when A is an integral domain.

Now recall two standard affine facts:

$$\operatorname{Spec} A \text{ is reduced} \iff A \text{ is reduced,}$$

and

$$\operatorname{Spec} A \text{ is irreducible} \iff \sqrt{(0)} \text{ is a prime ideal.}$$

Here

$$\sqrt{(0)} = \sqrt{(0)}$$

is the nilradical of A , the ideal of all nilpotent elements of A .

If A is reduced, then

$$\sqrt{(0)} = (0).$$

Therefore, if A is reduced and $\operatorname{Spec} A$ is irreducible, then

$$(0)$$

is a prime ideal.

But (0) is prime exactly when A is an integral domain.

Hence, in the affine case, we have

$A \text{ is an integral domain} \iff A \text{ is reduced and } \operatorname{Spec} A \text{ is irreducible.}$
--

From affine schemes to arbitrary schemes

For an arbitrary scheme X , the theorem is global:

$$X \text{ integral} \iff X \text{ reduced and irreducible.}$$

The reason is that every scheme is covered by affine open subsets.

If X is reduced and irreducible, then every nonempty affine open subset

$$U = \operatorname{Spec} A \subseteq X$$

is also reduced and irreducible. Hence A is an integral domain.

Thus locally, all affine pieces of X are spectra of domains.

Conversely, if X is integral, then every ring of functions on an open subset is an integral domain. Since every integral domain is reduced, X is reduced. Also, if X could be decomposed into two proper closed pieces, then one could find disjoint nonempty open subsets, producing zero divisors in a suitable ring of functions. That contradicts integrality.

Geometric intuition

Reduced removes nilpotents.

Irreducible removes multiple components.

Integral means both conditions at once:

An integral scheme is one genuine algebraic component with no nilpotent fuzz.

Typical examples are:

Scheme	Reduced?	Irreducible?	Integral?
$\text{Spec } k[x]$	yes	yes	yes
$\text{Spec } k[x]/(x^2)$	no	yes	no
$\text{Spec } k[x, y]/(xy)$	yes	no	no
$\text{Spec } k[x, y]/(xy, x^2)$	no	no	no

4.3 Theory for Problem 4: Locally Noetherian Schemes

Problem 4 proves that

X is locally Noetherian $\iff$ every affine open $U = \text{Spec } A \subseteq X$ has A Noetherian.

This is an affine-locality theorem.

Noetherian rings

A ring A is Noetherian if every ideal of A is finitely generated.

Equivalently, every ascending chain of ideals

$$I_1 \subseteq I_2 \subseteq I_3 \subseteq \cdots$$

eventually stabilizes.

That means there exists some N such that

$$I_N = I_{N+1} = I_{N+2} = \cdots.$$

So Noetherian rings are algebraically finite objects.

Important examples include polynomial rings over fields:

$$k[x_1, \dots, x_n],$$

which are Noetherian by Hilbert's basis theorem.

Also, localizations of Noetherian rings are Noetherian:

$$A \text{ Noetherian} \implies A_f \text{ Noetherian.}$$

Geometrically,

$$A_f$$

is the coordinate ring of the distinguished open

$$D(f) \subseteq \operatorname{Spec} A.$$

Thus Noetherianity is preserved when passing to distinguished affine open subsets.

Locally Noetherian schemes

A scheme X is locally Noetherian if it has an affine open cover

$$X = \bigcup_i \operatorname{Spec} A_i$$

such that every A_i is Noetherian.

This means that near every point, X looks like the spectrum of a Noetherian ring.

So locally Noetherian schemes are schemes of finite algebraic complexity near every point.

Why the theorem is not trivial

The definition of locally Noetherian only says that there exists some affine open cover by Noetherian rings.

Problem 4 proves something stronger:

If one Noetherian affine cover exists, then every affine open is Noetherian.

This is important because it means that being locally Noetherian is not dependent on a chosen cover. It is an intrinsic property of the scheme.

The key algebra lemma

The proof relies on the following algebraic fact.

Let A be a ring, and suppose

$$f_1, \dots, f_r \in A$$

generate the unit ideal:

$$(f_1, \dots, f_r) = A.$$

If each localization

$$A_{f_i}$$

is Noetherian, then A is Noetherian.

In symbols,

$$(f_1, \dots, f_r) = A \text{ and all } A_{f_i} \text{ Noetherian} \implies A \text{ Noetherian.}$$

Geometrically, the condition

$$(f_1, \dots, f_r) = A$$

means that the distinguished opens cover $\text{Spec } A$:

$$\text{Spec } A = D(f_1) \cup \dots \cup D(f_r).$$

Therefore, if $\text{Spec } A$ can be covered by finitely many distinguished opens whose coordinate rings are Noetherian, then A itself is Noetherian.

Why finite covers appear

Every affine scheme $\text{Spec } A$ is quasi-compact.

This means that every open cover of $\text{Spec } A$ admits a finite subcover.

Therefore, if

$$U = \text{Spec } A$$

is covered by many small affine open subsets with Noetherian coordinate rings, then finitely many of them already cover U .

After refining to distinguished opens, we obtain a finite cover

$$U = D(f_1) \cup \dots \cup D(f_r).$$

Then the coordinate rings are

$$\mathcal{O}_X(D(f_i)) \cong A_{f_i}.$$

If each A_{f_i} is Noetherian and the opens $D(f_i)$ cover U , then

$$(f_1, \dots, f_r) = A.$$

The algebra lemma then implies:

$$A \text{ is Noetherian.}$$

Role of Problem 2 in Problem 4

Problem 2 is used to compare an arbitrary affine open subset with a fixed Noetherian affine cover.

Suppose X has a Noetherian affine cover

$$X = \bigcup_{\lambda} V_{\lambda}, \quad V_{\lambda} = \text{Spec } B_{\lambda},$$

where each B_{λ} is Noetherian.

Now take an arbitrary affine open subset

$$U = \text{Spec } A \subseteq X.$$

The intersections

$$U \cap V_\lambda$$

may not themselves be affine.

But Problem 2 says that each $U \cap V_\lambda$ can be covered by affine opens which are distinguished both in U and in V_λ .

Therefore U can be covered by distinguished opens

$$D(f_i) \subseteq U$$

which are also distinguished opens inside some Noetherian affine scheme V_λ .

Since distinguished opens inside Noetherian affine schemes have Noetherian coordinate rings, each

$$A_{f_i}$$

is Noetherian.

Because $U = \operatorname{Spec} A$ is quasi-compact, finitely many such distinguished opens cover U . Hence

$$U = D(f_1) \cup \cdots \cup D(f_r),$$

with each A_{f_i} Noetherian.

By the algebra lemma,

$$A$$

is Noetherian.

Thus every affine open subset of X has Noetherian coordinate ring.

4.4 Conceptual Summary

The three results form a useful local-to-global toolkit.

Result	Local statement	Global meaning
Affine overlap refinement	Overlaps of affine opens can be covered by common distinguished opens.	Different affine charts can be compared on smaller affine pieces.
Integral = reduced + irreducible	On affine opens, domains correspond to reduced irreducible spectra.	An integral scheme is one reduced algebraic component.
Locally Noetherian criterion	Noetherianity can be checked on arbitrary affine opens.	The property is intrinsic and independent of the chosen cover.

The deepest common principle is:

Schemes are controlled by affine opens, and affine opens are controlled by rings.

More specifically,

Local affine data determines global scheme-theoretic properties.

5 Examples for Problems 2–4

We now give concrete examples illustrating the three main themes:

affine overlaps, integral = reduced + irreducible, locally Noetherian schemes.

The purpose of these examples is to show how the abstract definitions become explicit ring computations.

5.1 Examples for Intersections of Affine Open Subschemes

Example 1: Distinguished opens in $\mathbb{A}_k^1$

Let

$$X = \mathbb{A}_k^1 = \operatorname{Spec} k[x].$$

Take the two distinguished open subsets

$$U = D(x), \quad V = D(x - 1).$$

Then

$$U = \operatorname{Spec} k[x]_x, \quad V = \operatorname{Spec} k[x]_{x-1}.$$

Their intersection is

$$U \cap V = D(x) \cap D(x - 1).$$

Since distinguished opens satisfy

$$D(f) \cap D(g) = D(fg),$$

we get

$$U \cap V = D(x(x - 1)).$$

Thus the overlap is itself one distinguished affine open:

$$U \cap V = \operatorname{Spec} k[x]_{x(x-1)}.$$

Inside $U = D(x)$, the same open set is obtained by additionally requiring $x - 1 \neq 0$:

$$U \cap V = D(x - 1) \subseteq U.$$

Inside $V = D(x - 1)$, the same open set is obtained by additionally requiring $x \neq 0$:

$$U \cap V = D(x) \subseteq V.$$

Therefore the same affine open subset is distinguished from both affine charts.

$$\boxed{D(x) \cap D(x - 1) = D(x(x - 1))}.$$

This is the simplest example of the phenomenon in Problem 2.

Example 2: The standard affine cover of $\mathbb{P}_k^1$

Let

$$X = \mathbb{P}_k^1.$$

The standard affine open cover is

$$\mathbb{P}_k^1 = U_0 \cup U_1,$$

where

$$U_0 = D_+(X_0), \quad U_1 = D_+(X_1).$$

We have

$$U_0 \cong \operatorname{Spec} k[t], \quad t = \frac{X_1}{X_0},$$

and

$$U_1 \cong \operatorname{Spec} k[s], \quad s = \frac{X_0}{X_1}.$$

On the overlap $U_0 \cap U_1$, both X_0 and X_1 are nonzero. Therefore

$$t = \frac{X_1}{X_0}$$

is nonzero, and also

$$s = \frac{X_0}{X_1}$$

is nonzero.

Moreover,

$$s = \frac{1}{t}.$$

Hence

$$U_0 \cap U_1 = D(t) \subseteq U_0 = \operatorname{Spec} k[t],$$

and also

$$U_0 \cap U_1 = D(s) \subseteq U_1 = \operatorname{Spec} k[s].$$

Therefore

$$U_0 \cap U_1 \cong \operatorname{Spec} k[t, t^{-1}] \cong \operatorname{Spec} k[s, s^{-1}].$$

This open subset is the multiplicative group

$$\mathcal{G}_m = \operatorname{Spec} k[t, t^{-1}].$$

So

$$\boxed{U_0 \cap U_1 = D(t) \subseteq \operatorname{Spec} k[t] = D(s) \subseteq \operatorname{Spec} k[s].}$$

This is the standard geometric example of an affine overlap which is distinguished from both coordinate charts.

Example 3: Two standard opens in the affine plane

Let

$$X = \mathbb{A}_k^2 = \operatorname{Spec} k[x, y].$$

Take

$$U = D(x), \quad V = D(y).$$

Then

$$U = \operatorname{Spec} k[x, y]_x, \quad V = \operatorname{Spec} k[x, y]_y.$$

Their intersection is

$$U \cap V = D(x) \cap D(y) = D(xy).$$

Thus

$$U \cap V \cong \operatorname{Spec} k[x, y]_{xy}.$$

Inside

$$U = \operatorname{Spec} k[x, y]_x,$$

we obtain the overlap by additionally inverting y :

$$U \cap V = D(y) \subseteq U.$$

Inside

$$V = \operatorname{Spec} k[x, y]_y,$$

we obtain the overlap by additionally inverting x :

$$U \cap V = D(x) \subseteq V.$$

Therefore

$$\boxed{D(x) \cap D(y) = D(xy).}$$

This example shows directly how intersections of affine opens are computed by localization.

5.2 Examples for Integral, Reduced, and Irreducible Schemes

Example 4: $\text{Spec } k[x]$ is integral

Let

$$X = \text{Spec } k[x].$$

The ring $k[x]$ is an integral domain. Therefore X is an integral affine scheme.

It is reduced because $k[x]$ has no nonzero nilpotent elements.

It is irreducible because the zero ideal

$$(0) \subseteq k[x]$$

is prime.

Indeed,

$$\text{Spec } A \text{ is irreducible} \iff \sqrt{(0)} \text{ is prime in } A.$$

Since $k[x]$ is reduced,

$$\sqrt{(0)} = (0),$$

and since $k[x]$ is a domain, (0) is prime.

Hence

$$\text{Spec } k[x]$$

is reduced, irreducible, and integral.

Geometrically, this is the affine line. It is one algebraic component and has no nilpotent thickening.

$\text{Spec } k[x] \text{ is integral.}$

Example 5: $\text{Spec } k[x]/(x^2)$ is irreducible but not reduced

Let

$$X = \text{Spec } k[x]/(x^2).$$

In the ring

$$A = k[x]/(x^2),$$

the element x is nonzero, but

$$x^2 = 0.$$

Therefore x is a nonzero nilpotent element. Hence A is not reduced, and so X is not reduced.

Now we check irreducibility.

The nilradical of A is

$$\sqrt{(0)} = (x).$$

The quotient by this ideal is

$$A/(x) = \frac{k[x]/(x^2)}{(x)} \cong k.$$

Since k is a field, it is an integral domain. Therefore (x) is a prime ideal of A .

Hence

$$\sqrt{(0)}$$

is prime, so

$$\operatorname{Spec} k[x]/(x^2)$$

is irreducible.

Thus this scheme is irreducible but not reduced.

$\operatorname{Spec} k[x]/(x^2) \text{ is irreducible but not reduced.}$

Geometrically, it is a doubled point: its underlying topological space is one point, but its structure sheaf contains infinitesimal thickening.

Therefore it is not integral.

Example 6: $\operatorname{Spec} k[x, y]/(xy)$ is reduced but reducible

Let

$$X = \operatorname{Spec} k[x, y]/(xy).$$

The equation

$$xy = 0$$

describes the union of the two coordinate axes:

$$x = 0 \quad \text{or} \quad y = 0.$$

Topologically,

$$V(xy) = V(x) \cup V(y).$$

Therefore the scheme is reducible.

Now consider the coordinate ring

$$A = k[x, y]/(xy).$$

The ideal $(xy) \subseteq k[x, y]$ is radical, so A is reduced. Hence X is reduced.

However, A is not an integral domain. In A , we have

$$x \cdot y = 0,$$

but

$$x \neq 0, \quad y \neq 0.$$

Therefore x and y are zero divisors.

So X is reduced but reducible, and hence not integral.

$\operatorname{Spec} k[x, y]/(xy)$ is reduced but reducible.

This example shows that reducedness alone does not imply integrality.

Example 7: $\operatorname{Spec} k[x, y]/(y^2 - x^3)$ is integral but singular

Let

$$X = \operatorname{Spec} k[x, y]/(y^2 - x^3).$$

The polynomial

$$y^2 - x^3$$

is irreducible in $k[x, y]$. Therefore the ideal

$$(y^2 - x^3)$$

is prime, and hence the quotient ring

$$k[x, y]/(y^2 - x^3)$$

is an integral domain.

Therefore

$$X$$

is an integral scheme.

Geometrically, this is the cuspidal cubic curve.

It has a singular point at the origin, but the coordinate ring is still a domain. Thus the scheme is reduced and irreducible, even though it is singular.

This example is important because it shows that singularity is a different concept from non-integrality.

Integral schemes may have singularities.

5.3 Examples for Locally Noetherian Schemes

Example 8: Affine space is locally Noetherian

Let

$$X = \mathbb{A}_k^n = \operatorname{Spec} k[x_1, \dots, x_n].$$

By Hilbert's basis theorem, the polynomial ring

$$k[x_1, \dots, x_n]$$

is Noetherian.

Therefore

$$\mathbb{A}_k^n$$

is locally Noetherian.

In fact, it is Noetherian, because it is covered by one affine open subset with Noetherian coordinate ring.

$\mathbb{A}_k^n$ is Noetherian.

Example 9: Projective space is Noetherian

Let

$$X = \mathbb{P}_k^n.$$

The standard affine open cover is

$$\mathbb{P}_k^n = D_+(X_0) \cup D_+(X_1) \cup \cdots \cup D_+(X_n).$$

Each standard open is isomorphic to affine n -space:

$$D_+(X_i) \cong \mathbb{A}_k^n.$$

Therefore each has Noetherian coordinate ring, because each coordinate ring is a polynomial ring in n variables over k .

Since the cover

$$\mathbb{P}_k^n = \bigcup_{i=0}^n D_+(X_i)$$

is finite, $\mathbb{P}_k^n$ is Noetherian.

Therefore it is also locally Noetherian.

Problem 4 tells us something stronger:

Every affine open subset of $\mathbb{P}_k^n$ has a Noetherian coordinate ring.

Example 10: An infinite disjoint union of points

Let

$$X = \coprod_{n \geq 1} \operatorname{Spec} k.$$

Each component

$$\operatorname{Spec} k$$

is affine and Noetherian.

Therefore X has an affine open cover by Noetherian affine schemes. Hence X is locally Noetherian.

However, X is not Noetherian in the finite-cover sense, because it cannot be covered by finitely many of its point-components.

Equivalently, X is not quasi-compact.

Thus

$$X = \coprod_{n \geq 1} \operatorname{Spec} k$$

is locally Noetherian but not Noetherian.

$\coprod_{n \geq 1} \operatorname{Spec} k \text{ is locally Noetherian but not Noetherian.}$
--

This example shows the difference between local finiteness and global finiteness.

Example 11: A non-locally Noetherian affine scheme

Let

$$A = k[x_1, x_2, x_3, \dots]$$

be the polynomial ring in infinitely many variables.

This ring is not Noetherian.

Indeed, consider the ascending chain of ideals

$$(x_1) \subset (x_1, x_2) \subset (x_1, x_2, x_3) \subset \cdots$$

This chain never stabilizes. Therefore A is not Noetherian.

Now let

$$X = \operatorname{Spec} A.$$

Since X itself is affine, if X were locally Noetherian, then by the criterion from Problem 4 every affine open subset of X would have Noetherian coordinate ring.

In particular, the affine open subset

$$X = \operatorname{Spec} A$$

itself would have Noetherian coordinate ring.

That would imply that A is Noetherian, which is false.

Therefore

$\operatorname{Spec} k[x_1, x_2, x_3, \dots] \text{ is not locally Noetherian.}$
--

5.4 Summary Table

Example	Scheme	Main lesson
1	$D(x), D(x-1) \subseteq \mathbb{A}_k^1$	Intersections of distinguished opens are distinguished.
2	$\mathbb{P}_k^1$	The standard projective affine overlap is a common distinguished open.
3	$D(x), D(y) \subseteq \mathbb{A}_k^2$	Affine overlaps are computed by localization.
4	$\operatorname{Spec} k[x]$	Basic example of an integral scheme.
5	$\operatorname{Spec} k[x]/(x^2)$	Irreducible but not reduced.
6	$\operatorname{Spec} k[x, y]/(xy)$	Reduced but reducible.
7	$\operatorname{Spec} k[x, y]/(y^2 - x^3)$	Integral schemes may still be singular.
8	$\mathbb{A}_k^n$	Affine space is Noetherian.
9	$\mathbb{P}_k^n$	Projective space is Noetherian.
10	$\coprod_{n \geq 1} \operatorname{Spec} k$	Locally Noetherian but not Noetherian.
11	$\operatorname{Spec} k[x_1, x_2, x_3, \dots]$	Not locally Noetherian.

5.5 Main Computational Patterns

The examples repeatedly use the following basic computations:

$$D(f) \cap D(g) = D(fg).$$

This describes intersections of distinguished opens.

$$D(f) \cong \operatorname{Spec} A_f.$$

This says that passing to a distinguished open corresponds to localization.

$$\operatorname{Spec} A \text{ is integral} \iff A \text{ is an integral domain.}$$

This translates integrality of affine schemes into a ring-theoretic condition.

$$\operatorname{Spec} A \text{ is reduced} \iff A \text{ is reduced.}$$

This translates absence of nilpotents into geometry.

$$\text{Spec } A \text{ is irreducible} \iff \sqrt{(0)} \text{ is prime.}$$

This relates irreducibility to the nilradical.

$$A \text{ Noetherian} \implies A_f \text{ Noetherian.}$$

This says that Noetherianity is preserved under localization.

Finally,

$$\text{Spec } A = D(f_1) \cup \cdots \cup D(f_r) \iff (f_1, \dots, f_r) = A.$$

This connects finite distinguished open covers with the algebraic condition that the functions $f_1, \dots, f_r$ generate the unit ideal.

The main lesson is:

In affine examples, scheme theory becomes explicit commutative algebra.

This is the basic mechanism of scheme theory:

$$\text{geometry of schemes} \longleftrightarrow \text{commutative algebra of rings.}$$

6 Charts, Diagrams, and Tables for Problems 2–4

This section collects visual summaries for the three main themes:

affine overlap refinement, integral = reduced+irreducible, locally Noetherian schemes.

The goal is to make the main logical structure visible.

6.1 Affine Overlap Refinement

Let X be a scheme and let

$$U = \text{Spec } A, \quad V = \text{Spec } B$$

be affine open subsets of X .


The overlap

$$U \cap V$$

need not be affine in general. However, it can be covered by affine open subschemes U_i such that

$$U_i = D(f_i) \subseteq U, \quad U_i = D(g_i) \subseteq V.$$

Thus each small open U_i is distinguished from both affine charts.

Diagram: Common Distinguished Refinement

The meaning is:

On each U_i , both coordinate systems are valid.

6.2 Distinguished Open Dictionary

Problem 2 is mostly about the behavior of distinguished open subsets.

Geometric object	Algebraic object	Meaning
$\text{Spec } A$	A	An affine scheme is controlled by its coordinate ring.
$D(f) \subseteq \text{Spec } A$	A_f	Restricting to $D(f)$ means inverting f .
$D(f) \cap D(g)$	A_{fg}	Being in both opens means inverting both f and g .
$D(f_1) \cup \cdots \cup D(f_r) = \text{Spec } A$	$(f_1, \dots, f_r) = A$	The functions $f_1, \dots, f_r$ have no common zero.

The central computational rule is:

$$D(f) \cap D(g) = D(fg).$$

6.3 Diagram: The Standard Cover of $\mathbb{P}_k^1$

The projective line has the standard affine cover

$$\mathbb{P}_k^1 = U_0 \cup U_1,$$

where

$$U_0 = D_+(X_0) \cong \text{Spec } k[t], \quad t = \frac{X_1}{X_0},$$

and

$$U_1 = D_+(X_1) \cong \operatorname{Spec} k[s], \quad s = \frac{X_0}{X_1}.$$

On the overlap,


$$s = \frac{1}{t}.$$

Therefore

$$U_0 \cap U_1 = D(t) \subseteq U_0$$

and also

$$U_0 \cap U_1 = D(s) \subseteq U_1.$$



Thus

$$U_0 \cap U_1 = D(t) \subseteq \operatorname{Spec} k[t] = D(s) \subseteq \operatorname{Spec} k[s].$$

This is the standard example of gluing affine charts along a common distinguished open subset.


6.4 Reduced, Irreducible, and Integral Schemes

Problem 3 proves:

$$X \text{ integral} \iff X \text{ reduced and irreducible.}$$

The two conditions have different meanings.

Property	Algebraic meaning	Geometric meaning
<i>Reduced</i>	<i>Nonzeronilpotents</i>	<i>No infinitesimal thickening</i>
<i>Irreducible</i>	<i>Nilradical is prime on a fine opens</i>	<i>One topological component</i>
<i>Integral</i>	<i>Coordinaterings are domains</i>	<i>One reduced algebraic component</i>

Diagram: Integral as Reduced plus Irreducible

So one should remember:

$$\boxed{\text{Integral} = \text{no nilpotents} + \text{one component.}}$$

6.5 Comparison Table: Reduced, Irreducible, Integral

Scheme	Reduced?	Irreducible?	Integral?
$\text{Spec } k[x]$	yes	yes	yes
$\text{Spec } k[x]/(x^2)$	no	yes	no
$\text{Spec } k[x, y]/(xy)$	yes	no	no
$\text{Spec } k[x, y]/(y^2 - x^3)$	yes	yes	yes

This table shows why both reducedness and irreducibility are necessary.

6.6 Table: Algebraic Meaning of Geometric Pathologies

Geometry	Algebra	Example	Effect
Nilpotent thickening	There exists $a \neq 0$ with $a^n = 0$.	$k[x]/(x^2)$	The scheme is not reduced.
Multiple components	There exist $a, b \neq 0$ with $ab = 0$.	$k[x, y]/(xy)$	The scheme is reducible and not a domain.
One reduced component	The zero ideal (0) is prime.	$k[x]$	The scheme is integral.
Singular but one component	The ring is a domain but not smooth.	$k[x, y]/(y^2 - x^3)$	The scheme is integral but singular.

This table separates three notions:

reduced, irreducible, smooth.

A singular scheme can still be integral.


6.7 Locally Noetherian Schemes

Problem 4 proves:

$$X \text{ locally Noetherian} \iff \text{every affine open } U = \operatorname{Spec} A \subseteq X \text{ has } A \text{ Noetherian.}$$

This means that being locally Noetherian is independent of the chosen affine cover.

Flowchart: Proof of the Locally Noetherian Criterion



The proof uses three ingredients:

$$\text{affine overlap refinement} + \text{quasi-compactness} + \text{localization lemma.}$$

6.8 Locally Noetherian versus Noetherian

Scheme	Locally Noetherian?	Noetherian?	Reason
$\operatorname{Spec} k[x_1, \dots, x_n]$	<i>yes</i>	<i>yes</i>	It has one Noetherian affine chart.
$\mathbb{P}_k^n$	<i>yes</i>	<i>yes</i>	It has a finite affine cover by copies of $\mathbb{A}_k^n$.
$\coprod_{n \geq 1} \operatorname{Spec} k$	<i>yes</i>	<i>no</i>	Each point is Noetherian, but there are infinitely many components.
$\operatorname{Spec} k[x_1, x_2, x_3, \dots]$	<i>no</i>	<i>no</i>	The coordinate ring is not Noetherian.

This table separates:

local finite algebraic complexity

from


global finite cover by such pieces.

6.9 Dependency Table for Problems 2–4

Problem	Key tool	What it proves
<i>Problem2</i>	Distinguished opens form a basis of affine schemes.	Affine overlaps can be refined by common distinguished opens.
<i>Problem3</i>	$\operatorname{Spec} A$ is integral if and only if A is a domain.	A scheme is integral exactly when it is reduced and irreducible.
<i>Problem4</i>	Problem 2, quasi-compactness of affine schemes, and the algebra lemma on localizations.	Locally Noetherian is independent of the chosen affine cover.

This shows that Problem 4 depends strongly on the overlap refinement from Problem 2.

6.10 Final Summary Diagram



The main principle is:

Schemes are controlled by affine opens, and affine opens are controlled by rings.

Thus, in these problems, scheme theory becomes explicit commutative algebra:

geometry of schemes $\longleftrightarrow$ commutative algebra of rings.

7 Problem 5: Locally of finite type on every affine open

Problem. A morphism $f : X \rightarrow Y$ is *locally of finite type* if there exists a covering of Y by open affine subsets

$$V_i = \operatorname{Spec} B_i$$

such that, for each i , $f^{-1}(V_i)$ can be covered by open affine subsets

$$U_{ij} = \operatorname{Spec} A_{ij},$$

where each A_{ij} is a finitely generated B_i -algebra.

The morphism is *of finite type* if the cover of $f^{-1}(V_i)$ above can be taken to be finite.

Show that a morphism $f : X \rightarrow Y$ is locally of finite type if and only if for every open affine subset

$$V = \operatorname{Spec} B$$

of Y , $f^{-1}(V)$ can be covered by open affine subsets

$$U_j = \operatorname{Spec} A_j,$$

where each A_j is a finitely generated B -algebra.

Solution.

We prove both directions.

First direction

Assume that $f : X \rightarrow Y$ is locally of finite type.

By definition, there exists an affine open covering

$$Y = \bigcup_i V_i, \quad V_i = \operatorname{Spec} B_i,$$

such that, for each i ,

$$f^{-1}(V_i) = \bigcup_j U_{ij}, \quad U_{ij} = \operatorname{Spec} A_{ij},$$

and each A_{ij} is a finitely generated B_i -algebra.

Let now

$$V = \operatorname{Spec} B$$

be an arbitrary open affine subset of Y . We must show that $f^{-1}(V)$ admits a covering by affine opens whose coordinate rings are finitely generated B -algebras.

Since the V_i cover Y , the open subsets

$$V \cap V_i$$

cover V . Although $V \cap V_i$ need not itself be affine in a general scheme, it can be covered by affine opens which are distinguished both in V and in V_i . Thus, for each i , we may cover $V \cap V_i$ by open affine subsets

$$W_{i\alpha}$$

such that

$$W_{i\alpha} = D(b_{i\alpha}) \subseteq V = \operatorname{Spec} B$$

and also

$$W_{i\alpha} = D(c_{i\alpha}) \subseteq V_i = \operatorname{Spec} B_i.$$

Therefore

$$W_{i\alpha} \cong \operatorname{Spec} B_{b_{i\alpha}}$$

as an open subset of V , and

$$W_{i\alpha} \cong \operatorname{Spec}(B_i)_{c_{i\alpha}}$$

as an open subset of V_i .

Now consider the inverse image

$$f^{-1}(W_{i\alpha}) \subseteq f^{-1}(V_i).$$

Since $f^{-1}(V_i)$ is covered by the affine opens

$$U_{ij} = \operatorname{Spec} A_{ij},$$

the sets

$$U_{ij} \cap f^{-1}(W_{i\alpha})$$

cover $f^{-1}(W_{i\alpha})$.

Inside $U_{ij} = \operatorname{Spec} A_{ij}$, the inverse image of the distinguished open

$$W_{i\alpha} = D(c_{i\alpha}) \subseteq \operatorname{Spec} B_i$$

is the distinguished open

$$D(\varphi(c_{i\alpha})) \subseteq \operatorname{Spec} A_{ij},$$

where

$$\varphi : B_i \rightarrow A_{ij}$$

is the ring map induced by the morphism on this affine piece.

Hence

$$U_{ij} \cap f^{-1}(W_{i\alpha}) \cong \operatorname{Spec}(A_{ij})_{\varphi(c_{i\alpha})}.$$

Since A_{ij} is a finitely generated B_i -algebra, its localization

$$(A_{ij})_{\varphi(c_{i\alpha})}$$

is a finitely generated $(B_i)_{c_{i\alpha}}$ -algebra.

But

$$(B_i)_{c_{i\alpha}} \cong B_{b_{i\alpha}},$$

because both are coordinate rings of the same affine open subset $W_{i\alpha}$.

Therefore

$$(A_{ij})_{\varphi(c_{i\alpha})}$$

is a finitely generated $B_{b_{i\alpha}}$ -algebra.

Finally,

$$B_{b_{i\alpha}}$$

is a finitely generated B -algebra, because

$$B_{b_{i\alpha}} \cong B[t]/(b_{i\alpha}t - 1).$$

Therefore any finitely generated $B_{b_{i\alpha}}$ -algebra is also a finitely generated B -algebra.

So each affine open

$$U_{ij} \cap f^{-1}(W_{i\alpha})$$

has coordinate ring finitely generated over B .

Since the opens $W_{i\alpha}$ cover V , their inverse images cover $f^{-1}(V)$. Therefore the affine opens

$$U_{ij} \cap f^{-1}(W_{i\alpha})$$

cover $f^{-1}(V)$, and each has coordinate ring finitely generated over B .

Thus, for every affine open

$$V = \operatorname{Spec} B \subseteq Y,$$

the inverse image $f^{-1}(V)$ can be covered by affine opens

$$\operatorname{Spec} A_j$$

with each A_j a finitely generated B -algebra.

Second direction

Conversely, assume that for every affine open subset

$$V = \operatorname{Spec} B \subseteq Y,$$

the inverse image $f^{-1}(V)$ can be covered by affine opens

$$U_j = \operatorname{Spec} A_j,$$

where each A_j is a finitely generated B -algebra.

Choose any affine open covering

$$Y = \bigcup_i V_i, \quad V_i = \operatorname{Spec} B_i.$$

By assumption, for each i , the inverse image

$$f^{-1}(V_i)$$

can be covered by affine opens

$$U_{ij} = \operatorname{Spec} A_{ij},$$

where each A_{ij} is a finitely generated B_i -algebra.

But this is exactly the definition of f being locally of finite type.

Therefore f is locally of finite type.

Conclusion

We have proved that $f : X \rightarrow Y$ is locally of finite type if and only if for every affine open subset

$$V = \operatorname{Spec} B \subseteq Y,$$

the inverse image $f^{-1}(V)$ can be covered by affine open subsets

$$U_j = \operatorname{Spec} A_j$$

such that each A_j is a finitely generated B -algebra.

8 Problem 6: Examples violating the properties

Problem. For each of the properties defined above of schemes or morphisms, give an example of a scheme or morphism which violates that property. Give an example of a morphism which is locally of finite type but not of finite type.

Solution.

We list standard examples.

A scheme which is not reduced

Let

$$X = \operatorname{Spec} k[\varepsilon]/(\varepsilon^2).$$

The ring

$$k[\varepsilon]/(\varepsilon^2)$$

has a nonzero nilpotent element, namely ε , because

$$\varepsilon \neq 0, \quad \varepsilon^2 = 0.$$

Therefore the ring is not reduced.

Hence

$$\operatorname{Spec} k[\varepsilon]/(\varepsilon^2)$$

is a scheme which is not reduced.

A scheme which is not irreducible

Let

$$X = \operatorname{Spec} k[x, y]/(xy).$$

Geometrically, this is the union of the two coordinate axes in the affine plane.

Indeed, in $\mathbb{A}_k^2 = \operatorname{Spec} k[x, y]$, the equation

$$xy = 0$$

means

$$x = 0 \quad \text{or} \quad y = 0.$$

Thus

$$\operatorname{Spec} k[x, y]/(xy)$$

is the union of the closed subsets

$$V(x) \quad \text{and} \quad V(y).$$

Both are proper closed subsets. Therefore X is reducible.

Hence

$$X = \operatorname{Spec} k[x, y]/(xy)$$

is not irreducible.

A scheme which is not integral

A scheme is integral if it is both reduced and irreducible.

Thus either a non-reduced scheme or a reducible scheme gives an example of a non-integral scheme.

For example,

$$X = \operatorname{Spec} k[x, y]/(xy)$$

is reduced but reducible. Therefore it is not integral.

Another example is

$$X = \operatorname{Spec} k[\varepsilon]/(\varepsilon^2),$$

which is not reduced. Hence it is not integral.

A morphism which is not locally of finite type

Consider the morphism

$$f : \operatorname{Spec} k[x_1, x_2, x_3, \dots] \longrightarrow \operatorname{Spec} k.$$

This morphism corresponds to the inclusion

$$k \longrightarrow k[x_1, x_2, x_3, \dots].$$

The ring

$$k[x_1, x_2, x_3, \dots]$$

is not a finitely generated k -algebra, because it has infinitely many algebraically independent variables.

Since the target is affine,

$$\operatorname{Spec} k,$$

the definition of locally finite type would require the source to be covered by affine opens whose coordinate rings are finitely generated k -algebras.

But the whole source is

$$\operatorname{Spec} k[x_1, x_2, x_3, \dots],$$

and the algebra contains infinitely many independent variables. This is the standard example of a morphism which is not locally of finite type.

Therefore

$$\operatorname{Spec} k[x_1, x_2, x_3, \dots] \longrightarrow \operatorname{Spec} k$$

is not locally of finite type.

A morphism which is locally of finite type but not of finite type

Let

$$X = \coprod_{n \geq 1} \mathbb{A}_k^1$$

be a countably infinite disjoint union of affine lines over k . Consider the natural morphism

$$f : X \longrightarrow \operatorname{Spec} k.$$

Each component is

$$\mathbb{A}_k^1 = \operatorname{Spec} k[t],$$

and $k[t]$ is a finitely generated k -algebra. Therefore each component is of finite type over $\operatorname{Spec} k$.

The inverse image of the only affine open of $\operatorname{Spec} k$ is

$$f^{-1}(\operatorname{Spec} k) = X.$$

This is covered by the affine opens

$$\mathbb{A}_k^1, \mathbb{A}_k^1, \mathbb{A}_k^1, \dots$$

one for each component. On each such open, the coordinate ring is $k[t]$, which is finitely generated over k .

Therefore f is locally of finite type.

However, X cannot be covered by finitely many of these connected components. Indeed, a finite union of components would miss all the remaining components. Hence X is not quasi-compact.

Since a morphism of finite type is, in particular, quasi-compact, this morphism cannot be of finite type.

Therefore

$$\coprod_{n \geq 1} \mathbb{A}_k^1 \longrightarrow \operatorname{Spec} k$$

is locally of finite type but not of finite type.

A scheme which is not a variety

In the language of the remark, a variety over an algebraically closed field k is an integral scheme of finite type over $\operatorname{Spec} k$.

Thus a scheme can fail to be a variety in several ways.

For example,

$$\operatorname{Spec} k[\varepsilon]/(\varepsilon^2)$$

is not a variety because it is not reduced.

Also,

$$\operatorname{Spec} k[x, y]/(xy)$$

is not a variety because it is reducible, hence not integral.

Finally,

$$\operatorname{Spec} k[x_1, x_2, x_3, \dots]$$

is not a variety because it is not of finite type over k .

Summary of examples

Property violated	Example
Reduced	$\operatorname{Spec} k[\varepsilon]/(\varepsilon^2)$
Irreducible	$\operatorname{Spec} k[x, y]/(xy)$
Integral	$\operatorname{Spec} k[x, y]/(xy)$
Locally of finite type	$\operatorname{Spec} k[x_1, x_2, x_3, \dots] \rightarrow \operatorname{Spec} k$
Finite type	$\coprod_{n \geq 1} \mathbb{A}_k^1 \rightarrow \operatorname{Spec} k$
Variety over k	$\operatorname{Spec} k[\varepsilon]/(\varepsilon^2)$

9 Problem 7: Generic point and function field of an integral scheme

Problem. Let X be an integral scheme. Show that there is a unique point η such that the closure of $\{\eta\}$ is X ; this is called the *generic point* of X . Show that the stalk of $\mathcal{O}_X$ at η is a field, called the *function field* of X , denoted by $K(X)$. Show that if

$$U = \operatorname{Spec} A$$

is any open affine subset of X , then $K(X)$ is the field of fractions of A .

Solution.

Since X is integral, it is nonempty, reduced, and irreducible. In particular, every nonempty open subset of X is dense in X , and every nonempty affine open subset has coordinate ring an integral domain.

Existence of the generic point

Choose a nonempty affine open subset

$$U = \operatorname{Spec} A \subseteq X.$$

Since X is integral, the open subscheme U is also integral. Hence its coordinate ring A is an integral domain.

Because A is a domain, the zero ideal

$$(0) \subseteq A$$

is a prime ideal. Therefore it determines a point

$$\eta = (0) \in \operatorname{Spec} A.$$

In the affine scheme $\operatorname{Spec} A$, the closure of the point corresponding to a prime ideal $\mathfrak{p}$ is

$$\overline{\{\mathfrak{p}\}} = V(\mathfrak{p}).$$

Thus

$$\overline{\{(0)\}} = V((0)) = \operatorname{Spec} A.$$

So (0) is the generic point of U .

We now show that this same point is generic for all of X .

Let $W \subseteq X$ be a nonempty open subset. Since X is irreducible, any two nonempty open subsets of X intersect. Therefore

$$W \cap U \neq \emptyset.$$

But $W \cap U$ is a nonempty open subset of $U = \operatorname{Spec} A$. Since (0) is the generic point of U , it lies in every nonempty open subset of U . Hence

$$\eta \in W \cap U \subseteq W.$$

Therefore η lies in every nonempty open subset of X .

Equivalently, the closure of $\{\eta\}$ is all of X :

$$\overline{\{\eta\}} = X.$$

Thus X has a generic point.

Uniqueness of the generic point

Suppose that η and η' are two points of X satisfying

$$\overline{\{\eta\}} = X \quad \text{and} \quad \overline{\{\eta'\}} = X.$$

Then both η and η' lie in every nonempty open subset of X .

Schemes are T_0 -spaces. This means that if two points are distinct, then there exists an open subset containing one of them but not the other.

Assume, for contradiction, that

$$\eta \neq \eta'.$$

Then there is an open subset $W \subseteq X$ containing one of the two points but not the other. Since W contains one of the points, it is nonempty. But every nonempty open subset of X must contain both η and η' , because both have dense closure equal to X . This is a contradiction.

Therefore

$$\eta = \eta'.$$

So the generic point of X is unique.

The stalk at the generic point is a field

Let

$$U = \operatorname{Spec} A$$

be an affine open neighborhood of η . Since X is integral, A is an integral domain.

Inside $U = \operatorname{Spec} A$, the point η corresponds to the prime ideal

$$(0) \subseteq A.$$

Therefore the stalk of the structure sheaf at η is

$$\mathcal{O}_{X,\eta} \cong \mathcal{O}_{U,\eta} \cong A_{(0)}.$$

But localizing an integral domain A at the zero prime ideal means inverting all nonzero elements of A . Hence

$$A_{(0)} = \operatorname{Frac}(A).$$

Therefore

$$\mathcal{O}_{X,\eta}$$

is a field.

This field is called the function field of X , and is denoted

$$K(X) := \mathcal{O}_{X,\eta}.$$

The function field on any affine open

Let

$$U = \operatorname{Spec} A$$

be any nonempty affine open subset of X .

Since η is the generic point of X , it lies in every nonempty open subset of X . In particular,

$$\eta \in U.$$

Since X is integral, the affine scheme U is integral, and therefore A is an integral domain.

In $\operatorname{Spec} A$, the generic point corresponds to the zero ideal

$$(0) \subseteq A.$$

Thus

$$K(X) = \mathcal{O}_{X,\eta} \cong \mathcal{O}_{U,\eta} \cong A_{(0)} = \operatorname{Frac}(A).$$

Therefore, for every nonempty affine open subset

$$U = \operatorname{Spec} A \subseteq X,$$

we have

$$K(X) \cong \operatorname{Frac}(A).$$

Conclusion

For an integral scheme X , there exists a unique point

$$\eta \in X$$

such that

$$\overline{\{\eta\}} = X.$$

This point is called the generic point of X .

The stalk at this point,

$$K(X) := \mathcal{O}_{X,\eta},$$

is a field, called the function field of X .

Moreover, for every nonempty affine open subset

$$U = \operatorname{Spec} A \subseteq X,$$

there is a canonical identification

$$K(X) \cong \operatorname{Frac}(A).$$

10 Theory for Problems 5–7

This section collects the main theory behind Problems 5–7. The common theme is that many important properties of schemes and morphisms are *local on affine opens*. In particular, finite generation of coordinate rings controls morphisms of finite type, while integrality controls the existence of a generic point and the construction of the function field.

10.1 Locally finite type and finite type morphisms

Let

$$f : X \rightarrow Y$$

be a morphism of schemes.

Definition. The morphism f is called *locally of finite type* if there exists an affine open covering

$$Y = \bigcup_i V_i, \quad V_i = \operatorname{Spec} B_i,$$

such that for each i , the inverse image $f^{-1}(V_i)$ can be covered by affine open subsets

$$U_{ij} = \operatorname{Spec} A_{ij},$$

where each A_{ij} is a finitely generated B_i -algebra.

Equivalently, locally on the target and locally on the source, the morphism looks like a ring map

$$B_i \rightarrow A_{ij}$$

where A_{ij} is generated over B_i by finitely many elements.

Definition. The morphism f is called *of finite type* if, in the definition above, the covering of each $f^{-1}(V_i)$ by affine open subsets

$$U_{ij} = \operatorname{Spec} A_{ij}$$

can be chosen to be finite.

Thus:

$$\text{finite type} = \text{locally finite type} + \text{finite covering condition.}$$

More conceptually, finite type morphisms are the geometric analogue of finitely generated algebras.

10.2 Affine algebraic meaning

The basic affine model is this.

Let

$$f : \operatorname{Spec} A \rightarrow \operatorname{Spec} B$$

be a morphism of affine schemes. Such a morphism corresponds contravariantly to a ring homomorphism

$$B \rightarrow A.$$

Then f is of finite type precisely when A is a finitely generated B -algebra.

This means that there exist elements

$$a_1, \dots, a_n \in A$$

such that

$$A = B[a_1, \dots, a_n].$$

Equivalently, there is a surjective B -algebra homomorphism

$$B[x_1, \dots, x_n] \twoheadrightarrow A.$$

So the affine schemes of finite type over $\operatorname{Spec} B$ are exactly those built from finitely many algebraic coordinates over B .

10.3 Why the condition is local

The property of being locally of finite type is local on the target.

This means that it is enough to check the property on one affine open covering of Y , and then it automatically holds on every affine open subset of Y .

The key reason is that affine opens can be refined by smaller affine opens. More precisely, if

$$V = \operatorname{Spec} B$$

is an affine open subset of Y , and $V_i = \operatorname{Spec} B_i$ is another affine open subset, then the intersection

$$V \cap V_i$$

can be covered by smaller affine open subsets. Around any point of $V \cap V_i$, one can choose an affine open subset contained in both V and V_i .

In practice, one refines the intersection by distinguished opens. For example, inside

$$V = \operatorname{Spec} B,$$

a distinguished open has the form

$$D(b) = \operatorname{Spec} B_b.$$

The coordinate ring B_b is a finitely generated B -algebra because

$$B_b \cong B[t]/(bt - 1).$$

This simple fact is crucial: localizing at one element does not destroy finite generation over the original base ring.

10.4 Localization preserves finite generation

A basic algebraic fact used repeatedly is the following.

Lemma. Let A be a finitely generated B -algebra, and let $S \subseteq B$ be a multiplicative subset. Then

$$S^{-1}A$$

is a finitely generated $S^{-1}B$ -algebra.

Reason. If

$$A = B[a_1, \dots, a_n],$$

then after localization,

$$S^{-1}A = (S^{-1}B)[a_1/1, \dots, a_n/1].$$

Thus the same generators still generate after localization.

In particular, if $b \in B$, then

$$A_b$$

is a finitely generated B_b -algebra.

Also, since

$$B_b \cong B[t]/(bt - 1),$$

the ring B_b is a finitely generated B -algebra. Hence a finitely generated B_b -algebra is also finitely generated over B .

This explains the mechanism behind Problem 5.

10.5 Geometric meaning of finite type

A morphism

$$f : X \rightarrow Y$$

of finite type should be thought of as a family of geometric objects over Y which can be described using finitely many algebraic coordinates and finitely many affine charts.

For example, over a field k , a scheme of finite type over $\operatorname{Spec} k$ is locally described by rings of the form

$$k[x_1, \dots, x_n]/I.$$

Thus finite type schemes over k are the schemes that behave like algebraic sets cut out in finite-dimensional affine space.

By contrast,

$$\operatorname{Spec} k[x_1, x_2, x_3, \dots]$$

is not of finite type over k , because its coordinate ring needs infinitely many independent variables.

10.6 Locally finite type versus finite type

The difference between *locally of finite type* and *of finite type* is a finiteness condition on the number of affine pieces needed.

A morphism may be locally finite type because every small affine piece is finitely generated over the base, but it may fail to be finite type because infinitely many such pieces are needed.

The standard example is

$$X = \coprod_{n \geq 1} \mathbb{A}_k^1$$

with the natural morphism

$$X \rightarrow \operatorname{Spec} k.$$

Each component

$$\mathbb{A}_k^1 = \operatorname{Spec} k[t]$$

is of finite type over k , so the morphism is locally of finite type.

However, the source is an infinite disjoint union, so it is not quasi-compact. Therefore it cannot be of finite type.

Thus:

$$\coprod_{n \geq 1} \mathbb{A}_k^1 \rightarrow \operatorname{Spec} k$$

is locally of finite type but not of finite type.

10.7 Reduced, irreducible, and integral schemes

The examples in Problem 6 depend on the basic properties of schemes.

Reduced schemes. A scheme X is reduced if, for every open subset $U \subseteq X$, the ring

$$\mathcal{O}_X(U)$$

has no nonzero nilpotent elements.

Affine-locally, this means that if

$$X = \operatorname{Spec} A,$$

then X is reduced precisely when A is a reduced ring.

For example,

$$\operatorname{Spec} k[\varepsilon]/(\varepsilon^2)$$

is not reduced because

$$\varepsilon \neq 0, \quad \varepsilon^2 = 0.$$

Irreducible schemes. A topological space X is irreducible if it cannot be written as the union of two proper closed subsets.

Equivalently, every two nonempty open subsets of X intersect.

For affine schemes,

$$\operatorname{Spec} A$$

is irreducible if and only if the nilradical of A is a prime ideal. In particular, if A is a domain, then $\operatorname{Spec} A$ is irreducible.

For example,

$$\operatorname{Spec} k[x, y]/(xy)$$

is reducible because it is the union of the two coordinate axes:

$$V(xy) = V(x) \cup V(y).$$

Integral schemes. A scheme X is integral if it is both reduced and irreducible.

Affine-locally, an affine scheme

$$\operatorname{Spec} A$$

is integral if and only if A is an integral domain.

So:

$$\operatorname{Spec} A \text{ integral} \iff A \text{ is a domain.}$$

10.8 Varieties in this language

In the language of these exercises, a variety over an algebraically closed field k is an integral scheme of finite type over

$$\operatorname{Spec} k.$$

Thus a k -variety must satisfy two kinds of conditions:

Condition	Meaning
Integral	Geometrically one reduced irreducible piece
Finite type over k	Described by finitely many algebraic coordinates over k

Therefore the following are not varieties:

$$\operatorname{Spec} k[\varepsilon]/(\varepsilon^2),$$

because it is not reduced;

$$\operatorname{Spec} k[x, y]/(xy),$$

because it is reducible;

and

$$\operatorname{Spec} k[x_1, x_2, x_3, \dots],$$

because it is not of finite type over k .

10.9 Generic points

Let X be an integral scheme.

Because X is irreducible, it has a unique dense irreducible component: namely X itself.

A point

$$\eta \in X$$

is called a *generic point* of X if

$$\overline{\{\eta\}} = X.$$

For an affine integral scheme

$$X = \operatorname{Spec} A,$$

where A is a domain, the generic point is the point corresponding to the zero ideal:

$$\eta = (0).$$

Indeed, since A is a domain, (0) is a prime ideal. Its closure is

$$\overline{\{(0)\}} = V((0)) = \operatorname{Spec} A.$$

Thus (0) is dense in $\operatorname{Spec} A$.

10.10 Generic point of an integral scheme

For a general integral scheme X , choose a nonempty affine open subset

$$U = \operatorname{Spec} A \subseteq X.$$

Since X is integral, the ring A is a domain. Therefore U has generic point

$$(0) \in \operatorname{Spec} A.$$

This point is also the generic point of all of X .

The reason is that X is irreducible. Hence every nonempty open subset of X intersects U . Since (0) lies in every nonempty open subset of U , it lies in every nonempty open subset of X .

Therefore

$$\overline{\{\eta\}} = X.$$

The generic point is unique because schemes are T_0 -spaces: two distinct points can be topologically distinguished by an open set. If two points both had closure equal to X , then both would lie in every nonempty open subset, which would contradict the T_0 property unless the two points were equal.

10.11 The stalk at the generic point

Let X be an integral scheme with generic point η .

The stalk

$$\mathcal{O}_{X,\eta}$$

is the local ring of X at the generic point.

If

$$U = \operatorname{Spec} A$$

is an affine open neighborhood of η , then η corresponds to the prime ideal

$$(0) \subseteq A.$$

Therefore

$$\mathcal{O}_{X,\eta} \cong A_{(0)}.$$

Since A is a domain, localizing at (0) means inverting every nonzero element of A . Hence

$$A_{(0)} = \operatorname{Frac}(A).$$

Therefore

$$\mathcal{O}_{X,\eta}$$

is a field.

This field is called the *function field* of X , and is denoted

$$K(X).$$

Thus

$$K(X) := \mathcal{O}_{X,\eta}.$$

10.12 Function field on affine opens

If

$$U = \operatorname{Spec} A$$

is any nonempty affine open subset of an integral scheme X , then $\eta \in U$, because the generic point lies in every nonempty open subset.

Inside $U = \operatorname{Spec} A$, the point η corresponds to the prime ideal

$$(0) \subseteq A.$$

Hence

$$K(X) = \mathcal{O}_{X,\eta} \cong A_{(0)} = \operatorname{Frac}(A).$$

Therefore, for every nonempty affine open subset

$$U = \operatorname{Spec} A \subseteq X,$$

we have

$$K(X) \cong \operatorname{Frac}(A).$$

This is why the function field of an integral scheme may be computed from any affine chart.

10.13 Geometric meaning of the function field

If X is an integral variety over a field k , then $K(X)$ is the field of rational functions on X .

For example, if

$$X = \mathbb{A}_k^n = \operatorname{Spec} k[x_1, \dots, x_n],$$

then

$$K(X) = \operatorname{Frac}(k[x_1, \dots, x_n]) = k(x_1, \dots, x_n).$$

If

$$X = \operatorname{Spec} k[x, y]/(y^2 - x^3 - x)$$

is an integral affine curve, then its function field is

$$K(X) = \operatorname{Frac}\left(k[x, y]/(y^2 - x^3 - x)\right).$$

Elements of $K(X)$ are not necessarily regular everywhere on X . They are rational functions, meaning that they are regular on some nonempty open subset of X .

10.14 Conceptual summary

The three problems are connected as follows:

Topic	Algebraic condition	Geometric meaning
Locally finite type	locally finitely generated algebras	finite-dimensional algebraic behavior
Finite type	finite affine cover by finitely generated algebras	globally finite algebraic behavior
Reduced	no nilpotents	no infinitesimal thickening
Irreducible	one topological component	one algebraic piece
Integral	domain on affine charts	reduced and irreducible
Generic point	$(0) \in \operatorname{Spec} A$ for a domain A	point whose closure is all of X
Function field	$\operatorname{Frac}(A)$	rational functions on X

10.15 Main takeaways

Locally finite type is checked on affine opens and survives refinement.

Finite type adds a global finiteness condition to local finite generation.

Integral schemes have a unique generic point.

$K(X) = \mathcal{O}_{X,\eta} \cong \operatorname{Frac}(A)$ for every nonempty affine open $U = \operatorname{Spec} A \subseteq X$.

11 Examples for Problems 5–7

This section gives concrete examples illustrating the notions from Problems 5–7: locally finite type morphisms, finite type morphisms, reduced schemes, irreducible schemes, integral schemes, varieties, generic points, and function fields.

11.1 Example 1: An affine morphism of finite type

Let k be a field, and consider

$$X = \operatorname{Spec} k[x, y]/(y^2 - x^3 - x)$$

with the natural morphism

$$f : X \rightarrow \operatorname{Spec} k.$$

The coordinate ring of X is

$$A = k[x, y]/(y^2 - x^3 - x).$$

This is a finitely generated k -algebra, because it is generated by the images of x and y . Indeed,

$$A = k[\bar{x}, \bar{y}].$$

Equivalently, there is a surjective k -algebra homomorphism

$$k[x, y] \twoheadrightarrow k[x, y]/(y^2 - x^3 - x).$$

Therefore the morphism

$$\mathrm{Spec} k[x, y]/(y^2 - x^3 - x) \rightarrow \mathrm{Spec} k$$

is of finite type.

Geometrically, this is an affine algebraic curve over k .

11.2 Example 2: A morphism locally of finite type

Let

$$X = \mathbb{A}_k^2 = \mathrm{Spec} k[x, y],$$

and consider the natural morphism

$$f : \mathbb{A}_k^2 \rightarrow \mathrm{Spec} k.$$

Since

$$k[x, y]$$

is a finitely generated k -algebra, the morphism is of finite type. Hence it is also locally of finite type.

Now take a distinguished open subset

$$D(x) \subseteq \mathbb{A}_k^2.$$

Its coordinate ring is

$$k[x, y]_x.$$

This is still finitely generated over k , because

$$k[x, y]_x \cong k[x, y, t]/(xt - 1).$$

Thus localization preserves the finite type nature of the morphism.

This example illustrates the idea from Problem 5: when we restrict to smaller affine opens, we still get finitely generated algebras.

11.3 Example 3: Checking locally finite type on another affine open

Let

$$Y = \mathrm{Spec} k[s],$$

and let

$$X = \mathrm{Spec} k[s, t].$$

Consider the morphism

$$f : X \rightarrow Y$$

induced by the inclusion

$$k[s] \hookrightarrow k[s, t].$$

This morphism is of finite type because

$$k[s, t] = k[s][t],$$

so $k[s, t]$ is generated over $k[s]$ by one element, namely t .

Now take the distinguished open subset

$$V = D(s) \subseteq Y.$$

Then

$$V = \operatorname{Spec} k[s]_s.$$

The inverse image of V is

$$f^{-1}(D(s)) = D(s) \subseteq X.$$

Its coordinate ring is

$$k[s, t]_s.$$

But

$$k[s, t]_s \cong k[s]_s[t].$$

So $k[s, t]_s$ is a finitely generated $k[s]_s$ -algebra.

Therefore the morphism remains locally of finite type after passing to the affine open subset $D(s)$.

This is exactly the behavior described in Problem 5.

11.4 Example 4: A morphism not locally of finite type

Consider

$$X = \operatorname{Spec} k[x_1, x_2, x_3, \dots]$$

and the natural morphism

$$f : X \rightarrow \operatorname{Spec} k.$$

The coordinate ring

$$k[x_1, x_2, x_3, \dots]$$

is the polynomial ring in infinitely many variables.

This ring is not finitely generated as a k -algebra. Indeed, if it were generated by finitely many elements, then only finitely many variables could appear in those generators. But then variables outside that finite list could not be produced.

Therefore

$$k[x_1, x_2, x_3, \dots]$$

is not a finitely generated k -algebra.

Hence the morphism

$$\mathrm{Spec} k[x_1, x_2, x_3, \dots] \rightarrow \mathrm{Spec} k$$

is not locally of finite type.

This example shows that infinite-dimensional affine space is not of finite type over k .

11.5 Example 5: Locally of finite type but not finite type

Let

$$X = \coprod_{n \geq 1} \mathbb{A}_k^1$$

be the infinite disjoint union of copies of the affine line over k . Consider the natural morphism

$$f : X \rightarrow \mathrm{Spec} k.$$

Each component is

$$\mathbb{A}_k^1 = \mathrm{Spec} k[t],$$

and $k[t]$ is a finitely generated k -algebra.

Therefore every point of X has an affine open neighborhood mapping to $\mathrm{Spec} k$ by a finite type morphism. Hence

$$f : X \rightarrow \mathrm{Spec} k$$

is locally of finite type.

However, X is an infinite disjoint union of nonempty open subsets. It is not quasi-compact, because no finite collection of components covers all of X .

A finite type morphism is quasi-compact. Therefore

$$f : X \rightarrow \mathrm{Spec} k$$

is not of finite type.

Thus this is the standard example of a morphism which is locally of finite type but not finite type.

11.6 Example 6: A non-reduced scheme

Let

$$X = \mathrm{Spec} k[\varepsilon]/(\varepsilon^2).$$

In the ring

$$k[\varepsilon]/(\varepsilon^2),$$

the element ε satisfies

$$\varepsilon^2 = 0,$$

but

$$\varepsilon \neq 0.$$

Thus ε is a nonzero nilpotent element.

Therefore

$$k[\varepsilon]/(\varepsilon^2)$$

is not reduced, and so

$$X = \operatorname{Spec} k[\varepsilon]/(\varepsilon^2)$$

is not a reduced scheme.

Geometrically, this scheme is an infinitesimal thickening of a point. It has the same underlying topological space as $\operatorname{Spec} k$, but it has extra nilpotent structure.

11.7 Example 7: A reducible scheme

Let

$$X = \operatorname{Spec} k[x, y]/(xy).$$

The equation

$$xy = 0$$

means that either

$$x = 0$$

or

$$y = 0.$$

Thus X is the union of the two coordinate axes in the affine plane:

$$X = V(x) \cup V(y).$$

Both $V(x)$ and $V(y)$ are proper closed subsets of X . Hence X is reducible.

Therefore

$$\operatorname{Spec} k[x, y]/(xy)$$

is not irreducible.

This example is reduced if k is a field, because the ideal (xy) is radical in $k[x, y]$. However, it is not integral because it is reducible.

11.8 Example 8: A scheme that is irreducible but not reduced

Let

$$X = \operatorname{Spec} k[x]/(x^2).$$

The ring

$$k[x]/(x^2)$$

has a nilpotent element, namely the image of x , because

$$x^2 = 0.$$

So X is not reduced.

However, the underlying topological space has only one prime ideal, namely

$$(x)/(x^2).$$

Therefore X is irreducible as a topological space.

Thus

$$\operatorname{Spec} k[x]/(x^2)$$

is irreducible but not reduced.

Consequently, it is not integral.

11.9 Example 9: A reduced scheme that is not integral

Let

$$X = \operatorname{Spec} k[x, y]/(xy).$$

As above, this scheme is the union of two coordinate axes:

$$V(xy) = V(x) \cup V(y).$$

The ring

$$k[x, y]/(xy)$$

is reduced, because (xy) is a radical ideal in $k[x, y]$. But it is not a domain, since

$$\bar{x} \cdot \bar{y} = 0$$

while

$$\bar{x} \neq 0, \quad \bar{y} \neq 0.$$

Therefore the scheme is reduced but not integral.

This example shows that reducedness alone is not enough for integrality. We also need irreducibility.

11.10 Example 10: A variety over an algebraically closed field

Let k be algebraically closed, and consider

$$X = \operatorname{Spec} k[x, y]/(y - x^2).$$

The ring

$$k[x, y]/(y - x^2)$$

is isomorphic to

$$k[x],$$

because the relation $y = x^2$ allows us to eliminate y .

Indeed,

$$k[x, y]/(y - x^2) \cong k[x].$$

Since $k[x]$ is a domain, X is integral.

Since $k[x]$ is finitely generated over k , the morphism

$$X \rightarrow \operatorname{Spec} k$$

is of finite type.

Therefore X is a variety over k .

Geometrically, X is the parabola

$$y = x^2$$

in the affine plane.

11.11 Example 11: A scheme that is not a variety because it is not reduced

Let k be algebraically closed and let

$$X = \operatorname{Spec} k[\varepsilon]/(\varepsilon^2).$$

Although this scheme is finite type over k , it is not reduced. Since a variety in this setting means an integral scheme of finite type over k , X is not a variety.

Thus

$$\operatorname{Spec} k[\varepsilon]/(\varepsilon^2)$$

is a finite type k -scheme, but not a variety.

11.12 Example 12: A scheme that is not a variety because it is not irreducible

Let

$$X = \operatorname{Spec} k[x, y]/(xy).$$

This scheme is finite type over k , and it is reduced. However, it is reducible, because it is the union of the two coordinate axes.

Therefore X is not integral.

Hence

$$\operatorname{Spec} k[x, y]/(xy)$$

is not a variety in the sense of these exercises.

11.13 Example 13: A scheme that is not a variety because it is not finite type

Let

$$X = \operatorname{Spec} k[x_1, x_2, x_3, \dots].$$

The coordinate ring

$$k[x_1, x_2, x_3, \dots]$$

is a domain, so X is integral.

However, it is not finitely generated as a k -algebra.

Therefore the morphism

$$X \rightarrow \operatorname{Spec} k$$

is not of finite type.

Hence X is not a variety over k , even though it is integral.

11.14 Example 14: Generic point of affine space

Let

$$X = \mathbb{A}_k^n = \operatorname{Spec} k[x_1, \dots, x_n].$$

The ring

$$k[x_1, \dots, x_n]$$

is a domain. Therefore X is integral.

The generic point of X is the point corresponding to the zero ideal:

$$\eta = (0).$$

The closure of $\{\eta\}$ is

$$\overline{\{\eta\}} = V((0)) = \operatorname{Spec} k[x_1, \dots, x_n] = X.$$

Therefore (0) is the generic point of affine n -space.

The function field is

$$K(X) = \operatorname{Frac}(k[x_1, \dots, x_n]).$$

Hence

$$K(\mathbb{A}_k^n) = k(x_1, \dots, x_n).$$

11.15 Example 15: Generic point and function field of the affine line

Let

$$X = \mathbb{A}_k^1 = \operatorname{Spec} k[t].$$

Since $k[t]$ is a domain, X is integral.

The generic point is

$$\eta = (0) \in \operatorname{Spec} k[t].$$

The function field is

$$K(X) = \mathcal{O}_{X,\eta}.$$

Since η corresponds to the prime ideal (0) , we have

$$\mathcal{O}_{X,\eta} \cong k[t]_{(0)}.$$

Because $k[t]$ is a domain,

$$k[t]_{(0)} = \operatorname{Frac}(k[t]) = k(t).$$

Thus

$$K(\mathbb{A}_k^1) = k(t).$$

11.16 Example 16: Generic point and function field of an affine curve

Let

$$X = \operatorname{Spec} k[x, y]/(y^2 - x^3 - x).$$

Assume that

$$y^2 - x^3 - x$$

is irreducible in $k[x, y]$. Then

$$A = k[x, y]/(y^2 - x^3 - x)$$

is a domain.

Therefore $X = \operatorname{Spec} A$ is integral.

Its generic point is the point corresponding to the zero ideal:

$$\eta = (0) \in \operatorname{Spec} A.$$

The function field is

$$K(X) = \operatorname{Frac}(A).$$

Thus

$$K(X) = \operatorname{Frac}\left(k[x, y]/(y^2 - x^3 - x)\right).$$

This is the field of rational functions on the curve

$$y^2 = x^3 + x.$$

11.17 Example 17: Function field does not depend on the affine open

Let

$$X = \mathbb{A}_k^1 = \operatorname{Spec} k[t].$$

The function field of X is

$$K(X) = k(t).$$

Now consider the distinguished open subset

$$U = D(t) \subseteq X.$$

Then

$$U = \operatorname{Spec} k[t]_t.$$

The coordinate ring of U is

$$k[t]_t = k[t, t^{-1}].$$

The field of fractions of this ring is

$$\operatorname{Frac}(k[t, t^{-1}]) = k(t).$$

Therefore

$$K(U) = \operatorname{Frac}(k[t, t^{-1}]) = k(t) = K(X).$$

This illustrates the statement from Problem 7: for any nonempty affine open subset $U = \operatorname{Spec} A$ of an integral scheme X , we have

$$K(X) \cong \operatorname{Frac}(A).$$

11.18 Example 18: Function field of projective line

Let

$$X = \mathbb{P}_k^1.$$

The projective line is integral. It has standard affine open cover

$$U_0 = \operatorname{Spec} k[t], \quad U_1 = \operatorname{Spec} k[s],$$

where on the overlap one has

$$s = t^{-1}.$$

Using the affine open

$$U_0 = \operatorname{Spec} k[t],$$

we get

$$K(\mathbb{P}_k^1) = \operatorname{Frac}(k[t]) = k(t).$$

Using the affine open

$$U_1 = \operatorname{Spec} k[s],$$

we also get

$$K(\mathbb{P}_k^1) = \text{Frac}(k[s]) = k(s).$$

Since $s = t^{-1}$, the fields

$$k(s) \quad \text{and} \quad k(t)$$

are naturally the same field.

Thus

$$K(\mathbb{P}_k^1) = k(t).$$

This example shows that even for a non-affine integral scheme, the function field can be computed from any affine open chart.

11.19 Example 19: Why integrality is needed for the function field

Let

$$X = \text{Spec } k[x, y]/(xy).$$

This scheme is not integral, because

$$\bar{x} \cdot \bar{y} = 0$$

while neither $\bar{x}$ nor $\bar{y}$ is zero.

The scheme has two irreducible components:

$$V(x) \quad \text{and} \quad V(y).$$

Each component has its own generic point. Therefore X does not have a unique generic point whose closure is all of X .

Consequently, there is no single function field $K(X)$ in the sense of Problem 7.

Instead, each irreducible component has its own function field. Both components are isomorphic to affine lines, so each component has function field

$$k(t).$$

This example shows why Problem 7 assumes that X is integral.

11.20 Example 20: Summary table

Example	Property illustrated	Reason
$\text{Spec } k[x, y]/(y^2 - x^3 - x)$	finite type over k	coordinate ring finitely generated over k
$\text{Spec } k[x_1, x_2, \dots]$	not locally finite type over k	infinitely many algebra generators needed
$\coprod_{n \geq 1} \mathbb{A}_k^1$	locally finite type, not finite type	infinitely many affine components
$\text{Spec } k[\varepsilon]/(\varepsilon^2)$	not reduced	$\varepsilon^2 = 0, \varepsilon \neq 0$
$\text{Spec } k[x, y]/(xy)$	reducible	$V(xy) = V(x) \cup V(y)$
$\text{Spec } k[x, y]/(xy)$	not integral	$\overline{xy} = 0$
$\mathbb{A}_k^n$	generic point (0)	$k[x_1, \dots, x_n]$ is a domain
$\mathbb{A}_k^1$	$K(X) = k(t)$	$\text{Frac}(k[t]) = k(t)$
$\mathbb{P}_k^1$	$K(X) = k(t)$	computed from affine chart $\text{Spec } k[t]$

12 Charts, Diagrams, and Tables for Problems 5–7

This section summarizes the main ideas of Problems 5–7 using diagrams, comparison tables, and conceptual charts. The goal is to make the relationships between local finite type morphisms, finite type morphisms, reduced schemes, irreducible schemes, integral schemes, generic points, and function fields visually clear.

12.1 Local finite type refinement diagram

Problem 5 says that if a morphism is locally of finite type with respect to one affine open cover of the target, then it is locally of finite type with respect to every affine open subset of the target.

Suppose

$$f : X \rightarrow Y$$

is locally of finite type. We start with an affine open cover

$$Y = \bigcup_i V_i, \quad V_i = \text{Spec } B_i,$$

and affine opens above each V_i ,

$$U_{ij} = \text{Spec } A_{ij} \subseteq f^{-1}(V_i),$$

where each A_{ij} is a finitely generated B_i -algebra.

Now let

$$V = \text{Spec } B \subseteq Y$$

be another affine open subset. We refine the intersections $V \cap V_i$ by smaller affine opens

$$W_{i\alpha} \subseteq V \cap V_i.$$

The geometric picture is:

$$\begin{array}{ccccc}
U_{ij} \cap f^{-1}(W_{i\alpha}) & \subseteq & U_{ij} & \subseteq & f^{-1}(V_i) \\
\downarrow & & \downarrow & & \downarrow \\
W_{i\alpha} & \subseteq & V_i & \subseteq & Y
\end{array}$$

Here $W_{i\alpha}$ is chosen small enough so that it is affine both as an open subset of V and as an open subset of V_i . For example, locally one may take distinguished opens:

$$W_{i\alpha} = D(b_{i\alpha}) \subseteq V = \operatorname{Spec} B,$$

and also

$$W_{i\alpha} = D(c_{i\alpha}) \subseteq V_i = \operatorname{Spec} B_i.$$

12.2 Coordinate ring diagram for refinement

The corresponding algebraic diagram is:

$$\begin{array}{ccc}
B_i & \longrightarrow & A_{ij} \\
\downarrow & & \downarrow \\
(B_i)_{c_{i\alpha}} & \longrightarrow & (A_{ij})_{\varphi(c_{i\alpha})}
\end{array}$$

The original finite generation condition is

$$A_{ij} \text{ is finitely generated over } B_i.$$

After localization, we get

$$(A_{ij})_{\varphi(c_{i\alpha})} \text{ is finitely generated over } (B_i)_{c_{i\alpha}}.$$

Since

$$(B_i)_{c_{i\alpha}} \cong B_{b_{i\alpha}},$$

the localized coordinate ring is finitely generated over the coordinate ring of the smaller affine open in V .

Finally,

$$B_{b_{i\alpha}} \cong B[t]/(b_{i\alpha}t - 1),$$

so $B_{b_{i\alpha}}$ is a finitely generated B -algebra.

Therefore the affine opens above V still have coordinate rings finitely generated over B .

12.3 Algebraic mechanism behind Problem 5

Geometric step	Algebraic step	Reason
$V_i = \operatorname{Spec} B_i$	B_i	affine chart on Y
$U_{ij} = \operatorname{Spec} A_{ij}$	$B_i \rightarrow A_{ij}$	affine chart above V_i
A_{ij} finite type over B_i	$A_{ij} = B_i[a_1, \dots, a_n]$	definition of locally finite type
$W = D(c) \subseteq V_i$	$(B_i)_c$	distinguished affine open
$U_{ij} \cap f^{-1}(W)$	$(A_{ij})_{\varphi(c)}$	preimage of a distinguished open
$(A_{ij})_{\varphi(c)}$	$(B_i)_c[a_1/1, \dots, a_n/1]$	localization preserves generators
$D(b) \subseteq \operatorname{Spec} B$	$B_b \cong B[t]/(bt - 1)$	localization is finitely generated

The important algebraic principle is:

$$A = B[a_1, \dots, a_n] \implies A_b = B_b[a_1/1, \dots, a_n/1].$$

Thus localization preserves finite generation.

12.4 Finite type versus locally finite type

The distinction between locally finite type and finite type is a global finiteness condition.

Property	Local algebra condition	Global finiteness condition
Locally of finite type	A_{ij} finitely generated over B_i	possibly infinitely many affine pieces
Of finite type	A_{ij} finitely generated over B_i	finitely many affine pieces over each V_i

In words:


$\text{finite type} = \text{locally finite type} + \text{quasi-compactness}$

A standard example is

$$\coprod_{n \geq 1} \mathbb{A}_k^1 \longrightarrow \operatorname{Spec} k.$$

Each component is finite type over k , but infinitely many components are needed to cover the source.

12.5 Flow chart for checking finite type



Thus local finite generation gives local finite type, while adding a finite covering condition gives finite type.

12.6 Examples violating properties

Property	Example violating it	Reason
Reduced	$\operatorname{Spec} k[\varepsilon]/(\varepsilon^2)$	$\varepsilon^2 = 0, \varepsilon \neq 0$
Irreducible	$\operatorname{Spec} k[x, y]/(xy)$	$V(xy) = V(x) \cup V(y)$
Integral	$\operatorname{Spec} k[x, y]/(xy)$	$\bar{x}\bar{y} = 0, \bar{x}, \bar{y} \neq 0$
Locally of finite type	$\operatorname{Spec} k[x_1, x_2, \dots] \rightarrow \operatorname{Spec} k$	infinitely many algebra generators
Finite type	$\coprod_{n \geq 1} \mathbb{A}_k^1 \rightarrow \operatorname{Spec} k$	not quasi-compact
Variety over k	$\operatorname{Spec} k[\varepsilon]/(\varepsilon^2)$	not integral

12.7 Reduced, irreducible, and integral schemes

Scheme property	Affine algebra condition	Geometric meaning
Reduced	A has no nonzero nilpotents	no infinitesimal thickening
Irreducible	$\sqrt{(0)}$ is prime	one topological component
Integral	A is a domain	one reduced irreducible piece

For affine schemes, the most important equivalence is:

$$\boxed{\operatorname{Spec} A \text{ is integral} \iff A \text{ is a domain.}}$$

12.8 Diagram of a non-reduced point

The scheme

$$\mathrm{Spec} k[\varepsilon]/(\varepsilon^2)$$

has one underlying topological point, but with nilpotent thickening.

ordinary reduced point



non-reduced thickened point



The nilpotent element is

$$\varepsilon^2 = 0, \quad \varepsilon \neq 0.$$

Thus the scheme has infinitesimal structure invisible in the underlying topological space.

12.9 Diagram of the reducible scheme $xy = 0$


For

$$X = \mathrm{Spec} k[x, y]/(xy),$$

the equation is

$$xy = 0.$$

Thus either $x = 0$ or $y = 0$. Geometrically, this is the union of the two coordinate axes.



Algebraically,

$$V(xy) = V(x) \cup V(y).$$

Also,

$$\bar{x}\bar{y} = 0, \quad \bar{x} \neq 0, \quad \bar{y} \neq 0.$$

Therefore the coordinate ring is not a domain, and the scheme is not integral.

12.10 Implication chart for affine schemes

For an affine scheme $X = \operatorname{Spec} A$, the implications are:

$$\begin{array}{ccccc} A \text{ is a domain} & \implies & A \text{ is reduced} & \implies & \operatorname{Spec} A \text{ is reduced} \\ & & \downarrow & & \\ & & \operatorname{Spec} A \text{ is irreducible} & & \end{array}$$

Hence

$$A \text{ is a domain} \implies \operatorname{Spec} A \text{ is integral.}$$

But the separate converses fail:

$$\operatorname{Spec} k[x]/(x^2)$$

is irreducible but not reduced, while

$$\operatorname{Spec} k[x, y]/(xy)$$

is reduced but not irreducible.

12.11 Generic point diagram

Let

$$X = \operatorname{Spec} A$$

where A is a domain. Then the generic point is the point corresponding to the zero ideal:

$$\eta = (0).$$

The closure of this point is the whole space:

$$\overline{\{\eta\}} = V((0)) = \operatorname{Spec} A.$$

Diagrammatically:

$$\begin{array}{c} \eta = (0) \\ \downarrow \text{ closure} \\ X = \operatorname{Spec} A \end{array}$$

The generic point is not usually a closed point. It is a point whose closure is the entire irreducible scheme.

12.12 Generic point versus closed points

For

$$X = \mathbb{A}_k^1 = \operatorname{Spec} k[t],$$

we have two important types of points:

Type of point	Prime ideal	Closure
generic point	(0)	$\mathbb{A}_k^1$
closed point	$(t - a)$	$\{a\}$

Thus the point (0) is dense, while the point $(t - a)$ is closed.

The generic point represents the whole irreducible affine line.

12.13 Function field diagram

Let X be an integral scheme with generic point η . Choose a nonempty affine open subset

$$U = \operatorname{Spec} A \subseteq X.$$

Then $\eta \in U$, and inside U it corresponds to the zero ideal:

$$\eta = (0) \in \operatorname{Spec} A.$$

Therefore

$$\mathcal{O}_{X,\eta} \cong A_{(0)} = \operatorname{Frac}(A).$$

The diagram is:

$$\begin{array}{ccc} U = \operatorname{Spec} A & \subseteq & X \\ \eta = (0) & \in & U \\ \downarrow & & \\ \mathcal{O}_{X,\eta} & \cong & A_{(0)} = \operatorname{Frac}(A) \end{array}$$

Thus

$$\boxed{K(X) = \mathcal{O}_{X,\eta} \cong \operatorname{Frac}(A).}$$

12.14 Function field examples

Scheme	Affine coordinate ring	Function field
$\mathbb{A}_k^1$	$k[t]$	$k(t)$
$\mathbb{A}_k^n$	$k[x_1, \dots, x_n]$	$k(x_1, \dots, x_n)$
$\operatorname{Spec} k[x, y]/(y - x^2)$	$k[x, y]/(y - x^2) \cong k[x]$	$k(x)$
$\operatorname{Spec} k[x, y]/(y^2 - x^3 - x)$	$k[x, y]/(y^2 - x^3 - x)$	$\operatorname{Frac}(k[x, y]/(y^2 - x^3 - x))$
$\mathbb{P}_k^1$	chart $\operatorname{Spec} k[t]$	$k(t)$

12.15 Function field from different affine charts

For the projective line

$$X = \mathbb{P}_k^1,$$

we have two standard affine charts:

$$U_0 = \operatorname{Spec} k[t], \quad U_1 = \operatorname{Spec} k[s].$$

On the overlap,

$$s = t^{-1}.$$

The function field can be computed from either chart:

$$U_0 = \operatorname{Spec} k[t] \longrightarrow \operatorname{Frac}(k[t]) = k(t)$$

$$U_1 = \operatorname{Spec} k[s] \longrightarrow \operatorname{Frac}(k[s]) = k(s)$$

Since

$$s = t^{-1},$$

the fields are naturally identified:

$$k(s) = k(t).$$

Therefore

$$K(\mathbb{P}_k^1) = k(t).$$

This shows that the function field of an integral scheme does not depend on the affine chart used to compute it.

12.16 Concept map for finite type morphisms

finitely generated algebra $\iff$ finite type affine morphism $\implies$ locally finite type morphism

$$\begin{array}{c} \downarrow \\ \text{finite affine cover} \\ \downarrow \\ \text{finite type morphism} \end{array}$$

In affine form:

$$\text{Spec } A \rightarrow \text{Spec } B$$

is of finite type if and only if

$$A = B[a_1, \dots, a_n].$$

12.17 Concept map for integral schemes and function fields

A is a domain $\iff$ $\text{Spec } A$ is integral $\implies$ unique generic point

$$\begin{array}{c} \downarrow \\ K(X) = \mathcal{O}_{X,\eta} \\ \downarrow \\ K(X) = \text{Frac}(A) \end{array}$$

For every nonempty affine open

$$U = \text{Spec } A \subseteq X,$$

we have

$$K(X) \cong \text{Frac}(A).$$

12.18 Final summary table

Topic	Object	Algebra	Conclusion
Finite type	$\text{Spec } A \rightarrow \text{Spec } B$	$A = B[a_1, \dots, a_n]$	finite type
Not locally finite type	$\text{Spec } k[x_1, x_2, \dots] \rightarrow \text{Spec } k$	infinitely many generators	not locally finite type
Locally finite, not finite	$\coprod_{n \geq 1} \mathbb{A}_k^1 \rightarrow \text{Spec } k$	each $k[t]$ is finite type	not quasi-compact
Non-reduced	$\text{Spec } k[\varepsilon]/(\varepsilon^2)$	$\varepsilon^2 = 0$	nilpotent thickening
Reducible	$\text{Spec } k[x, y]/(xy)$	$xy = 0$	two components
Integral	$\text{Spec } A$	A is a domain	generic point (0)
Function field	$U = \text{Spec } A \subseteq X$	$A_{(0)} = \text{Frac}(A)$	$K(X) = \text{Frac}(A)$

13 Problem 8. Normalization

Problem. A scheme is normal if all its local rings are integrally closed domains. Let X be an integral scheme. For each open affine subset $U = \operatorname{Spec} A$ of X , let $\overline{A}$ be the integral closure of A in its quotient field, and let

$$\tilde{U} = \operatorname{Spec} \overline{A}.$$

Show that one can glue the schemes $\tilde{U}$ to obtain a normal integral scheme $\widetilde{X}$, called the normalization of X . Show that there is a morphism

$$\pi : \widetilde{X} \longrightarrow X$$

with the following universal property: for every normal integral scheme Z , and for every dominant morphism $f : Z \rightarrow X$, the morphism f factors uniquely through $\widetilde{X}$.

Solution.

Let X be an integral scheme. If $U = \operatorname{Spec} A \subseteq X$ is affine, then A is an integral domain. Its function field is

$$K = \operatorname{Frac}(A),$$

and $\overline{A}$ denotes the integral closure of A in K .

The basic compatibility needed for gluing is the following localization fact.

Lemma. If $S \subseteq A$ is a multiplicative subset, then the integral closure of $S^{-1}A$ in $\operatorname{Frac}(A)$ is

$$S^{-1}\overline{A}.$$

Proof of the lemma. Every element of $S^{-1}\overline{A}$ is integral over $S^{-1}A$, because integrality is preserved under localization.

Conversely, suppose $x \in \operatorname{Frac}(A)$ is integral over $S^{-1}A$. Then x satisfies a monic equation

$$x^n + \frac{a_1}{s_1}x^{n-1} + \cdots + \frac{a_n}{s_n} = 0, \quad a_i \in A, \quad s_i \in S.$$

After multiplying by a suitable element of S , one obtains that sx is integral over A for some $s \in S$. Hence

$$sx \in \overline{A},$$

and therefore

$$x = \frac{sx}{s} \in S^{-1}\overline{A}.$$

Thus the integral closure of $S^{-1}A$ is $S^{-1}\overline{A}$. $\square$

Now let $D(g) \subseteq U = \operatorname{Spec} A$ be a distinguished open subset. Then

$$D(g) = \operatorname{Spec} A_g.$$

By the lemma, the normalization of $D(g)$ is

$$\operatorname{Spec} \overline{A}_g = \operatorname{Spec}(\overline{A}_g).$$

But this is exactly the inverse image of $D(g)$ inside $\operatorname{Spec} \bar{A}$. Hence normalization is compatible with restriction to distinguished open subsets.

Since affine open subsets of a scheme have a basis of distinguished open subsets, these canonical identifications allow us to glue the affine schemes

$$\tilde{U} = \operatorname{Spec} \bar{A}$$

along overlaps. The cocycle condition is automatic because all identifications come from the same localization property above.

Thus the schemes $\tilde{U}$ glue to a scheme $\tilde{X}$.

The morphisms

$$A \longrightarrow \bar{A}$$

induce morphisms

$$\operatorname{Spec} \bar{A} \longrightarrow \operatorname{Spec} A.$$

These are compatible on overlaps, so they glue to a morphism

$$\pi : \tilde{X} \longrightarrow X.$$

We now check that $\tilde{X}$ is normal and integral. Normality is local on affine open subsets. Each affine piece of $\tilde{X}$ is of the form

$$\operatorname{Spec} \bar{A},$$

where $\bar{A}$ is integrally closed in its fraction field. Therefore every affine coordinate ring of $\tilde{X}$ is integrally closed. Hence $\tilde{X}$ is normal.

Also, since X is integral, the rings A are domains and their integral closures $\bar{A} \subseteq \operatorname{Frac}(A)$ are again domains. The affine pieces glue along nonempty open subsets in a compatible way, so $\tilde{X}$ is integral.

It remains to prove the universal property.

Let Z be a normal integral scheme, and let

$$f : Z \longrightarrow X$$

be a dominant morphism. Since X and Z are integral and f is dominant, f induces an inclusion of function fields

$$K(X) \hookrightarrow K(Z).$$

Let $U = \operatorname{Spec} A \subseteq X$ be affine, and let $W = \operatorname{Spec} B \subseteq f^{-1}(U)$ be affine. Since Z is normal, B is an integrally closed domain.

The morphism f gives a ring homomorphism

$$A \longrightarrow B.$$

Now take any element $\alpha \in \bar{A}$. By definition, α is integral over A . Its image in $K(Z)$ is therefore integral over the image of A , hence integral over B . Since B is integrally closed in its fraction field, the image of α actually lies in B .

Therefore the map $A \rightarrow B$ extends uniquely to a map

$$\overline{A} \longrightarrow B.$$

Equivalently, the morphism

$$W \longrightarrow U$$

lifts uniquely to a morphism

$$W \longrightarrow \tilde{U}.$$

These local lifts are compatible on overlaps, because they are all defined by the same induced map on function fields. Hence they glue to a global morphism

$$\tilde{f} : Z \longrightarrow \tilde{X}$$

such that

$$f = \pi \circ \tilde{f}.$$

Finally, uniqueness follows because on each affine open $\tilde{U} = \operatorname{Spec} \overline{A}$, the map

$$\overline{A} \longrightarrow \Gamma(W, \mathcal{O}_Z)$$

is forced by the induced map of function fields

$$K(X) \hookrightarrow K(Z).$$

Thus there is at most one such lift.

Therefore $\tilde{X}$ is the normalization of X , and $\pi : \tilde{X} \rightarrow X$ satisfies the desired universal property.

14 Problem 9. Fibres of Morphisms of Affine Schemes

Problem. Describe the fibres over all points of the target space for the following morphisms between affine schemes. In each case, the corresponding homomorphism of rings is the obvious one. Here k denotes a field. Which fibres are irreducible or reduced?

General principle. If

$$\varphi : A \longrightarrow B$$

is a ring homomorphism, then the corresponding morphism of affine schemes is

$$\operatorname{Spec} B \longrightarrow \operatorname{Spec} A.$$

For a point $\mathfrak{p} \in \operatorname{Spec} A$, the fibre over $\mathfrak{p}$ is

$$\operatorname{Spec} (B \otimes_A \kappa(\mathfrak{p})),$$

where

$$\kappa(\mathfrak{p}) = \operatorname{Frac}(A/\mathfrak{p})$$

is the residue field at $\mathfrak{p}$.

14.1 Part 1

Consider

$$\operatorname{Spec} k[T, U]/(TU - 1) \longrightarrow \operatorname{Spec} k[T].$$

Let $\mathfrak{p} \in \operatorname{Spec} k[T]$, and write

$$a = \overline{T} \in \kappa(\mathfrak{p}).$$

The fibre is

$$\operatorname{Spec} (\kappa(\mathfrak{p})[U]/(aU - 1)).$$

If $\mathfrak{p} = (T)$, then $a = 0$, so the fibre is

$$\operatorname{Spec} (k[U]/(-1)) = \operatorname{Spec}(0) = \emptyset.$$

If $\mathfrak{p} \neq (T)$, then $a \neq 0$ in $\kappa(\mathfrak{p})$, so

$$\kappa(\mathfrak{p})[U]/(aU - 1) \cong \kappa(\mathfrak{p}).$$

Hence the fibre is a single reduced point:

$$\operatorname{Spec} \kappa(\mathfrak{p}).$$

Thus:

Point of $\operatorname{Spec} k[T]$	Fibre	Reduced?	Irreducible?
(T)	$\emptyset$	yes	no, by the usual convention
$\mathfrak{p} \neq (T)$	$\operatorname{Spec} \kappa(\mathfrak{p})$	yes	yes

Geometrically, this is the morphism

$$\mathbb{A}_k^1 \setminus \{0\} \longrightarrow \mathbb{A}_k^1,$$

so the fibre over $T = 0$ is empty and every other fibre is one point.

14.2 Part 2

Consider

$$\operatorname{Spec} k[T, U]/(T^2 - U^2) \longrightarrow \operatorname{Spec} k[T].$$

Again let

$$a = \overline{T} \in \kappa(\mathfrak{p}).$$

The fibre is

$$\operatorname{Spec} (\kappa(\mathfrak{p})[U]/(a^2 - U^2)).$$

Equivalently,

$$a^2 - U^2 = (a - U)(a + U).$$

If $\mathfrak{p} = (T)$, then $a = 0$, so the fibre is

$$\operatorname{Spec} (k[U]/(U^2)).$$

This is one nonreduced point. Hence it is irreducible but not reduced.

Now suppose $\mathfrak{p} \neq (T)$, so $a \neq 0$.

If $\text{char } k \neq 2$, then $a \neq -a$, and

$$a^2 - U^2 = (a - U)(a + U)$$

has two distinct linear factors. Therefore

$$\kappa(\mathfrak{p})[U]/(a^2 - U^2) \cong \kappa(\mathfrak{p}) \times \kappa(\mathfrak{p}).$$

The fibre consists of two reduced points. It is reduced but not irreducible.

If $\text{char } k = 2$, then $a = -a$, and

$$a^2 - U^2 = (U - a)^2.$$

Hence the fibre is

$$\text{Spec } (\kappa(\mathfrak{p})[U]/((U - a)^2)).$$

This is one nonreduced point. It is irreducible but not reduced.

Thus, if $\text{char } k \neq 2$,

Point	Fibre	Reduced?	Irreducible?
(T)	$\text{Spec } k[U]/(U^2)$	no	yes
$\mathfrak{p} \neq (T)$	two reduced points	yes	no

If $\text{char } k = 2$, every fibre is a doubled point, hence irreducible but not reduced.

14.3 Part 3

Consider

$$\text{Spec } k[T, U, V, W]/((U + T)W, (U + T)(U^3 + U^2 + UV^2 - V^2)) \longrightarrow \text{Spec } k[T].$$

Let

$$a = \bar{T} \in \kappa(\mathfrak{p}).$$

The fibre is

$$\text{Spec } \frac{\kappa(\mathfrak{p})[U, V, W]}{((U + a)W, (U + a)(U^3 + U^2 + UV^2 - V^2))}.$$

Set

$$F(U, V) = U^3 + U^2 + UV^2 - V^2.$$

Then the fibre is cut out by

$$(U + a)W = 0, \quad (U + a)F(U, V) = 0.$$

Set-theoretically, this fibre is the union

$$\{U = -a\} \cup \{W = 0, F(U, V) = 0\}.$$

The first part is an affine plane:

$$\operatorname{Spec} \kappa(\mathfrak{p})[V, W].$$

The second part lies inside the plane $W = 0$ and is the curve

$$F(U, V) = 0.$$

Therefore every fibre is reducible: it contains the plane $U = -a$ and also the residual part $W = 0$, $F(U, V) = 0$.

Now discuss reducedness.

If $\operatorname{char} k \neq 2$, then $F(U, V)$ is square-free, and the factor $U + a$ is not a repeated factor of the defining equations. In this case the ideal

$$((U + a)W, (U + a)F)$$

is radical. Hence every fibre is reduced.

If $\operatorname{char} k = 2$, then

$$F(U, V) = U^3 + U^2 + UV^2 + V^2 = (U + 1)(U + V)^2.$$

Thus the equations contain a repeated factor. In this case every fibre is nonreduced.

Hence:

Characteristic of k	Reducedness of fibres	Irreducibility of fibres
$\operatorname{char} k \neq 2$	all fibres reduced	all fibres reducible
$\operatorname{char} k = 2$	all fibres nonreduced	all fibres reducible

14.4 Part 4

Consider

$$\operatorname{Spec} \mathbb{Z}[T] \longrightarrow \operatorname{Spec} \mathbb{Z}.$$

The points of $\operatorname{Spec} \mathbb{Z}$ are the generic point

$$(0)$$

and the closed points

$$(p),$$

where p is a prime number.

Over the generic point (0) , the residue field is

$$\kappa((0)) = \mathbb{Q}.$$

The fibre is

$$\operatorname{Spec} (\mathbb{Z}[T] \otimes_{\mathbb{Z}} \mathbb{Q}) = \operatorname{Spec} \mathbb{Q}[T] = \mathbb{A}_{\mathbb{Q}}^1.$$

Over a closed point (p) , the residue field is

$$\kappa((p)) = \mathbb{F}_p.$$

The fibre is

$$\operatorname{Spec}(\mathbb{Z}[T] \otimes_{\mathbb{Z}} \mathbb{F}_p) = \operatorname{Spec} \mathbb{F}_p[T] = \mathbb{A}_{\mathbb{F}_p}^1.$$

Thus:

Point of $\operatorname{Spec} \mathbb{Z}$	Fibre	Reduced?	Irreducible?
(0)	$\operatorname{Spec} \mathbb{Q}[T]$	yes	yes
(p)	$\operatorname{Spec} \mathbb{F}_p[T]$	yes	yes

So every fibre is an affine line over a field.

14.5 Part 5

Consider

$$\operatorname{Spec} \mathbb{Z}[T]/(T^2 + 1) \longrightarrow \operatorname{Spec} \mathbb{Z}.$$

Over the generic point (0), the fibre is

$$\operatorname{Spec}(\mathbb{Q}[T]/(T^2 + 1)).$$

Since $T^2 + 1$ is irreducible over $\mathbb{Q}$, we get

$$\mathbb{Q}[T]/(T^2 + 1) \cong \mathbb{Q}(i).$$

Thus the generic fibre is

$$\operatorname{Spec} \mathbb{Q}(i),$$

a single reduced point.

Now let p be a prime number. The fibre over (p) is

$$\operatorname{Spec}(\mathbb{F}_p[T]/(T^2 + 1)).$$

For $p = 2$, we have

$$T^2 + 1 = (T + 1)^2$$

in $\mathbb{F}_2[T]$. Hence the fibre is

$$\operatorname{Spec}(\mathbb{F}_2[T]/((T + 1)^2)),$$

a nonreduced one-point scheme. It is irreducible but not reduced.

For p odd, the polynomial $T^2 + 1$ splits over $\mathbb{F}_p$ if and only if -1 is a square modulo p . By the classical number theoretic criterion,

$$-1 \text{ is a square modulo } p \iff p \equiv 1 \pmod{4}.$$

Therefore:

If

$$p \equiv 1 \pmod{4},$$

then

$$T^2 + 1 = (T - a)(T + a)$$

for some $a \in \mathbb{F}_p$, and the fibre is

$$\operatorname{Spec}(\mathbb{F}_p \times \mathbb{F}_p).$$

It consists of two reduced points. Hence it is reduced but not irreducible.

If

$$p \equiv 3 \pmod{4},$$

then $T^2 + 1$ is irreducible over $\mathbb{F}_p$, so

$$\mathbb{F}_p[T]/(T^2 + 1) \cong \mathbb{F}_{p^2}.$$

The fibre is one reduced point:

$$\operatorname{Spec} \mathbb{F}_{p^2}.$$

Thus:

Point of $\operatorname{Spec} \mathbb{Z}$	Fibre	Reduced?	Irreducible?
(0)	$\operatorname{Spec} \mathbb{Q}(i)$	yes	yes
(2)	$\operatorname{Spec} \mathbb{F}_2[T]/((T+1)^2)$	no	yes
$(p), p \equiv 1 \pmod{4}$	two reduced points	yes	no
$(p), p \equiv 3 \pmod{4}$	$\operatorname{Spec} \mathbb{F}_{p^2}$	yes	yes

From the point of view of algebraic number theory, this is the behaviour of rational primes in the Gaussian integers $\mathbb{Z}[i]$:

Prime p	Behaviour in $\mathbb{Z}[i]$
$p = 2$	ramified
$p \equiv 1 \pmod{4}$	split
$p \equiv 3 \pmod{4}$	inert

14.6 Part 6

Consider

$$\operatorname{Spec} \mathbb{C} \longrightarrow \operatorname{Spec} \mathbb{Z}.$$

The corresponding ring homomorphism is

$$\mathbb{Z} \longrightarrow \mathbb{C}.$$

The scheme $\operatorname{Spec} \mathbb{C}$ has only one point, namely the zero ideal (0). Its inverse image in $\operatorname{Spec} \mathbb{Z}$ is

$$(0) \subseteq \mathbb{Z}.$$

Thus the image of the morphism is only the generic point of $\operatorname{Spec} \mathbb{Z}$.

The fibre over (0) is

$$\operatorname{Spec} (\mathbb{C} \otimes_{\mathbb{Z}} \mathbb{Q}) = \operatorname{Spec} \mathbb{C}.$$

This is one reduced point.

For a closed point (p) , the fibre is

$$\operatorname{Spec} (\mathbb{C} \otimes_{\mathbb{Z}} \mathbb{F}_p).$$

But

$$\mathbb{C} \otimes_{\mathbb{Z}} \mathbb{F}_p \cong \mathbb{C}/p\mathbb{C}.$$

Since p is invertible in $\mathbb{C}$, we have

$$p\mathbb{C} = \mathbb{C}.$$

Therefore

$$\mathbb{C}/p\mathbb{C} = 0.$$

Hence the fibre over (p) is empty.

Thus:

Point of $\text{Spec } \mathbb{Z}$	Fibre	Reduced?	Irreducible?
(0)	$\text{Spec } \mathbb{C}$	yes	yes
(p)	$\emptyset$	yes	no, by the usual convention

15 Problem 10. Cartesian Diagrams

Problem. A commutative diagram

$$\begin{array}{ccc} A & \longrightarrow & B \\ \downarrow & & \downarrow \\ C & \longrightarrow & D \end{array}$$

is Cartesian if the induced morphism

$$A \longrightarrow B \times_D C$$

is an isomorphism.

Show that the following diagrams are Cartesian in any category.

15.1 First diagram

We are given morphisms

$$X \longrightarrow S \longrightarrow T, \quad Y \longrightarrow S \longrightarrow T.$$

Then we have fibre products

$$X \times_S Y \quad \text{and} \quad X \times_T Y.$$

Also, the diagonal morphism

$$\Delta : S \longrightarrow S \times_T S$$

is induced by the two identity maps $S \rightarrow S$.

The diagram is

$$\begin{array}{ccc} X \times_S Y & \longrightarrow & S \\ \downarrow & & \downarrow \Delta \\ X \times_T Y & \longrightarrow & S \times_T S. \end{array}$$

We prove that it is Cartesian.

Let Q be any object. A morphism

$$Q \longrightarrow (X \times_T Y) \times_{S \times_T S} S$$

is the same as the following data:

$$x : Q \rightarrow X, \quad y : Q \rightarrow Y, \quad s : Q \rightarrow S,$$

such that:

$$x \text{ and } y \text{ agree after mapping to } T,$$

and

$$\Delta \circ s = (x_S, y_S),$$

where

$$x_S : Q \rightarrow X \rightarrow S, \quad y_S : Q \rightarrow Y \rightarrow S.$$

But the condition

$$\Delta \circ s = (x_S, y_S)$$

means precisely that

$$s = x_S = y_S.$$

Therefore the data above is exactly the same as a pair of morphisms

$$x : Q \rightarrow X, \quad y : Q \rightarrow Y$$

such that

$$x_S = y_S.$$

By the universal property of the fibre product, this is exactly the same as a morphism

$$Q \longrightarrow X \times_S Y.$$

Thus for every object Q , there is a natural bijection

$$\mathrm{Hom}(Q, X \times_S Y) \cong \mathrm{Hom}(Q, (X \times_T Y) \times_{S \times_T S} S).$$

By the Yoneda principle, this gives an isomorphism

$$X \times_S Y \cong (X \times_T Y) \times_{S \times_T S} S.$$

Therefore the diagram is Cartesian.

15.2 Second diagram

Now assume X and Y are objects over S , and let

$$f : X \longrightarrow Y$$

be a morphism over S .

The diagonal morphism is

$$\Delta : Y \longrightarrow Y \times_S Y.$$

The graph morphism of f is

$$\Gamma_f : X \longrightarrow X \times_S Y,$$

defined by

$$\Gamma_f = (\text{id}_X, f).$$

The diagram is

$$\begin{array}{ccc} X & \xrightarrow{f} & Y \\ \Gamma_f \downarrow & & \downarrow \Delta \\ X \times_S Y & \xrightarrow{f \times \text{id}_Y} & Y \times_S Y. \end{array}$$

We prove that it is Cartesian.

Let Q be any object. A morphism

$$Q \longrightarrow (X \times_S Y) \times_{Y \times_S Y} Y$$

is the same as the following data:

$$x : Q \rightarrow X, \quad y : Q \rightarrow Y, \quad z : Q \rightarrow Y,$$

such that (x, y) defines a morphism

$$Q \rightarrow X \times_S Y,$$

and

$$(f \circ x, y) = \Delta \circ z.$$

But

$$\Delta \circ z = (z, z).$$

Hence the condition becomes

$$(f \circ x, y) = (z, z).$$

Therefore

$$z = f \circ x \quad \text{and} \quad y = f \circ x.$$

Thus the whole data is uniquely determined by the single morphism

$$x : Q \rightarrow X.$$

Conversely, any morphism $x : Q \rightarrow X$ gives such a triple by setting

$$y = f \circ x, \quad z = f \circ x.$$

Therefore there is a natural bijection

$$\mathrm{Hom}(Q, X) \cong \mathrm{Hom}(Q, (X \times_S Y) \times_{Y \times_S Y} Y).$$

By the Yoneda principle,

$$X \cong (X \times_S Y) \times_{Y \times_S Y} Y.$$

Therefore the second diagram is Cartesian.

16 Problem 10. Cartesian Diagrams

Problem. A commutative diagram

$$\begin{array}{ccc} A & \longrightarrow & B \\ \downarrow & & \downarrow \\ C & \longrightarrow & D \end{array}$$

is Cartesian if the induced morphism

$$A \longrightarrow B \times_D C$$

is an isomorphism.

Show that the following diagrams are Cartesian in any category.

16.1 First diagram

We are given morphisms

$$X \longrightarrow S \longrightarrow T, \quad Y \longrightarrow S \longrightarrow T.$$

Then we have fibre products

$$X \times_S Y \quad \text{and} \quad X \times_T Y.$$

Also, the diagonal morphism

$$\Delta : S \longrightarrow S \times_T S$$

is induced by the two identity maps $S \rightarrow S$.

The diagram is

$$\begin{array}{ccc} X \times_S Y & \longrightarrow & S \\ \downarrow & & \downarrow \Delta \\ X \times_T Y & \longrightarrow & S \times_T S. \end{array}$$

We prove that it is Cartesian.

Let Q be any object. A morphism

$$Q \longrightarrow (X \times_T Y) \times_{S \times_T S} S$$

is the same as the following data:

$$x : Q \rightarrow X, \quad y : Q \rightarrow Y, \quad s : Q \rightarrow S,$$

such that:

$$x \text{ and } y \text{ agree after mapping to } T,$$

and

$$\Delta \circ s = (x_S, y_S),$$

where

$$x_S : Q \rightarrow X \rightarrow S, \quad y_S : Q \rightarrow Y \rightarrow S.$$

But the condition

$$\Delta \circ s = (x_S, y_S)$$

means precisely that

$$s = x_S = y_S.$$

Therefore the data above is exactly the same as a pair of morphisms

$$x : Q \rightarrow X, \quad y : Q \rightarrow Y$$

such that

$$x_S = y_S.$$

By the universal property of the fibre product, this is exactly the same as a morphism

$$Q \longrightarrow X \times_S Y.$$

Thus for every object Q , there is a natural bijection

$$\mathrm{Hom}(Q, X \times_S Y) \cong \mathrm{Hom}(Q, (X \times_T Y) \times_{S \times_T S} S).$$

By the Yoneda principle, this gives an isomorphism

$$X \times_S Y \cong (X \times_T Y) \times_{S \times_T S} S.$$

Therefore the diagram is Cartesian.

16.2 Second diagram

Now assume X and Y are objects over S , and let

$$f : X \longrightarrow Y$$

be a morphism over S .

The diagonal morphism is

$$\Delta : Y \longrightarrow Y \times_S Y.$$

The graph morphism of f is

$$\Gamma_f : X \longrightarrow X \times_S Y,$$

defined by

$$\Gamma_f = (\text{id}_X, f).$$

The diagram is

$$\begin{array}{ccc} X & \xrightarrow{f} & Y \\ \Gamma_f \downarrow & & \downarrow \Delta \\ X \times_S Y & \xrightarrow{f \times \text{id}_Y} & Y \times_S Y. \end{array}$$

We prove that it is Cartesian.

Let Q be any object. A morphism

$$Q \longrightarrow (X \times_S Y) \times_{Y \times_S Y} Y$$

is the same as the following data:

$$x : Q \rightarrow X, \quad y : Q \rightarrow Y, \quad z : Q \rightarrow Y,$$

such that (x, y) defines a morphism

$$Q \rightarrow X \times_S Y,$$

and

$$(f \circ x, y) = \Delta \circ z.$$

But

$$\Delta \circ z = (z, z).$$

Hence the condition becomes

$$(f \circ x, y) = (z, z).$$

Therefore

$$z = f \circ x \quad \text{and} \quad y = f \circ x.$$

Thus the whole data is uniquely determined by the single morphism

$$x : Q \rightarrow X.$$

Conversely, any morphism $x : Q \rightarrow X$ gives such a triple by setting

$$y = f \circ x, \quad z = f \circ x.$$

Therefore there is a natural bijection

$$\text{Hom}(Q, X) \cong \text{Hom}(Q, (X \times_S Y) \times_{Y \times_S Y} Y).$$

By the Yoneda principle,

$$X \cong (X \times_S Y) \times_{Y \times_S Y} Y.$$

Therefore the second diagram is Cartesian.