

Algebraic Geometry 6

Lecture notes with examples

July 29, 2026

Contents

1	Problem 5. Locally Finite Type Can Be Checked on Every Affine Open	3
2	Problem 6. Examples Violating the Properties Defined Above	5
3	Problem 7. Generic Point and Function Field of an Integral Scheme	10
4	Theory: Locally Finite Type, Finite Type, Generic Points, and Function Fields	13
4.1	Local Nature of Schemes	13
4.2	Locally of Finite Type Morphisms	15
4.3	Why Locally Finite Type Can Be Checked on Every Affine Open	15
4.4	Finite Type Morphisms	17
4.5	Noetherian Schemes	18
4.6	Generic Points	19
4.7	Function Fields	20
4.8	Function Field on Any Affine Open	20
4.9	Examples of Function Fields	21
4.10	Summary of the Main Ideas	21
4.11	Conceptual Picture	22
5	Examples: Locally Finite Type, Finite Type, Generic Points, and Function Fields	22
5.1	Examples of Locally Finite Type Morphisms	22
5.2	Examples Showing the Difference Between Locally Finite Type and Finite Type	24
5.3	Examples of Reduced, Irreducible, and Integral Schemes	25

5.4	Examples of Generic Points	27
5.5	Examples of Function Fields	28
5.6	Same Function Field on Different Affine Opens	30
5.7	Example Where the Function Field Is Not Defined in This Sense	30
5.8	Summary of Examples	31
6	Charts, Diagrams, and Tables: Locally Finite Type, Generic Points, and Function Fields	31
6.1	Chart: Local Properties of Schemes	31
6.2	Diagram: How Locally Finite Type Is Checked	31
6.3	Table: Locally Finite Type versus Finite Type	32
6.4	Conceptual Formula: Finite Type	32
6.5	Diagram: The Difference Between Locally Finite Type and Finite Type .	33
6.6	Table: Standard Counterexamples	33
6.7	Diagram: Generic Point of an Integral Affine Scheme	33
6.8	Example Diagram: Generic Point of $\mathbb{A}_k^1$	34
6.9	Diagram: Reducible Scheme with Two Generic Points	35
6.10	Chart: Function Field of an Integral Scheme	35
6.11	Table: Function Fields of Standard Schemes	36
6.12	Diagram: Computing $K(X)$ on Different Affine Opens	36
6.13	Table: Integral versus Reducible Case	37
6.14	Final Summary Diagram	37

1 Problem 5. Locally Finite Type Can Be Checked on Every Affine Open

Problem. A morphism $f : X \rightarrow Y$ is *locally of finite type* if there exists a covering of Y by open affine subsets

$$V_i = \operatorname{Spec} B_i$$

such that, for each i , the open subset $f^{-1}(V_i)$ can be covered by open affine subsets

$$U_{ij} = \operatorname{Spec} A_{ij},$$

where each A_{ij} is a finitely generated B_i -algebra.

The morphism is *of finite type* if the cover of $f^{-1}(V_i)$ above can be taken to be finite.

Show that a morphism $f : X \rightarrow Y$ is locally of finite type if and only if for every open affine subset

$$V = \operatorname{Spec} B \subseteq Y,$$

the inverse image $f^{-1}(V)$ can be covered by open affine subsets

$$U_j = \operatorname{Spec} A_j,$$

where each A_j is a finitely generated B -algebra.

Solution.

We prove both directions.

Step 1. The easy direction

Assume that for every open affine subset

$$V = \operatorname{Spec} B \subseteq Y,$$

the inverse image $f^{-1}(V)$ can be covered by open affine subsets

$$U_j = \operatorname{Spec} A_j$$

such that every A_j is a finitely generated B -algebra.

Then in particular this is true for the affine opens appearing in any affine open cover of Y . Therefore f is locally of finite type by definition.

Step 2. The nontrivial direction

Now assume that $f : X \rightarrow Y$ is locally of finite type.

By definition, there exists an affine open cover

$$Y = \bigcup_i V_i, \quad V_i = \operatorname{Spec} B_i,$$

such that each $f^{-1}(V_i)$ is covered by open affine subsets

$$U_{ij} = \operatorname{Spec} A_{ij},$$

where each A_{ij} is a finitely generated B_i -algebra.

We must show that the same property holds for an arbitrary affine open subset

$$V = \operatorname{Spec} B \subseteq Y.$$

Let $y \in V$. Since the V_i cover Y , there exists some i such that

$$y \in V_i.$$

Thus

$$y \in V \cap V_i.$$

The intersection $V \cap V_i$ is open both in V and in V_i . Since affine schemes have bases of distinguished open subsets, there exists an open subset

$$W \subseteq V \cap V_i$$

containing y such that W is distinguished open in both affine schemes. Hence we may write

$$W = D(b) \subseteq \operatorname{Spec} B$$

and also

$$W = D(c) \subseteq \operatorname{Spec} B_i.$$

Therefore

$$W \cong \operatorname{Spec} B_b$$

and also

$$W \cong \operatorname{Spec}(B_i)_c.$$

Now consider

$$f^{-1}(W) \subseteq f^{-1}(V_i).$$

Since $f^{-1}(V_i)$ is covered by the affine opens

$$U_{ij} = \operatorname{Spec} A_{ij},$$

the inverse image $f^{-1}(W)$ is covered by the intersections

$$U_{ij} \cap f^{-1}(W).$$

Because $W = D(c) \subseteq V_i$, the preimage of W inside

$$U_{ij} = \operatorname{Spec} A_{ij}$$

is a distinguished open subset. More precisely, if

$$B_i \longrightarrow A_{ij}$$

is the corresponding ring map and $c \in B_i$, then

$$U_{ij} \cap f^{-1}(W) = D(\varphi(c)) \subseteq \operatorname{Spec} A_{ij}.$$

Hence

$$U_{ij} \cap f^{-1}(W) \cong \operatorname{Spec}((A_{ij})_{\varphi(c)}).$$

Since A_{ij} is a finitely generated B_i -algebra, the localization

$$(A_{ij})_{\varphi(c)}$$

is a finitely generated $(B_i)_c$ -algebra.

But

$$(B_i)_c \cong B_b.$$

Therefore

$$(A_{ij})_{\varphi(c)}$$

is a finitely generated B_b -algebra.

Finally, B_b is itself a finitely generated B -algebra, because

$$B_b \cong B[t]/(bt - 1).$$

Thus $(A_{ij})_{\varphi(c)}$ is finitely generated as a B -algebra.

Since the sets W cover V , the affine opens

$$U_{ij} \cap f^{-1}(W)$$

cover $f^{-1}(V)$. Each of them has the form

$$\operatorname{Spec} A$$

where A is a finitely generated B -algebra.

Therefore, for every open affine subset

$$V = \operatorname{Spec} B \subseteq Y,$$

the inverse image $f^{-1}(V)$ can be covered by open affine subsets whose coordinate rings are finitely generated B -algebras.

Hence f is locally of finite type in the desired stronger sense.

$f : X \rightarrow Y$ is locally of finite type $\iff$ the condition holds over every affine open $V = \operatorname{Spec} B \subseteq Y$.

2 Problem 6. Examples Violating the Properties Defined Above

Problem. For each of the properties defined above of schemes or morphisms, give an example of a scheme or morphism which violates that property.

Give an example of a morphism which is locally of finite type but not of finite type.

Solution.

We collect standard examples. Throughout, let k be a field.

1. A scheme which is not reduced

Consider

$$X = \operatorname{Spec} k[\varepsilon]/(\varepsilon^2).$$

The ring

$$k[\varepsilon]/(\varepsilon^2)$$

has a nonzero nilpotent element, namely ε , because

$$\varepsilon \neq 0, \quad \varepsilon^2 = 0.$$

Therefore the ring is not reduced, and hence

$$X$$

is not a reduced scheme.

$\operatorname{Spec} k[\varepsilon]/(\varepsilon^2) \text{ is not reduced.}$

2. A scheme which is not irreducible

Consider

$$X = \operatorname{Spec}(k \times k).$$

The prime ideals of $k \times k$ are

$$k \times 0 \quad \text{and} \quad 0 \times k.$$

Thus

$$\operatorname{Spec}(k \times k)$$

has two closed points and is the disjoint union of two one-point spaces. Hence it is reducible.

Equivalently,

$$k \times k$$

has two nontrivial idempotents,

$$(1, 0) \quad \text{and} \quad (0, 1),$$

so the scheme splits into two pieces.

$\operatorname{Spec}(k \times k) \text{ is not irreducible.}$

3. A scheme which is not integral

A scheme is integral if it is both reduced and irreducible.

Consider

$$X = \operatorname{Spec} k[x, y]/(xy).$$

The ring

$$k[x, y]/(xy)$$

has zero divisors, because

$$x \cdot y = 0$$

in the quotient, while neither x nor y is zero.

Thus $k[x, y]/(xy)$ is not an integral domain. Therefore

$$X$$

is not integral.

Geometrically, this is the union of the two coordinate axes in the affine plane:

$$xy = 0.$$

It has two irreducible components:

$$x = 0 \quad \text{and} \quad y = 0.$$

$\operatorname{Spec} k[x, y]/(xy) \text{ is not integral.}$

4. A scheme which is not affine

The projective line

$$\mathbb{P}_k^1$$

is not affine.

Indeed, the global regular functions on $\mathbb{P}_k^1$ are only the constant functions:

$$\Gamma(\mathbb{P}_k^1, \mathcal{O}_{\mathbb{P}_k^1}) = k.$$

If $\mathbb{P}_k^1$ were affine, then it would have to be isomorphic to

$$\operatorname{Spec} k,$$

because an affine scheme is determined by its ring of global functions. But $\mathbb{P}_k^1$ is not a single point. Therefore $\mathbb{P}_k^1$ is not affine.

$\mathbb{P}_k^1 \text{ is not affine.}$

5. A scheme which is not Noetherian

Consider the affine scheme

$$X = \operatorname{Spec} k[x_1, x_2, x_3, \dots].$$

The ring

$$k[x_1, x_2, x_3, \dots]$$

is not Noetherian, because it has the infinite strictly increasing chain of ideals

$$(x_1) \subsetneq (x_1, x_2) \subsetneq (x_1, x_2, x_3) \subsetneq \dots.$$

Therefore

$$\operatorname{Spec} k[x_1, x_2, x_3, \dots]$$

is not a Noetherian scheme.

$\operatorname{Spec} k[x_1, x_2, x_3, \dots] \text{ is not Noetherian.}$

6. A morphism which is not locally of finite type

Consider the natural morphism

$$f : \operatorname{Spec} k[x_1, x_2, x_3, \dots] \longrightarrow \operatorname{Spec} k.$$

This corresponds to the inclusion

$$k \longrightarrow k[x_1, x_2, x_3, \dots].$$

The morphism is locally of finite type if the coordinate ring is locally covered by finitely generated k -algebras. But since

$$k[x_1, x_2, x_3, \dots]$$

is not finitely generated as a k -algebra, the affine morphism

$$\operatorname{Spec} k[x_1, x_2, x_3, \dots] \rightarrow \operatorname{Spec} k$$

is not of finite type.

Moreover, because the source is affine, if it were locally of finite type over $\operatorname{Spec} k$, then it would be covered by affine opens of finite type over k . Since the source is quasi-compact, finitely many of these affine opens would cover it, forcing the scheme to be of finite type over k , which it is not.

Thus

$$f$$

is not locally of finite type.

$\operatorname{Spec} k[x_1, x_2, x_3, \dots] \rightarrow \operatorname{Spec} k \text{ is not locally of finite type.}$

7. A morphism locally of finite type but not of finite type

Let

$$X = \coprod_{n=1}^{\infty} \operatorname{Spec} k$$

be the infinite disjoint union of countably many copies of $\operatorname{Spec} k$.

There is a natural morphism

$$f : X \longrightarrow \operatorname{Spec} k.$$

Each component

$$\operatorname{Spec} k \rightarrow \operatorname{Spec} k$$

is of finite type. Since X is covered by these open affine components, the morphism

$$f : X \rightarrow \operatorname{Spec} k$$

is locally of finite type.

However, f is not of finite type.

Indeed, if f were of finite type, then since the target $\operatorname{Spec} k$ is affine, the inverse image

$$f^{-1}(\operatorname{Spec} k) = X$$

would be covered by finitely many affine opens of finite type over k . In particular, X would be quasi-compact.

But

$$X = \coprod_{n=1}^{\infty} \operatorname{Spec} k$$

is not quasi-compact. The open cover by its connected components,

$$X = \bigcup_{n=1}^{\infty} \operatorname{Spec} k,$$

has no finite subcover.

Therefore f is locally of finite type but not of finite type.

$\coprod_{n=1}^{\infty} \operatorname{Spec} k \longrightarrow \operatorname{Spec} k$ is locally of finite type but not of finite type.

Summary table

Property	Counterexample	Reason
Reduced	$\operatorname{Spec} k[\varepsilon]/(\varepsilon^2)$	$\varepsilon \neq 0, \varepsilon^2 = 0$
Irreducible	$\operatorname{Spec}(k \times k)$	Two disconnected points
Integral	$\operatorname{Spec} k[x, y]/(xy)$	$x \cdot y = 0$
Affine	$\mathbb{P}_k^1$	Projective line is not affine
Noetherian	$\operatorname{Spec} k[x_1, x_2, x_3, \dots]$	Infinite ascending chain of ideals
Locally finite type	$\operatorname{Spec} k[x_1, x_2, x_3, \dots] \rightarrow \operatorname{Spec} k$	Infinitely many algebra generators
Finite type	$\coprod_{n=1}^{\infty} \operatorname{Spec} k \rightarrow \operatorname{Spec} k$	Not quasi-compact

3 Problem 7. Generic Point and Function Field of an Integral Scheme

Problem. Let X be an integral scheme.

Show that there is a unique point η such that

$$\overline{\{\eta\}} = X.$$

This point is called the *generic point* of X .

Show that the stalk of $\mathcal{O}_X$ at η is a field, called the *function field* of X , denoted

$$K(X).$$

Show that if

$$U = \operatorname{Spec} A$$

is any open affine subset of X , then

$$K(X)$$

is the field of fractions of A .

Solution.

Since X is integral, it is reduced and irreducible. In particular, the underlying topological space of X is irreducible.

Step 1. Existence of the generic point

Let

$$U = \operatorname{Spec} A$$

be a nonempty affine open subset of X .

Because X is integral, every nonempty affine open subset of X is also integral. Therefore A is an integral domain.

In the affine scheme

$$U = \operatorname{Spec} A,$$

the zero ideal

$$(0) \subseteq A$$

is a prime ideal, because A is a domain.

Thus (0) defines a point of U . Denote this point by

$$\eta.$$

Inside $U = \operatorname{Spec} A$, the closure of η is all of U , because

$$\overline{\{(0)\}} = V((0)) = \operatorname{Spec} A.$$

Thus

$$\overline{\{\eta\}}^U = U.$$

Since X is irreducible, every nonempty open subset of X is dense in X . Therefore

$$\overline{U}^X = X.$$

Since $\overline{\{\eta\}}^X$ contains U , it follows that

$$\overline{\{\eta\}}^X = X.$$

Thus X has a point η whose closure is all of X .

$$\boxed{\overline{\{\eta\}} = X.}$$

Step 2. Uniqueness of the generic point

Suppose $\eta' \in X$ is another point such that

$$\overline{\{\eta'\}} = X.$$

Then η and η' have the same closure.

Schemes are T_0 -spaces, meaning that two distinct points cannot have the same closure. Hence

$$\eta = \eta'.$$

Therefore the generic point of an integral scheme is unique.

$$\boxed{\text{An integral scheme has a unique generic point.}}$$

Step 3. The stalk at the generic point is a field

Let

$$U = \operatorname{Spec} A$$

be a nonempty affine open subset of X . As above, A is a domain, and the generic point η corresponds to the prime ideal

$$(0) \subseteq A.$$

The stalk of the structure sheaf at η can be computed on the affine open U :

$$\mathcal{O}_{X,\eta} = \mathcal{O}_{U,\eta}.$$

Since $U = \operatorname{Spec} A$ and $\eta = (0)$, we have

$$\mathcal{O}_{U,\eta} = A_{(0)}.$$

But localizing a domain A at the zero prime ideal gives its field of fractions:

$$A_{(0)} = \operatorname{Frac}(A).$$

Therefore

$$\mathcal{O}_{X,\eta} = \text{Frac}(A).$$

In particular, $\mathcal{O}_{X,\eta}$ is a field.

This field is called the *function field* of X and is denoted

$$K(X).$$

Thus

$$K(X) := \mathcal{O}_{X,\eta}.$$

$$\boxed{K(X) = \mathcal{O}_{X,\eta} \text{ is a field.}}$$

Step 4. Computing the function field on any affine open

Let

$$U = \text{Spec } A$$

be any nonempty affine open subset of X .

Since X is irreducible and η is the generic point, every nonempty open subset of X contains η . Therefore

$$\eta \in U.$$

Because X is integral, A is an integral domain. The point η corresponds to the zero ideal

$$(0) \subseteq A.$$

Hence

$$K(X) = \mathcal{O}_{X,\eta} = \mathcal{O}_{U,\eta} = A_{(0)} = \text{Frac}(A).$$

Therefore, for every nonempty affine open subset

$$U = \text{Spec } A \subseteq X,$$

we have

$$\boxed{K(X) = \text{Frac}(A).}$$

Geometric meaning

The function field $K(X)$ is the field of rational functions on X .

If

$$X = \text{Spec } A$$

is affine and integral, then

$$K(X) = \text{Frac}(A).$$

For example:

$$K(\mathbb{A}_k^1) = K(\text{Spec } k[x]) = k(x),$$

and

$$K(\mathbb{A}_k^n) = K(\operatorname{Spec} k[x_1, \dots, x_n]) = k(x_1, \dots, x_n).$$

Thus the generic point behaves like the “general point” of the scheme, and the stalk at that point consists of rational functions defined near the general point.

4 Theory: Locally Finite Type, Finite Type, Generic Points, and Function Fields

This section collects the main theoretical ideas behind Problems 5–7. The central theme is that many important properties of schemes and morphisms are *local*: they can be checked on affine open subsets. For schemes, affine opens reduce geometric questions to commutative algebra. For morphisms, inverse images of affine opens allow us to study the morphism by studying ring maps.

4.1 Local Nature of Schemes

A scheme is built by gluing affine schemes. An affine scheme has the form

$$\operatorname{Spec} A$$

for a commutative ring A . Therefore, many geometric properties of schemes can be tested on affine open subsets.

For example, if X is a scheme and

$$U = \operatorname{Spec} A \subseteq X$$

is an affine open subset, then the geometry of U is controlled by the algebra of A .

Reduced schemes

A ring A is *reduced* if it has no nonzero nilpotent elements. That is,

$$a^n = 0 \implies a = 0.$$

A scheme X is reduced if every affine open subset

$$U = \operatorname{Spec} A$$

has A reduced.

Equivalently, X is reduced if the structure sheaf $\mathcal{O}_X$ has no nonzero nilpotent sections locally.

A standard non-reduced example is

$$\operatorname{Spec} k[\varepsilon]/(\varepsilon^2).$$

Here

$$\varepsilon \neq 0, \quad \varepsilon^2 = 0.$$

So ε is a nonzero nilpotent element.

Irreducible schemes

A topological space X is *irreducible* if it cannot be written as a union of two proper closed subsets:

$$X \neq Z_1 \cup Z_2$$

with $Z_1, Z_2 \subsetneq X$ closed.

Equivalently, X is irreducible if every two nonempty open subsets intersect:

$$U \neq \emptyset, \quad V \neq \emptyset \quad \implies \quad U \cap V \neq \emptyset.$$

For an affine scheme

$$X = \operatorname{Spec} A,$$

the space X is irreducible if and only if the nilradical

$$\sqrt{(0)}$$

is a prime ideal.

If A is reduced, then $\sqrt{(0)} = (0)$. Hence, for a reduced affine scheme,

$$\operatorname{Spec} A \text{ is irreducible} \iff (0) \text{ is prime} \iff A \text{ is a domain.}$$

Integral schemes

A scheme X is *integral* if it is both reduced and irreducible.

On affine opens, this means the following:

$$X \text{ integral} \iff \text{every nonempty affine open } \operatorname{Spec} A \subseteq X \text{ has } A \text{ an integral domain.}$$

Thus integral schemes are the geometric analogues of integral domains.

A typical non-integral example is

$$\operatorname{Spec} k[x, y]/(xy).$$

In this ring,

$$x \cdot y = 0,$$

but neither x nor y is zero. Therefore the ring has zero divisors, so the scheme is not integral.

Geometrically, this is the union of the two coordinate axes:

$$xy = 0.$$

4.2 Locally of Finite Type Morphisms

Let

$$f : X \rightarrow Y$$

be a morphism of schemes.

The morphism f is *locally of finite type* if locally on the target and locally on the source, it is induced by finitely generated algebra maps.

More precisely, f is locally of finite type if there exists an affine open cover

$$Y = \bigcup_i V_i, \quad V_i = \operatorname{Spec} B_i,$$

such that each inverse image

$$f^{-1}(V_i)$$

can be covered by affine opens

$$U_{ij} = \operatorname{Spec} A_{ij},$$

where each A_{ij} is a finitely generated B_i -algebra.

So locally, the morphism is described by ring maps

$$B_i \rightarrow A_{ij}$$

such that

$$A_{ij} = B_i[a_1, \dots, a_n]$$

for finitely many algebra generators.

Geometric meaning

A morphism locally of finite type is one whose fibers and local coordinate rings are controlled by finitely many algebraic parameters.

For example,

$$\mathbb{A}_k^n = \operatorname{Spec} k[x_1, \dots, x_n]$$

is locally of finite type over

$$\operatorname{Spec} k$$

because

$$k[x_1, \dots, x_n]$$

is finitely generated as a k -algebra.

In contrast,

$$\operatorname{Spec} k[x_1, x_2, x_3, \dots] \rightarrow \operatorname{Spec} k$$

is not locally of finite type, because infinitely many variables are needed.

4.3 Why Locally Finite Type Can Be Checked on Every Affine Open

The definition of locally finite type only asks for *some* affine open cover of Y . However, the important theorem is that this automatically implies the same condition over *every* affine open subset of Y .

Theorem

Let

$$f : X \rightarrow Y$$

be a morphism of schemes. Then f is locally of finite type if and only if for every affine open subset

$$V = \operatorname{Spec} B \subseteq Y,$$

the inverse image

$$f^{-1}(V)$$

can be covered by affine open subsets

$$U_j = \operatorname{Spec} A_j$$

such that each A_j is a finitely generated B -algebra.

Idea of the proof

Assume f is locally of finite type. Then there is one good affine cover

$$Y = \bigcup_i \operatorname{Spec} B_i.$$

Now take any other affine open

$$V = \operatorname{Spec} B \subseteq Y.$$

We compare V with the chosen affine cover by looking at intersections

$$V \cap V_i.$$

Affine schemes have a basis of distinguished open subsets. Therefore, around any point of $V \cap V_i$, we can find an open subset

$$W$$

which is distinguished both inside V and inside V_i . Hence

$$W = D(b) \subseteq \operatorname{Spec} B$$

and

$$W = D(c) \subseteq \operatorname{Spec} B_i.$$

Thus

$$W \cong \operatorname{Spec} B_b$$

and

$$W \cong \operatorname{Spec}(B_i)_c.$$

If

$$U_{ij} = \operatorname{Spec} A_{ij}$$

lies over V_i , then its restriction over W is

$$U_{ij} \cap f^{-1}(W) \cong \operatorname{Spec}(A_{ij})_{\varphi(c)}.$$

Since A_{ij} is finitely generated over B_i , the localization

$$(A_{ij})_{\varphi(c)}$$

is finitely generated over

$$(B_i)_c.$$

But

$$(B_i)_c \cong B_b.$$

Also,

$$B_b \cong B[t]/(bt - 1),$$

so B_b is finitely generated over B . Therefore

$$(A_{ij})_{\varphi(c)}$$

is finitely generated over B .

Thus the property descends to every affine open $V = \operatorname{Spec} B \subseteq Y$.

4.4 Finite Type Morphisms

A morphism

$$f : X \rightarrow Y$$

is *of finite type* if it is locally of finite type and the affine covers of the inverse images of affine opens can be chosen to be finite.

Equivalently, a morphism is of finite type if it is locally of finite type and quasi-compact.

Informally:

locally of finite type = locally finitely many equations and variables,

while

finite type = locally finite type plus globally finitely many affine pieces over affine opens.

Difference between locally finite type and finite type

The difference is a compactness difference.

Consider

$$X = \coprod_{n=1}^{\infty} \operatorname{Spec} k.$$

There is a natural morphism

$$X \rightarrow \operatorname{Spec} k.$$

Each component

$$\operatorname{Spec} k \rightarrow \operatorname{Spec} k$$

is of finite type. Hence the morphism is locally of finite type.

But X is an infinite disjoint union of points. It is not quasi-compact, because the cover by its connected components has no finite subcover:

$$X = \bigcup_{n=1}^{\infty} \operatorname{Spec} k.$$

Therefore

$$\prod_{n=1}^{\infty} \operatorname{Spec} k \rightarrow \operatorname{Spec} k$$

is locally of finite type but not of finite type.

4.5 Noetherian Schemes

A ring A is *Noetherian* if every ideal of A is finitely generated. Equivalently, every ascending chain of ideals stabilizes:

$$I_1 \subseteq I_2 \subseteq I_3 \subseteq \cdots$$

eventually becomes constant.

A scheme X is *locally Noetherian* if it has an affine open cover

$$X = \bigcup_i \operatorname{Spec} A_i$$

where each A_i is Noetherian.

It is *Noetherian* if it is locally Noetherian and quasi-compact.

A standard non-Noetherian example is

$$\operatorname{Spec} k[x_1, x_2, x_3, \dots].$$

Indeed, the ring

$$k[x_1, x_2, x_3, \dots]$$

has the infinite strictly increasing chain of ideals

$$(x_1) \subsetneq (x_1, x_2) \subsetneq (x_1, x_2, x_3) \subsetneq \cdots.$$

Therefore it is not Noetherian.

4.6 Generic Points

Let X be an integral scheme.

Because X is irreducible, it should have one point representing the whole space. This is the *generic point*.

A point

$$\eta \in X$$

is called a generic point of X if

$$\overline{\{\eta\}} = X.$$

That is, η is a point whose closure is the entire space.

Existence

Let

$$U = \operatorname{Spec} A$$

be a nonempty affine open subset of X .

Since X is integral, A is an integral domain. Therefore the zero ideal

$$(0) \subseteq A$$

is prime.

Hence (0) defines a point of

$$\operatorname{Spec} A.$$

The closure of this point inside $\operatorname{Spec} A$ is

$$V((0)) = \operatorname{Spec} A.$$

Thus the point corresponding to (0) is dense in U .

Since X is irreducible, every nonempty open subset of X is dense in X . Therefore this point is dense in all of X .

Hence there exists a point

$$\eta \in X$$

such that

$$\overline{\{\eta\}} = X.$$

Uniqueness

Schemes are T_0 -spaces. This means distinct points have distinct closures.

Therefore, if two points η and η' both satisfy

$$\overline{\{\eta\}} = X \quad \text{and} \quad \overline{\{\eta'\}} = X,$$

then their closures are equal, so

$$\eta = \eta'.$$

Hence an integral scheme has a unique generic point.

4.7 Function Fields

Let X be an integral scheme with generic point η .

The stalk of the structure sheaf at η ,

$$\mathcal{O}_{X,\eta},$$

is called the *function field* of X . It is denoted

$$K(X).$$

Thus

$$K(X) := \mathcal{O}_{X,\eta}.$$

Why the stalk at the generic point is a field

Let

$$U = \operatorname{Spec} A$$

be a nonempty affine open subset of X .

Since X is integral, A is a domain. The generic point η corresponds to the zero prime ideal

$$(0) \subseteq A.$$

The stalk of the structure sheaf at this point is

$$\mathcal{O}_{X,\eta} = \mathcal{O}_{U,\eta} = A_{(0)}.$$

But localizing a domain at the zero prime gives its field of fractions:

$$A_{(0)} = \operatorname{Frac}(A).$$

Therefore

$$K(X) = \mathcal{O}_{X,\eta} = \operatorname{Frac}(A).$$

In particular, $K(X)$ is a field.

4.8 Function Field on Any Affine Open

Let

$$U = \operatorname{Spec} A$$

be any nonempty affine open subset of an integral scheme X .

The generic point η belongs to every nonempty open subset of X . Therefore

$$\eta \in U.$$

Inside $U = \operatorname{Spec} A$, the point η corresponds to the zero ideal

$$(0) \subseteq A.$$

Therefore

$$K(X) = \mathcal{O}_{X,\eta} = \mathcal{O}_{U,\eta} = A_{(0)} = \text{Frac}(A).$$

Thus, for every nonempty affine open subset

$$U = \text{Spec } A \subseteq X,$$

we have

$$\boxed{K(X) = \text{Frac}(A).}$$

4.9 Examples of Function Fields

For the affine line

$$\mathbb{A}_k^1 = \text{Spec } k[x],$$

we get

$$K(\mathbb{A}_k^1) = \text{Frac}(k[x]) = k(x).$$

For affine n -space

$$\mathbb{A}_k^n = \text{Spec } k[x_1, \dots, x_n],$$

we get

$$K(\mathbb{A}_k^n) = \text{Frac}(k[x_1, \dots, x_n]) = k(x_1, \dots, x_n).$$

For an irreducible affine variety

$$X = \text{Spec } k[x_1, \dots, x_n]/\mathfrak{p},$$

where $\mathfrak{p}$ is prime, we have

$$K(X) = \text{Frac}(k[x_1, \dots, x_n]/\mathfrak{p}).$$

This is the field of rational functions on X .

4.10 Summary of the Main Ideas

Concept	Affine algebraic meaning	Geometric meaning
Reduced	A has no nilpotents	No infinitesimal thickening
Irreducible	$\sqrt{(0)}$ is prime	One geometric component
Integral	A is a domain	Reduced and irreducible
Locally finite type	A locally finitely generated over B	Locally finitely many parameters
Finite type	Finite affine cover by finitely generated algebras	Locally finite type plus quasi-compact
Generic point	$(0) \in \text{Spec } A$ for a domain A	Dense point of an integral scheme
Function field	$\text{Frac}(A)$	Field of rational functions

4.11 Conceptual Picture

For an integral affine scheme

$$X = \operatorname{Spec} A,$$

where A is a domain, the points of X are prime ideals of A . The generic point is

$$(0) \in \operatorname{Spec} A.$$

Its closure is

$$\overline{\{(0)\}} = V((0)) = \operatorname{Spec} A.$$

So (0) lies “above” all other points in the specialization order. It represents the whole irreducible space.

The function field is obtained by looking at functions near this generic point:

$$K(X) = \mathcal{O}_{X,\eta}.$$

On an affine open

$$U = \operatorname{Spec} A,$$

this becomes

$$K(X) = A_{(0)} = \operatorname{Frac}(A).$$

Thus the slogan is:

On an integral scheme, rational functions are regular functions near the generic point.

5 Examples: Locally Finite Type, Finite Type, Generic Points, and Function Fields

This section gives concrete examples illustrating the theory from Problems 5–7.

5.1 Examples of Locally Finite Type Morphisms

Example 1. Affine space over a field

Let

$$X = \mathbb{A}_k^n = \operatorname{Spec} k[x_1, \dots, x_n]$$

and let

$$Y = \operatorname{Spec} k.$$

The natural morphism

$$f : \mathbb{A}_k^n \rightarrow \operatorname{Spec} k$$

corresponds to the ring map

$$k \rightarrow k[x_1, \dots, x_n].$$

The algebra

$$k[x_1, \dots, x_n]$$

is generated over k by finitely many elements:

$$x_1, \dots, x_n.$$

Therefore

$$k[x_1, \dots, x_n]$$

is a finitely generated k -algebra.

Hence

$$\mathbb{A}_k^n \rightarrow \operatorname{Spec} k \text{ is of finite type, and therefore locally of finite type.}$$

Example 2. A hypersurface over a field

Let

$$X = \operatorname{Spec} k[x, y]/(y^2 - x^3)$$

and let

$$Y = \operatorname{Spec} k.$$

The morphism

$$X \rightarrow Y$$

corresponds to the ring map

$$k \rightarrow k[x, y]/(y^2 - x^3).$$

The algebra

$$k[x, y]/(y^2 - x^3)$$

is generated over k by the images of x and y . Therefore it is a finitely generated k -algebra.

Hence

$$\operatorname{Spec} k[x, y]/(y^2 - x^3) \rightarrow \operatorname{Spec} k \text{ is of finite type.}$$

Geometrically, this is the affine cusp curve

$$y^2 = x^3.$$

Example 3. A distinguished open is locally of finite type

Let

$$X = \operatorname{Spec} A$$

and let

$$D(f) \subseteq X$$

be a distinguished open subset.

Then

$$D(f) \cong \operatorname{Spec} A_f.$$

The inclusion

$$D(f) \hookrightarrow X$$

corresponds to the localization map

$$A \rightarrow A_f.$$

But

$$A_f \cong A[t]/(ft - 1).$$

Therefore A_f is a finitely generated A -algebra, generated by t . Hence

$$D(f) \rightarrow \operatorname{Spec} A$$

is of finite type.

Thus

Distinguished open immersions are of finite type.

More generally, open immersions are locally of finite type.

5.2 Examples Showing the Difference Between Locally Finite Type and Finite Type

Example 4. Locally finite type but not finite type

Let

$$X = \coprod_{n=1}^{\infty} \operatorname{Spec} k$$

be the countably infinite disjoint union of copies of $\operatorname{Spec} k$.

Consider the natural morphism

$$f : X \rightarrow \operatorname{Spec} k.$$

Each component

$$\operatorname{Spec} k \rightarrow \operatorname{Spec} k$$

is of finite type. Therefore X is covered by affine opens over which the morphism is of finite type.

Hence

$$f : X \rightarrow \operatorname{Spec} k$$

is locally of finite type.

However, X is not quasi-compact. Indeed, the open cover

$$X = \bigcup_{n=1}^{\infty} \operatorname{Spec} k$$

has no finite subcover.

Since a finite type morphism is locally finite type and quasi-compact, f cannot be of finite type.

Therefore

$$\coprod_{n=1}^{\infty} \operatorname{Spec} k \rightarrow \operatorname{Spec} k \text{ is locally of finite type but not of finite type.}$$

Example 5. Not locally of finite type

Let

$$X = \operatorname{Spec} k[x_1, x_2, x_3, \dots]$$

and let

$$Y = \operatorname{Spec} k.$$

The morphism

$$X \rightarrow Y$$

corresponds to the ring map

$$k \rightarrow k[x_1, x_2, x_3, \dots].$$

The algebra

$$k[x_1, x_2, x_3, \dots]$$

is not finitely generated over k , because it requires infinitely many variables.

Moreover, it is not locally of finite type. The generic point corresponds to the zero ideal

$$(0) \subset k[x_1, x_2, x_3, \dots].$$

Every distinguished open neighbourhood of this generic point has the form

$$D(g) = \operatorname{Spec} k[x_1, x_2, x_3, \dots]_g.$$

The element g involves only finitely many variables. After localizing at g , infinitely many variables still remain. Hence

$$k[x_1, x_2, x_3, \dots]_g$$

is still not finitely generated as a k -algebra.

Therefore

$$\operatorname{Spec} k[x_1, x_2, x_3, \dots] \rightarrow \operatorname{Spec} k \text{ is not locally of finite type.}$$

5.3 Examples of Reduced, Irreducible, and Integral Schemes

Example 6. Reduced but not irreducible

Let

$$X = \operatorname{Spec} k[x, y]/(xy).$$

The ring

$$k[x, y]/(xy)$$

has no nilpotent elements if k is a field, so it is reduced.

However, it is not an integral domain, because

$$x \cdot y = 0$$

while neither x nor y is zero.

Geometrically,

$$xy = 0$$

is the union of the two coordinate axes.

Thus X is reduced but reducible.

Therefore

$\text{Spec } k[x, y]/(xy)$ is reduced but not irreducible, hence not integral.

Example 7. Irreducible but not reduced

Let

$$X = \text{Spec } k[\varepsilon]/(\varepsilon^2).$$

The topological space of X has only one point, because the only prime ideal is (ε) .

Therefore X is irreducible as a topological space.

However,

$$\varepsilon \neq 0, \quad \varepsilon^2 = 0.$$

So X is not reduced.

Therefore

$\text{Spec } k[\varepsilon]/(\varepsilon^2)$ is irreducible but not reduced, hence not integral.

Example 8. Integral affine line

Let

$$X = \mathbb{A}_k^1 = \text{Spec } k[x].$$

The ring

$$k[x]$$

is an integral domain. Therefore

$$\mathbb{A}_k^1$$

is integral.

The generic point is the point corresponding to the zero ideal:

$$\eta = (0) \in \text{Spec } k[x].$$

Its closure is

$$\overline{\{(0)\}} = V((0)) = \text{Spec } k[x].$$

Hence

$(0) \in \text{Spec } k[x]$ is the generic point of $\mathbb{A}_k^1$.

5.4 Examples of Generic Points

Example 9. Generic point of affine space

Let

$$X = \mathbb{A}_k^n = \operatorname{Spec} k[x_1, \dots, x_n].$$

Since

$$k[x_1, \dots, x_n]$$

is an integral domain, X is integral.

The generic point is

$$\eta = (0).$$

Indeed,

$$\overline{\{\eta\}} = V((0)) = \operatorname{Spec} k[x_1, \dots, x_n] = X.$$

Thus

$$\boxed{\eta = (0) \text{ is the generic point of } \mathbb{A}_k^n.}$$

Example 10. Generic point of an irreducible affine variety

Let

$$X = \operatorname{Spec} k[x_1, \dots, x_n]/\mathfrak{p},$$

where $\mathfrak{p}$ is a prime ideal.

Set

$$A = k[x_1, \dots, x_n]/\mathfrak{p}.$$

Since $\mathfrak{p}$ is prime, A is an integral domain. Hence X is integral.

The generic point of X is the zero ideal in A :

$$(0) \subseteq A.$$

Equivalently, as a prime ideal of $k[x_1, \dots, x_n]$, this point corresponds to

$$\mathfrak{p}.$$

Thus

$$\boxed{\text{The generic point of } \operatorname{Spec} k[x_1, \dots, x_n]/\mathfrak{p} \text{ corresponds to } \mathfrak{p}.}$$

Example 11. Generic points of a reducible scheme

Let

$$X = \operatorname{Spec} k[x, y]/(xy).$$

This scheme is reducible. It has two irreducible components:

$$V(x) \quad \text{and} \quad V(y).$$

The prime ideals

$$(x) \quad \text{and} \quad (y)$$

correspond to the generic points of these two components.

Thus X does not have one generic point whose closure is all of X . Instead, it has one generic point for each irreducible component.

Therefore

$\text{Spec } k[x, y]/(xy) \text{ has two component-generic points: } (x) \text{ and } (y).$

5.5 Examples of Function Fields

Example 12. Function field of the affine line

Let

$$X = \mathbb{A}_k^1 = \text{Spec } k[x].$$

The generic point is

$$\eta = (0).$$

The function field is

$$K(X) = \mathcal{O}_{X, \eta}.$$

Since

$$\mathcal{O}_{X, \eta} = k[x]_{(0)},$$

and localizing $k[x]$ at (0) gives its field of fractions, we obtain

$$K(\mathbb{A}_k^1) = \text{Frac}(k[x]) = k(x).$$

Thus

$K(\mathbb{A}_k^1) = k(x).$

Example 13. Function field of affine n -space

Let

$$X = \mathbb{A}_k^n = \text{Spec } k[x_1, \dots, x_n].$$

Since

$$k[x_1, \dots, x_n]$$

is a domain, X is integral.

Therefore

$$K(X) = \text{Frac}(k[x_1, \dots, x_n]).$$

Hence

$K(\mathbb{A}_k^n) = k(x_1, \dots, x_n).$

Example 14. Function field of an affine curve

Let

$$X = \operatorname{Spec} k[x, y]/(y^2 - x^3).$$

Assume that $y^2 - x^3$ is irreducible over k . Then

$$A = k[x, y]/(y^2 - x^3)$$

is an integral domain.

Therefore

$$X$$

is integral, and its function field is

$$K(X) = \operatorname{Frac}(A).$$

So

$$K(X) = \operatorname{Frac}\left(k[x, y]/(y^2 - x^3)\right).$$

Equivalently, this is the field generated by x and y subject to the relation

$$y^2 = x^3.$$

Thus

$$\boxed{K(X) = \operatorname{Frac}\left(k[x, y]/(y^2 - x^3)\right).}$$

Example 15. Function field of projective line

Let

$$X = \mathbb{P}_k^1.$$

Although $\mathbb{P}_k^1$ is not affine, it is integral.

Take the standard affine open

$$U_0 = \{X_0 \neq 0\} \cong \operatorname{Spec} k[t].$$

Since U_0 is a nonempty affine open subset of $\mathbb{P}_k^1$, the function field of $\mathbb{P}_k^1$ is

$$K(\mathbb{P}_k^1) = \operatorname{Frac}(k[t]).$$

Therefore

$$\boxed{K(\mathbb{P}_k^1) = k(t).}$$

This shows that the function field of a non-affine integral scheme can still be computed on any nonempty affine open.

5.6 Same Function Field on Different Affine Opens

Let

$$X = \mathbb{A}_k^1 = \operatorname{Spec} k[x].$$

The function field is

$$K(X) = k(x).$$

Now consider the distinguished open subset

$$D(x) \subseteq X.$$

We have

$$D(x) = \operatorname{Spec} k[x]_x.$$

Its coordinate ring is

$$k[x]_x.$$

The field of fractions of $k[x]_x$ is still

$$k(x).$$

Therefore

$$\operatorname{Frac}(k[x]) = \operatorname{Frac}(k[x]_x) = k(x).$$

Thus

$K(X) \text{ can be computed using } \operatorname{Spec} k[x] \text{ or using } D(x).$

This illustrates the theorem that for an integral scheme,

$$K(X) = \operatorname{Frac}(A)$$

for every nonempty affine open subset

$$U = \operatorname{Spec} A \subseteq X.$$

5.7 Example Where the Function Field Is Not Defined in This Sense

Let

$$X = \operatorname{Spec} k[x, y]/(xy).$$

This scheme is not integral, because

$$x \cdot y = 0$$

but neither x nor y is zero.

Since X is reducible, it does not have a unique generic point. It has two irreducible components:

$$V(x) \quad \text{and} \quad V(y).$$

Each component has its own function field:

$$K(V(x)) \cong k(y),$$

and

$$K(V(y)) \cong k(x).$$

But X itself is not integral, so there is no single function field $K(X)$ in the sense defined for integral schemes.

Therefore

$\text{Spec } k[x, y]/(xy)$ has component function fields, but no single function field as an integral scheme.

5.8 Summary of Examples

Example	Object	Conclusion
$\mathbb{A}_k^n$	$\text{Spec } k[x_1, \dots, x_n] \rightarrow \text{Spec } k$	Finite type
Infinite disjoint union	$\coprod_{n=1}^{\infty} \text{Spec } k \rightarrow \text{Spec } k$	Locally finite type, not finite type
Infinite variables	$\text{Spec } k[x_1, x_2, \dots] \rightarrow \text{Spec } k$	Not locally finite type
Dual numbers	$\text{Spec } k[\varepsilon]/(\varepsilon^2)$	Irreducible but not reduced
Two axes	$\text{Spec } k[x, y]/(xy)$	Reduced but reducible
Affine line	$\text{Spec } k[x]$	$K(X) = k(x)$
Affine n -space	$\text{Spec } k[x_1, \dots, x_n]$	$K(X) = k(x_1, \dots, x_n)$
Projective line	$\mathbb{P}_k^1$	$K(X) = k(t)$

6 Charts, Diagrams, and Tables: Locally Finite Type, Generic Points, and Function Fields


This section summarizes the material visually. The diagrams are designed to show the logical structure of the proofs and the relations between the main concepts.

6.1 Chart: Local Properties of Schemes

Property	Affine condition on A	Geometric meaning	Typical failure
Reduced	A has no nonzero nilpotents	No infinitesimal thickening	$k[\varepsilon]/(\varepsilon^2)$
Irreducible	$\sqrt{(0)}$ is prime	One topological component	$k \times k$
Integral	A is a domain	Reduced and irreducible	$k[x, y]/(xy)$
Noetherian	A is Noetherian	Finiteness of ideals locally	$k[x_1, x_2, x_3, \dots]$
Affine	$X \cong \text{Spec } A$	Globally described by one ring	$\mathbb{P}_k^1$

6.2 Diagram: How Locally Finite Type Is Checked

The definition starts with one affine cover of Y , but the theorem says the condition automatically holds over every affine open subset of Y .



6.3 Table: Locally Finite Type versus Finite Type

Property	Meaning	Example
Locally of finite type	Locally on source and target, coordinate rings are finitely generated	$\coprod_{n=1}^{\infty} \operatorname{Spec} k \rightarrow \operatorname{Spec} k$
Finite type	Locally of finite type + quasi-compactness	$\mathbb{A}_k^n \rightarrow \operatorname{Spec} k$
Not locally of finite type	Even locally, infinitely many algebra generators are needed	$\operatorname{Spec} k[x_1, x_2, x_3, \dots] \rightarrow \operatorname{Spec} k$

6.4 Conceptual Formula: Finite Type

$$\boxed{\text{finite type} = \text{locally finite type} + \text{quasi-compactness}}$$

More explicitly,

$$f : X \rightarrow Y \text{ finite type} \iff f \text{ locally of finite type and quasi-compact.}$$

6.5 Diagram: The Difference Between Locally Finite Type and Finite Type

Locally finite type	Finite type
Each small affine piece is controlled by finitely many generators	Only finitely many such affine pieces are needed over affine opens
$\coprod_{n=1}^{\infty} \operatorname{Spec} k \rightarrow \operatorname{Spec} k$	$\mathbb{A}_k^n \rightarrow \operatorname{Spec} k$
Locally finite type Not quasi-compact	Finite type Quasi-compact

6.6 Table: Standard Counterexamples

Failed property	Counterexample	Reason
Reduced	$\operatorname{Spec} k[\varepsilon]/(\varepsilon^2)$	$\varepsilon \neq 0, \varepsilon^2 = 0$
Irreducible	$\operatorname{Spec}(k \times k)$	Two disconnected points
Integral	$\operatorname{Spec} k[x, y]/(xy)$	$x \cdot y = 0$
Affine	$\mathbb{P}_k^1$	$\Gamma(\mathbb{P}_k^1, \mathcal{O}) = k$ but $\mathbb{P}_k^1 \neq \operatorname{Spec} k$
Noetherian	$\operatorname{Spec} k[x_1, x_2, x_3, \dots]$	$(x_1) \subsetneq (x_1, x_2) \subsetneq (x_1, x_2, x_3) \subsetneq \dots$
Locally finite type	$\operatorname{Spec} k[x_1, x_2, x_3, \dots] \rightarrow \operatorname{Spec} k$	Infinitely many generators remain locally
Finite type	$\coprod_{n=1}^{\infty} \operatorname{Spec} k \rightarrow \operatorname{Spec} k$	Not quasi-compact

6.7 Diagram: Generic Point of an Integral Affine Scheme

Let

$$X = \operatorname{Spec} A$$


where A is an integral domain. Then the generic point is

$$\eta = (0).$$

The closure of (0) is all of X :

$$\overline{\{(0)\}} = V((0)) = \operatorname{Spec} A.$$

So the picture is:



Thus the generic point is the most general point of the scheme.

6.8 Example Diagram: Generic Point of $\mathbb{A}_k^1$

For

$$\mathbb{A}_k^1 = \operatorname{Spec} k[x],$$

the points are prime ideals of $k[x]$.


The generic point is

$$(0).$$

Closed points correspond to maximal ideals such as

$$(x - a)$$

for $a \in k$, assuming k is algebraically closed.



The closure relation is

$$\overline{\{(0)\}} = \operatorname{Spec} k[x],$$

while

$$\overline{\{(x-a)\}} = \{(x-a)\}.$$

6.9 Diagram: Reducible Scheme with Two Generic Points

Consider

$$X = \operatorname{Spec} k[x, y]/(xy).$$

This is the union of two coordinate axes:

$$xy = 0.$$

The two irreducible components are

$$V(x) \quad \text{and} \quad V(y).$$

Their generic points are

$$(x) \quad \text{and} \quad (y).$$

$$\begin{array}{ccc} & \operatorname{Spec} k[x, y]/(xy) & \\ \swarrow & & \searrow \\ V(x) & & V(y) \\ \text{generic point } (x) & & \text{generic point } (y) \end{array}$$

So a reducible scheme does not have one generic point for the whole space. Instead, each irreducible component has its own generic point.

6.10 Chart: Function Field of an Integral Scheme

Let X be integral with generic point η . Then

$$K(X) = \mathcal{O}_{X, \eta}.$$

If

$$U = \operatorname{Spec} A \subseteq X$$

is any nonempty affine open subset, then

$$K(X) = \operatorname{Frac}(A).$$

The computation is:

$$\begin{aligned}
 U &= \operatorname{Spec} A \\
 &\Downarrow \\
 \eta &\leftrightarrow (0) \subseteq A \\
 &\Downarrow \\
 \mathcal{O}_{X,\eta} &= \mathcal{O}_{U,\eta} \\
 &\Downarrow \\
 \mathcal{O}_{U,\eta} &= A_{(0)} \\
 &\Downarrow \\
 A_{(0)} &= \operatorname{Frac}(A) \\
 &\Downarrow \\
 \boxed{K(X) &= \operatorname{Frac}(A)}
 \end{aligned}$$

6.11 Table: Function Fields of Standard Schemes

Scheme	Affine coordinate ring	Function field
$\mathbb{A}_k^1$	$k[x]$	$k(x)$
$\mathbb{A}_k^n$	$k[x_1, \dots, x_n]$	$k(x_1, \dots, x_n)$
$\operatorname{Spec} k[x, y]/(y^2 - x^3)$	$k[x, y]/(y^2 - x^3)$	$\operatorname{Frac}(k[x, y]/(y^2 - x^3))$
$\mathbb{P}_k^1$	computed on $U_0 \cong \operatorname{Spec} k[t]$	$k(t)$
$\operatorname{Spec} k[x_1, \dots, x_n]/\mathfrak{p}$	$k[x_1, \dots, x_n]/\mathfrak{p}$	$\operatorname{Frac}(k[x_1, \dots, x_n]/\mathfrak{p})$

6.12 Diagram: Computing $K(X)$ on Different Affine Opens

Let

$$X = \mathbb{A}_k^1 = \operatorname{Spec} k[x].$$

Take the distinguished open

$$D(x) = \operatorname{Spec} k[x]_x.$$

Then

$$\operatorname{Frac}(k[x]) = \operatorname{Frac}(k[x]_x) = k(x).$$

Thus both affine opens compute the same function field:

$$\begin{array}{ccc}
\mathrm{Spec} k[x] & \supseteq & D(x) = \mathrm{Spec} k[x]_x \\
\Downarrow & & \Downarrow \\
\mathrm{Frac}(k[x]) & = & \mathrm{Frac}(k[x]_x) \\
\Downarrow & & \Downarrow \\
k(x) & = & k(x)
\end{array}$$

6.13 Table: Integral versus Reducible Case

Scheme	Generic points	Function field
$\mathrm{Spec} k[x]$	One generic point (0)	$k(x)$
$\mathrm{Spec} k[x_1, \dots, x_n]$	One generic point (0)	$k(x_1, \dots, x_n)$
$\mathrm{Spec} k[x, y]/(xy)$	Two component-generic points $(x), (y)$	No single $K(X)$ as an integral scheme Component fields: $k(x), k(y)$
$\mathrm{Spec} k[\varepsilon]/(\varepsilon^2)$	One topological point	Not integral, so no $K(X)$ in this sense

6.14 Final Summary Diagram

$$\begin{array}{ccccc}
\text{Integral scheme } X & \implies & \text{unique generic point } \eta & \implies & K(X) = \mathcal{O}_{X,\eta} \\
\Downarrow & & \Downarrow & & \Downarrow \\
\text{Affine open } U = \mathrm{Spec} A & \implies & \eta = (0) \in \mathrm{Spec} A & \implies & K(X) = A_{(0)} = \mathrm{Frac}(A)
\end{array}$$

So the key slogan is:

On an integral scheme, rational functions are regular functions near the generic point.