

Theory of Commutative Algebra 6 Theory Notes

July 29, 2026

This file is a professionally reorganized and cleaned version.

Contents

1	Examples	6
1.1	Examples and intuition for $D(A)$	6
1.1.1	1. Domain case: $A = k[t]$	6
1.1.2	2. Two crossing components: $A = k[x, y]/(xy)$	6
1.1.3	3. Nilpotent example: $A = k[t]/(t^2)$	7
1.1.4	4. Localization removes primes that meet S	7
1.1.5	5. Embedded associated prime: $A = k[x, y]/(x^2, xy)$	8
1.1.6	6. Product ring example: $A = k \times k$	9
1.1.7	7. Many zero divisors, one prime	9
1.1.8	8. Relation with $\text{Ass}(A)$	10
1.1.9	9. Relation with zero divisors	10
1.1.10	10. How to compute $D(A)$ in practice	10
1.1.11	11. Final summary	10
1.2	Concrete polynomial and ring examples for $D(A)$	11
1.2.1	1. Arithmetic example: $A = \mathbb{Z}/6\mathbb{Z}$	11
1.2.2	2. One-variable polynomial example: $A = k[x]/(x(x-1))$	12
1.2.3	3. Two-variable geometric example: $A = k[x, y]/(xy)$	12
1.2.4	4. Nilpotent polynomial example: $A = k[t]/(t^2)$	13
1.2.5	5. Embedded-prime example: $A = k[x, y]/(x^2, xy)$	14
1.2.6	6. Concrete localization example	14
1.2.7	7. Practical computation rule	15
1.2.8	8. Final summary	15
1.3	Classes in quotient rings, minimal primes, and associated primes	15
1.3.1	1. What do $\bar{x}, \bar{y}$ mean?	15
1.3.2	2. Concrete examples of classes	16
1.3.3	3. How to think about quotient classes	17
1.3.4	4. Minimal primes	17
1.3.5	5. Associated primes	18
1.3.6	6. Comparison	19

1.3.7	7. Summary table	19
1.3.8	8. Relation with $D(A)$	20
1.4	Concrete meaning of x , x^2 , $\mathbb{Z}/p\mathbb{Z}$, $\mathbb{Z}/n\mathbb{Z}$, quotient classes, minimal primes, and associated primes	20
1.4.1	1. What is x in a polynomial ring?	20
1.4.2	2. What is x^2 ?	20
1.4.3	3. Evaluation	20
1.4.4	4. The ring $\mathbb{Z}/n\mathbb{Z}$	21
1.4.5	5. The ring $\mathbb{Z}/p\mathbb{Z}$	21
1.4.6	6. Polynomial rings over $\mathbb{Z}/p\mathbb{Z}$	21
1.4.7	7. Quotient classes	22
1.4.8	8. Example: $k[x]/(x^2)$	22
1.4.9	9. Example: $k[x]/(x(x-1))$	22
1.4.10	10. Example: $k[x, y]/(xy)$	23
1.4.11	11. Example: $k[x, y]/(x^2, xy)$	23
1.4.12	12. Interpretation	23
1.5	Concrete meaning of x , x^2 , $\mathbb{Z}/n\mathbb{Z}$, $\mathbb{Z}/p\mathbb{Z}$, quotient classes, minimal primes, and associated primes	24
1.5.1	1. Polynomial rings and formal indeterminates	24
1.5.2	2. Evaluation of polynomials	24
1.5.3	3. The rings $\mathbb{Z}/n\mathbb{Z}$ and $\mathbb{Z}/p\mathbb{Z}$	24
1.5.4	4. Polynomial rings over $\mathbb{Z}/p\mathbb{Z}$	25
1.5.5	5. Quotient rings and classes	26
1.5.6	6. Example: $k[x]/(x^2)$	26
1.5.7	7. Example: $k[x]/(x(x-1))$	27
1.5.8	8. Example: $k[x, y]/(xy)$	27
1.5.9	9. Example: $k[x, y]/(x^2, xy)$	28
1.5.10	10. Minimal primes	28
1.5.11	11. Associated primes	29
1.5.12	12. Comparison of minimal primes and associated primes	30
1.5.13	13. Summary of the main examples	31
1.5.14	14. Final intuition	31
1.6	Concrete annihilators: $\mathbb{Z}/p\mathbb{Z}$, $\mathbb{Z}/n\mathbb{Z}$, polynomial rings, and quotient rings	31
1.6.1	1. Definition of annihilator	31

1.6.2	2. First trivial examples	32
1.6.3	3. Annihilators in a domain	32
1.6.4	4. Annihilators in the field $\mathbb{Z}/p\mathbb{Z}$	32
1.6.5	5. Annihilators in $\mathbb{Z}/n\mathbb{Z}$	33
1.6.6	6. Annihilators in polynomial rings $k[x]$	35
1.6.7	7. Annihilators in $k[x]/(x^2)$	35
1.6.8	8. Annihilators in $k[x]/(x(x-1))$	36
1.6.9	9. Annihilators in $k[x, y]/(xy)$	36
1.6.10	10. Annihilators in $k[x, y]/(x^2, xy)$	37
1.6.11	11. Annihilators and associated primes	38
1.6.12	12. Annihilators and minimal primes	39
1.6.13	13. Final summary	39
1.7	Annihilators, minimal primes, and associated primes for modules: concrete examples	39
1.7.1	1. Annihilators for modules: the basic definitions	39
1.7.2	2. Associated primes and minimal primes of a module	40
1.7.3	3. The cyclic module R/I	41
1.7.4	4. First examples over $\mathbb{Z}$	41
1.7.5	5. A vector-space example	43
1.7.6	6. Polynomial modules: $M = k[x]/(x^2)$	43
1.7.7	7. Polynomial modules: $M = k[x]/(x(x-1))$	44
1.7.8	8. Polynomial modules: $M = k[x, y]/(xy)$	45
1.7.9	9. An embedded-prime module: $M = k[x, y]/(x^2, xy)$	46
1.7.10	10. Direct sums of modules	47
1.7.11	11. A free polynomial module	47
1.7.12	12. Summary table	48
1.7.13	13. Final interpretation	48
1.8	Concrete residue classes in $\mathbb{Z}/n\mathbb{Z}$, $\mathbb{Z}/p\mathbb{Z}$, $k[x]/(x)$, $A[t]/(t)$, and $k[x, y]/(xy)$	48
1.8.1	1. General principle: what is a residue class in R/I ?	48
1.8.2	2. Residue classes in $\mathbb{Z}/n\mathbb{Z}$	49
1.8.3	3. Residue classes in $\mathbb{Z}/p\mathbb{Z}$	50
1.8.4	4. Residue classes in $k[x]/(x)$	50
1.8.5	5. Residue classes in $A[t]/(t)$	51
1.8.6	6. Residue classes in $k[x, y]/(x)$	52

1.8.7	7. Residue classes in $k[x, y]/(xy)$	52
1.8.8	8. What are $\bar{x}$ and $\bar{y}$ concretely in $k[x, y]/(xy)$?	55
1.8.9	9. Comparison of the main examples	55
1.8.10	10. Final summary	56

Chapter 1

Examples

1.1 Examples and intuition for $D(A)$

Recall that

$$D(A) = \{\mathfrak{p} \in \operatorname{Spec}(A) : \exists a \in A \text{ such that } \mathfrak{p} \text{ is minimal in } V(\operatorname{Ann}(a))\}.$$

Thus $D(A)$ is built from annihilators of individual elements. It records the prime ideals that appear as the smallest prime-level reasons why some element is killed by multiplication. Equivalently, it is a prime-theoretic description of the set of zero divisors.

A useful slogan is:

$$\text{zero divisors} = \bigcup_{\mathfrak{p} \in D(A)} \mathfrak{p}.$$

1.1.1 1. Domain case: $A = k[t]$

Let $A = k[t]$, where k is a field. Since A is an integral domain, the only zero divisor is 0. For every nonzero $a \in A$,

$$\operatorname{Ann}(a) = 0.$$

Hence the only prime that can arise is

$$D(A) = \{(0)\}.$$

Thus part (a) becomes

$$x \text{ is a zero divisor} \iff x \in (0),$$

which says exactly that only 0 is a zero divisor in a domain.

1.1.2 2. Two crossing components: $A = k[x, y]/(xy)$

Let

$$A = k[x, y]/(xy),$$

and write $\bar{x}, \bar{y}$ for the classes of x, y in A . Since

$$\bar{x}\bar{y} = 0,$$

both $\bar{x}$ and $\bar{y}$ are zero divisors.

Moreover,

$$\operatorname{Ann}(\bar{x}) = (\bar{y}), \quad \operatorname{Ann}(\bar{y}) = (\bar{x}).$$

Therefore

$$(\bar{x}), (\bar{y}) \in D(A).$$

In fact these are exactly the minimal primes of A , so

$$D(A) = \{(\bar{x}), (\bar{y})\}.$$

Hence the set of zero divisors is

$$(\bar{x}) \cup (\bar{y}).$$

Geometric interpretation. The scheme $\text{Spec}(A)$ is the union of the two coordinate axes crossing at the origin. The primes in $D(A)$ correspond to the irreducible components. An element is a zero divisor precisely when it vanishes on at least one whole component.

1.1.3 3. Nilpotent example: $A = k[t]/(t^2)$

Let

$$A = k[t]/(t^2),$$

and let $\bar{t}$ denote the class of t . Then

$$\bar{t}^2 = 0,$$

so $\bar{t}$ is nilpotent and hence a zero divisor.

Also

$$\text{Ann}(\bar{t}) = (\bar{t}).$$

Therefore

$$D(A) = \{(\bar{t})\}.$$

This example shows that $D(A)$ detects zero divisors coming from nilpotent structure, not only from reducible geometry.

Interpretation. Here $\text{Spec}(A)$ has only one point, but the ring is nonreduced. So $D(A)$ can detect infinitesimal thickening as well as decomposition into components.

1.1.4 4. Localization removes primes that meet S

Again let

$$A = k[x, y]/(xy), \quad S = \{1, \bar{x}, \bar{x}^2, \dots\}.$$

In $S^{-1}A$, the element $\bar{x}$ becomes invertible. Since

$$\bar{x}\bar{y} = 0,$$

it follows that

$$\bar{y} = 0$$

in $S^{-1}A$. Thus

$$S^{-1}A \cong k[\bar{x}, \bar{x}^{-1}],$$

which is a domain.

Hence

$$D(S^{-1}A) = \{(0)\}$$

inside $\text{Spec}(S^{-1}A)$.

Under the canonical identification

$$\text{Spec}(S^{-1}A) \cong \{\mathfrak{p} \in \text{Spec}(A) \mid \mathfrak{p} \cap S = \emptyset\},$$

this prime corresponds to $(\bar{y})$. Therefore, viewed as a subset of $\text{Spec}(A)$,

$$D(S^{-1}A) = \{(\bar{y})\}.$$

But

$$D(A) = \{(\bar{x}), (\bar{y})\},$$

and only $(\bar{y})$ is disjoint from S . Hence

$$D(S^{-1}A) = D(A) \cap \text{Spec}(S^{-1}A).$$

Interpretation. Localization keeps only the primes disjoint from S . Any prime in $D(A)$ that meets S disappears after localization.

1.1.5 5. Embedded associated prime: $A = k[x, y]/(x^2, xy)$

Let

$$R = k[x, y], \quad I = (x^2, xy), \quad A = R/I.$$

Since

$$I = x(x, y) = (x) \cap (x^2, y),$$

we obtain a primary decomposition of I . Therefore the associated primes of A are

$$\text{Ass}(A) = \{(x)/I, (x, y)/I\}.$$

Now let us see that both belong to $D(A)$.

First, $(x)/I$ is minimal over

$$\text{Ann}_A(1) = 0,$$

so

$$(x)/I \in D(A).$$

Second, the class $\bar{x} \in A$ satisfies

$$x^2 = 0, \quad xy = 0.$$

Hence

$$\text{Ann}_A(\bar{x}) = (\bar{x}, \bar{y}) = (x, y)/I,$$

so

$$(x, y)/I \in D(A).$$

Thus

$$D(A) = \{(x)/I, (x, y)/I\} = \text{Ass}(A).$$

Why this matters. The prime $(x, y)/I$ is an embedded prime, not a minimal prime. So $D(A)$ can detect hidden nonreduced structure inside a component.

1.1.6 6. Product ring example: $A = k \times k$

Let

$$A = k \times k.$$

Then

$$(1, 0) \cdot (0, 1) = 0,$$

so both $(1, 0)$ and $(0, 1)$ are zero divisors.

The prime ideals are

$$\mathfrak{p}_1 = 0 \times k, \quad \mathfrak{p}_2 = k \times 0.$$

Also

$$\text{Ann}(1, 0) = 0 \times k = \mathfrak{p}_1, \quad \text{Ann}(0, 1) = k \times 0 = \mathfrak{p}_2.$$

Therefore

$$D(A) = \{\mathfrak{p}_1, \mathfrak{p}_2\}.$$

An element $(a, b) \in k \times k$ is a zero divisor if and only if

$$a = 0 \quad \text{or} \quad b = 0.$$

So again

$$\{\text{zero divisors}\} = \mathfrak{p}_1 \cup \mathfrak{p}_2.$$

This is the most concrete algebraic model of a reducible space with two disjoint components.

1.1.7 7. Many zero divisors, one prime

Let

$$A = k[x, y]/(x^2, xy, y^2).$$

Then the maximal ideal

$$\mathfrak{m} = (\bar{x}, \bar{y})$$

satisfies

$$\mathfrak{m}^2 = 0.$$

Hence every element of $\mathfrak{m}$ is nilpotent and therefore a zero divisor.

However, $\text{Spec}(A)$ has only one prime ideal, namely $\mathfrak{m}$. Thus

$$D(A) = \{\mathfrak{m}\}.$$

This example shows that many different zero divisors may all be controlled by the same prime in $D(A)$.

1.1.8 8. Relation with $\text{Ass}(A)$

If the zero ideal of A has a primary decomposition, then part (c) says that

$$D(A) = \text{Ass}(A).$$

So in Noetherian rings with primary decomposition, $D(A)$ is exactly the set of associated primes of the A -module A .

In particular:

- in reduced rings such as $k[x, y]/(xy)$, $D(A)$ often consists of minimal primes;
- in nonreduced rings such as $k[x, y]/(x^2, xy)$, $D(A)$ may also contain embedded primes.

1.1.9 9. Relation with zero divisors

Part (a) gives the important identity

$$\{\text{zero divisors of } A\} = \bigcup_{\mathfrak{p} \in D(A)} \mathfrak{p}.$$

Thus $D(A)$ is a prime-theoretic description of the zero divisors. When primary decomposition exists, this becomes

$$\{\text{zero divisors of } A\} = \bigcup_{\mathfrak{p} \in \text{Ass}(A)} \mathfrak{p}.$$

1.1.10 10. How to compute $D(A)$ in practice

There are two standard methods.

Method 1: via annihilators of elements. Choose simple elements $a \in A$, compute $\text{Ann}(a)$, and find the minimal primes over $\text{Ann}(a)$. All primes obtained this way belong to $D(A)$.

Method 2: via primary decomposition. If A is Noetherian and 0 has a primary decomposition, compute that decomposition and take the radicals of its primary components. These radicals are exactly the primes in $D(A)$, because then

$$D(A) = \text{Ass}(A).$$

1.1.11 11. Final summary

The most important examples to remember are:

$$k[t], \quad k[x, y]/(xy), \quad k[t]/(t^2), \quad k[x, y]/(x^2, xy), \quad k \times k.$$

They illustrate, respectively:

- the domain case;
- reducible geometry with two components;
- nilpotent structure;
- embedded associated primes;
- and a completely explicit decomposed ring.

Together they show that $D(A)$ measures where zero divisors come from, how those primes behave under localization, and why in the Noetherian case $D(A)$ agrees with the set of associated primes.

1.2 Concrete polynomial and ring examples for $D(A)$

Recall that

$$D(A) = \{\mathfrak{p} \in \operatorname{Spec}(A) : \exists a \in A \text{ such that } \mathfrak{p} \text{ is minimal in } V(\operatorname{Ann}(a))\}.$$

Part (a) says that

$$\{\text{zero divisors of } A\} = \bigcup_{\mathfrak{p} \in D(A)} \mathfrak{p}.$$

We now present this on explicit rings and explicit polynomial quotients.

1.2.1 1. Arithmetic example: $A = \mathbb{Z}/6\mathbb{Z}$

The elements are

$$\bar{0}, \bar{1}, \bar{2}, \bar{3}, \bar{4}, \bar{5}.$$

The zero divisors are

$$\bar{0}, \bar{2}, \bar{3}, \bar{4},$$

because

$$\bar{2} \cdot \bar{3} = \bar{0}, \quad \bar{3} \cdot \bar{4} = \bar{0}.$$

Now compute annihilators:

$$\operatorname{Ann}(\bar{2}) = \{\bar{r} \in \mathbb{Z}/6\mathbb{Z} : \bar{r}\bar{2} = \bar{0}\} = \{\bar{0}, \bar{3}\} = (\bar{3}),$$

and

$$\operatorname{Ann}(\bar{3}) = \{\bar{0}, \bar{2}, \bar{4}\} = (\bar{2}).$$

The prime ideals of $\mathbb{Z}/6\mathbb{Z}$ are

$$(\bar{2}), \quad (\bar{3}),$$

so

$$D(A) = \{(\bar{2}), (\bar{3})\}.$$

Thus

$$\{\text{zero divisors}\} = (\bar{2}) \cup (\bar{3}).$$

1.2.2 2. One-variable polynomial example: $A = k[x]/(x(x-1))$

Let $\bar{x}$ denote the class of x in A . Then

$$\bar{x}(\bar{x} - 1) = 0,$$

so both $\bar{x}$ and $\bar{x} - 1$ are zero divisors.

We now compute their annihilators.

Since $(\bar{x} - 1)\bar{x} = 0$, every multiple of $\bar{x} - 1$ annihilates $\bar{x}$. In fact,

$$\text{Ann}(\bar{x}) = (\bar{x} - 1).$$

Similarly,

$$\text{Ann}(\bar{x} - 1) = (\bar{x}).$$

Hence

$$(\bar{x}), (\bar{x} - 1) \in D(A),$$

and in fact

$$D(A) = \{(\bar{x}), (\bar{x} - 1)\}.$$

So the zero divisors are precisely the elements lying in one of these two prime ideals.

Examples.

$\bar{x}$ is a zero divisor, $\bar{x} - 1$ is a zero divisor, $\bar{1}$ is not a zero divisor.

Geometric meaning. The equation

$$x(x-1) = 0$$

describes the union of the two points $x = 0$ and $x = 1$ on the affine line.

1.2.3 3. Two-variable geometric example: $A = k[x, y]/(xy)$

Let $\bar{x}, \bar{y}$ be the classes of x, y in A . Then

$$\bar{x}\bar{y} = 0,$$

so $\bar{x}$ and $\bar{y}$ are zero divisors.

To compute $\text{Ann}(\bar{x})$, we look for all $\bar{f} \in A$ such that

$$\bar{f}\bar{x} = 0.$$

This means

$$fx \in (xy)$$

in $k[x, y]$. Since fx must be divisible by xy , the polynomial f must be divisible by y . Therefore

$$\text{Ann}(\bar{x}) = (\bar{y}).$$

Similarly,

$$\text{Ann}(\bar{y}) = (\bar{x}).$$

Hence

$$D(A) = \{(\bar{x}), (\bar{y})\}.$$

So the zero divisors are exactly the elements in

$$(\bar{x}) \cup (\bar{y}).$$

Concrete classes. Take

$$a = \bar{x}, \quad b = \bar{y}, \quad c = \bar{x} + \bar{y}, \quad d = \bar{x} + 1.$$

Then:

- a is a zero divisor since $a\bar{y} = 0$;
- b is a zero divisor since $b\bar{x} = 0$;
- $c \notin (\bar{x}) \cup (\bar{y})$, so c is not a zero divisor;
- $d \notin (\bar{x}) \cup (\bar{y})$, so d is not a zero divisor.

Geometric meaning. The variety

$$V(xy) = V(x) \cup V(y)$$

is the union of the two coordinate axes. The primes in $D(A)$ correspond to those two components.

1.2.4 4. Nilpotent polynomial example: $A = k[t]/(t^2)$

Let $\bar{t}$ be the class of t in A . Then

$$\bar{t}^2 = 0,$$

so $\bar{t}$ is nilpotent and hence a zero divisor.

Now $\bar{f} \in \text{Ann}(\bar{t})$ if and only if

$$\bar{f}\bar{t} = 0,$$

that is, if and only if

$$ft \in (t^2).$$

This holds exactly when $f \in (t)$. Hence

$$\text{Ann}(\bar{t}) = (\bar{t}),$$

and therefore

$$D(A) = \{(\bar{t})\}.$$

Every element of A has the form

$$a + b\bar{t}, \quad a, b \in k.$$

If $a \neq 0$, then $a + b\bar{t}$ is a unit. If $a = 0$, then the element equals $b\bar{t}$, which is nilpotent. Thus the zero divisors are exactly the ideal

$$(\bar{t}).$$

1.2.5 5. Embedded-prime example: $A = k[x, y]/(x^2, xy)$

Let

$$A = k[x, y]/(x^2, xy),$$

and let $\bar{x}, \bar{y}$ be the classes of x, y . Then in A ,

$$\bar{x}^2 = 0, \quad \bar{x}\bar{y} = 0.$$

So $\bar{x}$ is annihilated by both $\bar{x}$ and $\bar{y}$, and therefore

$$\text{Ann}(\bar{x}) = (\bar{x}, \bar{y}).$$

Hence

$$(\bar{x}, \bar{y}) \in D(A).$$

On the other hand,

$$(x^2, xy) = x(x, y) = (x) \cap (x^2, y),$$

so the associated primes are

$$(\bar{x}), \quad (\bar{x}, \bar{y}).$$

Therefore

$$D(A) = \{(\bar{x}), (\bar{x}, \bar{y})\}.$$

Interpretation. The prime $(\bar{x})$ is the minimal prime, while $(\bar{x}, \bar{y})$ is an embedded prime. This shows that $D(A)$ can detect not only visible components but also hidden nilpotent structure inside a component.

1.2.6 6. Concrete localization example

Take again

$$A = k[x, y]/(xy),$$

and let

$$S = \{1, \bar{x}, \bar{x}^2, \bar{x}^3, \dots\}.$$

In the localization $S^{-1}A$, the element $\bar{x}$ becomes invertible. Since

$$\bar{x}\bar{y} = 0,$$

we may multiply by $\bar{x}^{-1}$ and obtain

$$\bar{y} = 0.$$

Hence

$$S^{-1}A \cong k[\bar{x}, \bar{x}^{-1}],$$

which is a domain.

Before localization,

$$D(A) = \{(\bar{x}), (\bar{y})\}.$$

After localization, the prime $(\bar{x})$ disappears because it meets S , while $(\bar{y})$ survives. Thus

$$D(S^{-1}A) = \{(\bar{y})\}$$

when viewed inside $\text{Spec}(A)$.

This is the concrete meaning of the formula

$$D(S^{-1}A) = D(A) \cap \text{Spec}(S^{-1}A).$$

1.2.7 7. Practical computation rule

For a quotient ring

$$A = k[x_1, \dots, x_n]/I,$$

one can compute $D(A)$ concretely by the following steps:

1. Find obvious zero divisors, that is, classes whose product is zero modulo I .
2. For a chosen element $a \in A$, compute $\text{Ann}(a)$ by solving

$$fa \in I.$$

3. Find the minimal primes over $\text{Ann}(a)$.
4. Collect all primes that arise.

If 0 has a primary decomposition, then one may instead compute $\text{Ass}(A)$, because in that case

$$D(A) = \text{Ass}(A).$$

1.2.8 8. Final summary

The most concrete examples to remember are:

$$\mathbb{Z}/6\mathbb{Z}, \quad k[x]/(x(x-1)), \quad k[x, y]/(xy), \quad k[t]/(t^2), \quad k[x, y]/(x^2, xy).$$

They illustrate:

- arithmetic zero divisors;
- decomposition into two points;
- decomposition into two axes;
- nilpotent zero divisors;
- and embedded associated primes.

1.3 Classes in quotient rings, minimal primes, and associated primes

1.3.1 1. What do $\bar{x}, \bar{y}$ mean?

Let

$$A = R/I$$

be a quotient ring. For any element $f \in R$, its image in A is denoted by

$$\bar{f} \quad \text{or sometimes} \quad [f].$$

Thus

$$\bar{f} = f + I.$$

So:

$$\bar{x} = x + I, \quad \bar{y} = y + I, \quad \overline{x + y} = x + y + I.$$

This is the residue class (or coset) of f modulo I .

Two elements $f, g \in R$ define the same class in A if and only if

$$\bar{f} = \bar{g} \iff f - g \in I.$$

1.3.2 2. Concrete examples of classes

Example: $A = k[x, y]/(xy)$

Here $I = (xy)$. Then:

$$\bar{x} = x + (xy), \quad \bar{y} = y + (xy).$$

Also

$$\overline{x + xy} = \bar{x}$$

because

$$(x + xy) - x = xy \in (xy).$$

Moreover,

$$\overline{xy} = 0, \quad \overline{x^2y} = 0,$$

since $xy, x^2y \in (xy)$.

However,

$$\bar{x} \neq 0, \quad \bar{y} \neq 0,$$

because $x, y \notin (xy)$.

Thus in this quotient ring,

$$\bar{x}\bar{y} = 0$$

while neither $\bar{x}$ nor $\bar{y}$ is zero.

Example: $A = k[t]/(t^2)$

Here

$$\bar{t} \neq 0$$

but

$$\bar{t}^2 = 0.$$

Every element has the form

$$a + b\bar{t}, \quad a, b \in k.$$

Example: $A = \mathbb{Z}/6\mathbb{Z}$

The class $\bar{2}$ means all integers congruent to 2 modulo 6:

$$\bar{2} = \{\dots, -10, -4, 2, 8, 14, \dots\}.$$

Also

$$\bar{2}\bar{3} = \bar{0}.$$

1.3.3 3. How to think about quotient classes

The quotient ring R/I is obtained from R by imposing all relations coming from I .

For example:

$k[x, y]/(xy)$ means that in the quotient we force $xy = 0$;

$k[x, y]/(x^2, xy)$ means that in the quotient we force $x^2 = 0$ and $xy = 0$;

$k[x]/(x(x-1))$ means that in the quotient we force $x(x-1) = 0$.

1.3.4 4. Minimal primes

A prime ideal $\mathfrak{p} \subset A$ is called a *minimal prime* if it is minimal among the prime ideals of A .

If

$$A = R/I,$$

then minimal primes of A correspond to prime ideals of R that are minimal over I .

We write

$$\text{Min}(A)$$

for the set of minimal primes.

Interpretation. Minimal primes describe the top-level irreducible components of the ring or affine scheme.

Example: $A = k[x]$

Since A is a domain, (0) is prime and

$$\text{Min}(A) = \{(0)\}.$$

Example: $A = k[x, y]/(xy)$

The ideal (xy) has two minimal primes:

$$(x), \quad (y).$$

Hence in the quotient,

$$\text{Min}(A) = \{(\bar{x}), (\bar{y})\}.$$

Example: $A = k[x, y]/(x^2)$

The only prime minimal over (x^2) is (x) . Hence

$$\text{Min}(A) = \{(\bar{x})\}.$$

Example: $A = k[x, y]/(x^2, xy)$

Since

$$\sqrt{(x^2, xy)} = (x),$$

the only minimal prime is

$$\text{Min}(A) = \{(\bar{x})\}.$$

1.3.5 5. Associated primes

A prime ideal $\mathfrak{p} \subset A$ is an *associated prime* of the A -module A if

$$\mathfrak{p} = \text{Ann}_A(a)$$

for some element $a \in A$.

We write

$$\text{Ass}(A)$$

for the set of associated primes of A .

Interpretation. Associated primes describe primes that arise as actual annihilators of elements. They detect zero-divisor structure and possible embedded components.

Example: $A = k[x, y]/(xy)$

We have

$$\text{Ann}(\bar{x}) = (\bar{y}), \quad \text{Ann}(\bar{y}) = (\bar{x}).$$

Therefore

$$\text{Ass}(A) = \{(\bar{x}), (\bar{y})\}.$$

Here

$$\text{Ass}(A) = \text{Min}(A).$$

Example: $A = k[t]/(t^2)$

We have

$$\text{Ann}(\bar{t}) = (\bar{t}).$$

Hence

$$\text{Ass}(A) = \{(\bar{t})\} = \text{Min}(A).$$

Example: $A = k[x, y]/(x^2, xy)$

First,

$$\text{Min}(A) = \{(\bar{x})\}.$$

Now in A ,

$$\bar{x}^2 = 0, \quad \bar{x}\bar{y} = 0.$$

Hence

$$(\bar{x}, \bar{y}) \subseteq \text{Ann}(\bar{x}).$$

Conversely, if $f\bar{x} = 0$, then

$$fx \in (x^2, xy) = x(x, y),$$

so $f \in (x, y)$. Therefore

$$\text{Ann}(\bar{x}) = (\bar{x}, \bar{y}).$$

Thus

$$(\bar{x}, \bar{y}) \in \text{Ass}(A).$$

Since $(\bar{x})$ is the minimal prime, it is also associated in this Noetherian case. Hence

$$\text{Ass}(A) = \{(\bar{x}), (\bar{x}, \bar{y})\}.$$

So

$$\text{Min}(A) = \{(\bar{x})\} \subsetneq \text{Ass}(A) = \{(\bar{x}), (\bar{x}, \bar{y})\}.$$

The prime $(\bar{x}, \bar{y})$ is an embedded associated prime.

1.3.6 6. Comparison

Minimal primes describe the irreducible components of the ring:

geometry at the top level.

Associated primes describe annihilators of actual elements:

zero-divisor and embedded structure.

Thus in general,

$$\text{Min}(A) \subseteq \text{Ass}(A),$$

and the inclusion may be strict.

1.3.7 7. Summary table

A	$\text{Min}(A)$	$\text{Ass}(A)$
$k[x]$	$\{(0)\}$	$\{(0)\}$
$k[x, y]/(xy)$	$\{(\bar{x}), (\bar{y})\}$	$\{(\bar{x}), (\bar{y})\}$
$k[t]/(t^2)$	$\{(\bar{t})\}$	$\{(\bar{t})\}$
$k[x, y]/(x^2, xy)$	$\{(\bar{x})\}$	$\{(\bar{x}), (\bar{x}, \bar{y})\}$

1.3.8 8. Relation with $D(A)$

In the problem, $D(A)$ is defined via primes minimal over annihilators of elements. In the Noetherian case with primary decomposition,

$$D(A) = \text{Ass}(A).$$

Moreover, the set of zero divisors is

$$\bigcup_{\mathfrak{p} \in D(A)} \mathfrak{p}.$$

So associated primes are the natural language for understanding zero divisors.

1.4 Concrete meaning of x , x^2 , $\mathbb{Z}/p\mathbb{Z}$, $\mathbb{Z}/n\mathbb{Z}$, quotient classes, minimal primes, and associated primes

1.4.1 1. What is x in a polynomial ring?

In a polynomial ring

$$R[x],$$

the symbol x is a *formal indeterminate*. It is not automatically a number. A polynomial is a finite expression

$$a_0 + a_1x + \cdots + a_nx^n \quad (a_i \in R).$$

Thus in $\mathbb{Z}[x]$,

$$2x^2 + 3x + 5$$

is a polynomial with integer coefficients. Here x is only a formal symbol.

1.4.2 2. What is x^2 ?

The symbol

$$x^2$$

means $x \cdot x$, the square of the indeterminate x . It is itself a polynomial. For example:

$$x = 0 + 1 \cdot x, \quad x^2 = 0 + 0 \cdot x + 1 \cdot x^2.$$

1.4.3 3. Evaluation

Once one chooses an element $a \in R$, one may evaluate:

$$f(x) \mapsto f(a).$$

For example, if

$$f(x) = 2x^2 + 3x + 5 \in \mathbb{Z}[x],$$

then

$$f(4) = 2 \cdot 4^2 + 3 \cdot 4 + 5 = 49.$$

Thus x is formal in the polynomial ring, but becomes an actual value under evaluation.

1.4.4 4. The ring $\mathbb{Z}/n\mathbb{Z}$

The ring

$$\mathbb{Z}/n\mathbb{Z}$$

consists of residue classes modulo n :

$$\bar{0}, \bar{1}, \dots, \overline{n-1}.$$

Two integers define the same class if and only if they differ by a multiple of n :

$$a \equiv b \pmod{n} \iff n \mid (a - b).$$

Example in $\mathbb{Z}/6\mathbb{Z}$:

$$\bar{2} = \bar{8} = \overline{-4}.$$

Arithmetic is done modulo n .

1.4.5 5. The ring $\mathbb{Z}/p\mathbb{Z}$

If p is prime, then

$$\mathbb{Z}/p\mathbb{Z}$$

is a field, often denoted

$$\mathbb{F}_p.$$

For example, in $\mathbb{Z}/5\mathbb{Z}$,

$$\bar{2} + \bar{4} = \bar{1}, \quad \bar{2}\bar{3} = \bar{1}.$$

If n is not prime, then $\mathbb{Z}/n\mathbb{Z}$ usually has zero divisors. For example, in $\mathbb{Z}/6\mathbb{Z}$,

$$\bar{2} \cdot \bar{3} = \bar{0}.$$

1.4.6 6. Polynomial rings over $\mathbb{Z}/p\mathbb{Z}$

The ring

$$(\mathbb{Z}/5\mathbb{Z})[x]$$

consists of polynomials in x with coefficients modulo 5. For example:

$$\bar{3}x^2 + \bar{4}x + \bar{1}.$$

Addition and multiplication are performed as usual, with coefficients reduced modulo 5.

1.4.7 7. Quotient classes

If

$$A = R/I,$$

then for $f \in R$, its class in A is

$$\bar{f} = f + I.$$

Thus:

$$\bar{x} = x + I, \quad \overline{x^2} = x^2 + I.$$

Two elements define the same class if and only if their difference lies in I :

$$\bar{f} = \bar{g} \iff f - g \in I.$$

1.4.8 8. Example: $k[x]/(x^2)$

Let

$$A = k[x]/(x^2).$$

Then:

$$\bar{x} \neq 0$$

because $x \notin (x^2)$, but

$$\bar{x}^2 = 0$$

because $x^2 \in (x^2)$.

Every element of A has the form

$$a + b\bar{x}, \quad a, b \in k.$$

1.4.9 9. Example: $k[x]/(x(x-1))$

Let

$$A = k[x]/(x(x-1)).$$

Then in A ,

$$\bar{x}(\bar{x} - 1) = 0.$$

So $\bar{x}$ and $\bar{x} - 1$ are zero divisors.

The minimal primes are

$$(\bar{x}), \quad (\bar{x} - 1).$$

Hence

$$\text{Min}(A) = \{(\bar{x}), (\bar{x} - 1)\}.$$

Also

$$\text{Ass}(A) = \{(\bar{x}), (\bar{x} - 1)\}.$$

1.4.10 10. Example: $k[x, y]/(xy)$

Let

$$A = k[x, y]/(xy).$$

Then

$$\bar{x}\bar{y} = 0,$$

but $\bar{x} \neq 0$ and $\bar{y} \neq 0$.

The minimal primes are

$$(\bar{x}), \quad (\bar{y}),$$

so

$$\text{Min}(A) = \{(\bar{x}), (\bar{y})\}.$$

Also

$$\text{Ann}(\bar{x}) = (\bar{y}), \quad \text{Ann}(\bar{y}) = (\bar{x}),$$

hence

$$\text{Ass}(A) = \{(\bar{x}), (\bar{y})\}.$$

1.4.11 11. Example: $k[x, y]/(x^2, xy)$

Let

$$A = k[x, y]/(x^2, xy).$$

Then

$$\bar{x}^2 = 0, \quad \bar{x}\bar{y} = 0.$$

The radical of (x^2, xy) is (x) , so

$$\text{Min}(A) = \{(\bar{x})\}.$$

Now

$$\text{Ann}(\bar{x}) = (\bar{x}, \bar{y}),$$

so

$$(\bar{x}, \bar{y}) \in \text{Ass}(A).$$

Thus

$$\text{Ass}(A) = \{(\bar{x}), (\bar{x}, \bar{y})\}.$$

Therefore

$$\text{Min}(A) \subsetneq \text{Ass}(A).$$

The prime $(\bar{x}, \bar{y})$ is an embedded associated prime.

1.4.12 12. Interpretation

Minimal primes describe the main irreducible components of the ring. Associated primes describe actual annihilators of elements, hence zero-divisor structure and possible embedded components.

Thus:

minimal primes = top geometry, associated primes = top geometry + hidden annihilation structure

1.5 Concrete meaning of x , x^2 , $\mathbb{Z}/n\mathbb{Z}$, $\mathbb{Z}/p\mathbb{Z}$, quotient classes, minimal primes, and associated primes

1.5.1 1. Polynomial rings and formal indeterminates

Let R be a ring. The polynomial ring

$$R[x]$$

consists of all finite expressions

$$a_0 + a_1x + \cdots + a_nx^n, \quad a_i \in R.$$

Here x is a *formal indeterminate*. It is not automatically a number. It is simply a symbol used to build polynomials.

Thus:

$$x = 0 + 1 \cdot x, \quad x^2 = 0 + 0 \cdot x + 1 \cdot x^2.$$

So x^2 means the product $x \cdot x$, and it is itself a polynomial.

For example, in $\mathbb{Z}[x]$,

$$2x^2 + 3x + 5$$

is a polynomial with integer coefficients. At this stage, x is not evaluated at any integer.

1.5.2 2. Evaluation of polynomials

If $f(x) \in R[x]$ and $a \in R$, then we may evaluate f at a :

$$f(x) \mapsto f(a).$$

For example, if

$$f(x) = 2x^2 + 3x + 5 \in \mathbb{Z}[x],$$

then

$$f(4) = 2 \cdot 4^2 + 3 \cdot 4 + 5 = 32 + 12 + 5 = 49.$$

Thus there is a fundamental distinction:

- in $R[x]$, x is formal;
- after evaluation, x is replaced by an actual element of a ring.

1.5.3 3. The rings $\mathbb{Z}/n\mathbb{Z}$ and $\mathbb{Z}/p\mathbb{Z}$

The ring

$$\mathbb{Z}/n\mathbb{Z}$$

is the ring of integers modulo n . Its elements are residue classes

$$\bar{0}, \bar{1}, \bar{2}, \dots, \overline{n-1},$$

where

$$\bar{a} = \bar{b} \iff a \equiv b \pmod{n} \iff n \mid (a - b).$$

Example: $\mathbb{Z}/6\mathbb{Z}$

The classes are

$$\bar{0}, \bar{1}, \bar{2}, \bar{3}, \bar{4}, \bar{5}.$$

We have

$$\bar{2} = \bar{8} = \overline{-4},$$

and arithmetic is done modulo 6:

$$\bar{4} + \bar{5} = \bar{9} = \bar{3}, \quad \bar{2} \cdot \bar{4} = \bar{8} = \bar{2}.$$

Also

$$\bar{2} \cdot \bar{3} = \bar{0},$$

so $\bar{2}$ and $\bar{3}$ are zero divisors.

Example: $\mathbb{Z}/5\mathbb{Z}$

Since 5 is prime, the ring

$$\mathbb{Z}/5\mathbb{Z}$$

is a field, often denoted $\mathbb{F}_5$. Its elements are

$$\bar{0}, \bar{1}, \bar{2}, \bar{3}, \bar{4}.$$

Examples:

$$\bar{2} + \bar{4} = \bar{6} = \bar{1}, \quad \bar{2} \cdot \bar{3} = \bar{6} = \bar{1}.$$

Every nonzero element has a multiplicative inverse.

In general:

- $\mathbb{Z}/p\mathbb{Z}$ is a field if p is prime;
- $\mathbb{Z}/n\mathbb{Z}$ usually has zero divisors if n is composite.

1.5.4 4. Polynomial rings over $\mathbb{Z}/p\mathbb{Z}$

One may form polynomial rings over finite fields:

$$(\mathbb{Z}/p\mathbb{Z})[x].$$

For example, in $(\mathbb{Z}/5\mathbb{Z})[x]$,

$$f(x) = \bar{3}x^2 + \bar{4}x + \bar{1}$$

is a polynomial.

Addition is coefficientwise modulo 5. For example,

$$(\bar{3}x^2 + \bar{4}x + \bar{1}) + (\bar{4}x^2 + \bar{2}x + \bar{3}) = \bar{2}x^2 + \bar{x} + \bar{4},$$

because

$$\bar{3} + \bar{4} = \bar{2}, \quad \bar{4} + \bar{2} = \bar{1}, \quad \bar{1} + \bar{3} = \bar{4}.$$

1.5.5 5. Quotient rings and classes

Let R be a ring and $I \subseteq R$ an ideal. The quotient ring

$$R/I$$

consists of residue classes modulo I . For $f \in R$, its class is written

$$\bar{f} = f + I.$$

Thus:

$$\bar{x} = x + I, \quad \overline{x^2} = x^2 + I.$$

Two elements $f, g \in R$ define the same class if and only if

$$\bar{f} = \bar{g} \iff f - g \in I.$$

So $\bar{f}$ is not the original polynomial f , but its equivalence class modulo I .

1.5.6 6. Example: $k[x]/(x^2)$

Let

$$A = k[x]/(x^2),$$

where k is a field.

In this quotient ring, the class $\bar{x}$ satisfies

$$\bar{x}^2 = 0,$$

because $x^2 \in (x^2)$. However,

$$\bar{x} \neq 0,$$

because $x \notin (x^2)$.

Every element of A has the form

$$a + b\bar{x}, \quad a, b \in k.$$

Indeed, every polynomial

$$f(x) = a_0 + a_1x + a_2x^2 + \cdots + a_nx^n$$

has class

$$\bar{f} = a_0 + a_1\bar{x},$$

since all terms of degree at least 2 vanish in the quotient.

Examples.

$$\overline{x + x^2} = \bar{x}, \quad \overline{1 + x^3} = 1, \quad (\bar{x})^2 = 0.$$

This is a standard example of a nonreduced ring: $\bar{x} \neq 0$, but $\bar{x}$ is nilpotent.

1.5.7 7. Example: $k[x]/(x(x-1))$

Let

$$A = k[x]/(x(x-1)).$$

Then in A ,

$$\bar{x}(\bar{x}-1) = 0.$$

Hence $\bar{x}$ and $\bar{x}-1$ are zero divisors.

This ring encodes the relation

$$x(x-1) = 0.$$

Geometrically, it corresponds to two points on the affine line: $x = 0$ and $x = 1$.

Examples.

$$\bar{x} \neq 0, \quad \bar{x}-1 \neq 0, \quad \bar{x}(\bar{x}-1) = 0.$$

1.5.8 8. Example: $k[x, y]/(xy)$

Let

$$A = k[x, y]/(xy).$$

Then

$$\bar{x}\bar{y} = 0,$$

but neither $\bar{x}$ nor $\bar{y}$ is zero. Indeed,

$$x \notin (xy), \quad y \notin (xy).$$

This is one of the most important examples:

$$\bar{x}\bar{y} = 0 \quad \text{with} \quad \bar{x} \neq 0, \quad \bar{y} \neq 0.$$

So $\bar{x}$ and $\bar{y}$ are nonzero zero divisors.

Examples of classes.

$$\overline{x+xy} = \bar{x}, \quad \overline{y+x^2y} = \bar{y}, \quad \overline{xy} = 0.$$

Annihilators. We compute:

$$\text{Ann}(\bar{x}) = (\bar{y}), \quad \text{Ann}(\bar{y}) = (\bar{x}).$$

Indeed, $\bar{f} \in \text{Ann}(\bar{x})$ means

$$\bar{f}\bar{x} = 0 \iff fx \in (xy),$$

which forces $f \in (y)$. Similarly for $\text{Ann}(\bar{y})$.

1.5.9 9. Example: $k[x, y]/(x^2, xy)$

Let

$$A = k[x, y]/(x^2, xy).$$

Then in A ,

$$\bar{x}^2 = 0, \quad \bar{x}\bar{y} = 0.$$

Thus $\bar{x}$ is killed by both $\bar{x}$ and $\bar{y}$.

We compute:

$$\text{Ann}(\bar{x}) = (\bar{x}, \bar{y}).$$

Indeed, if $f\bar{x} = 0$, then

$$fx \in (x^2, xy) = x(x, y),$$

so $f \in (x, y)$.

This example is very important because it has an embedded associated prime.

1.5.10 10. Minimal primes

Let $A = R/I$. A prime ideal $\mathfrak{p} \subseteq R$ is *minimal over I* if:

- $I \subseteq \mathfrak{p}$,
- $\mathfrak{p}$ is prime,
- there is no smaller prime ideal between I and $\mathfrak{p}$.

The corresponding primes in $A = R/I$ are called the *minimal primes* of A . We write

$$\text{Min}(A)$$

for the set of minimal primes.

Interpretation. Minimal primes describe the top-level irreducible components of the ring or affine variety.

Example: $k[x]$

Since $k[x]$ is a domain, the zero ideal is prime. Thus

$$\text{Min}(k[x]) = \{(0)\}.$$

Example: $k[x]/(x^2)$

The radical of (x^2) is (x) , so

$$\text{Min}(k[x]/(x^2)) = \{(\bar{x})\}.$$

Example: $k[x]/(x(x-1))$

The ideal $(x(x-1))$ has two minimal primes:

$$(x), \quad (x-1).$$

Hence

$$\text{Min}\left(k[x]/(x(x-1))\right) = \{(\bar{x}), (\bar{x}-1)\}.$$

Example: $k[x, y]/(xy)$

The ideal (xy) has two minimal primes:

$$(x), \quad (y).$$

Therefore

$$\text{Min}\left(k[x, y]/(xy)\right) = \{(\bar{x}), (\bar{y})\}.$$

Example: $k[x, y]/(x^2, xy)$

We have

$$\sqrt{(x^2, xy)} = (x),$$

so

$$\text{Min}\left(k[x, y]/(x^2, xy)\right) = \{(\bar{x})\}.$$

1.5.11 11. Associated primes

A prime ideal $\mathfrak{p} \subseteq A$ is an *associated prime* of the A -module A if

$$\mathfrak{p} = \text{Ann}_A(a)$$

for some $a \in A$.

We write

$$\text{Ass}(A)$$

for the set of associated primes.

Interpretation. Associated primes detect actual annihilators of elements. They describe zero-divisor structure and may reveal hidden embedded components.

Example: $k[x]/(x^2)$

We have

$$\text{Ann}(\bar{x}) = (\bar{x}),$$

so

$$\text{Ass}\left(k[x]/(x^2)\right) = \{(\bar{x})\}.$$

Example: $k[x]/(x(x-1))$

We have

$$\text{Ann}(\bar{x}) = (\bar{x} - 1), \quad \text{Ann}(\bar{x} - 1) = (\bar{x}).$$

Hence

$$\text{Ass}\left(k[x]/(x(x-1))\right) = \{(\bar{x}), (\bar{x} - 1)\}.$$

Example: $k[x, y]/(xy)$

We computed

$$\text{Ann}(\bar{x}) = (\bar{y}), \quad \text{Ann}(\bar{y}) = (\bar{x}),$$

so

$$\text{Ass}\left(k[x, y]/(xy)\right) = \{(\bar{x}), (\bar{y})\}.$$

Example: $k[x, y]/(x^2, xy)$

We have already seen that

$$\text{Ann}(\bar{x}) = (\bar{x}, \bar{y}).$$

Also the minimal prime $(\bar{x})$ is associated in this Noetherian case. Therefore

$$\text{Ass}\left(k[x, y]/(x^2, xy)\right) = \{(\bar{x}), (\bar{x}, \bar{y})\}.$$

Thus

$$\text{Min}(A) = \{(\bar{x})\} \subsetneq \text{Ass}(A) = \{(\bar{x}), (\bar{x}, \bar{y})\}.$$

The prime $(\bar{x}, \bar{y})$ is an *embedded associated prime*.

1.5.12 12. Comparison of minimal primes and associated primes

Minimal primes describe the irreducible components:

top-level geometry.

Associated primes describe annihilators of actual elements:

zero-divisor structure and embedded behavior.

In general, for Noetherian rings,

$$\text{Min}(A) \subseteq \text{Ass}(A),$$

and the inclusion may be strict.

1.5.13 13. Summary of the main examples

A	$\text{Min}(A)$	$\text{Ass}(A)$
$k[x]$	$\{(0)\}$	$\{(0)\}$
$k[x]/(x^2)$	$\{(\bar{x})\}$	$\{(\bar{x})\}$
$k[x]/(x(x-1))$	$\{(\bar{x}), (\bar{x}-1)\}$	$\{(\bar{x}), (\bar{x}-1)\}$
$k[x, y]/(xy)$	$\{(\bar{x}), (\bar{y})\}$	$\{(\bar{x}), (\bar{y})\}$
$k[x, y]/(x^2, xy)$	$\{(\bar{x})\}$	$\{(\bar{x}), (\bar{x}, \bar{y})\}$

1.5.14 14. Final intuition

The key points are:

- x, y are formal variables in polynomial rings;
- x^2, xy are formal products;
- quotient rings impose algebraic relations such as $x^2 = 0$ or $xy = 0$;
- $\bar{x}, \bar{y}$ are classes of x, y in the quotient;
- minimal primes describe the main geometric components;
- associated primes describe where actual annihilation occurs.

Thus one may think of:

minimal primes = geometry, associated primes = geometry plus hidden zero-divisor structure.

1.6 Concrete annihilators: $\mathbb{Z}/p\mathbb{Z}$, $\mathbb{Z}/n\mathbb{Z}$, polynomial rings, and quotient rings

1.6.1 1. Definition of annihilator

Let A be a ring and let $a \in A$. The annihilator of a is

$$\text{Ann}_A(a) = \{r \in A : ra = 0\}.$$

It is an ideal of A .

More generally, if $S \subseteq A$, then

$$\text{Ann}_A(S) = \{r \in A : rs = 0 \text{ for all } s \in S\}.$$

For a single element a , this is just $\text{Ann}_A(a)$.

Basic intuition. The annihilator of a measures which elements of the ring kill a by multiplication. So:

- if $\text{Ann}_A(a) = 0$, then no nonzero element kills a ;
- if $\text{Ann}_A(a) \neq 0$, then a is a zero divisor (unless $a = 0$);
- prime annihilators are exactly what lead to associated primes.

1.6.2 2. First trivial examples

For every ring A ,

$$\text{Ann}_A(0) = A,$$

because every element kills 0.

Also,

$$\text{Ann}_A(1) = 0,$$

because if $r \cdot 1 = 0$, then $r = 0$.

Thus in every nonzero ring:

$$\text{Ann}_A(0) = A, \quad \text{Ann}_A(1) = 0.$$

1.6.3 3. Annihilators in a domain

Suppose A is an integral domain. Then for every nonzero $a \in A$,

$$\text{Ann}_A(a) = 0.$$

Indeed, if $ra = 0$ and $a \neq 0$, then $r = 0$ because there are no zero divisors.

Examples. In $k[x]$, where k is a field,

$$\text{Ann}_{k[x]}(x) = 0, \quad \text{Ann}_{k[x]}(x^2 + 1) = 0, \quad \text{Ann}_{k[x]}(f) = 0$$

for every nonzero polynomial $f \in k[x]$.

So in a domain, annihilators of nonzero elements are always trivial.

1.6.4 4. Annihilators in the field $\mathbb{Z}/p\mathbb{Z}$

Let p be prime. Then

$$\mathbb{Z}/p\mathbb{Z}$$

is a field. Therefore every nonzero element is invertible, hence cannot have a nonzero annihilator.

So:

$$\text{Ann}_{\mathbb{Z}/p\mathbb{Z}}(\bar{0}) = \mathbb{Z}/p\mathbb{Z},$$

and for every $\bar{a} \neq \bar{0}$,

$$\text{Ann}_{\mathbb{Z}/p\mathbb{Z}}(\bar{a}) = 0.$$

Example: $\mathbb{Z}/5\mathbb{Z}$

The elements are

$$\bar{0}, \bar{1}, \bar{2}, \bar{3}, \bar{4}.$$

Then:

$$\text{Ann}(\bar{0}) = \mathbb{Z}/5\mathbb{Z},$$

but

$$\text{Ann}(\bar{1}) = 0, \quad \text{Ann}(\bar{2}) = 0, \quad \text{Ann}(\bar{3}) = 0, \quad \text{Ann}(\bar{4}) = 0.$$

Indeed, for example, if $\bar{r}\bar{2} = \bar{0}$, then multiply by the inverse of $\bar{2}$, namely $\bar{3}$, to get $\bar{r} = \bar{0}$.

Conclusion. Fields have no nonzero zero divisors, and therefore all nonzero annihilators are trivial.

1.6.5 5. Annihilators in $\mathbb{Z}/n\mathbb{Z}$

Now let

$$A = \mathbb{Z}/n\mathbb{Z}.$$

This is much more interesting when n is composite.

For $\bar{a} \in \mathbb{Z}/n\mathbb{Z}$,

$$\text{Ann}(\bar{a}) = \{\bar{r} \in \mathbb{Z}/n\mathbb{Z} : \bar{r}\bar{a} = \bar{0}\}.$$

This means

$$ra \equiv 0 \pmod{n},$$

or equivalently,

$$n \mid ra.$$

Let

$$d = \gcd(n, a).$$

Then one has the concrete formula

$$\text{Ann}_{\mathbb{Z}/n\mathbb{Z}}(\bar{a}) = \left(\frac{\bar{n}}{\bar{d}}\right).$$

Why? Write

$$n = dn_1, \quad a = da_1, \quad \gcd(n_1, a_1) = 1.$$

Then

$$n \mid ra \iff dn_1 \mid rda_1 \iff n_1 \mid ra_1.$$

Since $\gcd(n_1, a_1) = 1$, this is equivalent to

$$n_1 \mid r.$$

So r must be a multiple of n/d , exactly as claimed.

Example: $\mathbb{Z}/6\mathbb{Z}$

Annihilator of $\bar{2}$. Here

$$\gcd(6, 2) = 2, \quad \frac{6}{2} = 3.$$

So

$$\text{Ann}(\bar{2}) = (\bar{3}) = \{\bar{0}, \bar{3}\}.$$

Indeed,

$$\bar{0} \cdot \bar{2} = \bar{0}, \quad \bar{3} \cdot \bar{2} = \bar{6} = \bar{0}.$$

Annihilator of $\bar{3}$. Here

$$\gcd(6, 3) = 3, \quad \frac{6}{3} = 2.$$

So

$$\text{Ann}(\bar{3}) = (\bar{2}) = \{\bar{0}, \bar{2}, \bar{4}\}.$$

Indeed,

$$\bar{2} \cdot \bar{3} = \bar{0}, \quad \bar{4} \cdot \bar{3} = \bar{12} = \bar{0}.$$

Annihilator of $\bar{4}$. Since $\gcd(6, 4) = 2$, we get

$$\text{Ann}(\bar{4}) = (\bar{3}) = \{\bar{0}, \bar{3}\}.$$

Annihilator of $\bar{1}$. Since $\gcd(6, 1) = 1$,

$$\text{Ann}(\bar{1}) = (\bar{6}) = 0.$$

Thus the nonzero elements $\bar{2}, \bar{3}, \bar{4}$ have nontrivial annihilators, while $\bar{1}, \bar{5}$ do not.

Example: $\mathbb{Z}/12\mathbb{Z}$

Annihilator of $\bar{3}$.

$$\gcd(12, 3) = 3, \quad \frac{12}{3} = 4.$$

Hence

$$\text{Ann}(\bar{3}) = (\bar{4}) = \{\bar{0}, \bar{4}, \bar{8}\}.$$

Annihilator of $\bar{4}$.

$$\gcd(12, 4) = 4, \quad \frac{12}{4} = 3.$$

Hence

$$\text{Ann}(\bar{4}) = (\bar{3}) = \{\bar{0}, \bar{3}, \bar{6}, \bar{9}\}.$$

Annihilator of $\bar{6}$.

$$\gcd(12, 6) = 6, \quad \frac{12}{6} = 2.$$

Hence

$$\text{Ann}(\bar{6}) = (\bar{2}) = \{\bar{0}, \bar{2}, \bar{4}, \bar{6}, \bar{8}, \bar{10}\}.$$

So in $\mathbb{Z}/n\mathbb{Z}$, annihilators can be computed explicitly using $\gcd$.

Deep point. In $\mathbb{Z}/n\mathbb{Z}$,

$$\bar{a} \text{ is a zero divisor} \iff \gcd(n, a) \neq 1 \iff \text{Ann}(\bar{a}) \neq 0.$$

1.6.6 6. Annihilators in polynomial rings $k[x]$

Let k be a field. Since $k[x]$ is a domain, every nonzero polynomial has trivial annihilator:

$$\text{Ann}_{k[x]}(f) = 0 \quad \text{for all } f \neq 0.$$

Examples

$$\text{Ann}(x) = 0, \quad \text{Ann}(x^2) = 0, \quad \text{Ann}(x^2 + 1) = 0, \quad \text{Ann}((x - 1)(x + 2)) = 0.$$

So in ordinary polynomial rings over a field, annihilators do not become interesting until we pass to quotients.

1.6.7 7. Annihilators in $k[x]/(x^2)$

Let

$$A = k[x]/(x^2).$$

Then $\bar{x} \neq 0$, but

$$\bar{x}^2 = 0.$$

So $\bar{x}$ has a nontrivial annihilator.

Annihilator of $\bar{x}$. We compute

$$\text{Ann}_A(\bar{x}) = \{\bar{f} : \bar{f}\bar{x} = 0\}.$$

This means

$$fx \in (x^2).$$

So f must be divisible by x . Therefore

$$\text{Ann}_A(\bar{x}) = (\bar{x}).$$

General element. Every element of A has the form

$$a + b\bar{x}, \quad a, b \in k.$$

Now:

- if $a \neq 0$, then $a + b\bar{x}$ is a unit, so

$$\text{Ann}(a + b\bar{x}) = 0;$$

- if $a = 0$ and $b \neq 0$, then $a + b\bar{x} = b\bar{x}$, and

$$\text{Ann}(b\bar{x}) = (\bar{x}).$$

So the annihilators in this ring are:

$$\text{Ann}(0) = A, \quad \text{Ann}(\bar{x}) = (\bar{x}), \quad \text{Ann}(1) = 0, \quad \text{Ann}(1 + \bar{x}) = 0.$$

Interpretation. This ring has only one nonzero prime ideal, namely $(\bar{x})$, and $\bar{x}$ is exactly the nilpotent element killed by that prime.

1.6.8 8. Annihilators in $k[x]/(x(x-1))$

Let

$$A = k[x]/(x(x-1)).$$

Then

$$\bar{x}(\bar{x}-1) = 0.$$

Annihilator of $\bar{x}$. Since $(\bar{x}-1)\bar{x} = 0$, we have

$$(\bar{x}-1) \subseteq \text{Ann}(\bar{x}).$$

Conversely, if $f\bar{x} = 0$, then

$$fx \in (x(x-1)).$$

So fx is divisible by $x(x-1)$, hence f is divisible by $x-1$. Therefore

$$\text{Ann}(\bar{x}) = (\bar{x}-1).$$

Annihilator of $\bar{x}-1$. Similarly,

$$\text{Ann}(\bar{x}-1) = (\bar{x}).$$

Thus:

$$\text{Ann}(\bar{x}) = (\bar{x}-1), \quad \text{Ann}(\bar{x}-1) = (\bar{x}).$$

Interpretation. Each component kills the other one. This is the algebraic reflection of the decomposition into the two points $x = 0$ and $x = 1$.

1.6.9 9. Annihilators in $k[x, y]/(xy)$

Let

$$A = k[x, y]/(xy).$$

Then

$$\bar{x}\bar{y} = 0.$$

Annihilator of $\bar{x}$. We want

$$\bar{f}\bar{x} = 0 \iff fx \in (xy).$$

This means fx is divisible by xy , so f must be divisible by y . Hence

$$\text{Ann}(\bar{x}) = (\bar{y}).$$

Annihilator of $\bar{y}$. Similarly,

$$\text{Ann}(\bar{y}) = (\bar{x}).$$

Annihilator of $\bar{x} + \bar{y}$. Now ask whether $\bar{x} + \bar{y}$ has a nontrivial annihilator. Suppose

$$\bar{f}(\bar{x} + \bar{y}) = 0.$$

Then

$$f(x + y) \in (xy).$$

But $x + y$ is not divisible by x or by y , so the only way this product lies in (xy) is that $f \in (xy)$, i.e. $\bar{f} = 0$. Therefore

$$\text{Ann}(\bar{x} + \bar{y}) = 0.$$

So in this ring:

$$\text{Ann}(\bar{x}) = (\bar{y}), \quad \text{Ann}(\bar{y}) = (\bar{x}), \quad \text{Ann}(\bar{x} + \bar{y}) = 0.$$

Concrete lesson. Not every nonunit-looking element is a zero divisor. The elements $\bar{x}$ and $\bar{y}$ are killed by nonzero ideals, but $\bar{x} + \bar{y}$ is not.

1.6.10 10. Annihilators in $k[x, y]/(x^2, xy)$

Let

$$A = k[x, y]/(x^2, xy).$$

Then

$$\bar{x}^2 = 0, \quad \bar{x}\bar{y} = 0.$$

Annihilator of $\bar{x}$. Since both $\bar{x}$ and $\bar{y}$ kill $\bar{x}$, we have

$$(\bar{x}, \bar{y}) \subseteq \text{Ann}(\bar{x}).$$

Conversely, if $\bar{f}\bar{x} = 0$, then

$$fx \in (x^2, xy) = x(x, y).$$

So $f \in (x, y)$. Therefore

$$\text{Ann}(\bar{x}) = (\bar{x}, \bar{y}).$$

Annihilator of $\bar{y}$. Suppose $\bar{f}\bar{y} = 0$. Then

$$fy \in (x^2, xy).$$

Since x^2 and xy are both divisible by x , and fy contains no forced factor x , the only possibility is that $f \in (x)$ and the remaining factor still vanishes. In fact one checks that no nonzero class kills $\bar{y}$, so

$$\text{Ann}(\bar{y}) = 0.$$

Annihilator of $\bar{x} + \bar{y}$. Again, $\bar{y}$ cannot be killed nontrivially, so $\bar{x} + \bar{y}$ is not killed by all of $(\bar{x}, \bar{y})$. In fact

$$\text{Ann}(\bar{x} + \bar{y}) = 0.$$

Thus:

$$\text{Ann}(\bar{x}) = (\bar{x}, \bar{y}), \quad \text{Ann}(\bar{y}) = 0, \quad \text{Ann}(\bar{x} + \bar{y}) = 0.$$

Interpretation. The element $\bar{x}$ is very fragile: it is killed by the maximal ideal $(\bar{x}, \bar{y})$. But $\bar{y}$ behaves much more like an element in a domain.

1.6.11 11. Annihilators and associated primes

A prime ideal $\mathfrak{p} \subseteq A$ is associated if

$$\mathfrak{p} = \text{Ann}_A(a)$$

for some $a \in A$.

So annihilators are the direct source of associated primes.

Examples

In $k[x]/(x^2)$:

$$\text{Ann}(\bar{x}) = (\bar{x}),$$

so

$$\text{Ass}\left(k[x]/(x^2)\right) = \{(\bar{x})\}.$$

In $k[x]/(x(x-1))$:

$$\text{Ann}(\bar{x}) = (\bar{x} - 1), \quad \text{Ann}(\bar{x} - 1) = (\bar{x}),$$

hence

$$\text{Ass}\left(k[x]/(x(x-1))\right) = \{(\bar{x}), (\bar{x} - 1)\}.$$

In $k[x, y]/(xy)$:

$$\text{Ann}(\bar{x}) = (\bar{y}), \quad \text{Ann}(\bar{y}) = (\bar{x}),$$

hence

$$\text{Ass}\left(k[x, y]/(xy)\right) = \{(\bar{x}), (\bar{y})\}.$$

In $k[x, y]/(x^2, xy)$:

$$\text{Ann}(\bar{x}) = (\bar{x}, \bar{y}),$$

so $(\bar{x}, \bar{y})$ is associated. Also the minimal prime $(\bar{x})$ is associated in this Noetherian example. Therefore

$$\text{Ass}\left(k[x, y]/(x^2, xy)\right) = \{(\bar{x}), (\bar{x}, \bar{y})\}.$$

1.6.12 12. Annihilators and minimal primes

Minimal primes describe the main geometric components. Associated primes come from annihilators and may include embedded primes.

Comparison table

A	sample element a	$\text{Ann}_A(a)$	relevant primes
$\mathbb{Z}/5\mathbb{Z}$	2	0	no nonzero annihilator
$\mathbb{Z}/6\mathbb{Z}$	$\bar{2}$	$(\bar{3})$	$(\bar{3})$
$k[x]/(x^2)$	$\bar{x}$	$(\bar{x})$	$(\bar{x})$
$k[x]/(x(x-1))$	$\bar{x}$	$(\bar{x}-1)$	$(\bar{x}), (\bar{x}-1)$
$k[x, y]/(xy)$	$\bar{x}$	$(\bar{y})$	$(\bar{x}), (\bar{y})$
$k[x, y]/(x^2, xy)$	$\bar{x}$	$(\bar{x}, \bar{y})$	$(\bar{x}), (\bar{x}, \bar{y})$

1.6.13 13. Final summary

The annihilator is one of the most concrete and useful constructions in commutative algebra:

$$\text{Ann}_A(a) = \{r \in A : ra = 0\}.$$

Its behavior is easy to understand in important examples:

- in a field such as $\mathbb{Z}/p\mathbb{Z}$, every nonzero annihilator is trivial;
- in $\mathbb{Z}/n\mathbb{Z}$, annihilators are computed by gcd:

$$\text{Ann}(\bar{a}) = \left(\frac{\overline{n}}{\gcd(n, a)} \right);$$

- in a domain such as $k[x]$, every nonzero annihilator is trivial;
- in quotient rings such as $k[x]/(x^2)$ or $k[x, y]/(xy)$, annihilators become nontrivial and reveal zero-divisor structure;
- prime annihilators are exactly the source of associated primes.

Thus annihilators form the bridge between concrete multiplication computations and the abstract prime-ideal structure of the ring.

1.7 Annihilators, minimal primes, and associated primes for modules: concrete examples

1.7.1 1. Annihilators for modules: the basic definitions

Let R be a ring and let M be an R -module.

For an element $m \in M$, the annihilator of m is

$$\text{Ann}_R(m) = \{r \in R : rm = 0\}.$$

This is an ideal of R .

The annihilator of the whole module M is

$$\text{Ann}_R(M) = \{r \in R : rM = 0\}.$$

Equivalently,

$$\text{Ann}_R(M) = \bigcap_{m \in M} \text{Ann}_R(m).$$

Interpretation.

- $\text{Ann}_R(m)$ tells us which scalars kill one specific element m .
- $\text{Ann}_R(M)$ tells us which scalars kill the entire module.

These two ideals are usually different.

1.7.2 2. Associated primes and minimal primes of a module

A prime ideal $\mathfrak{p} \subseteq R$ is an *associated prime* of the R -module M if

$$\mathfrak{p} = \text{Ann}_R(m)$$

for some $m \in M$.

We write

$$\text{Ass}_R(M)$$

for the set of associated primes of M .

If M is finitely generated over a Noetherian ring, one defines the minimal primes of M as the minimal primes in the support of M , equivalently the minimal primes over $\text{Ann}_R(M)$. We write

$$\text{Min}_R(M)$$

for this set.

Interpretation.

- $\text{Min}_R(M)$ describes the main geometric components seen by the module.
- $\text{Ass}_R(M)$ records prime annihilators of actual elements of the module.

In the Noetherian finitely generated setting,

$$\text{Min}_R(M) \subseteq \text{Ass}_R(M),$$

and the inclusion may be strict.

1.7.3 3. The cyclic module R/I

A very important case is the cyclic module

$$M = R/I.$$

Then the element $\bar{1} = 1 + I \in R/I$ satisfies

$$\text{Ann}_R(\bar{1}) = I.$$

More generally, for $\bar{f} = f + I \in R/I$,

$$\text{Ann}_R(\bar{f}) = \{r \in R : rf \in I\} = (I : f),$$

the colon ideal.

So cyclic modules give a very concrete way to compute annihilators.

1.7.4 4. First examples over $\mathbb{Z}$

Example 1: $M = \mathbb{Z}$ as a $\mathbb{Z}$ -module

Take $R = \mathbb{Z}$ and $M = \mathbb{Z}$.

For any nonzero integer m ,

$$\text{Ann}_{\mathbb{Z}}(m) = 0,$$

because if $rm = 0$ in $\mathbb{Z}$, then $r = 0$.

Thus

$$\text{Ann}_{\mathbb{Z}}(\mathbb{Z}) = 0.$$

Since (0) is prime in $\mathbb{Z}$, we get

$$\text{Ass}_{\mathbb{Z}}(\mathbb{Z}) = \{(0)\}, \quad \text{Min}_{\mathbb{Z}}(\mathbb{Z}) = \{(0)\}.$$

So the free module $\mathbb{Z}$ behaves like a torsion-free object.

Example 2: $M = \mathbb{Z}/12\mathbb{Z}$

Now let

$$M = \mathbb{Z}/12\mathbb{Z}.$$

As a $\mathbb{Z}$ -module, this is a cyclic module:

$$M = \mathbb{Z}/(12).$$

First,

$$\text{Ann}_{\mathbb{Z}}(M) = (12),$$

because 12 kills every class modulo 12, and no smaller ideal does.

Now compute annihilators of specific elements.

Element $\bar{1}$.

$$\text{Ann}(\bar{1}) = (12).$$

Element $\bar{2}$. We want all $r \in \mathbb{Z}$ such that

$$r\bar{2} = \bar{0} \text{ in } \mathbb{Z}/12\mathbb{Z}.$$

This means

$$12 \mid 2r,$$

so r must be divisible by 6. Hence

$$\text{Ann}(\bar{2}) = (6).$$

Element $\bar{3}$. We need

$$12 \mid 3r,$$

so r must be divisible by 4. Thus

$$\text{Ann}(\bar{3}) = (4).$$

Element $\bar{4}$. We need

$$12 \mid 4r,$$

so r must be divisible by 3. Hence

$$\text{Ann}(\bar{4}) = (3).$$

Element $\bar{6}$. We need

$$12 \mid 6r,$$

so r must be divisible by 2. Hence

$$\text{Ann}(\bar{6}) = (2).$$

So the prime annihilators that occur are

$$(2), \quad (3).$$

Therefore

$$\text{Ass}_{\mathbb{Z}}(\mathbb{Z}/12\mathbb{Z}) = \{(2), (3)\}.$$

Also the minimal primes over (12) are exactly

$$(2), \quad (3),$$

so

$$\text{Min}_{\mathbb{Z}}(\mathbb{Z}/12\mathbb{Z}) = \{(2), (3)\}.$$

General formula. For $M = \mathbb{Z}/n\mathbb{Z}$ and $\bar{a} \in M$,

$$\text{Ann}_{\mathbb{Z}}(\bar{a}) = \left(\frac{n}{\gcd(n, a)} \right).$$

1.7.5 5. A vector-space example

Let k be a field, and let $M = k^n$ be a k -vector space.

For every nonzero $v \in k^n$,

$$\text{Ann}_k(v) = 0,$$

because only $0 \in k$ kills a nonzero vector.

Thus

$$\text{Ann}_k(k^n) = 0, \quad \text{Ass}_k(k^n) = \{(0)\}, \quad \text{Min}_k(k^n) = \{(0)\}.$$

So over a field, modules behave very simply: nonzero elements have trivial annihilator.

1.7.6 6. Polynomial modules: $M = k[x]/(x^2)$

Let

$$R = k[x], \quad M = R/(x^2).$$

Then M is an R -module.

Since $M = R/I$ with $I = (x^2)$,

$$\text{Ann}_R(M) = (x^2).$$

Now compute annihilators of concrete elements.

Element $\bar{1}$.

$$\text{Ann}_R(\bar{1}) = (x^2).$$

Element $\bar{x}$. We need all $f \in k[x]$ such that

$$f\bar{x} = 0 \text{ in } R/(x^2).$$

This means

$$fx \in (x^2),$$

so f must be divisible by x . Hence

$$\text{Ann}_R(\bar{x}) = (x).$$

Element $\overline{1+x}$. Suppose

$$f(\overline{1+x}) = 0.$$

Then

$$f(1+x) \in (x^2).$$

But $1+x$ is not divisible by x , so the only way this can happen is $f \in (x^2)$. Therefore

$$\text{Ann}_R(\overline{1+x}) = (x^2).$$

Thus the prime annihilator that appears is

$$(x).$$

So

$$\text{Ass}_R(k[x]/(x^2)) = \{(x)\}.$$

Also the only minimal prime over (x^2) is

$$(x),$$

hence

$$\text{Min}_R(k[x]/(x^2)) = \{(x)\}.$$

1.7.7 7. Polynomial modules: $M = k[x]/(x(x-1))$

Let

$$R = k[x], \quad M = R/(x(x-1)).$$

Then

$$\text{Ann}_R(M) = (x(x-1)).$$

Now compute annihilators of elements.

Element $\bar{1}$.

$$\text{Ann}(\bar{1}) = (x(x-1)).$$

Element $\bar{x}$. We need

$$f\bar{x} = 0 \iff fx \in (x(x-1)).$$

So fx must be divisible by $x(x-1)$, hence f must be divisible by $x-1$. Therefore

$$\text{Ann}(\bar{x}) = (x-1).$$

Element $\overline{x-1}$. Similarly,

$$\text{Ann}(\overline{x-1}) = (x).$$

Thus the associated primes are

$$\text{Ass}_R(k[x]/(x(x-1))) = \{(x), (x-1)\}.$$

These are also the minimal primes over $(x(x-1))$, so

$$\text{Min}_R(k[x]/(x(x-1))) = \{(x), (x-1)\}.$$

Interpretation. The module sees two components, corresponding to the two points $x = 0$ and $x = 1$.

1.7.8 8. Polynomial modules: $M = k[x, y]/(xy)$

Let

$$R = k[x, y], \quad M = R/(xy).$$

Then

$$\text{Ann}_R(M) = (xy).$$

Now compute element annihilators.

Element $\bar{x}$. We need

$$f\bar{x} = 0 \iff fx \in (xy).$$

Thus f must be divisible by y , so

$$\text{Ann}(\bar{x}) = (y).$$

Element $\bar{y}$. Similarly,

$$\text{Ann}(\bar{y}) = (x).$$

Element $\overline{x+y}$. Suppose

$$f(\overline{x+y}) = 0.$$

Then

$$f(x+y) \in (xy).$$

Since $x+y$ is not divisible by x or by y , the only possibility is

$$f \in (xy).$$

Therefore

$$\text{Ann}(\overline{x+y}) = (xy).$$

So the prime annihilators that occur are

$$(x), \quad (y).$$

Hence

$$\text{Ass}_R(k[x, y]/(xy)) = \{(x), (y)\}.$$

These are also the minimal primes over (xy) , so

$$\text{Min}_R(k[x, y]/(xy)) = \{(x), (y)\}.$$

Interpretation. The element $\bar{x}$ is killed by the y -axis, and $\bar{y}$ is killed by the x -axis.

1.7.9 9. An embedded-prime module: $M = k[x, y]/(x^2, xy)$

Let

$$R = k[x, y], \quad M = R/(x^2, xy).$$

Then

$$\text{Ann}_R(M) = (x^2, xy).$$

Now compute annihilators of elements.

Element $\bar{1}$.

$$\text{Ann}(\bar{1}) = (x^2, xy).$$

Element $\bar{x}$. We need

$$f\bar{x} = 0 \iff fx \in (x^2, xy) = x(x, y).$$

This is equivalent to $f \in (x, y)$. Hence

$$\text{Ann}(\bar{x}) = (x, y).$$

Element $\bar{y}$. We need

$$f\bar{y} = 0 \iff fy \in (x^2, xy).$$

Since fy belongs to $(x^2, xy) = x(x, y)$, it must be divisible by x . Because x and y are independent variables, this forces $f \in (x)$. Conversely, any multiple of x kills $\bar{y}$, because $xy \in (x^2, xy)$. Therefore

$$\text{Ann}(\bar{y}) = (x).$$

Thus we get two prime annihilators:

$$(x), \quad (x, y).$$

Hence

$$\text{Ass}_R(k[x, y]/(x^2, xy)) = \{(x), (x, y)\}.$$

Now

$$\sqrt{(x^2, xy)} = (x),$$

so the only minimal prime over $\text{Ann}_R(M)$ is

$$(x).$$

Thus

$$\text{Min}_R(k[x, y]/(x^2, xy)) = \{(x)\}.$$

Therefore

$$\text{Min}_R(M) = \{(x)\} \subsetneq \text{Ass}_R(M) = \{(x), (x, y)\}.$$

The prime (x, y) is an embedded associated prime.

1.7.10 10. Direct sums of modules

Let

$$R = k[x, y], \quad M = R/(x) \oplus R/(y).$$

Then

$$\text{Ann}_R(M) = \text{Ann}_R(R/(x)) \cap \text{Ann}_R(R/(y)) = (x) \cap (y) = (xy).$$

Now consider the two elements

$$m_1 = (\bar{1}, 0), \quad m_2 = (0, \bar{1}).$$

Their annihilators are:

$$\text{Ann}_R(m_1) = (x), \quad \text{Ann}_R(m_2) = (y).$$

So

$$\text{Ass}_R(M) = \{(x), (y)\}.$$

And the minimal primes over (xy) are again

$$(x), \quad (y).$$

Hence

$$\text{Min}_R(M) = \{(x), (y)\}.$$

This is a very concrete example showing that associated primes of a direct sum are the union of the associated primes of the summands.

1.7.11 11. A free polynomial module

Let

$$R = k[x], \quad M = R^2.$$

Then no nonzero polynomial kills all of R^2 , so

$$\text{Ann}_R(M) = 0.$$

Also, for a nonzero vector such as

$$m = (1, x),$$

if $f(1, x) = (0, 0)$, then $f = 0$. Hence

$$\text{Ann}_R(m) = 0.$$

Since $R = k[x]$ is a domain, (0) is prime. Therefore

$$\text{Ass}_R(R^2) = \{(0)\}, \quad \text{Min}_R(R^2) = \{(0)\}.$$

So free modules over domains behave like torsion-free modules.

1.7.12 12. Summary table

R	M	$\text{Ann}_R(M)$	$\text{Ass}_R(M)$	$\text{Min}_R(M)$
$\mathbb{Z}$	$\mathbb{Z}$	(0)	$\{(0)\}$	$\{(0)\}$
$\mathbb{Z}$	$\mathbb{Z}/12\mathbb{Z}$	(12)	$\{(2), (3)\}$	$\{(2), (3)\}$
$k[x]$	$k[x]/(x^2)$	(x^2)	$\{(x)\}$	$\{(x)\}$
$k[x]$	$k[x]/(x(x-1))$	$(x(x-1))$	$\{(x), (x-1)\}$	$\{(x), (x-1)\}$
$k[x, y]$	$k[x, y]/(xy)$	(xy)	$\{(x), (y)\}$	$\{(x), (y)\}$
$k[x, y]$	$k[x, y]/(x^2, xy)$	(x^2, xy)	$\{(x), (x, y)\}$	$\{(x)\}$
$k[x, y]$	$R/(x) \oplus R/(y)$	(xy)	$\{(x), (y)\}$	$\{(x), (y)\}$
$k[x]$	R^2	(0)	$\{(0)\}$	$\{(0)\}$

1.7.13 13. Final interpretation

For modules, one must always distinguish:

$$\text{Ann}_R(m) \quad \text{for one element } m \in M,$$

from

$$\text{Ann}_R(M) \quad \text{for the whole module.}$$

The annihilator of the whole module need not be prime and usually describes the global defining equations of the module. The annihilator of a specific element may be much smaller, and when it is prime, it contributes to the associated primes.

Thus:

$$\text{minimal primes of } M = \text{main geometric support of } M,$$

while

$$\text{associated primes of } M = \text{prime annihilators of actual elements of } M.$$

That is why associated primes are finer than minimal primes: they can detect embedded structure that the global annihilator $\text{Ann}_R(M)$ does not show by itself.

1.8 Concrete residue classes in $\mathbb{Z}/n\mathbb{Z}$, $\mathbb{Z}/p\mathbb{Z}$, $k[x]/(x)$, $A[t]/(t)$, and $k[x, y]/(xy)$

1.8.1 1. General principle: what is a residue class in R/I ?

Let R be a ring and $I \subseteq R$ an ideal. An element of the quotient ring

$$R/I$$

is a residue class

$$\bar{f} = f + I = \{f + h : h \in I\}.$$

Thus $\bar{f}$ is not just one element f , but the whole set of elements differing from f by something in I .

Two elements $f, g \in R$ define the same class if and only if

$$\bar{f} = \bar{g} \iff f - g \in I.$$

So the quotient R/I identifies all elements that differ by an element of I .

1.8.2 2. Residue classes in $\mathbb{Z}/n\mathbb{Z}$

The quotient

$$\mathbb{Z}/n\mathbb{Z}$$

consists of congruence classes modulo n . For an integer a ,

$$\bar{a} = a + n\mathbb{Z} = \{a + nk : k \in \mathbb{Z}\}.$$

So $\bar{a}$ is the set of all integers congruent to a modulo n .

Example: $\mathbb{Z}/5\mathbb{Z}$

We have

$$\bar{2} = 2 + 5\mathbb{Z} = \{\dots, -8, -3, 2, 7, 12, 17, \dots\}.$$

Also

$$\bar{2} = \bar{7} = \overline{-3},$$

because

$$7 - 2 = 5, \quad -3 - 2 = -5,$$

and both are divisible by 5.

Thus the elements of $\mathbb{Z}/5\mathbb{Z}$ are

$$\bar{0}, \bar{1}, \bar{2}, \bar{3}, \bar{4}.$$

Example: $\mathbb{Z}/6\mathbb{Z}$

Similarly,

$$\bar{4} = 4 + 6\mathbb{Z} = \{\dots, -8, -2, 4, 10, 16, \dots\}.$$

Also

$$\bar{4} = \bar{10} = \overline{-2}.$$

Thus in $\mathbb{Z}/n\mathbb{Z}$, a residue class is simply a set of integers having the same remainder modulo n .

1.8.3 3. Residue classes in $\mathbb{Z}/p\mathbb{Z}$

If p is prime, then

$$\mathbb{Z}/p\mathbb{Z}$$

is a field. The residue classes are defined exactly as above:

$$\bar{a} = a + p\mathbb{Z}.$$

Example: $\mathbb{Z}/7\mathbb{Z}$

$$\bar{3} = 3 + 7\mathbb{Z} = \{\dots, -11, -4, 3, 10, 17, 24, \dots\}.$$

So the only difference between $\mathbb{Z}/p\mathbb{Z}$ and $\mathbb{Z}/n\mathbb{Z}$ is not how classes are formed, but the algebraic structure afterward: when p is prime, every nonzero class has a multiplicative inverse.

1.8.4 4. Residue classes in $k[x]/(x)$

Now let k be a field. Consider

$$k[x]/(x).$$

An element is a class

$$\bar{f} = f(x) + (x).$$

Two polynomials $f(x)$ and $g(x)$ lie in the same class if and only if

$$f(x) - g(x) \in (x),$$

that is, if $f(x) - g(x)$ is divisible by x .

This means precisely that f and g have the same constant term.

Indeed, if

$$f(x) = a_0 + a_1x + \dots + a_nx^n,$$

then

$$f(x) - a_0 = x(a_1 + a_2x + \dots + a_nx^{n-1}) \in (x),$$

so

$$\bar{f} = \overline{a_0}.$$

Thus every class is determined by the constant term alone.

Class of 0

$$\bar{0} = (x) = \{xq(x) : q(x) \in k[x]\}.$$

So this class consists of all polynomials divisible by x :

$$x, \quad x^2, \quad 3x + 7x^2, \quad x(x^3 + 1), \dots$$

Class of 1

$$\bar{1} = 1 + (x) = \{1 + xq(x) : q(x) \in k[x]\}.$$

Examples:

$$1, \quad 1 + x, \quad 1 + 3x^2, \quad 1 + x + x^4, \dots$$

Class of $a \in k$

$$\bar{a} = a + (x) = \{a + xq(x) : q(x) \in k[x]\}.$$

Thus in $k[x]/(x)$, two polynomials are equal if and only if they have the same constant term. Therefore

$$k[x]/(x) \cong k.$$

1.8.5 5. Residue classes in $A[t]/(t)$

Let A be any commutative ring. Consider

$$A[t]/(t).$$

An element is a class

$$\bar{f} = f(t) + (t).$$

If

$$f(t) = a_0 + a_1t + \dots + a_nt^n,$$

then

$$f(t) - a_0 \in (t),$$

so

$$\bar{f} = \overline{a_0}.$$

Thus every class is again determined by the constant term.

Class of $a \in A$

$$\bar{a} = a + (t) = \{a + tq(t) : q(t) \in A[t]\}.$$

Class of 0

$$\bar{0} = (t) = \{tq(t) : q(t) \in A[t]\}.$$

So $A[t]/(t)$ behaves exactly like A , and in fact

$$A[t]/(t) \cong A.$$

Example

In $\mathbb{Z}[t]/(t)$,

$$\overline{7 + 3t + t^2} = \bar{7},$$

and the class $\bar{7}$ consists of all polynomials

$$7 + tq(t).$$

1.8.6 6. Residue classes in $k[x, y]/(x)$

Before treating $k[x, y]/(xy)$, it is useful to look at

$$k[x, y]/(x).$$

Here two polynomials are equivalent if and only if their difference is divisible by x .

So every polynomial becomes equal to what remains after setting $x = 0$. In other words, only the y -part survives.

If

$$f(x, y) = g(y) + xh(x, y),$$

then

$$\bar{f} = \overline{g(y)}.$$

Thus every class has a representative depending only on y , and

$$k[x, y]/(x) \cong k[y].$$

Example

$$x^2 + xy + y^3 + 2y + 5$$

has class

$$y^3 + 2y + 5.$$

So the class of $g(y)$ is

$$g(y) + (x) = \{g(y) + xh(x, y) : h(x, y) \in k[x, y]\}.$$

1.8.7 7. Residue classes in $k[x, y]/(xy)$

Now consider

$$B = k[x, y]/(xy).$$

An element is a class

$$\bar{f} = f(x, y) + (xy).$$

Two polynomials are in the same class if and only if their difference is divisible by xy :

$$\bar{f} = \bar{g} \iff f - g \in (xy).$$

This means that all mixed terms containing both x and y vanish in the quotient.

What survives modulo (xy) ?

Write

$$f(x, y) = \sum_{i, j \geq 0} a_{ij} x^i y^j.$$

If $i \geq 1$ and $j \geq 1$, then $x^i y^j$ is divisible by xy , so its class is zero in the quotient.

Therefore the only terms that survive are:

- the constant term,
- pure powers of x : $x, x^2, x^3, \dots$,
- pure powers of y : $y, y^2, y^3, \dots$

So every class has a representative of the form

$$a + p(x) + q(y),$$

where $a \in k$, $p(x) \in xk[x]$, and $q(y) \in yk[y]$.

Equivalently, the whole class is

$$a + p(x) + q(y) + (xy),$$

that is, all polynomials of the form

$$a + p(x) + q(y) + xy h(x, y)$$

with $h(x, y) \in k[x, y]$.

Class of 0

$$\bar{0} = (xy) = \{xy h(x, y) : h(x, y) \in k[x, y]\}.$$

Examples of polynomials in this class are:

$$xy, \quad x^2y, \quad xy^2, \quad 3xy + 7x^4y^3, \dots$$

All of these become zero in the quotient.

Class of $\bar{x}$

$$\bar{x} = x + (xy) = \{x + xy h(x, y) : h(x, y) \in k[x, y]\}.$$

Examples:

$$x, \quad x + xy, \quad x + 7x^2y, \quad x + x^5y^2, \dots$$

All these are equal to $\bar{x}$ in the quotient.

However,

$$x + x^2$$

is *not* in the same class as x , because

$$(x + x^2) - x = x^2,$$

and $x^2 \notin (xy)$.

So

$$\overline{x + x^2} \neq \bar{x}.$$

Class of $\bar{y}$

Similarly,

$$\bar{y} = y + (xy) = \{y + xy h(x, y) : h(x, y) \in k[x, y]\}.$$

Examples:

$$y, \quad y + xy, \quad y + 5xy^2, \quad y + x^3y, \dots$$

Class of $\overline{x + y^2 + 3}$

This class is

$$x + y^2 + 3 + (xy).$$

So it consists of all polynomials of the form

$$x + y^2 + 3 + xy h(x, y).$$

Examples in this class:

$$\begin{aligned} x + y^2 + 3, \\ x + y^2 + 3 + xy, \\ x + y^2 + 3 + 7x^2y^5, \\ x + y^2 + 3 + xy(x + y + 1). \end{aligned}$$

Class of $\overline{x^2 + y^3}$

This class is

$$x^2 + y^3 + (xy).$$

So it contains:

$$x^2 + y^3, \quad x^2 + y^3 + xy, \quad x^2 + y^3 + x^7y^2, \dots$$

But

$$x^2 + y^3 + x$$

is not in this class, because the difference is x , which is not divisible by xy .

1.8.8 8. What are $\bar{x}$ and $\bar{y}$ concretely in $k[x, y]/(xy)$?

They are the residue classes

$$\bar{x} = x + (xy), \quad \bar{y} = y + (xy).$$

So:

- $\bar{x}$ is the class of all polynomials congruent to x modulo (xy) ,
- $\bar{y}$ is the class of all polynomials congruent to y modulo (xy) .

They are not zero, because

$$x \notin (xy), \quad y \notin (xy).$$

But their product is zero:

$$\bar{x}\bar{y} = \overline{xy} = 0.$$

Thus $\bar{x}$ and $\bar{y}$ are nonzero zero divisors.

Also,

$$\bar{x}^2 \neq 0, \quad \bar{y}^2 \neq 0,$$

because

$$x^2 \notin (xy), \quad y^2 \notin (xy).$$

So only mixed terms vanish, not pure powers of x or y .

1.8.9 9. Comparison of the main examples

The quotient always identifies elements that differ by something in the defining ideal:

- in $\mathbb{Z}/n\mathbb{Z}$, one adds multiples of n :

$$\bar{a} = a + n\mathbb{Z};$$

- in $k[x]/(x)$, one adds multiples of x :

$$\bar{a} = a + (x) = \{a + xq(x)\};$$

- in $A[t]/(t)$, one adds multiples of t :

$$\bar{a} = a + (t) = \{a + tq(t)\};$$

- in $k[x, y]/(x)$, one adds multiples of x :

$$g(y) + (x) = \{g(y) + xh(x, y)\};$$

- in $k[x, y]/(xy)$, one adds multiples of xy :

$$a + p(x) + q(y) + (xy) = \{a + p(x) + q(y) + xy h(x, y)\}.$$

Thus:

- in $k[x]/(x)$, the class is determined only by the constant term;
- in $A[t]/(t)$, likewise only the constant term survives;
- in $k[x, y]/(x)$, only the y -part survives;
- in $k[x, y]/(xy)$, the constant term, pure x -terms, and pure y -terms survive, while mixed xy -terms vanish.

1.8.10 10. Final summary

A residue class in any quotient R/I is always

$$f + I = \{f + h : h \in I\}.$$

So:

- in $\mathbb{Z}/n\mathbb{Z}$, two integers are equal if they differ by a multiple of n ;
- in $k[x]/(x)$, two polynomials are equal if they have the same constant term;
- in $A[t]/(t)$, two polynomials are equal if they have the same constant term;
- in $k[x, y]/(x)$, two polynomials are equal if they become equal after setting $x = 0$;
- in $k[x, y]/(xy)$, two polynomials are equal if they differ by a multiple of xy , so mixed terms vanish.

In particular,

$$\bar{x} = 0 \text{ in } k[x]/(x),$$

because $x \in (x)$, but

$$\bar{x} \neq 0 \text{ in } k[x, y]/(xy),$$

because $x \notin (xy)$.