

Chapter 10 Equality constrained minimization

Last update on 2024-05-21 16:50

Table of contents

Equality constrained minimization

Eliminating equality constraints

Newton's method with equality constraints

Infeasible start Newton method

Equality constrained minimization

Eliminating equality constraints

Newton's method with equality constraints

Infeasible start Newton method

Equality constrained minimization

equality constrained minimization problem

$$\begin{array}{ll}\text{minimize} & f(x) \\ \text{subject to} & Ax = b\end{array}$$

- ▶ f convex and twice continuously differentiable
- ▶ $A \in \mathbb{R}^{p \times n}$ with **rank** $A = p$
- ▶ assume optimal value p^* is finite and attained

optimality condition (review)

$$x^* \text{ is optimal} \quad \Longleftrightarrow \quad \begin{array}{l} x^* \in \text{dom } f, \quad Ax^* = b, \\ \text{there exists } \nu^* \text{ such that } \nabla f(x^*) + A^T \nu^* = 0 \end{array}$$

equality constrained quadratic minimization (with $P \in \mathbb{S}_+^n$)

$$\begin{array}{ll} \text{minimize} & (1/2)x^T Px + q^T x + r \\ \text{subject to} & Ax = b \end{array}$$

optimality condition

$$\begin{bmatrix} P & A^T \\ A & 0 \end{bmatrix} \begin{bmatrix} x^* \\ \nu^* \end{bmatrix} = \begin{bmatrix} -q \\ b \end{bmatrix}$$

- ▶ coefficient matrix is called KKT matrix
- ▶ KKT matrix is nonsingular if and only if

$$Ax = 0, \quad x \neq 0 \quad \implies \quad x^T Px > 0$$

- ▶ equivalent condition for nonsingularity

$$P + A^T A \succ 0$$

Equality constrained minimization

Eliminating equality constraints

Newton's method with equality constraints

Infeasible start Newton method

Eliminating equality constraints

represent solutions of $\{x \mid Ax = b\}$ as

$$\{x \mid Ax = b\} = \{Fz + \hat{x} \mid z \in \mathbb{R}^{n-p}\}$$

- ▶ \hat{x} is any particular solution
- ▶ range of $F \in \mathbb{R}^{n \times (n-p)}$ is nullspace of A

reduced or eliminated problem

$$\text{minimize} \quad f(Fz + \hat{x})$$

- ▶ unconstrained problem with variable $z \in \mathbb{R}^{n-p}$
- ▶ from solution z^* , obtain x^* and ν^* as

$$x^* = Fz^* + \hat{x}, \quad \nu^* = -(AA^T)^{-1} A \nabla f(x^*)$$

example optimal allocation with resource constraint

$$\begin{array}{ll}\text{minimize} & f_1(x_1) + \cdots + f_n(x_n) \\ \text{subject to} & x_1 + \cdots + x_n = b\end{array}$$

eliminate $x_n = b - x_1 - \cdots - x_{n-1}$, namely, choose

$$\hat{x} = be_n, \quad F = \begin{bmatrix} I \\ -\mathbf{1}^T \end{bmatrix} \in \mathbb{R}^{n \times (n-1)}$$

reduced problem

$$\text{minimize} \quad f_1(x_1) + \cdots + f_{n-1}(x_{n-1}) + f_n(b - x_1 - \cdots - x_{n-1})$$

Equality constrained minimization

Eliminating equality constraints

Newton's method with equality constraints

Infeasible start Newton method

Newton step

Newton step Δx_{nt} of f at feasible x is given by the solution v of

$$\begin{bmatrix} \nabla^2 f(x) & A^T \\ A & 0 \end{bmatrix} \begin{bmatrix} v \\ w \end{bmatrix} = \begin{bmatrix} -\nabla f(x) \\ 0 \end{bmatrix}$$

interpretations

- ▶ Δx_{nt} solves second order approximation (with variable v)

$$\begin{array}{ll} \text{minimize} & \hat{f}(x+v) = f(x) + \nabla f(x)^T v + (1/2)v^T \nabla^2 f(x)v \\ \text{subject to} & A(x+v) = b \end{array}$$

- ▶ Δx_{nt} equations follow from linearizing optimality conditions

$$\nabla f(x+v) + A^T w \approx \nabla f(x) + \nabla^2 f(x)v + A^T w = 0, \quad A(x+v) = b$$

$$\lambda(x) = (\Delta x_{\text{nt}}^T \nabla^2 f(x) \Delta x_{\text{nt}})^{1/2} = (-\nabla f(x)^T \Delta x_{\text{nt}})^{1/2}$$

interpretations

- ▶ gives an estimate of $f(x) - p^*$ using quadratic approximation \hat{f}

$$f(x) - \inf_{Ay=b} \hat{f}(y) = (1/2)\lambda(x)^2$$

- ▶ directional derivative in Newton direction

$$\left. \frac{d}{dt} f(x + t\Delta x_{\text{nt}}) \right|_{t=0} = -\lambda(x)^2$$

- ▶ in general $\lambda(x) \neq (\nabla f(x)^T \nabla^2 f(x)^{-1} \nabla f(x))^{1/2}$

Newton's method with equality constraints

given starting point $x \in \text{dom } f$ with $Ax = b$, tolerance $\epsilon > 0$

repeat

1. Compute the Newton step and decrement $\Delta x_{\text{nt}}, \lambda(x)$
 2. *Stopping criterion.* **quit** if $\lambda^2/2 \leq \epsilon$
 3. *Line search.* Choose step size t by backtracking line search
 4. *Update.* $x := x + t\Delta x_{\text{nt}}$
-

- ▶ feasible descent method: $x^{(k)}$ feasible and $f(x^{(k+1)}) < f(x^{(k)})$
- ▶ affine invariant

Newton's method for reduced problem

$$\text{minimize } \tilde{f}(z) = f(Fz + \hat{x})$$

- ▶ $z \in \mathbb{R}^{n-p}$ are variables, \hat{x} satisfies $A\hat{x} = b$, range of F is the nullspace of A
- ▶ Newton's method for \tilde{f} starts at $z^{(0)}$, generates iterates $z^{(k)}$

relation to Newton's method with equality constraints

when starting at $x^{(0)} = Fz^{(0)} + \hat{x}$, iterates are

$$x^{(k)} = Fz^{(k)} + \hat{x}$$

hence no separate convergence analysis is needed

Equality constrained minimization

Eliminating equality constraints

Newton's method with equality constraints

Infeasible start Newton method

Newton step at infeasible points

Newton step Δx_{nt} of f at infeasible x is given by the solution of

$$\begin{bmatrix} \nabla^2 f(x) & A^T \\ A & 0 \end{bmatrix} \begin{bmatrix} \Delta x_{\text{nt}} \\ w \end{bmatrix} = - \begin{bmatrix} \nabla f(x) \\ Ax - b \end{bmatrix}$$

interpretation

- ▶ Δx_{nt} equations follow from linearizing optimality conditions

$$\nabla f(x + v) + A^T w \approx \nabla f(x) + \nabla^2 f(x)v + A^T w = 0, \quad A(x + v) = b$$

primal-dual interpretation

- ▶ write optimality condition as $r(y) = 0$ where

$$y = (x, \nu), \quad r(y) = (\nabla f(x) + A^T \nu, Ax - b)$$

- ▶ linearizing $r(y) = 0$ gives

$$r(y + \Delta y) \approx r(y) + Dr(y)\Delta y = 0$$

which is equivalent to

$$\begin{bmatrix} \nabla^2 f(x) & A^T \\ A & 0 \end{bmatrix} \begin{bmatrix} \Delta x_{\text{nt}} \\ \Delta \nu_{\text{nt}} \end{bmatrix} = - \begin{bmatrix} \nabla f(x) + A^T \nu \\ Ax - b \end{bmatrix}$$

same as the above equation with $w = \nu + \Delta \nu_{\text{nt}}$

Infeasible start Newton method

given starting point $x \in \mathbf{dom} f$, ν , tolerance $\epsilon > 0$, $\alpha \in (0, 1/2)$, $\beta \in (0, 1)$

repeat

1. Compute primal and dual Newton steps $\Delta x_{\text{nt}}, \Delta \nu_{\text{nt}}$
2. *Backtracking line search on $\|r\|_2$.* $t := 1$.
while $\|r(x + t\Delta x_{\text{nt}}, \nu + t\Delta \nu_{\text{nt}})\|_2 > (1 - \alpha t)\|r(x, \nu)\|_2$, $t := \beta t$
3. *Update.* $x := x + t\Delta x_{\text{nt}}, \nu := \nu + t\Delta \nu_{\text{nt}}$

until $Ax = b$ and $\|r(x, \nu)\|_2 \leq \epsilon$

- ▶ not a descent method: $f(x^{(k+1)}) > f(x^{(k)})$ is possible
- ▶ directional derivative of $\|r(y)\|_2$ in direction $\Delta y = (\Delta x_{\text{nt}}, \Delta \nu_{\text{nt}})$ is

$$\left. \frac{d}{dt} \|r(y + t\Delta y)\|_2 \right|_{t=0} = -\|r(y)\|_2$$