

Publication status: Preprint has been published in a journal as an article
DOI of the published article: <https://doi.org/10.1590/2236-8906-08/2021>

Topsoil physical properties under a riparian forest in Central Brazil: infiltration and penetration resistance

Johnny Rodrigues de Melo Murta, Gleicon Queiroz de Brito, Sergio Fernandes Mendonça-Filho,
Luiz Felipe Salemi

<https://doi.org/10.1590/2236-8906-62/2020>

Submitted on: 2022-03-10

Posted on: 2022-03-11 (version 1)
(YYYY-MM-DD)

Topsoil physical properties under a riparian forest in Central Brazil: infiltration and penetration resistance

Johnny Rodrigues de Melo Murta¹, Gleicon Queiroz de Brito¹, Sergio Fernandes Mendonça Filho¹
and Luiz Felipe Salemi^{1,2}

Johnny Rodrigues de Melo Murta: <https://orcid.org/0000-0002-5379-5666>

Gleicon Queiroz de Brito: <https://orcid.org/0000-0002-5491-2334>

Sergio Fernandes Mendonça Filho: <https://orcid.org/0000-0002-1264-2866>

Luiz Felipe Salemi: <https://orcid.org/0000-0003-2271-5712>

Título resumido: Topsoil physical properties under a riparian forest in Central Brazil

1. Universidade de Brasília, Faculdade de Planaltina, Núcleo de Estudos e Pesquisas Ambientais e Limnológicas (NEPAL), Villa Nossa Senhora de Fátima 73345-010, Planaltina, Distrito Federal, Brazil
2. Corresponding author: lfsalemi@unb.br

ABSTRACT – (Topsoil physical properties under a riparian forest in Central Brazil: infiltration and penetration resistance). Plant composition, diversity and structure of riparian forests of Central Brazil are well known. However, little is known about soil physical properties under these forests. This knowledge is important as a baseline for biodiversity restoration and ecosystem services that occur in riparian zones. In order to bridge this gap, here we assessed the infiltration capacity and soil penetration resistance in a plinthic soil under gallery forest in Planaltina, Distrito Federal, Brazil. We measured infiltration capacity (Mini-Disk infiltrometer) and soil penetration resistance (Stof penetrometer) following linear transects. The plinthic soil had high infiltration capacity and low penetration resistance. Our infiltration estimate is in the middle range when compared to other permeability studies in tropical forests. Like their counterparts, high biological activity along with the lack of disturbance are the likely explanations for such high topsoil permeability to water.

Keywords: buffer zones, soil conservation, streamside áreas, tropical ecosystems, water management

RESUMO – (Propriedades físicas da camada superficial do solo em florestas ripária no Brasil central: resistência à infiltração e penetração). As florestas de galeria são bem descritas em termos de composição, diversidade e estrutura da vegetação. No entanto, pouco se sabe sobre as propriedades físicas do solo sob essas florestas que são importantes para a restauração de zonas ripárias. A fim de preencher essa lacuna, no presente trabalho, avaliou-se a capacidade de infiltração e a resistência do solo à penetração em um solo plíntico sob floresta ripária em Planaltina, Distrito Federal, Brasil. Mediu-se a capacidade de infiltração (infiltrômetro de Mini-Disk) e a resistência à penetração no solo (penetrômetro de Stolf) seguindo transectos lineares. O solo plíntico apresentou alta capacidade de infiltração e baixa resistência à penetração. Nossa estimativa de infiltração está na faixa intermediária em comparação com outros estudos de permeabilidade em florestas tropicais. Mesmo assim, a mediana da capacidade de infiltração foi superior à intensidade de chuva de alto período de retorno o que descarta a possibilidade de escoamento superficial Hortoniano. Como em outras florestas tropicais, a alta atividade biológica junto com a ausência de perturbação do solo são as razões prováveis para essa alta permeabilidade do solo à água.

Palavras-chave: zonas tampão, conservação do solo, áreas ribeirinhas, ecossistemas tropicais, manejo da água.

Introduction

Gallery forests, also known as riparian forests, occur along small streams in the Cerrado biome of Brazil (Ribeiro and Walter, 1998). These forests are well known in terms of plant species composition, diversity and structure (Felfili 1994, Felfili 1995, Nóbrega et al. 2001, Silva Júnior 2004, Silva Júnior 2005, Moretti et al. 2013, Cabacinha & Fontes 2014, Pio 2018). Generally, they present a high biological diversity. One of the possible explanations for such diversity might be attributed to heterogeneity of soil types over which these forests occur which include both well and poorly drained soils (Haridasan 1998). For example, they might occur on oxisols (latossolos) and histosols (organossolos) (Resck & Silva 1998). Since these soil types present a clearly distinct hydric behaviour, they not only affect plant community but are also expected to influence the ecosystem service of buffering zones that riparian forests play. For instance, riparian forests under histosols might not reduce surface runoff from uplands in the wet season since these soils would be water saturated during this period following the runoff generation (as described in Dunne & Black 1970a,b). So, it is important to understand hydrological functions of these forests under different soil types.

Like many riparian forests elsewhere, such ecosystems are expected to carry out important buffer functions including retaining sediment, nutrients and human-made chemicals such as pesticides. The assumption behind is that riparian soil under these forests present high topsoil permeability which, in turn, diminish or extinguish surface runoff from uplands (Salemi et al. 2011). To date, there is no documentation on basic topsoil permeability for these riparian forests. Understanding such properties under specific physical settings may allow to infer whether they carry out the expected functions they are expected to. Moreover, such information may serve as a baseline which, in turn, may help scientists and managers to predict the impacts of inappropriate riparian occupation and restoration (see Brito et al. 2019; Pereira et al. 2021).

In this context, the objective of the present paper was to characterize a plinthic soil under a riparian forest in terms of infiltration capacity and soil penetration resistance. We expected that, like other tropical rainforests, these riparian forests would present high permeability to water.

Methods

Study área - We selected a riparian forest located in Planaltina, Distrito Federal, Brazil. The Köppen-Geiger climatic classification is Aw, with two markedly defined seasons (dry and wet) (Alvares et al. 2013). Annual rainfall is 1,393 mm with nearly 100% of it falling in the wet season (October to April) [Digite aqui]

(Malaquias et al., 2010). Average temperature is 20.7°C. The soil type is a clay Plintossolo (Plinthosols), according to the Brazilian Soil Classification System (Embrapa, 2018).

Riparian forests in the study area usually have high plant species diversity. Trees density and basal area are around, respectively, 1.300-1.900 individuals ha⁻¹ and 32-38 m² ha⁻¹ (Nóbrega et al. 2001, Silva Júnior 2004, Silva Júnior 2005). Common species found in these forests are *Protium spruceanum* (Benth.) Engl. (Burseraceae), *Copaifera langsdorffii* Desf. (Fabaceae), *Tapirira guianensis* Aubl. (Anacardiaceae), *Inga alba* (Sw.) Willd. (Fabaceae), *Cheiloclinium cognatum* (Miers) A.C.Sm. (Celastraceae), *Pseudolmedia guaranitica* Hasler (Moraceae), *Euterpe edulis* Mart. (Arecaceae) and *Talauma ovata* A. St.-Hill. (Magnoliaceae) (Nóbrega et al. 2001, Silva Júnior, 2004, Silva Júnior, 2005).

Variables and sampling design - We measure both infiltration capacity and soil resistance to penetration using, respectively, Mini-Disk infiltrometer (Decagon) and Stof Impact Penetrometer (Kamaq) during the dry season of 2018.

The Mini-disk infiltrometer is a two-chamber device in which the Mariotte principle is applied to control soil suction. Such equipment uses the analytical solution proposed by Zhang (1997). Similar to Ghimire et al. (2013), to increase the contact area between the infiltrometer and the soil, we carried out two steps: (i) we carefully removed the litter layer and (ii) we used a thin (< 1 mm) of fine sand. These procedures ensure optimum contact between the infiltrometer and the soil. To capture the highest range of pores, we set a suction pressure of 0 cm which corresponds to Ko, that is, the saturated hydraulic conductivity (Reichardt & Timm 2019). The rates of water discharge through the Mini-Disk, as inferred from changes in the water levels in the storage chamber, were recorded until steady-state flow was reached. Infiltration was measured using three linear transects randomly distributed within the riparian forest (figure 2). To randomize linear transects, we used a randomizer software (randomizer.org) which selected 3 out of 50 grid transects. Ten infiltration samples were taken from each of these transects. Transects were established in parallel to stream channel. When measuring infiltration, it is important to minimize the effect of soil moisture on infiltration measurements since substantial variation can occur from one month to the next (see Pereira et al. 2021). Thus, we carried out two steps: (i) measurements were made in August (dry season) when soil moisture in the region is minimal (Oliveira et al. 2005), and (ii) we set 0 cm of suction pressure in the Mariotte bottle.

In order to predict the effectiveness of the riparian forest in accommodating storms of the region, we compared our median of infiltration capacity with 5-minutes rainfall intensities of different return periods (100, 50 and 25-years) available in Souza (2014).

Soil penetration resistance was measured for ten times following the linear transects described above (figure 2). We arbitrarily selected four impacts to determine soil penetration resistance. This

[Digite aqui]

include the first impact which is the inherent weight of the equipment followed by the other three impacts which resulted from the impact of a standard 2 kg weight into the soil. To minimize the effect of soil moisture on our penetration measurements, we carried out the measurements in the dry season (August) when soil moisture is minimal (Oliveira et al. 2005).

Data analysis - Sample sufficiency was verified for infiltration capacity using a simple plot of mean and standard deviation versus number of samples. An adequate sample size was considered met when central tendency statistics stabilized for at least three times consecutively (Salemi et al. 2020).

Residuals of soil penetration resistance were subjected to Shapiro-Wilk test which indicated a lack of gaussian distribution. Therefore, we used Kruskal-Wallis test followed by the post hoc Dunn test to check significant differences between impacts. Statistical analyses were carried out in PAST software at $\alpha < 0.05$.

Results and Discussion

The number of samples used for infiltration was considered appropriate given the relative stabilization of the median statistics (figure 3). The median infiltration capacity was 893 mm hr⁻¹ (figure 4). Mean (\pm standard deviation) of soil penetration resistance for all four impacts were, respectively, as follows: 0.63 ± 0.21 MPa, 0.66 ± 0.32 MPa, 0.74 ± 0.22 MPa and 0.76 ± 0.22 MPa (figure 5). There was a significant increase in soil penetration resistance with depth. Such increase has previously been documented for a forest within the Brazilian savanna (see Pereira et al. 2021). Our resistance to penetration was lower compared to a nearby savanna (regionally known as “Cerrado típico”) in the same area (Murta et al. 2021). Such difference of soil penetration resistance in the riparian forest compared to a savanna might be attributed to the higher and more continuous addition of litter year-round in the forest compared to the savanna (Costa et al. 2020). For example, Aquino et al. (2016) measured an average of 6 ton ha⁻¹ year⁻¹ in a riparian forest whereas Valenti et al. (2008) found 0.6 ton ha⁻¹ year⁻¹ in a savanna.

We found infiltration capacity of the plinthic soil under riparian forest to be in the middle range compared to estimates carried out in other tropical forests such as the Amazon and Atlantic Forest (table 1). Generally, high soil biological activity (e.g. continuous litter production, root growth and decay and soil fauna) along with the lack of disturbance (e.g. human and animal trampling, use of heavy machinery) leads to high infiltration in forest soils (Bruijnzeel 1990, Salemi et al. 2011).

Our median infiltration capacity estimate was higher compared to the rain intensity of various storms in the region (figure 4). Thus, we rule out the occurrence of infiltration excess overland-flow in this forest. However, our penetration resistance indicated a likely reduction of these permeability with depth which was, to some extent, expected due the presence of the plinthic horizon. A previous [Digite aqui]

study on plinthic soils showed the formation of a perched water table above such layer (Moraes et al. 2006). Furthermore, riparian forests, like the one studied here, generally might occur in shallow water table zones (Ribeiro & Walter 2008). In case the water table (perched or not) intersects the soil surface, both return-flow and overland-flow are triggered (Dunne & Black 1970a, b, Elsenbeer & Vertessy 2000). Thus, saturation overland-flow can occur. In this case, unlike the generally expected role of surface runoff sinks, these zones might act as surface runoff sources instead. More studies are needed to clarify percolation capacities of these soil and their implications for runoff processes.

Acknowledgments

We thank Maurício Rigon Hoffmann of Elo Florestal Inkora farm, for his permission to work on his rural property. This study was partially financed by the Coordenação de Aperfeiçoamento de Pessoal de Nível Superior - Brasil (CAPES) - Finance Code - 001.

Author Contributions

Johnny Rodrigues de Melo Murta: sampling design, sampling, data analyses, manuscript preparation.

Gleicon Queiroz de Brito: sampling, manuscript preparation.

Sergio Fernandes Mendonça Filho: sampling, manuscript preparation.

Luiz Felipe Salemi: sampling design, sampling, data analyses, manuscript preparation.

Conflict of interest

The authors declare no conflict of interest.

Literatura citada

Alvares, C. A., Stape, J. L., Sentelhas, P. C., de Moraes Gonçalves, J. L. & Sparovek, G. 2013.

Köppen's climate classification map for Brazil. *Meteorologische Zeitschrift* 22(6): 711-728.

Aquino, P.S.R., Nappo, M.E., Rodrigues, M.S., Pereira, I.M., Matricardi, E.A.T. & Pelá, G.M.

2016. Análise espacial da produtividade de serapilheira em uma mata de galeria. *Ciência Florestal* 26(2): 489-500.

Brito, G.Q., Sampaio, J.A.G., Luiz, G.P., Melo, A.C.A., Simões, V.P., Martins, P.R., Neves, G., Murta, J.R.M., Mendonça Filho, S.F., Couto Júnior, A.F., Vieira, L.C.G., Teixeira, A.C.O.

[Digite aqui]

- & Salemi, L.F.** 2019. Efeitos da conversão de floresta riparia em pastagem sobre as propriedades físico-hídricas do solo. *Revista Espaço e Geografia* 22: 75-91.
- Bruijnzeel, L.A.** 1990. Hydrology of moist tropical forests and effects of conversion: a state of knowledge review. Amsterdam, International Hydrology Programme.
- Cabacinha, C.D. & Fontes, M.A.L.** 2014. Caracterização florística e estrutural de fragments de matas de galeria da bacia do Alto Araguaia. *Ciência Florestal* 24(2): 379-390.
- Costa, A.N., Souza, J.R., Alves, K.M., Penna-Oliveira, A., Paula-Silva, G., Becker, I.S., Marinho-Vieira, K., Bonfim, A.L., Bartimachi, A. & Vieira-Neto, E.H.M.** 2020. Linking the spatial temporal variation of litterfall to standing vegetation biomass in Brazilian savannas. *Journal of Plant Ecology* 13(5): 517-524.
- Dunne, T. & Black, R.** 1970a. An investigation of runoff production in permeable soils. *Water Resources Research* 6(2): 179-191.
- Dunne, T. & Black, R.** 1970b. Partial are contribution to storm runoff in a small New England Watershed. *Water Resources Research* 6(5): 1296-1311.
- Elsenbeer, H. & Vertessy, R.A.** 2000. Stormflow Generation and flowpaths characteristics in an Amazoninan rainforest catchment. *Hydrological Processes*, 14(14): 2367-2381.
- Felfili, J.M.** 1994. Floristic composition and phytosociology of the gallery forest alongside Gama Stream in Brasília, DF, Brazil. *Revista Brasileira de Botânica* 7(1): 1-11.
- Felfili, J.M.** 1995. Diversity, strcuture and dynamics of gallery forest in central Brazil. *Vegetatio* 117: 1-15.
- Ghimire, C.P., Bonell, M., Bruijnzeel, L.A., Coles, N.A. & Lubczynski, M.W.** 2013. Reforesting severelydegraded grassland in the Lesser Himalaya of Nepal: effects on soil hydraulic donuctivity and overland-flow production. *Journal of Geophysical Research: Earth Surface* 118: 2528-2545.
- Haridasan, M.** 1998. Solos de matas de galeria e nutrição mineral de espécies arbóreas em condições naturais. *In: Ribeiro, J.F. (ed.). Cerrado: matas de galeria. EMBRAPA-CPAC, Planaltina*, pp. 17-28.
- Lozano-Baez, S.E., Cooper, M., Ferraz, S.F.B., Rodrigues, R.R., Lassabatere, L., Castellini, M. & Di Prima, S.** 2020. Assessing water infiltration and soil water repellency in Brazilian Atlantic Forest Soils. *Applied Sciences* 10: 1-14.
- Malaquias, J. V., Silva, F. A. M. & Evangelista, B. A.** 2010. Precipitação pluviométrica em Planaltina, DF: análise de dados da estação principal da Embrapa Cerrados, 1974 a 2003. Planaltina, Embrapa Cerrados.
- Moraes, J.M., Schuler, A.E., Dunne, T., Figueiredo, R.O. & Victoria, R.L.** 2006. Water storage and runoff processes in plinthic soils under forest and pasture in Eastern Amazonia. *Hydrological Processes* 20: 2509-2526.

- Moretti, M.S., Ribeiro, E.S., Cabral, R.S.O., Barros, L.S. & Soares, G.S.** 2013. Levantamento fitossociológico de mata de galeria e cerradão no município de Poconé, Mato Grosso. *Revista Eletrônica Geoaraguaia*, 1: 193-208.
- Murta, J.R.M., Brito, G.Q., Mendonça Filho, S.F., Hoffmann, M.R. & Salemi, L.F.** 2020. Understanding the effect of an agroforestry system with high litter input on topsoil permeability. *Soil Use and Management*, 37(4): 802-809.
- Oliveira, R. S., Bezerra, L., Davidson, E. A., Pinto, F., Klink, C. A., Nepstad, D. C. & Moreira, A.** 2005. Deep root function in soil water dynamics in cerrado savannas of central Brazil. *Functional Ecology* 19(4): 574-581.
- Pereira, L.C., Balbinot, L., Matus, G.N., Dias, H.C.T. & Tonello, K.C.** 2021. Aspects of forest restoration and hydrology: linking passive restoration and soil-water recovery in Brazilian Cerrado. *Journal of Forestry Research* (2021).
- Pio, A.D.** 2018. Estrutura e diversidade de florestas estacionais em áreas do domínio do Cerrado. Dissertação de Mestrado, Instituto Federal de Goiano, Urutaí.
- Reichardt, K. & Timm, L.C.** 2019. Soil, plant and atmosphere. Springer.
- Resck, D.V.S. & Silva, J.E.** 1998. A importância da mata de galeria no ciclo hidrológico de uma bacia hidrográfica. *In: Ribeiro, J.F. (ed.). Cerrado: matas de galeria. EMBRAPA-CPAC, Planaltina*, pp. 29-50.
- Ribeiro, J. F. & Walter, B.M.T.** 1998. Fitofisionomias do bioma Cerrado. *In: Sano, S. M.; Almeida, S. P. (Ed.) Cerrado: ambiente e flora. EMBRAPA-CPAC, Planaltina*. pp. 89-168.
- Salemi, L.F., Groppo, J.D., Trevisan, R., Moraes, J.M., Lima, W.P. & Martinelli, L.A.** 2011. Aspectos hidrológicos da recuperação florestal de áreas de preservação permanente ao longo dos corpos de água. *Revista do Instituto Florestal* 23(1): 69-80.
- Salemi, L.F., Fernandes, R.P., Silva, R.W.C., Garcia, L.G., Moraes, J.M., Groppo, J.D. & Martinelli, L.A.** 2020. Soil hydraulic properties: a simple and practical approach to estimate the number of samples. *Eurasian Journal of Soil Science*, 9(1): 18-23.
- Scheffler, R., Neill, C., Krusche, A.V. & Elsenbeer, H.** 2011. Soil hydraulic response to land-use change associated with the recente soybean expansion at the Amazon agricultural frontier. *Agriculture, Ecosystems & Environment* 144(1): 281-289.
- Silva Júnior, M.C.** 2004. Fitossociologia e estrutura diamétrica da mata de galeria do Taquara, na Reserva Ecológica do IBGE, DF. *Revista Árvore* 28(3): 419-428.
- Silva Júnior, M.C.** 2005. Fitossociologia e estrutura diamétrica na Mata de Galeria do Pitoco, na Reserva Ecológica do IBGE, DF. *Cerne* 11(2): 147-158.
- Souza, F.P.** 2014. Monitoramento e modelagem hidrológica da sub-bacia do lago Paranoá - Brasília/DF e avaliação de bacia de retenção. Dissertação de Mestrado, Universidade de Brasília, Brasília.

- Valenti, M.W., Cianciaruso, M.V. & Batalha, M.A.** 2008. Seasonality of litterfall and leaf decomposition in a cerrado site. *Brazilian Journal of Biology* 68(3): 459-465.
- Zhang, R.** 1997. Determination of soil sorptivity and hydraulic conductivity from disk infiltrometer. *Soil Science Society of America Journal* 61(4): 1024-1030.
- Zimmermann, B., Elsenbeer, H. & De Moraes, J.M.** 2006. The influence of land-use changes on soil hydraulic properties: implication for runoff generation. *Forest Ecology and Management* 222: 29-38.

Editora Associada: Ana Paula Souza Caetano

Submissão: 07/06/2020

Aceito: 25/02/2022




Figure 1. Study area located in Brazil (a), Distrito Federal State (b), Planaltina (c). Photograph shows the forest structure near the floor (d).




Figure 2. Design used for soil sampling in the riparian forest in Planaltina, Brasília, Distrito Federal State, Brazil. White lines represent infiltration transects. White arrows represent soil penetration resistance sampling points.




Figure 3. Box-Jitter plot showing infiltration capacity as function of the number of samples. Horizontal lines within the box represent the median. x represent the mean. Horizontal boundaries of the boxes represent the first and third quatiles. Tips of the vertical lines represent maximum (upper) and minimum (lower) values.




Figure 4. Box-plot of infiltration capacity of a plinthic soil under riparian forest. Horizontal line within the box represent the median. x represent the mean. Horizontal boundaries of the boxes represent the first and third quartiles. Tips of the vertical lines represent the maximum (upper) and minimum (lower) values. Notches indicate the 95% confidence interval of the median. Dotted lines 1, 2 and 3 indicate rainfall intensities with, respectively, 100 (300 mm hr⁻¹), 50 (260 mm hr⁻¹), and 25 (220 mm hr⁻¹) years return periods available in Souza (2014).




Figure 5. Soil penetration resistance (cm) in a plinthic soil under riparian forest. Bars represent the mean. Error bars represent the standard deviation.

Table 1. Central tendency statistics of infiltration capacity found in different studies under tropical forests.

Infiltration capacity (mm h ⁻¹)	Method	Soil type	Number of samples	Vegetation type	Location	Reference
550*	Guelph permeameter	Plinthic soil	21	Tropical rain forest	Pará, Brazil	Moraes et al. (2006)
1533	Hood infiltrometer	Ultisol	25	Open tropical rain forest	Rondônia, Brazil	Zimmermann et al. (2006)
1200	Hood infiltrometer	Oxisol	75	Tropical rain forest	Mato Grosso, Brazil	Scheffler et al. (2011)
557	Mini-Disk	Gleysol	10	Tropical riparian forest	Mato Grosso, Brazil	Brito et al. (2019)
200	Single disk	Ultisol	42	Semideciduous tropical forest	São Paulo, Brazil	Lozano-Baez et al. (2020)
893	Mini-Disk	Plinthsol	30	Tropical riparian forest	Distrito Federal, Brazil	Present study

*saturated hydraulic conductivity at 0.15 m


CARTA DE AUTORIZAÇÃO DE PUBLICAÇÃO NO PORTAL DE PREPRINTS DO SCIELO

Ao Comitê Editorial de HOEHNEA

Declaro, em meu próprio nome e nos dos demais Autores, que concordo com a publicação do Artigo Aceito pelo Corpo Editorial de Hoehnea, intitulado “Topsoil physical properties under a riparian forest in Central Brazil: infiltration and penetration resistance” de autoria de Johnny Rodrigues de Melo Murta, Gleicon Queiroz de Brito, Sergio Fernandes Mendonça Filho e Luiz Felipe Salemi no Portal de Preprints do SciELO Brasil (<https://preprints.scielo.org/index.php/scielo/preprints/section/biological>).

Declaro, ainda, que o referido artigo é original, sendo que o conteúdo não foi ou não está sendo considerado para publicação em outro periódico, quer seja no formato impresso e/ou eletrônico.

Brasília, 05 de março de 2022.



Luiz Felipe Salemi

This preprint was submitted under the following conditions:

- The authors declare that they are aware that they are solely responsible for the content of the preprint and that the deposit in SciELO Preprints does not mean any commitment on the part of SciELO, except its preservation and dissemination.
- The authors declare that the necessary Terms of Free and Informed Consent of participants or patients in the research were obtained and are described in the manuscript, when applicable.
- The authors declare that the preparation of the manuscript followed the ethical norms of scientific communication.
- The authors declare that the data, applications, and other content underlying the manuscript are referenced.
- The deposited manuscript is in PDF format.
- The authors declare that the research that originated the manuscript followed good ethical practices and that the necessary approvals from research ethics committees, when applicable, are described in the manuscript.
- The authors declare that once a manuscript is posted on the SciELO Preprints server, it can only be taken down on request to the SciELO Preprints server Editorial Secretariat, who will post a retraction notice in its place.
- The authors agree that the approved manuscript will be made available under a [Creative Commons CC-BY](#) license.
- The submitting author declares that the contributions of all authors and conflict of interest statement are included explicitly and in specific sections of the manuscript.
- The authors declare that the manuscript was not deposited and/or previously made available on another preprint server or published by a journal.
- If the manuscript is being reviewed or being prepared for publishing but not yet published by a journal, the authors declare that they have received authorization from the journal to make this deposit.
- The submitting author declares that all authors of the manuscript agree with the submission to SciELO Preprints.