

Publication status: Preprint has been published in a journal as an article
DOI of the published article: <https://doi.org/10.1590/pboci.2020.152>

Characterization of dental surgeons of Pernambuco in the COVID-19 pandemic context: preliminary data.

Gabriela da Silveira Gaspar, Nilcema Figueiredo, Edson Hilan Gomes de Lucena, Cindy Avani
Silva Ceissler, Ronald Pereira Cavalcanti, Paulo Sávio Angeiras de Goes

<https://doi.org/10.1590/SciELOPreprints.832>

Submitted on: 2020-06-22

Posted on: 2020-06-23 (version 1)
(YYYY-MM-DD)

Characterization of dental surgeons of Pernambuco in the COVID-19 pandemic context: preliminary data.

Gabriela da Silveira Gaspar (author of correspondence)

(81) 9.9147-3749/gabrielasgaspar@gmail.com

Public Health Center, Academic Center of Vitória, Federal University of Pernambuco, Vitória de Santo Antão, Pernambuco, Brazil.

ORCID: 0000-0001-8391-7524

Nilcema Figueiredo

(81)9.9975-1015/nilcema.figueiredo@ufpe.br

Academic Area of Social Medicine, Center of Medical Sciences, Federal University of Pernambuco, Recife, Pernambuco, Brazil.

ORCID: 0000-0001-6181-8728

Edson Hilan Gomes de Lucena

(81) 9.9770-0510/ehglucena@gmail.com

Department of Clinical and Social Dentistry, Center of Health Sciences, Federal University of Paraíba, João Pessoa, Paraíba, Brazil.

ORCID: 0000-0003-3431-115X

Cindy Avani Silva Ceissler

(81) 9.9875-4900/ cindyceissler@gmail.com

Federal University of Pernambuco, Oral health observatory, Recife, Pernambuco, Brazil.

ORCID: 0000-0002-4077-9245

Ronald Pereira Cavalcanti

(81) 9.9949-1632/ronald.cavalcanti@ufpe.br

Public Health Center, Academic Center of Vitória, Federal University of Pernambuco, Vitória de Santo Antão, Pernambuco, Brazil.

ORCID: 0000-0002-4088-0765

Paulo Sávio Angeiras de Goes

(81) 9.9173-5200/paulosaviogoes@gmail.com

Postgraduate Program in Child and Adolescent Health, Federal University of Pernambuco, Recife, Pernambuco, Brazil.

ORCID: 0000-0002-6708-0450

Author's contributions:

GSG contributed to the study design, data collection, analysis and interpretation and writing of the manuscript, NF contributed to the data collection and analysis and writing of the manuscript, EHGL contributed to the data collection and interpretation and writing of the manuscript, CASC performed data analysis and interpretation and wrote the manuscript, RPC collected and interpreted the data and wrote the manuscript, PSAG contributed to the study design, data analysis and interpretation and to the critical review of the manuscript.

ABSTRACT

This study aimed to present preliminary research results about the impact of COVID-19 on the professional practice of dentists in Pernambuco. This is a cross-sectional, descriptive, and exploratory study, whose population was composed of dental surgeons with active enrollment in the Regional Dentistry Council of Pernambuco. Data collection was performed using an electronic form and included characterization of professionals (gender, age, time since graduation, marital status, family income and field of work) and health status (vaccination schedule, presence of comorbidities, biosafety knowledge and testing for COVID-19). Preliminary data correspond to the first week of collection, which were analyzed from the frequency, proportions, and measures of central tendency distributions. Of the 363 dental surgeons, for the field of work, 38.6% work in both the public and private sectors. Comorbidities related to the worsening of COVID-19 were identified in 35.0% of participants, 24.5% are not up to date with influenza and hepatitis vaccines, and 79.3% have not been tested for COVID-19. Regarding the biosafety instructions for COVID-19, 30.7% received no training. It is necessary to immunize dental surgeons to prevent from immunological diseases and expansion of the testing capacity for COVID-19, especially for professionals belonging to the risk group. In addition to guaranteeing the offer of qualification courses on biosafety, which is essential for the safe resumption of activities.

Keywords: Oral Health; Dentistry; Dentists; Prevention and Control; COVID-19

INTRODUCTION

The first cases of SARS-CoV-2 infection (Severe Acute Respiratory Syndrome Coronavirus 2), the causative agent of COVID-19 (Coronavirus Disease), appeared in the city of Wuhan, Hubei province, China (Chen N et al, 2020). The disease spread rapidly through Chinese territory and, subsequently, cases were identified in other countries such as Japan, Thailand, and the Republic of Korea. With the significant global expansion of cases, the World Health Organization (WHO) declared the disease as a pandemic in March 2020 (WHO, 2020).

The first case of COVID-19 in Brazil was registered on February 26, 2020. And, a month later, community transmission was acknowledged throughout the territory. In the context of a public health emergency of national importance, the Secretariat of Health Surveillance of the Ministry of Health (SVS) adapted the Surveillance System for Acute Respiratory Syndromes, aiming to guide the National Health Surveillance System for simultaneous circulation of coronavirus, influenza and other respiratory viruses (Brasil, 2020).

According to the Epidemiological Bulletin number 18, of the Ministry of Health, of June 15, 2020, in Brazil there were 888 thousand cases confirmed by COVID-19 and 43,959 deaths (Brasil, 2020). However, these data do not reflect the real prevalence of COVID-19 in the population, since, in the country, the tests are almost exclusively performed on people with symptoms, especially in the most severe cases, that is, it fails to diagnose the great majority of those contaminated, who present mild symptoms, or even, are asymptomatic (HALLAL et al., 2020).

The state of Pernambuco, according to the Epidemiological Bulletin of the State Health Secretariat, 06/16/2020, registered 46,427 confirmed cases, of which 17,405 were serious and 29,022 were mild, placing it in second place in the Northeast Region, with the highest number of deaths of the disease. So far, confirmed severe cases of the disease are distributed in 168 (90.8%) municipalities in Pernambuco, in addition to the Fernando de Noronha Archipelago (PERNAMBUCO, 2020).

Once the virus is present within oral tissue cells, this causes the aerosol generated during dental procedures to be potentially contaminated. Thus, dental practice, although indispensable for maintaining general health, has been considered to be at high risk of contagion (PENG et al, 2020). The fear of contamination has led some people to avoid attending the dental service, considered after the reopening of health services, which tends to increase the number of emergencies or more complex procedures. Another effect to be analyzed will consider possible needs for adaptations in the dental office to reduce or eliminate the viruses present in aerosols and on the surfaces of the service environment, and also, dentists need to adapt to this new routine, preserving their health and that of the patient.

Due to the profound impacts of the pandemic on the provision of dental services, investigating mechanisms to adapt this practice may contribute to guarantee the provision of this essential service to society. As it is a topic of national and international interest, the results obtained with this study intend to generate evidence about the risk that dental activity presents during the pandemic period for professionals in the area.

Thus, the objective is to quantify this risk with protective measures, identification of dentists with comorbidities, testing for COVID-19 and through information received by the professionals.

MATERIAL AND METHODS

The present study has a quantitative, cross-sectional, descriptive, and exploratory approach. For the communication, respondents were considered in the first eight days of the survey, from June 9th to 16th, 2020.

The population consisted of all 9,793 dental surgeons in the state of Pernambuco with active enrollment in Regional Dentistry Council of Pernambuco (CRO-PE). All dental surgeons with active enrollment in the Regional Dentistry Council of Pernambuco were included until the beginning of the research. Those who did not have an e-mail, who showed any inconsistency in the e-mail or telephone number, who did not have the Whatsapp messaging app, which did not make access to the form possible, and professionals with unsigned or suspended registrations were excluded.

The instrument for data collection was developed and applied in an electronic form, using Google® Forms. These were sent through the official email list of the CRO-PE, through the message transmission list of the Whatsapp messaging app, via text messaging type messages and posted on the social media networks of that entity. This protocol was established to minimize losses and enhance results in order to have a high response rate and due to the need for social distancing, which would make it impossible to collect data face-to-face.

The characterization of the investigated professionals considered the socioeconomic profile: gender, age, time since graduation, marital status, family income and field of work (public and/or private). To analyze the health situation, the variables: updated vaccination schedule, knowledge about biosafety for COVID-19, testing for COVID-19 and presence of comorbidities were analyzed.

The data were exported to the Excel program, version 2000 (Microsoft Corp, United States), and subsequently analyzed from the distribution of frequencies, percentages, mean and median. To guarantee the respondent's identity, the professional's registration number provided by the CRO-PE and the one registered by the respondent in the questionnaire were crossed. However, to guarantee the confidentiality of the participant, all participants were identified by a number, thus guaranteeing confidentiality, and making it impossible for the professional to be recognized.

The research followed all the precepts of the Resolution 466/12 of the National Health Council, related to research in human beings, and was approved by the Research Ethics Committee of the Academic Center of Vitória (CEP/CAV) of the Federal University of Pernambuco, under CAAE n°. 32362120.3.0000.9430, and report n°. 4.068.546.

RESULTS

One week after the start, the survey received responses from 363 dental surgeons. 72.5% of these (n = 263) are female, 58.1% (n = 211) are married or in a common-law marriage and 33.9% (n = 123) are single. As for family income, 66.1% (n = 240) is equal to or greater than five minimum wages, 23.1% (n = 84) is 3 to 4 minimum wages and 10.7% (n = 39) corresponds to a maximum of 2 minimum wages. On average, the participants have 14.6 years of graduation. Regarding postgraduate education, 55.9% (n = 203) have specialization or residency as the highest degree and 22% (n = 80) do not have postgraduate degrees. 38.6% (n = 140) work professionally in both public and private sectors.

Analyzing the health situation of these professionals, it was identified that 35.0% (n = 127) has some type of comorbidities related to the worsening of the symptoms of COVID-19. Among the main ones, arterial hypertension (14.3%), pulmonary disease, respiratory failure or asthma (9.6%), obesity (8.8%), and diabetes (4.1%) stand out. Regarding the vaccination schedule, 24.5% are not up to date with one of the influenza or hepatitis vaccines, 79.3% (n = 288) of the dentists were not tested for COVID-19, among those who performed testing, 70.6% (n = 53) had negative results. As for training for clinical care, 30.3% (n = 110) of the respondents said they had not received any instruction through training or a course on biosafety for COVID-19 (table1).

Table 1 - Characterization of dental surgeons according to socioeconomic profile and health situation, Pernambuco, 2020.

		Variables	Frequency	Percentage
Socioeconomic Profile	Gender	Female	263	72,5%
		Male	100	27,5%
	Marital status	Married/Common-law marriage	211	50,1%
		Divorced	25	6,9%
		Single	124	34,1%
		Widow	3	0,8%
	Family Income (consider the minimum wage in the amount of R\$ 1,045.00):	Less than 1 minimum wage	3	0,8%
		1 to 2 minimum wages	36	9,9%
		3 to 4 minimum wages	84	23,1%
		Equal to or greater than 5 minimum wages	240	66,1%
	Mean of graduation time (in years)		14,6	
	Highest degree, in case of postgraduate education	Doctorate	39	10,7%
		Masters	41	11,3%
		Specilization/residency	203	55,9%
		No postgraduate degree	80	22,0%
	Field of work you carry out professional activities	Only in the private sector	123	33,9%
		Only in the public sector	92	25,3%
		In both (private and public sectors)	140	38,6%
		Does not practice clinical dentistry	8	2,2%

Health situation	Presence of comorbidity	Yes	127	35,0%
		No, none	236	65,0%
	Testing	Was tested for COVID-19 and the result was negative	53	14,6%
		Was tested for COVID-19 and the result was positive	22	6,1%
		Was not tested for COVID-19	288	79,3%
	Updated vaccination schedule	No, for both	18	5,0%
		Yes, for both (hepatitis and influenza)	274	75,5%
		Yes, only for influenza (2020 vaccine)	25	6,9%
		Yes, only for hepatitis B	46	12,7%
	Received instructions on biosafety for COVID-19	Não	110	30,3%
		Sim	253	69,7%

DISCUSSION

The present work is the first report that seeks to trace the impact of the COVID-19 pandemic on the oral health team on different aspects; including the establishment of a situational picture of the health conditions of dentists, which are the financial and emotional impact, and change in the work process. Their findings will help to reflect on the necessary changes to be adopted by this professional group in the post-pandemic period.

The research findings point to a predominance of women in accordance with the literature on the participation of this gender in Dentistry (MORITA; HADDAD; ARAÚJO, 2010). Therefore, it is important to reflect on gender issues within the discussions of the impact of COVID-19 and its economic and social effects for Dentistry, considering the greater risk of women becoming infected because they are the majority, but also the consequences of informal occupational bonds, greater economic vulnerability, greater burden of responsibility for family care within the domestic environment and little occupation in the representation spaces and leadership (WENHAM; SMITH; MORGAN, 2020).

The National Oral Health Policy - Smiling Brazil ("Brasil Sorridente") opened jobs in the public service (CAYETANO et al., 2019) and thus brought dentist's aspiration for occupation in the public sector, aiming at financial stability, easing the uncertainty of practicing in the private sector. Data from 2014 indicate that one third of

the country's dental surgeons work in the Unified Health System - SUS (CASCAES; DOTTO; BOMFIM, 2018), however they also seek to work in the private sector (CARDOSO; VIEIRA; MACHADO, 2010), corroborating the results of this study, in which most respondents reported working in both sectors. Thus, discussions about the transformations in the routine of dental work imposed by the pandemic must consider this double relation in the economy fields, since it is the same professional practicing in different workplaces.

Health professionals are immersed in a background where occupational risk is part of their daily lives. These risks and, consequently, the fear of becoming ill and/or its realization directly interfere in the performance of workers with repercussions on their quality of life (NORONHA, 2013). A study carried out with dental surgeons in India shows that only 10% are confident with the return of post-pandemic care. Among the factors considered for those who reported being hesitant are their personal health and financial issues (KINARIWALA et al., 2020).

Personal health is an important issue to be considered in the face of the risk of contamination, since the findings of this research identified an important number of professionals with chronic diseases that are risk factors for COVID-19. This health condition can keep them away from work for fear of getting worse in case of contamination. The literature reports the association between the presence of comorbidities and the worsening of the outcomes of patients with the infection of the new coronavirus, in addition to pointing out that the increase in the number of these comorbidities increases the severity of the outcomes (YE et al., 2020).

It was observed that almost one third of the professionals did not receive any instructions through training or a course on biosafety for COVID-19, which can result in contamination. An alternative to fill this gap may be the use of virtual courses, being a possible educational tool in periods of social isolation (QUIJANO-ESCATTE et al., 2020).

To minimize occupational risk, the vaccination schedule needs to be updated, since immunization promotes protection for professionals and users. In addition to being safe and efficient, it is available in the Unified Health System (SUS). There is evidence that points to the effectiveness of the influenza vaccine in reducing the incidence of cases and absence from work among health workers (IMAI et al., 2018). For hepatitis B, similar findings were identified in dental surgeons in the public health system in São Paulo (GARBIN et al., 2019).

The small amount of testing that is being carried out in all regions of Brazil is reflected in the preliminary results of this study, where less than a quarter of the professionals took the test for COVID-19. The expansion of the testing capacity of the population and health professionals is an imperative measure to generate the information and indicators necessary to support the best strategies to develop projections that can reflect the social and demographic diversity of the society, allowing a safe return to normal and minimizing the possibility of subsequent epidemic outbreaks (BARRETO et al., 2020).

This study has the limitation of presenting preliminary data, however, it is possible to reflect on the health situation of dental surgeons in Pernambuco in the face of the greatest public health crisis of this century. It is noteworthy that data collection will continue to be carried out and with a larger number of the sample, analytical measures can be evaluated and inferences about the financial and emotional impact, and change in the work process, in addition to enabling the expansion of the analysis to other categories of oral health professionals, such as oral health assistants and technicians.

CONCLUSION

The performance of the dental surgeon is essential for people's health and should be based on safe clinical practice for the oral health team and users of health services. The new coronavirus has brought about new demands for dentistry, both immediately and long term, and for its solution, partnerships between professionals, professional entities, government and universities are necessary to propose, implement and monitor actions that subsidize them.

The research of the impact of COVID-19 in the professional practice of dental surgeons presents, as preliminary results, the need for immunization of these professionals for immunological diseases, the expansion of the testing capacity for COVID-19 to improve the accuracy of the actions to control the disease, including special attention to professionals belonging to the risk group. Another verified demand was qualification on biosafety, which is fundamental for the safe resumption of activities. These results are expected to be used for the planning and reorganization of health services in the public and private sectors.

ACKNOWLEDGEMENTS

We thank the support of the Regional Council of Dentistry and the Pernambuco Health Secretariat through the State Coordination of Oral Health.

CONFLICT OF INTERESTS

All authors have declared that there is no conflict of interest related to the manuscript.

REFERENCES

Aquilante AG, Aciole GG. Oral health care after the National Policy on Oral Health - "Smiling Brazil": a case study. *Ciêns Saúde Coletiva* 2015 Jan;20(1):239-48.

<https://doi.org/10.1590/1413-81232014201.21192013>

Barreto ML, Barros AJD, Carvalho MS, Codeço CT, Hallal PRC, Medronho RA et al. O que é urgente e necessário para subsidiar as políticas de enfrentamento da pandemia de COVID-19 no Brasil? *Rev. bras. epidemiol.* 2020 Apr;23:e200032.

<https://doi.org/10.1590/1980-549720200032>

Brasil. Ministério da Saúde. Secretaria de Vigilância à Saúde. Painel de casos de doença pelo coronavírus 2019 (COVID-19) no Brasil. 2020. <https://covid.saude.gov.br/>.

Cardoso AL, Vieira ALS, Machado MH. Mercado de Trabalho dos Odontólogos no Brasil. *Divulg Saúde Debate.* 2010 May;45(5):71-9.

Cascaes AM, Dotto L, Bomfim RA. Trends in the dental surgeon workforce in Brazil 2007-2014: a time series study using data from the National Registry of Health Services. *Epidemiol. Serv. Saúde* [Internet]. 2018 [cited 2020 June 17]; 27(1):e201723615. http://www.scielo.br/scielo.php?script=sci_arttext&pid=S2237-

96222018000100313&lng=en. Epub Mar 05, 2018. <https://doi.org/10.5123/s1679-49742018000100015>

Cayetano MH, Carrer FC, Gabriel M, Martins FC, Pucca Jr. GA, Araujo MA. Política Nacional de Saúde Bucal Brasileira (Brasil Sorridente): Um resgate da história, aprendizados e futuro. *Univ Odontol.* 2019 ene-jun; 38(80).
<https://doi.org/10.11144/Javeriana.uo38-80.pnsb>

Chen, N, Zhou M, Dong X, Qu J, Gong F, Han Y, Qiu Y, Wang J, Liu Y, Wei Y, Xia J, Yu T, Zhang X, Zhang L. Epidemiological and clinical characteristics of 99 cases of 2019 novel coronavirus pneumonia in Wuhan, China: a descriptive study. *Lancet* 395, 507–513 (2020).

Garbin CAS, Vanzo KLT, Moimaz SAS, Garbin AJI, Chiba FY. Vaccination coverage and immunity against hepatitis B in public health dentists. *Rev. Soc. Bras. Med. Trop.* 2019;52:e20180534. <https://doi.org/10.1590/0037-8682-0534-2018>

Garbin CAS, Wakayama B, Saliba TA, Saliba Junior OA, Garbin AJI. A cross-sectional study on dental surgeons' immune status against hepatitis B virus in the Public Health System. *Rev. Inst. Med. trop. São Paulo.* 2020;62:e18. <https://doi.org/10.1590/s1678-9946202062018>

Hallal PC, Horta BL, Barros AJD, Dellagostin OA, Hartwig FP, Silveira MF et al. Evolução da prevalência de infecção por COVID-19 no Rio Grande do Sul: inquéritos sorológicos seriados. *Ciência e Saúde Coletiva.* 2020 Abr;65.
<https://preprints.scielo.org/index.php/scielo/preprint/view/41>>.

Imai C, Toizumi M, Hall L, Lambert S, Halton K, Merollini K. A systematic review and meta-analysis of the direct epidemiological and economic effects of seasonal influenza vaccination on healthcare workers. *PLoS One.* 2018 Jun 7;13(6):e0198685.
doi:10.1371/journal.pone.0198685

Kinariwala N, Perera Samaranayake L, Perera I, Patel Z. Concerns and Fears of Indian Dentists on Professional Practice During the Coronavirus Disease-2019 (COVID-19) Pandemic [published online ahead of print, 2020 Jun 7]. *Oral Dis.* 2020;10.1111/odi.13459. doi:10.1111/odi.13459

Morita MC, Haddad AE, Araújo ME. Perfil atual e tendências do cirurgião-dentista brasileiro. 1st ed. Maringá: Dental Press International; 2010.

Noronha MSM. Riscos ocupacionais e patologias em cirurgiões-dentistas de centro de especialidades odontológicas. *RFO UPF [online].* 2013; 18(3):316-320.

Peng X, Xu X, Li Y, Chend L, Zhou X, Ren B. Transmission routes of 2019-nCoV and controls in dental practice. *Int J Oral Sci.* 2020 Mar; 12(1):9.

Pernambuco. Secretaria de Saúde de Pernambuco. Secretaria Executiva de Vigilância à Saúde. Centro de Informações Estratégicas de Vigilância em Saúde de Pernambuco. Informe Epidemiológico nº 107/2020. <https://www.cievspe.com/novo-coronavirus-2019-ncov>

Quijano-Escate R, Rebatta-Acuña A, Garayar-Peceros H, Gutierrez-Flores KE, Bendezu-Quispe G. Aprendizaje en tiempos de aislamiento social: cursos masivos

abiertos en línea sobre la COVID-19. Rev Peru Med Exp Salud Publica. 2020; 37(2):375-7. doi: <https://doi.org/10.17843/rpmesp.2020.372.5478>

Wenham C, Smith J, Morgan R. COVID-19: the gendered impacts of the outbreak. The Lancet. 2020 Mar; 395(10227):846-848. [https://doi.org/10.1016/S0140-6736\(20\)30526-2](https://doi.org/10.1016/S0140-6736(20)30526-2)

World Health Organization, Coronavirus disease (COVID-19) pandemic [Internet]. Novel coronavirus(2019-nCoV) Situation Report - 52. 2020. https://www.who.int/docs/default-source/coronaviruse/situation-reports/20200312-sitrep-52-covid-19.pdf?sfvrsn=e2bfc9c0_4

Ye C, Zhang S, Zhang X, Cai H, Gu J, Lian J et al. Impact of comorbidities on patients with COVID- 19: A large retrospective study in Zhejiang, China. J Med Virol. 2020 Jun Accepted Author Manuscript. doi:10.1002/jmv.26183.

This preprint was submitted under the following conditions:

- The authors declare that they are aware that they are solely responsible for the content of the preprint and that the deposit in SciELO Preprints does not mean any commitment on the part of SciELO, except its preservation and dissemination.
- The authors declare that the necessary Terms of Free and Informed Consent of participants or patients in the research were obtained and are described in the manuscript, when applicable.
- The authors declare that the preparation of the manuscript followed the ethical norms of scientific communication.
- The authors declare that the data, applications, and other content underlying the manuscript are referenced.
- The deposited manuscript is in PDF format.
- The authors declare that the research that originated the manuscript followed good ethical practices and that the necessary approvals from research ethics committees, when applicable, are described in the manuscript.
- The authors declare that once a manuscript is posted on the SciELO Preprints server, it can only be taken down on request to the SciELO Preprints server Editorial Secretariat, who will post a retraction notice in its place.
- The authors agree that the approved manuscript will be made available under a [Creative Commons CC-BY](#) license.
- The submitting author declares that the contributions of all authors and conflict of interest statement are included explicitly and in specific sections of the manuscript.
- The authors declare that the manuscript was not deposited and/or previously made available on another preprint server or published by a journal.
- If the manuscript is being reviewed or being prepared for publishing but not yet published by a journal, the authors declare that they have received authorization from the journal to make this deposit.
- The submitting author declares that all authors of the manuscript agree with the submission to SciELO Preprints.