

Mathematical Induction

Mala Rani Barman

Lecturer

Dept. of CSE

Sheikh Hasina University

- **Mathematical induction**, is a technique for proving results or establishing statements for natural numbers. This part illustrates the method through a variety of examples.
- Definition
- **Mathematical Induction** is a mathematical technique which is used to prove a statement, a formula or a theorem is true for every natural number.
- The technique involves two steps to prove a statement, as stated below –
- **Step 1(Base step)** – It proves that a statement is true for the initial value.
- **Step 2(Inductive step)** – It proves that if the statement is true for the n^{th} iteration (or number n), then it is also true for $(n+1)^{\text{th}}$ iteration (or number $n+1$).

- **Procedure**

- **Step 1** – Consider an initial value for which the statement is true. It is to be shown that the statement is true for $n = \text{initial value}$.
- **Step 2** – Assume the statement is true for any value of $n = k$. Then prove the statement is true for $n = k+1$. We actually break $n = k+1$ into two parts, one part is $n = k$ (which is already proved) and try to prove the other part.

Example-1

$$1 + 3 + 5 + \dots + (2n - 1) = n^2 \quad \text{for } n = 1, 2, \dots$$

Solution

Step 1 – For $n = 1$, $1 = 1^2$, Hence, step 1 is satisfied.

Step 2 – Let us assume the statement is true for $n = k$.

Hence, $1 + 3 + 5 + \dots + (2k - 1) = k^2$ is true (It is an assumption)

We have to prove that $1 + 3 + 5 + \dots + (2(k + 1) - 1) = (k + 1)^2$ also holds

$$1 + 3 + 5 + \dots + (2(k + 1) - 1)$$

$$= 1 + 3 + 5 + \dots + (2k + 2 - 1)$$

Example-1

$$= 1 + 3 + 5 + \cdots + (2k + 2 - 1)$$

$$= 1 + 3 + 5 + \cdots + (2k + 1)$$

$$= 1 + 3 + 5 + \cdots + (2k - 1) + (2k + 1)$$

$$= k^2 + (2k + 1)$$

$$= (k + 1)^2$$

So, $1 + 3 + 5 + \cdots + (2(k + 1) - 1) = (k + 1)^2$ hold which satisfies the step 2.

Hence, $1 + 3 + 5 + \cdots + (2n - 1) = n^2$ is proved.

Example-2

$3^n - 1$ is a multiple of 2 for $n = 1, 2, \dots$

Solution

Step 1 – For $n = 1, 3^1 - 1 = 3 - 1 = 2$ which is a multiple of 2

Step 2 – Let us assume $3^n - 1$ is true for $n = k$, Hence, $3^k - 1$ is true (It is an assumption)

We have to prove that $3^{k+1} - 1$ is also a multiple of 2

$$3^{k+1} - 1 = 3 \times 3^k - 1 = (2 \times 3^k) + (3^k - 1)$$

The first part (2×3^k) is certain to be a multiple of 2 and the second part $(3^k - 1)$ is also true as our previous assumption.

Hence, $3^{k+1} - 1$ is a multiple of 2.

So, it is proved that $3^n - 1$ is a multiple of 2.

Example-3

Prove that $(ab)^n = a^n b^n$ is true for every natural number n

Solution

Step 1 – For $n = 1$, $(ab)^1 = a^1 b^1 = ab$, Hence, step 1 is satisfied.

Step 2 – Let us assume the statement is true for $n = k$, Hence, $(ab)^k = a^k b^k$ is true (It is an assumption).

We have to prove that $(ab)^{k+1} = a^{k+1} b^{k+1}$ also hold

Given, $(ab)^k = a^k b^k$

Or, $(ab)^k (ab) = (a^k b^k)(ab)$ [Multiplying both side by 'ab']

Example-3

$$\text{Or, } (ab)^{k+1} = (aa^k)(bb^k)$$

$$\text{Or, } (ab)^{k+1} = (a^{k+1}b^{k+1})$$

Hence, step 2 is proved.

So, $(ab)^n = a^n b^n$ is true for every natural number n .