

# Sheikh Hasina University, Netrokona

Course:

MATH-3105 (Multivariable Calculus & Geometry)

Textbook:

Calculus, Early Transcendentals

By Anton, Bivens, Davis (10<sup>th</sup> Edition)

Course Teacher:

Dr. Md. Tauhedul Azam

3<sup>rd</sup> Year 1<sup>st</sup> Semester

1

## Vectors–Valued Functions

2

## Chapter 12.1

# Introduction to Vector-Valued Functions

MATH-3105

Dept. of CSE

3

3

## Parametric Curves in 3-Space

If  $f$ ,  $g$ , and  $h$  are well-behaved functions, then the parametric equations

$$x = f(t), y = g(t), z = h(t)$$

generate a curve in 3-space that is traced in a specific direction as  $t$  increases. This direction is called the orientation or direction of increasing parameter, and the curve together with its orientation is called the graph of the parametric equations or the parametric curve represented by the equations. If no restrictions are stated explicitly or are implied by the equations, then it will be understood that  $t$  varies over the interval  $(-\infty, \infty)$ .

MATH-3105



Dept. of CSE

4

4

## Parametric Curves in 3-Space

Example (p842-843)



The parametric equation  
 $x = 1 - t, y = 3t, z = 3t$   
 represents a line in 3-space.

The parametric equation  
 $x = a \cos t, y = a \sin t, z = ct$   
 represent a curve called a circular helix.

MATH-3105

Dept. of CSE

5

5

## Vector-Valued Functions

### Definition

The parametric equation  $x = t, y = t^2, z = t^3$  is called **twisted cubic** is the set of points of the form  $(t, t^2, t^3)$  for real values of  $t$ . If we view each of these points as a terminal point for a vector  $\underline{r}$  whose initial point is at the origin,

$$\underline{r} = (x, y, z) = (t, t^2, t^3) = t \hat{i} + t^2 \hat{j} + t^3 \hat{k}$$

then we obtain  $\underline{r}$  as a function of the parameter  $t$ , that is,  $\underline{r} = \underline{r}(t)$ . Since this function produces a vector, we say that  $\underline{r} = \underline{r}(t)$  defines  $\underline{r}$  as a **vector-valued function of a real variable**, or more simply, a **vector-valued function**.

MATH-3105

Dept. of CSE

6

6

## Vector-Valued Functions

### Definition

If  $\underline{\mathbf{r}}(t)$  is a vector-valued function in 3-space, then for each allowable value of  $t$  the vector  $\underline{\mathbf{r}} = \underline{\mathbf{r}}(t)$  can be represented in terms of components as

$$\underline{\mathbf{r}} = \underline{\mathbf{r}}(t) = (x(t), y(t), z(t)) = x(t) \hat{\mathbf{i}} + y(t) \hat{\mathbf{j}} + z(t) \hat{\mathbf{k}}$$

The functions  $x(t)$ ,  $y(t)$ , and  $z(t)$  are called the **component functions** or the **components** of  $\underline{\mathbf{r}}(t)$ .

MATH-3105

Dept. of CSE

7

7

## Vector-Valued Functions

### Domain

The **domain** of a vector-valued function  $\underline{\mathbf{r}}(t)$  is the set of allowable values for  $t$ .

If  $\underline{\mathbf{r}}(t)$  is defined in terms of component functions and the domain is not specified explicitly, then it will be understood that the domain is the intersection of the natural domains of the component functions; this is called the **natural domain** of  $\underline{\mathbf{r}}(t)$ .

MATH-3105

Dept. of CSE

8

8

## Vector-Valued Functions

### Homework

Example 1 – 6 (p842 –845)

## Chapter 12.1

### Homework

Exercise Set 12.1 (p845 –847)

## Chapter 12.2

# Calculus of Vector-Valued Functions

MATH-3105

Dept. of CSE

11

11

## Limits and Continuity

### Definition 12.2.1

Let  $\underline{\mathbf{r}}(t)$  be a vector-valued function that is defined for all  $t$  in some open interval containing the number  $a$ , except that  $\underline{\mathbf{r}}(t)$  need not be defined at  $a$ . We will write

$$\lim_{t \rightarrow a} \underline{\mathbf{r}}(t) = \underline{\mathbf{L}}$$

if and only if

$$\lim_{t \rightarrow a} \|\underline{\mathbf{r}}(t) - \underline{\mathbf{L}}\| = 0$$


MATH-3105

Dept. of CSE


12

12

## Limits and Continuity



$\mathbf{r}(t)$  approaches  $\mathbf{L}$  in length and direction if  $\lim_{t \rightarrow a} \mathbf{r}(t) = \mathbf{L}$ .



$\|\mathbf{r}(t) - \mathbf{L}\|$  is the distance between terminal points for vectors  $\mathbf{r}(t)$  and  $\mathbf{L}$  when positioned with the same initial points.

MATH-3105

Dept. of CSE

13

13

## Limits and Continuity

### Theorem 12.2.2

(a) If  $\underline{\mathbf{r}} = \underline{\mathbf{r}}(t) = (x(t), y(t)) = x(t) \hat{\mathbf{i}} + y(t) \hat{\mathbf{j}}$ , then

$$\lim_{t \rightarrow a} \underline{\mathbf{r}}(t) = \left( \lim_{t \rightarrow a} x(t), \lim_{t \rightarrow a} y(t) \right) = \lim_{t \rightarrow a} x(t) \hat{\mathbf{i}} + \lim_{t \rightarrow a} y(t) \hat{\mathbf{j}}$$

provided the limits of the component functions exist. Conversely, the limits of the component functions exist provided  $\underline{\mathbf{r}}(t)$  approaches a limiting vector as  $t$  approaches  $a$ .

MATH-3105

Dept. of CSE

14

14

## Limits and Continuity

### Theorem 12.2.2

(b) If  $\underline{\mathbf{r}} = \underline{\mathbf{r}}(t) = (x(t), y(t), z(t)) = x(t) \hat{\mathbf{i}} + y(t) \hat{\mathbf{j}} + z(t) \hat{\mathbf{k}}$  then

$$\begin{aligned} \lim_{t \rightarrow a} \underline{\mathbf{r}}(t) &= \left( \lim_{t \rightarrow a} x(t), \lim_{t \rightarrow a} y(t), \lim_{t \rightarrow a} z(t) \right) \\ &= \lim_{t \rightarrow a} x(t) \hat{\mathbf{i}} + \lim_{t \rightarrow a} y(t) \hat{\mathbf{j}} + \lim_{t \rightarrow a} z(t) \hat{\mathbf{k}} \end{aligned}$$

provided the limits of the component functions exist. Conversely, the limits of the component functions exist provided  $\underline{\mathbf{r}}(t)$  approaches a limiting vector as  $t$  approaches  $a$ .

MATH-3105

Dept. of CSE

15

15

## Limits and Continuity

### Example 1 (Textbook p849)

Let  $\underline{\mathbf{r}}(t) = t^2 \hat{\mathbf{i}} + e^t \hat{\mathbf{j}} - (2 \cos \pi t) \hat{\mathbf{k}}$ . Then

$$\begin{aligned} \lim_{t \rightarrow 0} \underline{\mathbf{r}}(t) &= \left( \lim_{t \rightarrow 0} t^2 \right) \hat{\mathbf{i}} + \left( \lim_{t \rightarrow 0} e^t \right) \hat{\mathbf{j}} - \left( \lim_{t \rightarrow 0} 2 \cos \pi t \right) \hat{\mathbf{k}} \\ &= (0) \hat{\mathbf{i}} + (1) \hat{\mathbf{j}} - (2) \hat{\mathbf{k}} \\ &= \hat{\mathbf{j}} - 2 \hat{\mathbf{k}} \end{aligned}$$

MATH-3105

Dept. of CSE

16

16



# Continuity

## Definition

We define a vector-valued function  $\underline{\mathbf{r}}(t)$  to be continuous at  $t = a$  if

$$\lim_{t \rightarrow a} \underline{\mathbf{r}}(t) = \underline{\mathbf{r}}(a)$$

That is,  $\underline{\mathbf{r}}(a)$  is defined, the limit of  $\underline{\mathbf{r}}(t)$  as  $t \rightarrow a$  exists, and the two are equal.

MATH-3105

Dept. of CSE

17

17

# Continuity

## Definition

We say that  $\underline{\mathbf{r}}(t)$  is continuous on an interval  $I$  if it is continuous at each point of  $I$  [with the understanding that at an endpoint in  $I$  the two-sided limit is replaced by the appropriate one-sided limit].

It follows that a vector-valued function is continuous at  $t = a$  if and only if its component functions are continuous at  $t = a$ .

MATH-3105

Dept. of CSE

18

18

## Derivatives

### Definition

If  $\underline{\mathbf{r}}(t)$  is a vector-valued function, we define the **derivative of  $\underline{\mathbf{r}}$  with respect to  $t$**  to be the vector-valued function  $\underline{\mathbf{r}}'$  given by

$$\underline{\mathbf{r}}'(t) = \lim_{h \rightarrow 0} \frac{\underline{\mathbf{r}}(t+h) - \underline{\mathbf{r}}(t)}{h}$$

The domain of  $\underline{\mathbf{r}}'$  consists of all values of  $t$  in the domain of  $\underline{\mathbf{r}}(t)$  for which the limit exists.

The function  $\underline{\mathbf{r}}(t)$  is **differentiable** at  $t$  if the limit exists.

MATH-3105

Dept. of CSE

19

19

## Derivatives

### Notes:

- The derivative of  $\underline{\mathbf{r}}(t)$  can be expressed as

$$\frac{d}{dt}[\underline{\mathbf{r}}(t)], \quad \frac{d\underline{\mathbf{r}}}{dt}, \quad \underline{\mathbf{r}}'(t), \quad \underline{\mathbf{r}}'$$

- It is important to keep in mind that  $\underline{\mathbf{r}}'(t)$  is a vector, not a number, and hence has a magnitude and a direction for each value of  $t$  [except if  $\underline{\mathbf{r}}'(t) = \mathbf{0}$ , in which case  $\underline{\mathbf{r}}'(t)$  has magnitude zero but no specific direction].

MATH-3105

Dept. of CSE


20

20

# Derivatives

## Geometric Interpretation of the Derivative

Suppose that  $C$  is the graph of a vector-valued function  $\underline{\mathbf{r}}(t)$  in 2-space or 3-space and that  $\underline{\mathbf{r}}'(t)$  exists and is nonzero for a given value of  $t$ . If the vector  $\underline{\mathbf{r}}'(t)$  is positioned with its initial point at the terminal point of the radius vector  $\underline{\mathbf{r}}(t)$ , then  $\underline{\mathbf{r}}'(t)$  is tangent to  $C$  and points in the direction of increasing parameter.



MATH-3105

Dept. of CSE

21

21

# Derivatives

## Theorem 12.2.5

If  $\underline{\mathbf{r}}(t)$  is a vector-valued function, then  $\underline{\mathbf{r}}$  is differentiable at  $t$  if and only if each of its component functions is differentiable at  $t$ , in which case the component functions of  $\underline{\mathbf{r}}'(t)$  are the derivatives of the corresponding component functions of  $\underline{\mathbf{r}}(t)$ . If  $\underline{\mathbf{r}}(t) = x(t) \hat{\mathbf{i}} + y(t) \hat{\mathbf{j}} + z(t) \hat{\mathbf{k}}$  then

$$\underline{\mathbf{r}}'(t) = \lim_{h \rightarrow 0} \frac{\underline{\mathbf{r}}(t+h) - \underline{\mathbf{r}}(t)}{h} = x'(t) \hat{\mathbf{i}} + y'(t) \hat{\mathbf{j}} + z'(t) \hat{\mathbf{k}}$$

MATH-3105

Dept. of CSE

22

22

## Derivatives

### Example 2 (Textbook p850)

Let  $\underline{\mathbf{r}}(t) = t^2 \hat{\mathbf{i}} + e^t \hat{\mathbf{j}} - (2 \cos \pi t) \hat{\mathbf{k}}$ . Then

$$\begin{aligned}\underline{\mathbf{r}}'(t) &= \frac{d}{dt}(t^2) \hat{\mathbf{i}} + \frac{d}{dt}(e^t) \hat{\mathbf{j}} - \frac{d}{dt}(2 \cos \pi t) \hat{\mathbf{k}} \\ &= 2t \hat{\mathbf{i}} + e^t \hat{\mathbf{j}} + (2\pi \sin \pi t) \hat{\mathbf{k}}\end{aligned}$$

MATH-3105

Dept. of CSE

23

23

## Derivatives

**12.2.6 THEOREM (Rules of Differentiation)** Let  $\mathbf{r}(t)$ ,  $\mathbf{r}_1(t)$ , and  $\mathbf{r}_2(t)$  be differentiable vector-valued functions that are all in 2-space or all in 3-space, and let  $f(t)$  be a differentiable real-valued function,  $k$  a scalar, and  $\mathbf{c}$  a constant vector (that is, a vector whose value does not depend on  $t$ ). Then the following rules of differentiation hold:

- (a)  $\frac{d}{dt}[\mathbf{c}] = \mathbf{0}$
- (b)  $\frac{d}{dt}[k\mathbf{r}(t)] = k \frac{d}{dt}[\mathbf{r}(t)]$
- (c)  $\frac{d}{dt}[\mathbf{r}_1(t) + \mathbf{r}_2(t)] = \frac{d}{dt}[\mathbf{r}_1(t)] + \frac{d}{dt}[\mathbf{r}_2(t)]$
- (d)  $\frac{d}{dt}[\mathbf{r}_1(t) - \mathbf{r}_2(t)] = \frac{d}{dt}[\mathbf{r}_1(t)] - \frac{d}{dt}[\mathbf{r}_2(t)]$
- (e)  $\frac{d}{dt}[f(t)\mathbf{r}(t)] = f(t) \frac{d}{dt}[\mathbf{r}(t)] + \frac{d}{dt}[f(t)]\mathbf{r}(t)$

MATH-3105

Dept. of CSE


24

24

## Tangent Lines

### Definition

Let  $P$  be a point on the graph of a vector-valued function  $\underline{\mathbf{r}}(t)$ , and let  $\underline{\mathbf{r}}(t_0)$  be the radius vector from the origin to  $P$  (Figure). If  $\underline{\mathbf{r}}'(t_0)$  exists and  $\underline{\mathbf{r}}'(t_0) \neq \mathbf{0}$ , then we call  $\underline{\mathbf{r}}'(t_0)$  a tangent vector to the graph of  $\underline{\mathbf{r}}(t)$  at  $\underline{\mathbf{r}}(t_0)$ , and we call the line through  $P$  that is parallel to the tangent vector the tangent line to the graph of  $\underline{\mathbf{r}}(t)$  at  $\underline{\mathbf{r}}(t_0)$ .



MATH-3105

Dept. of CSE

25

25

## Tangent Lines

### Definition

Let  $\underline{\mathbf{r}}_0 = \underline{\mathbf{r}}(t_0)$  and  $\underline{\mathbf{v}}_0 = \underline{\mathbf{r}}'(t_0)$ . Then the tangent line to the graph of  $\underline{\mathbf{r}}(t)$  at  $\underline{\mathbf{r}}_0$  is given by the vector equation

$$\underline{\mathbf{r}} = \underline{\mathbf{r}}_0 + t \underline{\mathbf{v}}_0$$

MATH-3105

Dept. of CSE

26

26

## Tangent Lines

### Example 3 (p851)

Find parametric equations of the tangent line to the circular helix

$$x = \cos t, \quad y = \sin t, \quad z = t$$

where  $t = t_0$ , and use that result to find parametric equations for the tangent line at the point where  $t = \pi$ .

MATH-3105

Dept. of CSE

27

27

## Tangent Lines

### Solution

The vector equation of the helix is

$$\underline{\mathbf{r}}(t) = \cos t \, \hat{\mathbf{i}} + \sin t \, \hat{\mathbf{j}} + t \, \hat{\mathbf{k}}$$

so we have

$$\underline{\mathbf{r}}_0 = \underline{\mathbf{r}}(t_0) = \cos t_0 \, \hat{\mathbf{i}} + \sin t_0 \, \hat{\mathbf{j}} + t_0 \, \hat{\mathbf{k}}$$

$$\underline{\mathbf{v}}_0 = \underline{\mathbf{r}}'(t_0) = (-\sin t_0) \hat{\mathbf{i}} + \cos t_0 \, \hat{\mathbf{j}} + \hat{\mathbf{k}}$$

MATH-3105

Dept. of CSE

28

28

## Tangent Lines

### Solution

Therefore, the vector equation of the tangent line at  $t = t_0$  is

$$\begin{aligned}\underline{\mathbf{r}}(t) &= \underline{\mathbf{r}}_0 + t \underline{\mathbf{v}}_0 \\ &= \cos t_0 \hat{\mathbf{i}} + \sin t_0 \hat{\mathbf{j}} + t_0 \hat{\mathbf{k}} + t[(-\sin t_0)\hat{\mathbf{i}} + \cos t_0 \hat{\mathbf{j}} + \hat{\mathbf{k}}] \\ &= (\cos t_0 - t \sin t_0)\hat{\mathbf{i}} + (\sin t_0 + t \cos t_0) \hat{\mathbf{j}} + (t_0 + t)\hat{\mathbf{k}}\end{aligned}$$

MATH-3105

Dept. of CSE

29

29

## Tangent Lines

### Solution

Thus the parametric equations of the tangent line at  $t = t_0$  are

$$x = \cos t_0 - t \sin t_0, \quad y = \sin t_0 + t \cos t_0, \quad z = t_0 + t$$

In particular, the tangent line at  $t = \pi$  has parametric equations

$$x = -1, \quad y = -t, \quad z = \pi + t$$

MATH-3105

Dept. of CSE

30

30

## Tangent Lines

### Example 4 (p852)

Let,

$$\underline{\mathbf{r}}_1(t) = (\tan^{-1} t) \hat{\mathbf{i}} + (\sin t) \hat{\mathbf{j}} + t^2 \hat{\mathbf{k}}$$

and

$$\underline{\mathbf{r}}_2(t) = (t^2 - t) \hat{\mathbf{i}} + (2t - 2) \hat{\mathbf{j}} + (\ln t) \hat{\mathbf{k}}$$

The graphs of  $\underline{\mathbf{r}}_1(t)$  and  $\underline{\mathbf{r}}_2(t)$  intersect at the origin. Find the degree measure of the acute angle between the tangent lines to the graphs of  $\underline{\mathbf{r}}_1(t)$  and  $\underline{\mathbf{r}}_2(t)$  at the origin.

MATH-3105

Dept. of CSE

31

31

## Tangent Lines

### Solution

At the origin,

$$\tan^{-1} t = 0, \sin t = 0, t^2 = 0 \implies t = 0 \text{ for } \underline{\mathbf{r}}_1(t).$$

The graphs of  $\underline{\mathbf{r}}_1(t)$  passes through the origin at  $t = 0$ , where its tangent vector is

$$\underline{\mathbf{r}}_1'(0) = \left( \frac{1}{1+t^2}, \cos t, 2t \right) \Big|_{t=0} = (1, 1, 0)$$

MATH-3105

Dept. of CSE

32

32



## Tangent Lines

### Solution

At the origin,

$$t^2 - t = 0, \quad 2t - t = 0, \quad \ln t = 0 \Rightarrow t = 1 \text{ for } \underline{\mathbf{r}}_2(t).$$

The graphs of  $\underline{\mathbf{r}}_2(t)$  passes through the origin at  $t = 1$ , where its tangent vector is

$$\underline{\mathbf{r}}'_2(1) = \left( 2t - 1, 2, \frac{1}{t} \right) \Big|_{t=1} = (1, 2, 1)$$

MATH-3105

Dept. of CSE

33

33

## Tangent Lines

### Solution

The angle between two tangent vectors satisfies

$$\cos \theta = \frac{(1, 1, 0) \cdot (1, 2, 1)}{\|(1, 1, 0)\| \|(1, 2, 1)\|} = \frac{1 + 2 + 0}{\sqrt{2} \sqrt{6}} = \frac{3}{\sqrt{12}} = \frac{\sqrt{3}}{2}$$

$$\Rightarrow \theta = \cos^{-1} \left( \frac{\sqrt{3}}{2} \right)$$

$$= \frac{\pi}{6} \text{ radian}$$

$$= 30^\circ$$

MATH-3105

Dept. of CSE

34

34

## Derivatives of Dot & Cross Products

$$\frac{d}{dt} [\underline{\mathbf{r}}_1(t) \cdot \underline{\mathbf{r}}_2(t)] = \underline{\mathbf{r}}_1(t) \cdot \frac{d\underline{\mathbf{r}}_2}{dt} + \frac{d\underline{\mathbf{r}}_1}{dt} \cdot \underline{\mathbf{r}}_2(t)$$

$$\frac{d}{dt} [\underline{\mathbf{r}}_1(t) \times \underline{\mathbf{r}}_2(t)] = \underline{\mathbf{r}}_1(t) \times \frac{d\underline{\mathbf{r}}_2}{dt} + \frac{d\underline{\mathbf{r}}_1}{dt} \times \underline{\mathbf{r}}_2(t)$$

MATH-3105

Dept. of CSE

35

35

## Derivatives of Dot & Cross Products

### Theorem 12.2.8

If  $\underline{\mathbf{r}}(t)$  is a differentiable vector-valued function in 2-space or 3-space and  $\|\underline{\mathbf{r}}(t)\|$  is constant for all  $t$ , then

$$\underline{\mathbf{r}}(t) \cdot \underline{\mathbf{r}}'(t) = 0$$

that is,  $\underline{\mathbf{r}}(t)$  and  $\underline{\mathbf{r}}'(t)$  are orthogonal vectors for all  $t$ .

### Proof

Homework (p853).

MATH-3105

Dept. of CSE

36

36

## Definite Integrals of Vector-Valued F<sup>n</sup>s

If  $\underline{\mathbf{r}}(t)$  is a vector-valued function that is continuous on the interval  $a \leq t \leq b$ , then we define the definite integral of  $\underline{\mathbf{r}}(t)$  over this interval as

$$\int_a^b \underline{\mathbf{r}}(t) dt = \left( \int_a^b x(t) dt \right) \hat{\mathbf{i}} + \left( \int_a^b y(t) dt \right) \hat{\mathbf{j}} + \left( \int_a^b z(t) dt \right) \hat{\mathbf{k}}$$

MATH-3105

Dept. of CSE

37

37

## Definite Integrals of Vector-Valued F<sup>n</sup>s

### Example 6 (p854)

Let  $\underline{\mathbf{r}}(t) = t^2 \hat{\mathbf{i}} + e^t \hat{\mathbf{j}} - (2 \cos \pi t) \hat{\mathbf{k}}$ . Then

$$\begin{aligned} \int_0^1 \underline{\mathbf{r}}(t) dt &= \left( \int_0^1 t^2 dt \right) \hat{\mathbf{i}} + \left( \int_0^1 e^t dt \right) \hat{\mathbf{j}} - \left( \int_0^1 2 \cos \pi t dt \right) \hat{\mathbf{k}} \\ &= \frac{t^3}{3} \Big|_0^1 \hat{\mathbf{i}} + e^t \Big|_0^1 \hat{\mathbf{j}} - \frac{2}{\pi} \sin \pi t \Big|_0^1 \hat{\mathbf{k}} \\ &= \frac{1}{3} \hat{\mathbf{i}} + (e - 1) \hat{\mathbf{j}} \end{aligned}$$

MATH-3105

Dept. of CSE

38

38

## Rules of Integration

**12.2.9 THEOREM (Rules of Integration)** Let  $\mathbf{r}(t)$ ,  $\mathbf{r}_1(t)$ , and  $\mathbf{r}_2(t)$  be vector-valued functions in 2-space or 3-space that are continuous on the interval  $a \leq t \leq b$ , and let  $k$  be a scalar. Then the following rules of integration hold:

$$(a) \int_a^b k\mathbf{r}(t) dt = k \int_a^b \mathbf{r}(t) dt$$

$$(b) \int_a^b [\mathbf{r}_1(t) + \mathbf{r}_2(t)] dt = \int_a^b \mathbf{r}_1(t) dt + \int_a^b \mathbf{r}_2(t) dt$$

$$(c) \int_a^b [\mathbf{r}_1(t) - \mathbf{r}_2(t)] dt = \int_a^b \mathbf{r}_1(t) dt - \int_a^b \mathbf{r}_2(t) dt$$

MATH-3105

Dept. of CSE

39

39

## Antiderivatives of Vector-Valued Functions

An antiderivative for a vector-valued function  $\underline{\mathbf{r}}(t)$  is a vector-valued function  $\underline{\mathbf{R}}(t)$  such that

$$\underline{\mathbf{R}}'(t) = \underline{\mathbf{r}}(t)$$

We express this Equation using integral notation as

$$\int \underline{\mathbf{r}}(t) dt = \underline{\mathbf{R}}(t) + \underline{\mathbf{C}}$$

where  $\underline{\mathbf{C}}$  represents an arbitrary constant vector.

MATH-3105

Dept. of CSE

40

40

## Antiderivatives of Vector-Valued Functions

Example 7 (p855)

$$\begin{aligned}\int (2t \hat{\mathbf{i}} + 3t^2 \hat{\mathbf{j}}) dt &= \left( \int 2t dt \right) \hat{\mathbf{i}} + \left( \int 3t^2 dt \right) \hat{\mathbf{j}} \\ &= (t^2 + C_1) \hat{\mathbf{i}} + (t^3 + C_2) \hat{\mathbf{j}} \\ &= t^2 \hat{\mathbf{i}} + t^3 \hat{\mathbf{j}} + \underline{\mathbf{C}}\end{aligned}$$

where  $\underline{\mathbf{C}} = C_1 \hat{\mathbf{i}} + C_2 \hat{\mathbf{j}}$  is an arbitrary vector constant of integration.

MATH-3105

Dept. of CSE

41

41

## Antiderivatives of Vector-Valued Functions

$$\frac{d}{dt} \left[ \int \underline{\mathbf{r}}(t) dt \right] = \underline{\mathbf{r}}(t)$$

$$\int \underline{\mathbf{r}}'(t) dt = \underline{\mathbf{r}}(t) + \underline{\mathbf{C}}$$

$$\int_a^b \underline{\mathbf{r}}(t) dt = \underline{\mathbf{R}}(t) \Big|_a^b = \underline{\mathbf{R}}(b) - \underline{\mathbf{R}}(a)$$

MATH-3105

Dept. of CSE

42

42

## Antiderivatives of Vector-Valued Functions

### Example 8 (p855)

Evaluate the definite integral

$$\int_0^2 (2t \hat{i} + 3t^2 \hat{j}) dt.$$

### Solution

$$\begin{aligned} \int_0^2 (2t \hat{i} + 3t^2 \hat{j}) dt &= t^2 \hat{i} + t^3 \hat{j} \Big|_0^2 \\ &= 4 \hat{i} + 8 \hat{j} \end{aligned}$$

MATH-3105

Dept. of CSE

43

43

## Antiderivatives of Vector-Valued Functions

### Example 9 (p855)

Find  $\underline{r}(t)$  given that  $\underline{r}'(t) = (3, 2t)$  and  $\underline{r}(1) = (2, 5)$ .

### Solution

We know

$$\underline{r}(t) = \int \underline{r}'(t) dt = \int (3, 2t) dt = (3t, t^2) + \underline{C} \quad \dots \dots \dots (1)$$

where  $\underline{C}$  is a vector constant of integration.

MATH-3105

Dept. of CSE

44

44

## Antiderivatives of Vector-Valued Functions

### Solution

Substituting  $t = 1$  in equation (1) we get

$$\underline{\mathbf{r}}(1) = (3,1) + \underline{\mathbf{C}}$$

Given that  $\underline{\mathbf{r}}(1) = (2,5)$ , therefore comparing

$$(3,1) + \underline{\mathbf{C}} = (2,5)$$

$$\Rightarrow \underline{\mathbf{C}} = (2,5) - (3,1) = (-1,4)$$

Thus from equation (1) we get

$$\underline{\mathbf{r}}(t) = (3t, t^2) + (-1, 4) = (3t - 1, t^2 + 4)$$

## Chapter 12.2

### Homework

Exercise Set 12.2 (p856 –857)

## Chapter 12.3

# Change of Parameter; Arc Length

MATH-3105

Dept. of CSE

47

47

## Smooth Parametrizations

### Definition

We will say that a curve represented by  $\underline{\mathbf{r}}(t)$  is **smoothly parametrized** by  $\underline{\mathbf{r}}(t)$ , or that  $\underline{\mathbf{r}}(t)$  is a **smooth function** of  $t$  if  $\underline{\mathbf{r}}'(t)$  is continuous and  $\underline{\mathbf{r}}'(t) \neq \mathbf{0}$  for any allowable value of  $t$ .

Geometrically, this means that a smoothly parametrized curve can have no abrupt changes in direction as the parameter increases.

MATH-3105

Dept. of CSE

48

48



## Smooth Parametrizations

### Example 1 (p858)

Determine whether the following vector-valued functions are smooth.

(a)  $\underline{\mathbf{r}}(t) = a \cos t \, \hat{\mathbf{i}} + a \sin t \, \hat{\mathbf{j}} + ct \, \hat{\mathbf{k}} \quad (a > 0, c > 0)$

(b)  $\underline{\mathbf{r}}(t) = t^2 \, \hat{\mathbf{i}} + t^3 \, \hat{\mathbf{j}}$

MATH-3105

Dept. of CSE

49

49


## Smooth Parametrizations

### Solution (a)

We have

$$\underline{\mathbf{r}}'(t) = -a \sin t \, \hat{\mathbf{i}} + a \cos t \, \hat{\mathbf{j}} + c \, \hat{\mathbf{k}}$$

The components are continuous functions of  $t$ , and there is no value of  $t$  for which all three of them are zero, so  $\underline{\mathbf{r}}(t)$  is a smooth function.



MATH-3105

Dept. of CSE

50

50


## Smooth Parametrizations

### Solution (b)

We have

$$\underline{\mathbf{r}}'(t) = 2t \hat{\mathbf{i}} + 3t^2 \hat{\mathbf{j}}$$

Although the components are continuous functions, they are both equal to zero if  $t = 0$ , so  $\underline{\mathbf{r}}(t)$  is not a smooth function.



MATH-3105

Dept. of CSE

51

51

## Arc Length from the Vector Viewpoint

### Arc Length for Parametric Curve

The arc length  $L$  of a parametric curve

$$x = x(t), \quad y = y(t), \quad z = z(t), \quad (a \leq t \leq b)$$

in 3-space is given by the formula

$$L = \int_a^b \sqrt{\left(\frac{dx}{dt}\right)^2 + \left(\frac{dy}{dt}\right)^2 + \left(\frac{dz}{dt}\right)^2} dt$$

MATH-3105

Dept. of CSE

52

52

## Arc Length from the Vector Viewpoint

### Vector Form

Let us consider a vector-valued function

$$\underline{\mathbf{r}}(t) = x(t) \hat{\mathbf{i}} + y(t) \hat{\mathbf{j}} + z(t) \hat{\mathbf{k}}$$

It follows that

$$\frac{d\underline{\mathbf{r}}}{dt} = \frac{dx}{dt} \hat{\mathbf{i}} + \frac{dy}{dt} \hat{\mathbf{j}} + \frac{dz}{dt} \hat{\mathbf{k}}$$

and hence

$$\left\| \frac{d\underline{\mathbf{r}}}{dt} \right\| = \sqrt{\left( \frac{dx}{dt} \right)^2 + \left( \frac{dy}{dt} \right)^2 + \left( \frac{dz}{dt} \right)^2}$$

MATH-3105

Dept. of CSE

53

53

## Arc Length from the Vector Viewpoint

### Theorem 12.3.1

If  $\mathcal{C}$  is the graph of a smooth vector-valued function  $\underline{\mathbf{r}}(t)$ , then its arc length  $L$  from  $t = a$  to  $t = b$  is

$$L = \int_a^b \left\| \frac{d\underline{\mathbf{r}}}{dt} \right\| dt$$

MATH-3105

Dept. of CSE

54

54

## Arc Length from the Vector Viewpoint

### Example 2 (p859)

Find the arc length of that portion of the circular helix

$$x = \cos t, \quad y = \sin t, \quad z = t$$

from  $t = 0$  to  $t = \pi$ .

MATH-3105

Dept. of CSE

55

55

## Arc Length from the Vector Viewpoint

### Solution

Let,  $\underline{\mathbf{r}}(t) = \cos t \, \hat{\mathbf{i}} + \sin t \, \hat{\mathbf{j}} + t \, \hat{\mathbf{k}} = (\cos t, \sin t, t)$ .

$$\therefore \underline{\mathbf{r}}'(t) = (-\sin t, \cos t, 1)$$

$$\Rightarrow \|\underline{\mathbf{r}}'(t)\| = \sqrt{(-\sin t)^2 + (\cos t)^2 + 1^2} = \sqrt{2}$$

Then the arc length of the helix is

$$L = \int_0^\pi \left\| \frac{d\underline{\mathbf{r}}}{dt} \right\| dt = \int_0^\pi \sqrt{2} \, dt = \sqrt{2}\pi$$

MATH-3105

Dept. of CSE

56


56

## Arc Length As a Parameter

**Step 1:** Select an arbitrary point on the curve  $C$  to serve as a **reference point**.

**Step 2:** Starting from the reference point, choose one direction along the curve to be the **positive direction** and the other to be the **negative direction**.

**Step 3:** If  $P$  is a point on the curve, let  $s$  be the “signed” arc length along  $C$  from the reference point to  $P$ , where  $s$  is positive if  $P$  is in the positive direction from the reference point and  $s$  is negative if  $P$  is in the negative direction. Figure illustrates this idea.



MATH-3105

Dept. of CSE

57

57

## Arc Length As a Parameter

By this procedure, a unique point  $P$  on the curve is determined when a value for  $s$  is given.

Let us now treat  $s$  as a variable. As the value of  $s$  changes, the corresponding point  $P$  moves along  $C$  and the coordinates of  $P$  become functions of  $s$ . Thus, the coordinates of  $P$  are  $(x(s), y(s), z(s))$ . Therefore, the curve  $C$  is given by the parametric equations

$$x = x(s), \quad y = y(s), \quad z = z(s)$$

MATH-3105

Dept. of CSE

58

58

## Arc Length As a Parameter

A parametric representation of a curve with arc length as the parameter is called an **arc length parametrization** of the curve.

Note that a given curve will generally have infinitely many different arc length parametrizations, since the reference point and orientation can be chosen arbitrarily.

MATH-3105

Dept. of CSE

59

59

## Arc Length As a Parameter

### Example 3 (p860)

Find the arc length parametrization of the circle  $x^2 + y^2 = a^2$  with counterclockwise orientation and  $(a, 0)$  as the reference point.

MATH-3105

Dept. of CSE

60

60


## Arc Length As a Parameter

### Solution

The circle with counterclockwise orientation can be represented by the parametric equations

$$x = a \cos t, y = a \sin t \quad (0 \leq t \leq 2\pi) \dots \dots \dots (1)$$

in which  $t$  can be interpreted as the angle in radian measure from the positive  $x$ -axis to the radius from the origin to the point  $P(x, y)$  (Figure).



MATH-3105

Dept. of CSE

61

61

## Arc Length As a Parameter

### Solution

If we take the positive direction for measuring the arc length to be counterclockwise, and we take  $(a, 0)$  to be the reference point, then  $s$  and  $t$  are related by

$$s = at \text{ or } t = s/a.$$

Making this change of variable in (1) and noting that  $s$  increases from  $0$  to  $2\pi a$  as  $t$  increases from  $0$  to  $2\pi$  yields the following arc length parametrization of the circle:

$$x = a \cos(s/a), \quad y = a \sin(s/a) \quad (0 \leq s \leq 2\pi a)$$

MATH-3105


Dept. of CSE

62

62

## Change of Parameter

A **change of parameter** in a vector-valued function  $\underline{\mathbf{r}}(t)$  is a substitution  $t = g(\tau)$  that produces a new vector-valued function  $\underline{\mathbf{r}}(g(\tau))$  having the same graph as  $\underline{\mathbf{r}}(t)$ , but possibly traced differently as the parameter  $\tau$  increases.



MATH-3105

Dept. of CSE

63

63

## Change of Parameter

### Example 4 (p861)

Find a change of parameter  $t = g(\tau)$  for the circle

$$\underline{\mathbf{r}}(t) = \cos t \, \hat{\mathbf{i}} + \sin t \, \hat{\mathbf{j}} \quad (0 \leq t \leq 2\pi)$$

such that

- (a) the circle is traced counterclockwise as  $\tau$  increases over the interval  $[0, 1]$ ;
- (b) the circle traced clockwise as  $\tau$  increases over the interval  $[0, 1]$ .

MATH-3105

Dept. of CSE

64

64



## Change of Parameter

### Solution (a)

The given circle is traced counterclockwise as  $t$  increases.

Thus, if we choose  $g$  to be an increasing function, then it will follow from the relationship  $t = g(\tau)$  that  $t$  increases when  $\tau$  increases, thereby ensuring that the circle will be traced counterclockwise as  $\tau$  increases.

We also want to choose  $g$  so that  $t$  increases from  $0$  to  $2\pi$  as  $\tau$  increases from  $0$  to  $1$ .

A simple choice of  $g$  that satisfies all of the required criteria is the line as follows

$$t = g(\tau) = 2\pi\tau$$

MATH-3105

Dept. of CSE

65

65

## Change of Parameter

### Solution (a)

A simple choice of  $g$  that satisfies all of the required criteria is the line as follows

$$t = g(\tau) = 2\pi\tau$$

The resulting representation of the circle in terms of the parameter  $\tau$  is

$$\underline{\mathbf{r}}(g(\tau)) = \cos(2\pi\tau) \hat{\mathbf{i}} + \sin(2\pi\tau) \hat{\mathbf{j}} \quad (0 \leq \tau \leq 1).$$

MATH-3105

Dept. of CSE

66

66

## Change of Parameter

### Solution (b)

To ensure that the circle is traced clockwise, we will choose  $g$  to be a decreasing function such that  $t$  decreases from  $2\pi$  to  $0$  as  $\tau$  increases from  $0$  to  $1$ . A simple choice of  $g$  that achieves this is the linear function

$$t = g(\tau) = 2\pi(1 - \tau).$$

The resulting representation of the circle in terms of the parameter  $\tau$  is

$$\underline{\mathbf{r}}(g(\tau)) = \cos(2\pi(1 - \tau)) \hat{\mathbf{i}} + \sin(2\pi(1 - \tau)) \hat{\mathbf{j}} \quad (0 \leq \tau \leq 1).$$

which simplifies to

$$\underline{\mathbf{r}}(g(\tau)) = \cos(2\pi\tau) \hat{\mathbf{i}} - \sin(2\pi\tau) \hat{\mathbf{j}} \quad (0 \leq \tau \leq 1).$$

MATH-3105

Dept. of CSE

67

67

## Chain Rule

### Theorem 12.3.2

Let  $\underline{\mathbf{r}}(t)$  be a vector-valued function that is differentiable with respect to  $t$ .

If  $t = g(\tau)$  is a change of parameter in which  $g$  is differentiable with respect to  $\tau$ , then  $\underline{\mathbf{r}}(g(\tau))$  is differentiable with respect to  $\tau$  and

$$\frac{d\underline{\mathbf{r}}}{d\tau} = \frac{d\underline{\mathbf{r}}}{dt} \frac{dt}{d\tau}$$

MATH-3105

Dept. of CSE

68

68

## Smooth Change of Parameter

A change of parameter  $t = g(\tau)$  in which  $\underline{r}(g(\tau))$  is smooth if  $\underline{r}(t)$  is smooth is called a **smooth change of parameter**.

It follows from (9) that  $t = g(\tau)$  will be a smooth change of parameter if  $\frac{dt}{d\tau}$  is continuous and  $\frac{dt}{d\tau} \neq 0$  for all values of  $\tau$ , since these conditions imply that  $\frac{d\mathbf{r}}{d\tau}$  is continuous and nonzero if  $\frac{d\mathbf{r}}{dt}$  is continuous and nonzero.

MATH-3105

Dept. of CSE

69

69

## Smooth Change of Parameter

Smooth changes of parameter fall into two categories—

those for which  $dt/d\tau > 0$  for all  $\tau$  (called positive changes of parameter) and

those for which  $dt/d\tau < 0$  for all  $\tau$  (called negative changes of parameter).

A positive change of parameter preserves the orientation of a parametric curve, and a negative change of parameter reverses it.

MATH-3105

Dept. of CSE

70

70


## Finding Arc Length Parametrizations

### Theorem 12.3.3

Let  $C$  be the graph of a smooth vector-valued function  $\underline{\mathbf{r}}(t)$ , and let  $\underline{\mathbf{r}}(t_0)$  be any point on  $C$ . Then the following formula defines a positive change of parameter from  $t$  to  $s$ , where  $s$  is an arc length parameter having  $\underline{\mathbf{r}}(t_0)$  as its reference point

$$s = \int_{t_0}^t \left\| \frac{d\underline{\mathbf{r}}}{du} \right\| du$$

Proof



Homework (p863).

MATH-3105

Dept. of CSE

71

71

## Finding Arc Length Parametrizations

### Example 6 (p863)

Find the arc length parametrization of the circular helix

$$\underline{\mathbf{r}}(t) = \cos t \, \hat{\mathbf{i}} + \sin t \, \hat{\mathbf{j}} + t \, \hat{\mathbf{k}}$$

that has reference point  $\underline{\mathbf{r}}(0) = (1, 0, 0)$  and the same orientation as the given helix.

MATH-3105

Dept. of CSE

72

72

## Finding Arc Length Parametrizations

### Solution (Example 6)

Replacing  $t$  by  $u$  in  $\underline{\mathbf{r}}$  for integration purpose, we have

$$\underline{\mathbf{r}}(t) = \cos u \, \hat{\mathbf{i}} + \sin u \, \hat{\mathbf{j}} + u \, \hat{\mathbf{k}}$$

$$\therefore \left\| \frac{d\underline{\mathbf{r}}}{du} \right\| = \sqrt{(-\sin u)^2 + \cos^2 u + 1} = \sqrt{2}$$

MATH-3105

Dept. of CSE

73

73

## Finding Arc Length Parametrizations

### Solution (Example 6)

Now taking  $t_0 = 0$

$$s = \int_0^t \left\| \frac{d\underline{\mathbf{r}}}{du} \right\| du = \int_0^t \sqrt{2} du = \sqrt{2}u \Big|_0^t = \sqrt{2}t$$

Thus,  $t = s/\sqrt{2}$ , so the given equation can be reparametrized in terms of  $s$  as

$$\underline{\mathbf{r}} = \cos\left(\frac{s}{\sqrt{2}}\right) \hat{\mathbf{i}} + \sin\left(\frac{s}{\sqrt{2}}\right) \hat{\mathbf{j}} + \left(\frac{s}{\sqrt{2}}\right) \hat{\mathbf{k}}$$

MATH-3105

Dept. of CSE

74

74

## Finding Arc Length Parametrizations

### Example 7 (p863)

A bug walks along the trunk of a tree following a path modeled by

$$\underline{\mathbf{r}}(t) = \cos t \, \hat{\mathbf{i}} + \sin t \, \hat{\mathbf{j}} + t \, \hat{\mathbf{k}}.$$

The bug starts at the reference point (1, 0, 0) and walks up the helix for a distance of 10 units. What are the bug's final coordinates?

MATH-3105

Dept. of CSE

75

75

## Finding Arc Length Parametrizations

### Solution (Example 7)

Replacing  $t$  by  $u$  in  $\underline{\mathbf{r}}$  for integration purpose, we have

$$\underline{\mathbf{r}}(t) = \cos u \, \hat{\mathbf{i}} + \sin u \, \hat{\mathbf{j}} + u \, \hat{\mathbf{k}}$$

$$\therefore \left\| \frac{d\underline{\mathbf{r}}}{du} \right\| = \sqrt{(-\sin u)^2 + \cos^2 u + 1} = \sqrt{2}$$

MATH-3105

Dept. of CSE

76

76

## Finding Arc Length Parametrizations

### Solution (Example 6)

Now taking  $t_0 = 0$

$$s = \int_0^t \left\| \frac{d\mathbf{r}}{du} \right\| du = \int_0^t \sqrt{2} du = \sqrt{2}u \Big|_0^t = \sqrt{2}t$$

Thus,  $t = s/\sqrt{2}$ , so the given equation can be reparametrized in terms of  $s$  as

$$\mathbf{r} = \cos\left(\frac{s}{\sqrt{2}}\right) \hat{\mathbf{i}} + \sin\left(\frac{s}{\sqrt{2}}\right) \hat{\mathbf{j}} + \left(\frac{s}{\sqrt{2}}\right) \hat{\mathbf{k}}$$

MATH-3105

Dept. of CSE

77

77

## Finding Arc Length Parametrizations

### Solution (Example 6)

Parametrically,

$$x = \cos\left(\frac{s}{\sqrt{2}}\right), \quad y = \sin\left(\frac{s}{\sqrt{2}}\right), \quad z = \frac{s}{\sqrt{2}}$$

Thus, at  $s = 10$  the coordinates are

$$(x, y, z) = \left( \cos\left(\frac{10}{\sqrt{2}}\right), \sin\left(\frac{10}{\sqrt{2}}\right), \frac{10}{\sqrt{2}} \right) \approx (0.705, 0.709, 7.07)$$

MATH-3105

Dept. of CSE

78

78

## Finding Arc Length Parametrizations

### Homework

Example 8 – 9 (p864)

## Chapter 12.3

### Homework

Exercise Set 12.3 (p866 –868)