

NaturalApproaches.com

Personalised Wellness Scan Report

Wellness screening summary

Patient details

Name: Test Test

Sex: Male

Age: 58

Height: 184.0 cm

Weight: 90.0 kg

Profile: Male

Scan time: 21/03/2026 21:03

This report is for wellness education and discussion only. It is not a medical diagnosis.

Report generated: 18/04/2026 09:58 BST

Eze Chiana
care@naturalapproaches.com
https://www.naturalapproaches.com

NAVIGATION

Table of contents

Cover

At a glance

Scan score overview

Priority category review

Body composition analysis

Clinical recommendations

Recommended support plan

Complete product recommendation mapping

Complete system marker tables

Appendix

OVERVIEW

At a glance

This section summarises the dominant patterns in the scan, the wider downstream themes they may be influencing, and the immediate priorities for follow-up.

Clinical snapshot

The scan is most suggestive of reduced digestive and absorptive efficiency, with likely downstream effects on repair, resilience, and wider function. There is also a secondary picture of barrier-related inflammatory stress and reduced energy-production efficiency.

Practitioner overview

Taken together, the findings point most strongly to reduced digestive and absorptive efficiency. Priority should be given to digestive support, absorptive support, and mucosal repair.

Why this matters

This matters because reduced digestive and absorptive efficiency can translate into poorer nutrient availability, slower repair, reduced resilience, and a tendency for other systems to become more reactive over time. It may also help explain the overlap with mitochondrial and energy-production strain and connective-tissue and repair pattern.

Primary pattern

Absorption and assimilation strain

This pattern points to reduced digestive and absorptive efficiency, most evident across digestive support, absorptive support, mucosal repair, and foundational nutrient repletion.

Follow-up focus: digestive support, absorptive support, mucosal repair, foundational nutrient repletion

Contributing patterns

- **Mitochondrial and energy-production strain** — Reduced energy-production efficiency, most evident across mitochondrial support, cofactor repletion, and energy-production support.
- **Connective-tissue and repair pattern** — Connective-tissue and repair pattern, most evident across repair support, connective-tissue support, and collagen support.
- **Toxic burden and detoxification strain** — Toxic-load pressure, most evident across exposure review, detoxification support, and antioxidant reserve.

Immediate priorities for follow-up

- From a clinical perspective, the most relevant next steps are to focus on digestive support, absorptive support, mucosal repair, foundational nutrient repletion, and mitochondrial support. Addressing these areas is likely to produce the greatest overall shift in the current pattern.

OVERVIEW

Health Index overview

This section provides a weighted overview of the scan, helping organise attention across major body systems before moving into more detailed interpretation.



Overall Scan Score

Generally Balanced

- Robust functional balance (85-100)
- Generally balanced (70-84)
- Needs focused support (55-69)
- Priority review recommended (0-54)

How to read this dashboard

This dashboard provides a weighted clinical overview of the scan. The Health Index and body-system cards are not based on a simple count of flagged markers. They are derived from the severity, distribution, and clustering of findings across related sections, with greater weighting given to patterns that may carry broader functional significance.

Each body-system card may include more than one underlying scan section, which is why the totals shown here may differ from the marker counts shown later on individual section pages. In general, lower scores suggest a more active pattern worth reviewing in follow-up, while higher scores suggest a more stable pattern on this dashboard view.

These scores are designed to help organise clinical attention. They are not a diagnosis and should always be interpreted alongside symptoms, history, and appropriate clinical assessment.

Renal & Fluid Balance



3 flagged · 1 within range

Includes: Kidney Function

Digestive



9 flagged · 10 within range

Includes: Gastrointestinal Function, Large Intestine Function, Liver Function, Gallbladder Function

Endocrine & Hormonal



13 flagged · 14 within range

Includes: Endocrine System, Thyroid, Prostate, Male Sexual Function, Sperm and semen, Male Hormone

Skin & Barrier



12 flagged · 6 within range

Includes: Skin, Eye

Cardiovascular



11 flagged · 18 within range

Includes: Cardiovascular and Cerebrovascular, Pulse of heart and brain, Blood lipids

Metabolic



24 flagged · 47 within range

Includes: Pancreatic Function, Blood Sugar, Trace Element, Vitamin, Amino Acid, Coenzyme, Essential Fatty Acid, Obesity, Lecithin, Fatty acid

Musculoskeletal



25 flagged · 27 within range

Includes: Bone Disease, Bone Mineral Density, Rheumatoid Bone Disease, Bone Growth Index, Collagen, Channels and collaterals

Immune & Inflammatory



14 flagged · 25 within range

Includes: Immune System, Human Toxin, Heavy Metal, Allergy, Human Immunity

Neurocognitive



7 flagged · 15 within range

Includes: Brain Nerve, Human Consciousness Level

Respiratory



2 flagged · 6 within range

Includes: Lung Function, Respiratory Function

Other



1 flagged · 4 within range

Includes: Basic Physical Quality

Priority overview

Prostate

Priority: Medium priority

Flagged markers: 3

Section Index: 56

Obesity

Priority: Medium priority

Flagged markers: 4

Section Index: 62

Gastrointestinal Function

Priority: Medium priority

Flagged markers: 3

Section Index: 65

Large Intestine Function

Priority: Medium priority

Flagged markers: 3

Section Index: 65

Male Sexual Function

Priority: Medium priority

Flagged markers: 3

Section Index: 67

Kidney Function

Priority: Low priority

Flagged markers: 3

Section Index: 72

RECOMMENDATIONS

Clinical recommendations

This section translates the scan findings into high-level clinical priorities, showing which functional areas appear most relevant for follow-up support.

Nutrient repletion

Given the dominant pattern, priority should be given to restoring nutrient sufficiency and absorptive capacity, particularly in the context of reduced digestive and absorptive efficiency.

Why this was suggested: Vitamin contains 6 flagged markers, led by Vitamin C, Vitamin E, Vitamin A.

Inflammation and barrier support

Given the broader scan picture, priority should be given to reducing inflammatory load while strengthening barrier integrity, particularly in the context of reduced digestive and absorptive efficiency.

Why this was suggested: Allergy contains 6 flagged markers, led by Hair dye allergy index, Injection allergy index, Chemical Sensitivity.

Visual and microvascular support

Support should focus on visual and microvascular function.

Why this was suggested: Visual and microvascular support contains 6 flagged markers, led by Edema, Collagen eye wrinkle, Dark circles.

Metabolic regulation and weight balance

Support should focus on metabolic balance and weight regulation, particularly in the context of reduced energy-production efficiency.

Why this was suggested: Metabolic regulation and weight balance contains 4 flagged markers, led by Abnormal lipid metabolism coefficient, Brown adipose tissue abnormalities coefficient, Triglyceride content of abnormal coefficient.

Gut function

Support digestive function, gut motility, and nutrient absorption so downstream inflammation and energy strain are not perpetuated.

Why this was suggested: Gastrointestinal Function contains 3 flagged markers, led by Gastric Absorption Function Coefficient, Small Intestine Motility, Small Intestine Absorption.

RECOMMENDATIONS

Recommended support plan

This section organises suggested support options into a staged plan, combining foundational and targeted measures based on the dominant scan patterns.

Absorption and assimilation strain Support Plan

This plan focuses on addressing the primary functional pattern, starting with foundational support before moving into targeted optimisation.

This plan is organised around reduced digestive and absorptive efficiency, with additional support for barrier-related inflammatory stress and reduced energy-production efficiency.

Phase 1 — Foundation

This phase focuses on core digestive, nutritional, and repair support.

Digestive Bitters

Natural Approaches

Core

Included to strengthen digestive support within the broader absorption and assimilation strain picture.

Gut Repair Complex

Natural Approaches

Core

Included to strengthen mucosal repair within the broader absorption and assimilation strain picture.

Mineral Repletion Blend

Natural Approaches

Included to strengthen nutrient repletion within the broader absorption and assimilation strain picture.

Phase 2 — Targeted Support

This phase adds more specific support aligned with the strongest secondary patterns.

Barrier Repair Complex

Natural Approaches

Core

Included to strengthen barrier support within the broader inflammatory and barrier dysfunction pattern picture.

Inflammation Support Formula

Natural Approaches

Core

Included to strengthen inflammation reduction within the broader inflammatory and barrier dysfunction pattern picture.

REFERENCE MAPPING

Complete product recommendation mapping

This section lists all products identified through the scan analysis. It includes full mapping across systems and markers to support practitioner review, product selection, and fulfilment planning.

Product	Source	Systems	Key markers
Barrier Repair Complex	Natural Approaches	Inflammatory and barrier dysfunction pattern, barrier support	
Digestive Bitters	Natural Approaches	Absorption and assimilation strain, digestive support	
Gut Repair Complex	Natural Approaches	Absorption and assimilation strain, mucosal repair	
Inflammation Support Formula	Natural Approaches	Inflammatory and barrier dysfunction pattern, inflammation reduction	
Mineral Repletion Blend	Natural Approaches	Absorption and assimilation strain, nutrient repletion	

Prostate

Medium priority

Priority interpretation

Prostate is one of the more active sections in this scan, with multiple flagged markers combining into a broader functional pattern.

Moderately and severely abnormal markers are prioritised here for faster clinical reading.

Flagged markers: 3 | Within range: 0 | Section Index: 56

Top findings

- Prostatitis Syndrome is moderately elevated
- Degree of Prostatic Hyperplasia is moderately elevated
- Degree of Prostatic Calcification is mildly elevated

Priority marker explanations

Prostatitis Syndrome

Result: 6.11 | **Reference range:** 2.213 - 2.717 | Moderately elevated

See Appendix for full clinical interpretation of this marker.

Degree of Prostatic Hyperplasia

Result: 5.03 | **Reference range:** 1.023 - 3.230 | Moderately elevated

See Appendix for full clinical interpretation of this marker.

Obesity

Medium priority

Priority interpretation

Obesity is one of the more active sections in this scan, with multiple flagged markers combining into a broader functional pattern.

Moderately and severely abnormal markers are prioritised here for faster clinical reading.

Flagged markers: 4 | Within range: 1 | Section Index: 62

Top findings

- Triglyceride content of abnormal coefficient is moderately elevated
- Brown adipose tissue abnormalities coefficient is moderately reduced
- Abnormal lipid metabolism coefficient is moderately reduced

Priority marker explanations

Triglyceride content of abnormal coefficient

Result: 5.222 | **Reference range:** 1.341 - 1.991 | Moderately elevated

See Appendix for full clinical interpretation of this marker.

Brown adipose tissue abnormalities coefficient

Result: 1.858 | **Reference range:** 2.791 - 4.202 | Moderately reduced

See Appendix for full clinical interpretation of this marker.

Abnormal lipid metabolism coefficient

Result: 0.89 | **Reference range:** 1.992 - 3.713 | Moderately reduced

See Appendix for full clinical interpretation of this marker.

Gastrointestinal Function

Medium priority

Priority interpretation

Gastrointestinal Function is one of the more active sections in this scan, with multiple flagged markers combining into a broader functional pattern.

Moderately and severely abnormal markers are prioritised here for faster clinical reading.

Flagged markers: 3 | Within range: 2 | Section Index: 65

Top findings

- Small Intestine Peristalsis Function Coefficient is moderately reduced
- Small Intestine Absorption Function Coefficient is moderately reduced
- Gastric Absorption Function Coefficient is moderately reduced

Priority marker explanations

Small Intestine Motility

Software marker: Small Intestine Peristalsis Function Coefficient

Result: 126.057 | **Reference range:** 133.437 - 140.476 | Moderately reduced

See Appendix for full clinical interpretation of this marker.

Small Intestine Absorption

Software marker: Small Intestine Absorption Function Coefficient

Result: 2.82 | **Reference range:** 3.572 - 6.483 | Moderately reduced

See Appendix for full clinical interpretation of this marker.

Gastric Absorption Function Coefficient

Result: 29.734 | **Reference range:** 34.367 - 35.642 | Moderately reduced

See Appendix for full clinical interpretation of this marker.

Large Intestine Function

Medium priority

Priority interpretation

Large Intestine Function is one of the more active sections in this scan, with multiple flagged markers combining into a broader functional pattern.

Moderately and severely abnormal markers are prioritised here for faster clinical reading.

Flagged markers: 3 | Within range: 1 | Section Index: 65

Top findings

- Large intestine peristalsis function coefficient is moderately reduced
- Intestinal bacteria coefficient is moderately reduced
- Intraluminal pressure coefficient is mildly elevated

Priority marker explanations

Colon Motility

Software marker: Large intestine peristalsis function coefficient

Result: 3.088 | **Reference range:** 4.572 - 6.483 | Moderately reduced

See Appendix for full clinical interpretation of this marker.

Gut Microbiome

Software marker: Intestinal bacteria coefficient

Result: 0.807 | **Reference range:** 1.734 - 2.621 | Moderately reduced

See Appendix for full clinical interpretation of this marker.

Male Sexual Function

Medium priority

Priority interpretation

Male Sexual Function contains a meaningful cluster of non-optimal markers, enough to warrant interpretation alongside symptoms, history, and related sections.

Moderately and severely abnormal markers are prioritised here for faster clinical reading.

Flagged markers: 3 | Within range: 0 | Section Index: 67

Top findings

- Testosterone is moderately reduced
- Gonadotropin is mildly reduced
- Erection Transmitter is mildly reduced

Priority marker explanations

Testosterone

Result: 2.492 | **Reference range:** 3.342 - 9.461 | Moderately reduced

See Appendix for full clinical interpretation of this marker.

Kidney Function

Low priority

Priority interpretation

Kidney Function contains a meaningful cluster of non-optimal markers, enough to warrant interpretation alongside symptoms, history, and related sections.

Moderately and severely abnormal markers are prioritised here for faster clinical reading.

Flagged markers: 3 | Within range: 1 | Section Index: 72

Top findings

- Uric acid Index is moderately elevated
- Urobilinogen Index is mildly elevated
- Proteinuria Index is mildly elevated

Priority marker explanations

Uric acid Index

Result: 2.689 | **Reference range:** 1.435 - 1.987 | Moderately elevated

See Appendix for full clinical interpretation of this marker.

DEDICATED SUMMARY

Body Composition & Metabolic Profile

This section summarises QRMA-derived body composition and metabolic profile indicators, including hydration, lean tissue, adiposity, and weight-related measures.

Within range

QRMA category: Element of Human

Section interpretation

This section summarises QRMA-derived body composition and metabolic profile indicators, including hydration, lean tissue, adiposity, and weight-related measures. Body fat percentage appears broadly reasonable for the profile shown. Where QRMA terminology is unconventional, clinically clearer labels are shown while preserving the original category reference.

- Hydration markers appear broadly balanced, without an obvious fluid-distribution concern on this page.
- Body fat percentage sits in a broadly reasonable range for a male profile.
- Body mass index is above the conventional range and may reflect increased metabolic load.
- Current weight sits materially above the QRMA target-weight estimate, so weight-related metrics should be interpreted alongside metabolic and inflammatory findings.
- Abdominal fat ratio is relatively high, which may align with greater central metabolic burden.

This section presents body-composition findings in a dedicated summary layout so the scan is easier to review clinically.

Fluid & Hydration

Intracellular fluid (QRMA)	21.6
Extracellular fluid (QRMA)	11.1

Lean Tissue & Structural Mass

Protein mass (QRMA)	8.58
Mineral mass (QRMA)	27
Muscle mass / muscle volume	41.3Kg
Body cell mass (BCM)	30.26 Kg
Muscle mass change target	0 Kg

Adiposity & Fat Distribution

Body fat mass (QRMA)	21.7
Body fat mass	21.7Kg
Body fat percentage	24.1%
Abdominal fat ratio	1.04
Fat mass change target	17.2 Kg

Weight & Metabolic Indicators

Height	184cm
Body weight	90kg
Obesity degree of body (QRMA)	124%
Body mass index (BMI)	26.6 kg/m²
Basal metabolic rate (BMR)	1825 kcal/day
Target weight	72.8 Kg
Weight change target	72.8 Kg

Other QRMA Body Composition Indices

Body shape / form assessment	63.3 Standard

TECHNICAL REFERENCE

Complete system marker tables

This section includes every parsed system and every marker across the report, including those within range, as the full explained reference layer.

Cardiovascular and Cerebrovascular

High priority

Complete system marker table — all markers and all severity grades are shown below.

Flagged markers: 4 | Within range: 11

Blood Viscosity

Within range

Cardiovascular and Cerebrovascular

Reference range: 48.264 - 65.371 | **Value:** 53.723

What this means: Reflects the thickness and flow resistance of blood, indicating how efficiently it can move through vessels and deliver oxygen and nutrients to tissues.

Why it matters: Elevated viscosity increases circulatory strain, impairs microvascular perfusion, and raises the workload on the heart while reducing tissue oxygen delivery.

Functional significance: Closely linked to hydration status, haematocrit balance, inflammatory proteins, lipid burden, red-cell aggregation, and endothelial function.

Common patterns: High viscosity -> reduced microcirculation -> lower tissue oxygenation -> fatigue, headaches, and cardiovascular stress; inflammatory and lipid burdens often intensify the pattern.

Your result in this scan: Blood Viscosity appears within the analyser's expected range in this scan.

Cholesterol Crystal

Within range

Cardiovascular and Cerebrovascular

Reference range: 56.749 - 67.522 | **Value:** 66.455

What this means: Reflects altered cholesterol handling and a tendency toward less soluble or more atherogenic lipid states within circulation.

Why it matters: Disturbed cholesterol dynamics can indicate vascular risk, inflammatory oxidation of lipids, impaired bile-dependent cholesterol disposal, and broader cardiometabolic strain.

Functional significance: Interacts with liver cholesterol metabolism, bile secretion, oxidative stress, thyroid signalling, and inflammatory modification of lipid particles.

Common patterns: Poor bile flow + liver congestion → weaker cholesterol disposal; oxidative stress can make lipid particles more damaging to vascular endothelium and amplify inflammatory risk.

Your result in this scan: Cholesterol Crystal appears within the analyser's expected range in this scan.

Blood Fat

Within range

Cardiovascular and Cerebrovascular

Reference range: 0.481 - 1.043 | **Value:** 0.902

What this means: Reflects the circulating fat burden within blood, indicating how effectively lipids are being transported, cleared, and metabolically handled.

Why it matters: Raised circulating fat is a functional marker of dyslipidaemia, insulin resistance, hepatic lipid overproduction, and increased vascular-inflammatory risk.

Functional significance: Linked to liver fat handling, insulin sensitivity, thyroid function, mitochondrial beta oxidation, and the balance between storage and utilisation of fuels.

Common patterns: Insulin resistance + hepatic lipogenesis -> elevated circulating fat -> endothelial stress and vascular burden; low thyroid and poor oxidation can worsen persistence.

Your result in this scan: Blood Fat appears within the analyser's expected range in this scan.

Vascular Resistance

Mildly elevated

Cardiovascular and Cerebrovascular

Reference range: 0.327 - 0.937 | **Value:** 1.226

What this means: Reflects the resistance encountered by blood as it moves through the vascular tree, indicating vessel tone, flexibility, and microvascular openness.

Why it matters: Higher vascular resistance forces the heart to work harder and reduces downstream tissue perfusion, increasing the risk of hypertension-type strain and poor oxygen delivery.

Functional significance: Linked to endothelial nitric oxide signalling, arterial elasticity, blood viscosity, autonomic tone, inflammation, and smooth muscle constriction.

Common patterns: Stress + endothelial dysfunction + high viscosity -> increased vascular resistance -> higher cardiac workload and lower tissue perfusion.

Your result in this scan: Indicates vasoconstrictive stress, endothelial dysfunction, elevated blood thickness, sympathetic overdrive, or reduced arterial flexibility.

Vascular Elasticity

Within range

Cardiovascular and Cerebrovascular

Reference range: 1.672 - 1.978 | **Value:** 1.934

What this means: Reflects the compliance and recoil capacity of blood vessels, indicating how well vascular connective tissue can expand and return without excessive stiffness.

Why it matters: Healthy vascular elasticity is essential for stable blood pressure, microcirculatory delivery, and protection against endothelial and cardiovascular strain.

Functional significance: Depends on collagen-elastin integrity, magnesium status, oxidative stress control, glycation burden, inflammatory tone, and endothelial nitric oxide signalling.

Common patterns: Oxidative stress, high blood sugar, and low magnesium can reduce vascular elasticity; collagen-elastin degradation increases cardiovascular burden and reduces tissue perfusion efficiency.

Your result in this scan: Vascular Elasticity appears within the analyser's expected range in this scan.

Myocardial Blood Demand

Within range

Cardiovascular and Cerebrovascular

Reference range: 0.192 - 0.412 | **Value:** 0.407

What this means: Reflects the oxygen and nutrient demand placed on cardiac muscle tissue during its workload.

Why it matters: When myocardial demand rises disproportionately, the heart becomes more vulnerable to ischemic strain, reduced efficiency, and fatigue under stress.

Functional significance: Linked to heart rate, pressure load, ventricular workload, oxygen delivery, mitochondrial energy production, and vascular supply adequacy.

Common patterns: High vascular resistance + reduced perfusion + elevated sympathetic tone -> increased myocardial demand with reduced reserve margin.

Your result in this scan: Myocardial Blood Demand appears within the analyser's expected range in this scan.

Myocardial Perfusion

Within range

Cardiovascular and Cerebrovascular

Reference range: 4.832 - 5.147 | **Value:** 5.132

What this means: Reflects the adequacy of blood flow reaching cardiac muscle tissue, indicating how well the myocardium is being nourished and oxygenated.

Why it matters: Cardiac muscle depends on continuous perfusion to maintain rhythmic contraction and metabolic stability; impaired perfusion increases the risk of ischemic stress and reduced output.

Functional significance: Linked to coronary circulation, blood viscosity, perfusion pressure, endothelial function, and the balance between myocardial demand and supply.

Common patterns: Low perfusion + high myocardial demand -> cardiac strain; viscosity, endothelial dysfunction, or reduced coronary pressure can all reduce perfusion volume.

Your result in this scan: Myocardial Perfusion appears within the analyser's expected range in this scan.

Myocardial Oxygen Consumption

Within range

Cardiovascular and Cerebrovascular

Reference range: 3.321 - 4.244 | **Value:** 3.382

What this means: Reflects the oxygen utilisation requirement of heart muscle tissue during its pumping activity.

Why it matters: Cardiac muscle has a very high metabolic demand, so increased oxygen consumption without matched delivery can indicate strain and reduced energetic efficiency.

Functional significance: Linked to myocardial workload, mitochondrial ATP generation, coronary perfusion, heart rate, and oxygen transport capacity.

Common patterns: High workload + reduced coronary supply -> mismatch between oxygen consumption and delivery; stress and vascular resistance often increase this mismatch.

Your result in this scan: Myocardial Oxygen Consumption appears within the analyser's expected range in this scan.

Stroke Volume

Mildly reduced

Cardiovascular and Cerebrovascular

Reference range: 1.338 - 1.672 | **Value:** 1.29

What this means: Reflects the volume of blood ejected by the heart with each beat, indicating pumping efficiency and circulatory support capacity.

Why it matters: Stroke volume is central to tissue perfusion, exercise tolerance, and the efficiency with which the heart can meet systemic demands.

Functional significance: Linked to preload, myocardial contractility, afterload, ventricular filling, and overall cardiac-muscular energy availability.

Common patterns: High resistance + weak contractility -> reduced stroke volume; low stroke volume can contribute to fatigue, poor exercise tolerance, and reduced tissue perfusion.

Your result in this scan: Suggests weaker pumping efficiency, reduced ventricular filling or contractility, or increased afterload impairing forward flow.

LV Ejection Impedance

Within range

Cardiovascular and Cerebrovascular

Reference range: 0.669 - 1.544 | **Value:** 1.456

What this means: Reflects the resistance faced by the left ventricle when ejecting blood into the arterial system, indicating afterload-related strain.

Why it matters: Greater ejection impedance increases myocardial workload and can reduce effective output while contributing to long-term ventricular stress.

Functional significance: Linked to arterial stiffness, vascular resistance, blood pressure patterns, endothelial tone, and left-ventricular workload.

Common patterns: Arterial stiffness + vascular resistance -> higher ejection impedance -> increased myocardial demand and reduced pump efficiency.

Your result in this scan: LV Ejection Impedance appears within the analyser's expected range in this scan.

LV Pump Power

Moderately reduced

Cardiovascular and Cerebrovascular

Reference range: 1.554 - 1.988 | **Value:** 0.991

What this means: Reflects the effective mechanical power generated by the left ventricle to sustain forward blood flow into systemic circulation.

Why it matters: This marker helps indicate whether the heart is generating adequate force to support circulation without excessive strain or inefficiency.

Functional significance: Linked to myocardial contractility, ventricular energy production, afterload, perfusion support, and systemic circulatory demand.

Common patterns: Reduced perfusion + higher afterload can lower effective pump power; mitochondrial and oxygen supply constraints may also impair ventricular performance.

Your result in this scan: Suggests reduced effective pump strength, weaker contractility, or reduced efficiency under afterload stress.

Coronary Artery Elasticity

Within range

Cardiovascular and Cerebrovascular

Reference range: 1.553 - 2.187 | **Value:** 1.66

What this means: Reflects the structural flexibility and recoil quality of coronary arterial tissue supplying the heart muscle.

Why it matters: Coronary elasticity influences myocardial perfusion reserve, blood flow responsiveness, and resilience against cardiovascular stress.

Functional significance: Closely linked to collagen-elastin quality, lipid oxidation, endothelial function, blood pressure regulation, and mineral balance.

Common patterns: Inflammation, dyslipidaemia, and glycation can reduce coronary elasticity and increase myocardial strain; low magnesium and endothelial stress may worsen the pattern.

Your result in this scan: Coronary Artery Elasticity appears within the analyser's expected range in this scan.

Coronary Perfusion Pressure

Moderately elevated

Cardiovascular and Cerebrovascular

Reference range: 11.719 - 18.418 | **Value:** 19.312

What this means: Reflects the pressure gradient driving blood flow through the coronary circulation to nourish heart muscle tissue.

Why it matters: Adequate coronary perfusion pressure is essential for delivering oxygen to the myocardium, especially during increased workload or stress.

Functional significance: Linked to aortic pressure, vascular tone, coronary flow dynamics, myocardial demand, and the adequacy of cardiac oxygen supply.

Common patterns: Low coronary perfusion pressure + high myocardial demand -> ischemic stress risk; vascular stiffness and endothelial dysfunction may impair effective coronary flow despite pressure changes.

Your result in this scan: Generally indicates stronger perfusion pressure support, although it must be interpreted in the context of vascular condition and myocardial demand.

Cerebral Blood Vessel Elasticity

Within range

Cardiovascular and Cerebrovascular

Reference range: 0.708 - 1.942 | **Value:** 1.021

What this means: Reflects the compliance and recoil of cerebral vessels, indicating how well connective tissue within brain circulation can adapt to flow and pressure changes.

Why it matters: Cerebrovascular elasticity is important for stable perfusion, oxygen delivery, and protection against pressure-related strain on neural tissue.

Functional significance: Interacts with vascular collagen-elastin integrity, blood viscosity, oxidative stress, magnesium status, and endothelial regulation.

Common patterns: High blood sugar, oxidative stress, and vascular inflammation can reduce cerebral vessel elasticity and contribute to headaches, brain fog, or perfusion instability.

Your result in this scan: Cerebral Blood Vessel Elasticity appears within the analyser's expected range in this scan.

Brain Tissue Blood Supply Status

Within range

Cardiovascular and Cerebrovascular

Reference range: 6.138 - 21.396 | **Value:** 20.196

What this means: Reflects the adequacy of cerebral perfusion reaching brain tissue, indicating how well oxygen and nutrients are being delivered to support neural metabolism.

Why it matters: The brain is highly dependent on uninterrupted blood flow, so this marker is relevant to cognition, concentration, autonomic regulation, and fatigue patterns.

Functional significance: Linked to vascular elasticity, blood viscosity, oxygen transport, microcirculation, endothelial function, and mitochondrial support within neural tissue.

Common patterns: High viscosity + poor vascular compliance -> reduced cerebral blood supply; low perfusion can amplify fatigue, poor concentration, and autonomic instability.

Your result in this scan: Brain Tissue Blood Supply Status appears within the analyser's expected range in this scan.

Gastrointestinal Function

High priority

Complete system marker table — all markers and all severity grades are shown below.

Flagged markers: 3 | Within range: 2

Pepsin Secretion Coefficient

Within range

Gastrointestinal Function

Reference range: 59.847 - 65.234 | **Value:** 59.915

What this means: Reflects gastric proteolytic capacity through pepsin secretion and activation, indicating how effectively the stomach can initiate protein digestion after adequate hydrochloric acid exposure.

Why it matters: Pepsin activity is essential for early-stage protein breakdown, liberation of amino acids, and preparation of nutrients for downstream pancreatic and intestinal digestion. Weak function can contribute to putrefaction, bloating, and poor protein assimilation.

Functional significance: Depends on sufficient gastric acidification, mucosal secretory integrity, vagal stimulation, and coordinated gastric phase digestion. It also influences antigen load reaching the intestine and subsequent immune activation.

Common patterns: Low hydrochloric acid → poor pepsin activation → incomplete protein digestion → intestinal fermentation/putrefaction → dysbiosis, bloating, and increased antigenic burden; this can secondarily increase hepatic detox and immune load.

Your result in this scan: Pepsin Secretion Coefficient appears within the analyser's expected range in this scan.

Gastric Peristalsis Function Coefficient

Within range

Gastrointestinal Function

Reference range: 58.425 - 61.213 | **Value:** 58.71

What this means: Reflects the motor efficiency of the stomach, including churning, mixing, and regulated emptying of gastric contents into the duodenum.

Why it matters: Effective gastric motility is required for mechanical breakdown of food, proper contact with acid and enzymes, and timely gastric emptying. Dysfunction contributes to fullness, reflux tendency, nausea, and upper abdominal distension.

Functional significance: Regulated by vagal tone, smooth muscle function, electrolyte balance, and pyloric coordination. It interfaces with reflux patterns, pancreatic timing, and small intestinal digestive readiness.

Common patterns: Low vagal tone or stress → reduced gastric motility → prolonged food retention → fermentation, reflux, and bloating; poor gastric emptying can also disturb pancreatic and biliary timing, worsening downstream digestion.

Your result in this scan: Gastric Peristalsis Function Coefficient appears within the analyser's expected range in this scan.

Gastric Absorption Function Coefficient

Moderately reduced

Gastrointestinal Function

Reference range: 34.367 - 35.642 | **Value:** 29.734

What this means: Reflects the stomach's limited but important absorptive and transport role, including handling of water, alcohol, certain medications, and support functions linked to intrinsic factor-dependent B12 assimilation downstream.

Why it matters: Although the stomach is not the primary absorptive organ, impaired gastric mucosal integrity can signal wider dysfunction in secretory and barrier roles that compromise nutrient handling and increase inflammatory sensitivity.

Functional significance: Closely linked to parietal cell function, intrinsic factor production, mucosal integrity, acid secretion, and the quality of transition into duodenal digestion and absorption.

Common patterns: Parietal cell dysfunction → low acid and low intrinsic factor → impaired B12 status → fatigue, neuropathy, and methylation compromise; mucosal injury can also increase sensitivity to reflux, gastritis, and upper GI inflammation.

Your result in this scan: Suggests gastric mucosal compromise, reduced intrinsic factor/parietal cell support, poor secretory integrity, or inflammatory impairment of the gastric lining.

Small Intestine Motility

Moderately reduced

Gastrointestinal Function

Reference range: 133.437 - 140.476 | **Value:** 126.057

What this means: Reflects coordinated small bowel motility, including segmental mixing and propulsive movement required for digestion, nutrient contact, and microbial regulation.

Why it matters: Adequate small intestinal motility prevents stasis, supports enzyme-substrate contact, and helps protect against bacterial overgrowth. Dysfunction is strongly linked to bloating, malabsorption, and SIBO-type presentations.

Functional significance: Depends on enteric nervous system signalling, vagal modulation, electrolyte balance, thyroid status, and smooth muscle responsiveness. It strongly influences transit time and absorptive efficiency.

Common patterns: Reduced small bowel motility → stasis → bacterial overgrowth → fermentation gases, mucosal irritation, and malabsorption; hypermotility can shorten nutrient contact time and contribute to loose stools and depletion.

Your result in this scan: Suggests stagnation, impaired migrating motor complex activity, autonomic dysfunction, hypothyroid tendency, or post-inflammatory motility suppression.

Small Intestine Absorption

Moderately reduced

Gastrointestinal Function

Reference range: 3.572 - 6.483 | **Value:** 2.82

What this means: Reflects absorptive efficiency of the small intestine across macronutrients, micronutrients, electrolytes, and water-soluble compounds, indicating the functional quality of villous and mucosal transport processes.

Why it matters: The small intestine is the main site of nutrient assimilation. Impaired function can produce broad downstream effects including fatigue, deficiency states, immune dysregulation, poor repair, and endocrine compromise.

Functional significance: Depends on villous integrity, brush border enzyme activity, bile and pancreatic support, mucosal permeability control, and healthy interaction with the microbiome.

Common patterns: Low bile or pancreatic enzyme output → incomplete breakdown → poor absorption; intestinal inflammation or permeability → nutrient deficiency + immune activation; malabsorption can secondarily impair thyroid, adrenal, mitochondrial, and detox pathways.

Your result in this scan: Suggests malabsorption, villous or brush-border compromise, inflammatory mucosal injury, pancreatic-biliary insufficiency, or dysbiosis-related impairment.

Large Intestine Function

High priority

Complete system marker table — all markers and all severity grades are shown below.

Flagged markers: 3 | Within range: 1

Colon Motility

Moderately reduced

Large Intestine Function

Reference range: 4.572 - 6.483 | **Value:** 3.088

What this means: Reflects colonic motility and propulsive function, including transit dynamics required for stool formation, waste movement, and timely elimination.

Why it matters: Healthy colonic transit reduces toxin reabsorption, supports microbiome balance, and prevents stagnation-associated constipation, bloating, and inflammatory burden.

Functional significance: Influenced by hydration status, magnesium and electrolyte balance, autonomic tone, fibre-fermentation dynamics, thyroid function, and pelvic floor or smooth muscle coordination.

Common patterns: Slow colonic transit → prolonged toxin and oestrogen metabolite contact → reabsorption and hepatic burden; constipation also shifts microbial ecology toward dysbiosis and inflammatory metabolite production.

Your result in this scan: Suggests sluggish transit, constipation tendency, autonomic suppression, dehydration, low magnesium status, or hypothyroid-associated bowel slowing.

Colon Absorption

Within range

Large Intestine Function

Reference range: 2.946 - 3.815 | **Value:** 2.952

What this means: Reflects absorptive activity of the colon, particularly water, electrolytes, and microbiome-derived metabolites such as short-chain fatty acids.

Why it matters: Balanced colonic absorption is important for stool consistency, mucosal energy supply, electrolyte stability, and maintenance of a favourable microbial environment.

Functional significance: Closely tied to microbiome fermentation quality, mucosal integrity, transit time, inflammation, and hydration-electrolyte regulation.

Common patterns: Slow transit + excessive fluid reabsorption → dry stools and constipation; dysbiosis → reduced butyrate production → weaker mucosal repair and higher inflammatory susceptibility.

Your result in this scan: Colon Absorption appears within the analyser's expected range in this scan.

Gut Microbiome

Moderately reduced

Large Intestine Function

Reference range: 1.734 - 2.621 | **Value:** 0.807

What this means: Reflects the functional state and relative balance of intestinal microbial ecology, including commensal support, fermentative activity, and microbial pressure within the gut lumen.

Why it matters: The intestinal microbiome shapes digestion, vitamin production, immune education, barrier integrity, oestrogen metabolism, and inflammatory signalling throughout the body.

Functional significance: Interacts directly with bile acids, fibre fermentation, mucosal immunity, short-chain fatty acid generation, endotoxin production, and enterohepatic circulation.

Common patterns: Dysbiosis → impaired barrier function → endotoxin translocation → hepatic and immune activation; altered beta-glucuronidase activity → oestrogen recirculation; microbial imbalance also influences mood, skin, and metabolic signalling.

Your result in this scan: Suggests reduced microbial diversity or beneficial flora support, often associated with antimicrobial exposure, low fibre intake, poor bile flow, or chronic stress.

Intraluminal pressure coefficient

Mildly elevated

Large Intestine Function

Reference range: 1.173 - 2.297 | **Value:** 2.639

What this means: Reflects pressure dynamics within the intestinal lumen, influenced by gas production, motility coordination, stool transit, muscular tone, and bowel distension patterns.

Why it matters: Abnormal intraluminal pressure is commonly associated with bloating, pain, altered transit, spasm, constipation, and fermentative bowel states.

Functional significance: Links mechanical bowel function with motility, microbial gas generation, stool burden, visceral sensitivity, and autonomic regulation of the gut.

Common patterns: Poor motility + dysbiosis → fermentation gas accumulation → raised luminal pressure → bloating, discomfort, and altered peristalsis; chronic stool retention can further increase pressure and toxin reabsorption.

Your result in this scan: Suggests bowel distension, trapped gas, constipation, spasm, fermentation excess, or inflammatory-irritative pressure states.

Liver Function

Medium priority

Complete system marker table — all markers and all severity grades are shown below.

Flagged markers: 1 | Within range: 4

Protein Metabolism

Within range

Liver Function

Reference range: 116.34 - 220.621 | **Value:** 116.569

What this means: Represents the liver's role in processing, transforming, and regulating amino acid metabolism and nitrogen balance.

Why it matters: The liver is central to protein metabolism, influencing tissue repair, enzyme production, and detoxification through the urea cycle.

Functional significance: Linked to amino acid turnover, hepatic enzyme activity, urea cycle function, and systemic nitrogen balance.

Common patterns: Inflammation + metabolic stress -> increased amino acid turnover -> altered hepatic protein metabolism.

Your result in this scan: Protein Metabolism appears within the analyser's expected range in this scan.

Energy Production Function

Within range

Liver Function

Reference range: 0.713 - 0.992 | **Value:** 0.881

What this means: Reflects hepatic contribution to systemic energy production through glucose regulation, glycogen storage, and metabolic processing.

Why it matters: The liver plays a central role in maintaining energy stability via gluconeogenesis, glycogen storage, and metabolic flexibility.

Functional significance: Linked to mitochondrial function, glucose regulation, glycogen dynamics, and hepatic metabolic pathways.

Common patterns: Insulin resistance + mitochondrial dysfunction -> impaired hepatic energy regulation -> metabolic instability.

Your result in this scan: Energy Production Function appears within the analyser's expected range in this scan.

Detoxification Function

Within range

Liver Function

Reference range: 0.202 - 0.991 | **Value:** 0.977

What this means: Represents the liver’s integrated capacity to process and eliminate endogenous and exogenous toxins.

Why it matters: Hepatic detoxification is essential for clearing hormones, metabolic waste, and environmental toxins, influencing systemic health.

Functional significance: Linked to Phase I and Phase II pathways, glutathione status, and hepatobiliary excretion.

Common patterns: Toxin exposure + nutrient depletion -> impaired detox pathways -> increased oxidative and inflammatory stress.

Your result in this scan: Detoxification Function appears within the analyser's expected range in this scan.

Bile Secretion Function

Within range

Liver Function

Reference range: 0.432 - 0.826 | **Value:** 0.717

What this means: Reflects the liver’s ability to produce and regulate bile flow for digestion and elimination.

Why it matters: Bile is critical for fat digestion, cholesterol metabolism, and toxin excretion.

Functional significance: Linked to hepatobiliary flow, gallbladder function, and fat-soluble nutrient absorption.

Common patterns: Liver congestion + impaired flow -> fat malabsorption and toxin recirculation.

Your result in this scan: Bile Secretion Function appears within the analyser's expected range in this scan.

Liver Fat Content

Moderately elevated

Liver Function

Reference range: 0.097 - 0.419 | **Value:** 0.636

What this means: Represents the degree of lipid accumulation within hepatic tissue.

Why it matters: Excess liver fat impairs metabolic regulation, insulin sensitivity, and detoxification capacity.

Functional significance: Linked to insulin resistance, lipid metabolism, and mitochondrial function.

Common patterns: High carbohydrate intake + insulin resistance -> hepatic fat accumulation -> metabolic dysfunction.

Your result in this scan: Indicates increased fat accumulation, often associated with metabolic syndrome or NAFLD.

Gallbladder Function

Medium priority

Complete system marker table — all markers and all severity grades are shown below.

Flagged markers: 2 | Within range: 3

Serum Globulin (A/G)

Within range

Gallbladder Function

Reference range: 126 - 159 | **Value:** 126.767

What this means: Reflects the albumin-to-globulin relationship within the source gallbladder-bile-flow cluster, functioning as a broad marker of protein balance, hepatic synthetic context, immune burden, and chronic inflammatory load rather than a bile-specific analyte alone.

Why it matters: The albumin-globulin relationship matters because it can help reveal whether liver synthetic function, protein status, immune activation, or chronic inflammatory burden is altering the terrain in which bile production and hepatobiliary flow occur.

Functional significance: Linked to hepatic protein synthesis, chronic inflammation, immune activation, gut-derived antigenic burden, nutritional protein status, and the liver's broader metabolic and detoxification workload.

Common patterns: Gut-derived inflammation + chronic immune activation + liver synthetic strain -> altered A/G balance; when this occurs alongside bile stagnation or raised bilirubin markers, the wider hepatobiliary picture becomes more significant.

Your result in this scan: Serum Globulin (A/G) appears within the analyser's expected range in this scan.

Total Bilirubin (TBIL)

Within range

Gallbladder Function

Reference range: 0.232 - 0.686 | **Value:** 0.542

What this means: Reflects total bilirubin burden across conjugated and unconjugated fractions, indicating the liver's handling of haem breakdown products and the effectiveness of bile-dependent excretion pathways.

Why it matters: Bilirubin is one of the clearest hepatobiliary flow markers because elevation can indicate impaired conjugation, reduced bile transport, cholestatic tendency, or increased haem turnover overwhelming clearance capacity.

Functional significance: Linked to red-cell turnover, hepatic conjugation capacity, bile secretion, biliary drainage, gallbladder flow, and the distinction between pre-hepatic, hepatic, and post-hepatic contributors to bilirubin burden.

Common patterns: Haem breakdown + sluggish hepatic conjugation + poor bile flow -> higher TBIL; gallbladder stasis or biliary obstruction can push levels higher when drainage is impaired.

Your result in this scan: Total Bilirubin (TBIL) appears within the analyser's expected range in this scan.

Alkaline Phosphatase (ALP)

Within range

Gallbladder Function

Reference range: 0.082 - 0.342 | **Value:** 0.316

What this means: Reflects alkaline phosphatase activity, a marker especially relevant to cholestatic stress, bile-duct irritation, and hepatobiliary drainage resistance, while also requiring distinction from bone-derived ALP sources.

Why it matters: ALP matters because it often rises when bile cannot move freely through the canalicular and ductal system, making it clinically useful in detecting cholestatic or obstructive hepatobiliary patterns.

Functional significance: Linked to bile-duct epithelium, canalicular flow, gallbladder drainage, hepatic congestion, cholestasis, and - when interpreted carefully - bone turnover as a non-biliary confounder.

Common patterns: Cholestasis + bile-duct irritation + reduced gallbladder emptying -> raised ALP; when combined with bilirubin or bile-acid elevation, biliary flow impairment becomes more likely.

Your result in this scan: Alkaline Phosphatase (ALP) appears within the analyser's expected range in this scan.

Serum Total Bile Acid (TBA

Mildly elevated

Gallbladder Function

Reference range: 0.317 - 0.695 | **Value:** 0.704

What this means: Reflects the circulating burden of bile acids, indicating how effectively the liver synthesises, secretes, recycles, and clears bile within the enterohepatic system.

Why it matters: Total bile acids are highly relevant to hepatobiliary flow because they rise when bile secretion, transport, or recirculation becomes impaired, making them a sensitive marker of cholestatic stress.

Functional significance: Linked to hepatocyte bile synthesis, canalicular export, gallbladder emptying, intestinal reabsorption, enterohepatic circulation, microbiome transformation of bile acids, and overall bile-flow efficiency.

Common patterns: Liver congestion + gallbladder stasis + impaired biliary export -> raised bile acids; gut dysbiosis and poor intestinal reabsorption patterns may further distort enterohepatic balance.

Your result in this scan: Higher total bile acids indicate impaired bile flow, cholestasis, hepatic transport strain, or disrupted enterohepatic circulation affecting detoxification and fat handling.

Bilirubin (DBIL)

Mildly elevated

Gallbladder Function

Reference range: 0.218 - 0.549 | **Value:** 0.56

What this means: Reflects direct (conjugated) bilirubin, indicating the fraction that has already been processed by the liver and is awaiting effective excretion through bile pathways.

Why it matters: Direct bilirubin is particularly useful in identifying cholestatic or post-conjugation excretory problems because elevation suggests that conjugation may be occurring but biliary drainage is impaired.

Functional significance: Linked to hepatic conjugation quality, canalicular export, bile-duct patency, gallbladder-biliary flow, and the distinction between processing failure and drainage failure within the hepatobiliary system.

Common patterns: Conjugation preserved + bile export impaired -> higher DBIL; when paired with raised ALP and bile acids, the pattern points more strongly to cholestasis or biliary obstruction-type stress.

Your result in this scan: Higher direct bilirubin indicates impaired biliary excretion, cholestatic stress, or drainage resistance after bilirubin has already been conjugated by the liver.

Pancreatic Function

Within range

Complete system marker table — all markers and all severity grades are shown below.

Flagged markers: 0 | Within range: 3

Insulin

Within range

Pancreatic Function

Reference range: 2.845 - 4.017 | **Value:** 3.036

What this means: Reflects insulin output within the pancreatic endocrine axis, indicating beta-cell signalling involved in glucose uptake, blood-sugar regulation, and anabolic storage physiology.

Why it matters: Insulin is central to glucose homeostasis, fat storage, protein anabolism, and metabolic flexibility. Disturbance can indicate insulin resistance, beta-cell strain, reactive dysglycaemia, or broader metabolic dysfunction.

Functional significance: Linked to pancreatic beta-cell function, glucose exposure, insulin sensitivity, hepatic glucose output, muscle glucose uptake, adipose storage signalling, and inflammatory-metabolic stress.

Common patterns: High refined-carbohydrate load + visceral adiposity + inflammation -> insulin resistance -> compensatory insulin elevation; later-stage beta-cell fatigue may eventually reduce output.

Your result in this scan: Insulin appears within the analyser's expected range in this scan.

Pancreatic Polypeptide (PP)

Within range

Pancreatic Function

Reference range: 3.210 - 6.854 | **Value:** 3.26

What this means: Reflects pancreatic polypeptide signalling, a regulatory peptide involved in pancreatic secretory balance, gastrointestinal signalling, appetite modulation, and autonomic-pancreatic coordination.

Why it matters: Pancreatic polypeptide helps regulate exocrine pancreatic output, gastrointestinal motility, and neuroendocrine digestive signalling. Disturbance may reflect altered pancreatic regulation, vagal dysfunction, or broader digestive-endocrine imbalance.

Functional significance: Linked to pancreatic islet signalling, vagal tone, postprandial digestive regulation, exocrine pancreatic coordination, appetite control, and gut-pancreas autonomic communication.

Common patterns: Digestive dysregulation + vagal imbalance + pancreatic stress -> altered PP signalling; this may coexist with appetite irregularity, enzyme dysregulation, or broader gut-pancreas axis disturbance.

Your result in this scan: Pancreatic Polypeptide (PP) appears within the analyser's expected range in this scan.

Glucagon

Within range

Pancreatic Function

Reference range: 2.412 - 2.974 | **Value:** 2.666

What this means: Reflects glucagon output within the pancreatic alpha-cell axis, indicating counter-regulatory signalling that raises blood glucose during fasting or increased energy demand.

Why it matters: Glucagon is essential for metabolic flexibility because it mobilises stored fuel, supports fasting glucose maintenance, and counterbalances insulin. Dysregulation can contribute to unstable blood sugar, hepatic glucose excess, or poor fasting adaptation.

Functional significance: Linked to pancreatic alpha-cell function, hepatic glycogenolysis, gluconeogenesis, fasting adaptation, insulin-glucagon balance, and overall glucose-energy homeostasis.

Common patterns: Insulin resistance + hepatic metabolic dysregulation -> excessive glucagon influence -> higher fasting or post-absorptive glucose output; low glucagon may align with poor fasting tolerance.

Your result in this scan: Glucagon appears within the analyser's expected range in this scan.

Kidney Function

Medium priority

Complete system marker table — all markers and all severity grades are shown below.

Flagged markers: 3 | Within range: 1

Urobilinogen Index

Mildly elevated

Kidney Function

Reference range: 2.762 - 5.424 | **Value:** 6.215

What this means: Reflects the burden of urobilinogen within the renal-urinary context, indicating downstream handling of bilirubin metabolism and possible interaction between liver processing, gut conversion, and urinary excretion.

Why it matters: Urobilinogen patterns can provide indirect insight into haem breakdown, liver conjugation efficiency, bile flow, gut microbial conversion, and renal excretion of downstream pigment metabolites.

Functional significance: Linked to bilirubin turnover, hepatobiliary function, intestinal transformation of bilirubin, enterohepatic handling, and urinary elimination dynamics.

Common patterns: Liver stress + altered bile flow + gut conversion changes -> shifted urobilinogen output; interpretation should consider bilirubin and hepatobiliary markers rather than kidney function alone.

Your result in this scan: Indicates increased urobilinogen burden, potentially reflecting enhanced haem breakdown, altered hepatobiliary handling, or increased gut-derived conversion with downstream urinary excretion.

Uric acid Index

Moderately elevated

Kidney Function

Reference range: 1.435 - 1.987 | **Value:** 2.689

What this means: Reflects uric-acid burden within renal handling, indicating purine metabolism load and the kidney's role in excreting this metabolic end product.

Why it matters: Uric acid is clinically relevant to oxidative stress, metabolic syndrome, renal burden, vascular risk, and crystal-related inflammatory tendencies when clearance is impaired or production is excessive.

Functional significance: Linked to purine turnover, fructose burden, renal excretion efficiency, insulin resistance, dehydration status, and inflammatory-metabolic terrain.

Common patterns: Fructose excess + insulin resistance + dehydration + renal under-clearance -> higher uric-acid burden; this may overlap with metabolic syndrome and inflammatory joint patterns.

Your result in this scan: Indicates increased uric-acid burden, potentially associated with higher purine load, reduced renal clearance, insulin resistance, dehydration, or crystal-forming inflammatory risk.

BUN

Within range

Kidney Function

Reference range: 4.725 - 8.631 | **Value:** 4.766

What this means: Reflects the blood burden of urea nitrogen as a marker of protein-waste handling across the liver-kidney axis, especially renal excretion of nitrogenous end products.

Why it matters: BUN helps indicate whether protein metabolism, hydration state, hepatic urea production, and renal waste clearance are functioning in coordinated balance.

Functional significance: Linked to protein intake and breakdown, hepatic urea-cycle output, renal filtration, hydration status, and catabolic stress states.

Common patterns: Dehydration + catabolic stress + reduced renal filtration -> higher BUN burden; interpretation should distinguish pre-renal concentration effects from true renal under-clearance.

Your result in this scan: BUN appears within the analyser's expected range in this scan.

Proteinuria Index

Mildly elevated

Kidney Function

Reference range: 1.571 - 4.079 | **Value:** 4.761

What this means: Reflects urinary protein leakage, indicating compromise of the renal filtration barrier or altered glomerular permeability within the kidney axis.

Why it matters: Proteinuria is one of the most clinically important renal warning signals because it can indicate glomerular stress, endothelial dysfunction, inflammatory injury, or metabolic-kidney damage before more obvious decline appears.

Functional significance: Linked to glomerular basement membrane integrity, podocyte stability, blood-pressure load, endothelial health, insulin resistance, and inflammatory renal stress.

Common patterns: Hypertension + hyperglycaemia + endothelial damage -> protein leakage -> rising renal risk; proteinuria often parallels broader cardiovascular-metabolic stress.

Your result in this scan: Indicates increased protein leakage across the glomerular barrier, suggesting renal stress, endothelial dysfunction, inflammatory injury, or metabolic-kidney burden.

Lung Function

Low priority

Complete system marker table — all markers and all severity grades are shown below.

Flagged markers: 1 | Within range: 3

Vital Capacity

Within range

Lung Function

Reference range: 3348 - 3529 | **Value:** 3489

What this means: Reflects the maximal volume of air that can be exhaled after a full inhalation, providing a functional measure of lung expansion, ventilatory reserve, and chest-wall-respiratory muscle capacity.

Why it matters: Vital capacity helps indicate whether the lungs and thoracic cage can move adequate air. Reduced values often track with restrictive breathing patterns, respiratory muscle weakness, poor posture, chronic inflammation, or low exercise tolerance.

Functional significance: Linked to diaphragmatic function, chest-wall mobility, lung tissue elasticity, respiratory-muscle strength, and the degree to which inflammation, fibrosis, or deconditioning restrict expansion.

Common patterns: Poor posture + diaphragmatic weakness + inflammatory or fibrotic restriction -> lower VC; anxiety-breathing patterns and deconditioning can further reduce effective breathing capacity.

Your result in this scan: Vital Capacity appears within the analyser's expected range in this scan.

Total Lung Capacity

Mildly reduced

Lung Function

Reference range: 4301 - 4782 | **Value:** 4286

What this means: Reflects the total volume of air the lungs can contain at maximal inflation, indicating the overall size and expansion capacity of the pulmonary-thoracic system.

Why it matters: TLC helps differentiate restrictive from hyperinflation patterns and provides insight into whether lung tissue, pleural mechanics, or chest-wall movement are limiting respiration.

Functional significance: Linked to lung compliance, thoracic mobility, respiratory-muscle function, alveolar expansion, and the presence of restrictive tissue change or air trapping.

Common patterns: Restriction -> lower TLC; air trapping and obstructive patterns can raise TLC through hyperinflation rather than healthy expansion.

Your result in this scan: Suggests restrictive mechanics, reduced expansion capacity, fibrosis-type limitation, obesity-related restriction, or poor thoracic mobility.

Airway Resistance

Within range

Lung Function

Reference range: 1.374 - 1.709 | **Value:** 1.621

What this means: Reflects the degree of opposition to airflow through the bronchial tree, indicating how easily air can move through the conducting airways.

Why it matters: Airway resistance is a core determinant of breathing effort. Higher resistance increases work of breathing and may indicate bronchospasm, mucus burden, inflammation, or airway narrowing.

Functional significance: Linked to bronchial tone, smooth-muscle constriction, mucus load, airway calibre, autonomic regulation, and inflammatory-allergic respiratory reactivity.

Common patterns: Allergy + inflammation + autonomic overdrive -> higher airway resistance; mucus, irritant exposure, and poor respiratory regulation can sustain the pattern.

Your result in this scan: Airway Resistance appears within the analyser's expected range in this scan.

Arterial Oxygen Content

Within range

Lung Function

Reference range: 17.903 - 21.012 | **Value:** 20.78

What this means: Reflects a source-label gas-exchange marker best interpreted cautiously as arterial blood-gas balance, with the PaCO2 label pointing more directly to carbon-dioxide handling and ventilatory adequacy than to oxygen content.

Why it matters: Carbon-dioxide clearance is a critical part of respiratory physiology. Abnormal values can indicate hypoventilation, overbreathing, impaired gas exchange, altered acid-base balance, or respiratory compensation patterns.

Functional significance: Linked to alveolar ventilation, respiratory drive, acid-base buffering, airway patency, respiratory-muscle function, and the balance between oxygenation and carbon-dioxide clearance.

Common patterns: Poor ventilation + airway resistance + respiratory-muscle fatigue -> raised PaCO2-type burden; anxiety-hyperventilation can push values in the opposite direction.

Your result in this scan: Arterial Oxygen Content appears within the analyser's expected range in this scan.

Brain Nerve

Low priority

Complete system marker table — all markers and all severity grades are shown below.

Flagged markers: 1 | Within range: 4

Brain Tissue Blood Supply

Within range

Brain Nerve

Reference range: 143.37 - 210.81 | **Value:** 168.036

What this means: Reflects adequacy of cerebral perfusion to brain tissue, including microvascular blood flow, oxygen delivery, and nutrient availability required for neuronal metabolism.

Why it matters: The brain is highly dependent on continuous perfusion for ATP generation, neurotransmitter synthesis, and preservation of cognitive and autonomic function.

Functional significance: Integrates vascular tone, blood viscosity, oxygen transport, mitochondrial efficiency, endothelial integrity, and cerebrovascular autoregulation.

Common patterns: Poor vascular elasticity or increased blood viscosity → reduced microcirculatory flow → lower oxygen delivery → mitochondrial shortfall in neurons; chronic hypoperfusion can amplify brain fog, poor memory, headaches, and autonomic instability.

Your result in this scan: Brain Tissue Blood Supply appears within the analyser's expected range in this scan.

Cerebral Arteriosclerosis

Within range

Brain Nerve

Reference range: 0.103 - 0.642 | **Value:** 0.196

What this means: Reflects structural and functional hardening of cerebral vessels, indicating reduced arterial compliance, impaired endothelial regulation, and higher cerebrovascular resistance.

Why it matters: Loss of vascular flexibility in the brain can compromise perfusion reserve, oxygen delivery, and resilience to metabolic or cognitive demand.

Functional significance: Closely linked to lipid metabolism, oxidative stress, inflammatory vascular injury, hypertension patterns, and impaired nitric oxide signalling.

Common patterns: Inflammation + dyslipidaemia + oxidative stress → endothelial injury → vascular stiffening → reduced cerebral perfusion reserve; this can contribute to memory decline, headaches, and reduced stress tolerance.

Your result in this scan: Cerebral Arteriosclerosis appears within the analyser's expected range in this scan.

Cranial Nerve Function

Within range

Brain Nerve

Reference range: 0.253 - 0.659 | **Value:** 0.292

What this means: Reflects the functional integrity of cranial nerve signalling involved in sensory processing, autonomic regulation, facial motor function, swallowing, vestibular balance, and vagal communication.

Why it matters: Cranial nerve performance influences eye movement, hearing, balance, parasympathetic tone, speech, swallowing, and gut-brain regulation through vagal pathways.

Functional significance: Depends on myelin integrity, micronutrient sufficiency, stable blood supply, mitochondrial support, and low neuroinflammatory burden.

Common patterns: Low vagal tone → impaired parasympathetic regulation → poor digestion, reduced motility, and stress sensitivity; nerve irritation or nutrient depletion can also affect balance, sensory clarity, and autonomic stability.

Your result in this scan: Cranial Nerve Function appears within the analyser's expected range in this scan.

Sentiment Index

Within range

Brain Nerve

Reference range: 0.109 - 0.351 | **Value:** 0.262

What this means: Reflects functional emotional regulation capacity, incorporating neurochemical balance, limbic system tone, and stress-related affective responsiveness.

Why it matters: Mood regulation influences sleep, motivation, behavioural resilience, social function, and downstream endocrine-immune regulation through the stress axis.

Functional significance: Closely linked to serotonin, dopamine, GABA-glutamate balance, HPA axis tone, inflammatory cytokines, and gut-brain signalling.

Common patterns: Chronic stress or gut dysbiosis → altered neurotransmitter production + inflammatory cytokine signalling → low mood, irritability, and poor sleep; blood sugar instability can further aggravate affective volatility.

Your result in this scan: Sentiment Index appears within the analyser's expected range in this scan.

Memory

Mildly reduced

Brain Nerve

Reference range: 0.442 - 0.817 | **Value:** 0.379

What this means: Reflects the capacity for encoding, consolidation, and recall of information within the source cognitive-performance framework.

Why it matters: Memory is central to learning, academic performance, adaptive behaviour, and the ability to integrate past experience into current functioning.

Functional significance: Linked to sleep quality, hippocampal support, attention, methylation status, energy availability, and neuroinflammatory burden.

Common patterns: Poor sleep + inattentive intake + inflammatory fatigue -> weaker memory; energy and attentional support often improve this domain.

Your result in this scan: Suggests weaker encoding or recall, often associated with stress, fatigue, sleep disruption, or poor attentional intake.

Bone Disease

Within range

Complete system marker table — all markers and all severity grades are shown below.

Flagged markers: 0 | Within range: 7

Adhesion Degree of Shoulder Muscle

Unclear

Bone Disease

Reference range: — | **Value:** < u 0.2

What this means: Reflects fibrotic adhesion and reduced glide between muscle and fascial layers in the shoulder complex.

Why it matters: Adhesions restrict mobility, impair circulation, and prolong recovery after injury.

Functional significance: Linked to chronic inflammation, microtrauma, and impaired lymphatic drainage.

Common patterns: Inflammation + immobility -> fascial adhesion formation.

Your result in this scan: Adhesion Degree of Shoulder Muscle appears within the analyser's expected range in this scan.

Age of Ligament

Unclear

Bone Disease

Reference range: — | **Value:** 10%-40%

What this means: Reflects the functional ageing state of ligament tissue, indicating the degree of connective tissue resilience, elasticity, and structural wear within supportive joint tissues.

Why it matters: Ligament quality affects joint stability, injury risk, recovery capacity, and the overall biomechanical integrity of the musculoskeletal system.

Functional significance: Depends on collagen turnover, mineral support, hydration, inflammatory load, and cumulative mechanical stress on connective tissues.

Common patterns: Oxidative stress, low collagen support, and chronic inflammatory or mechanical load can accelerate ligament ageing and reduce tissue recovery capacity.

Your result in this scan: Age of Ligament appears within the analyser's expected range in this scan.

Bone Mineral Density

Medium priority

Complete system marker table — all markers and all severity grades are shown below.

Flagged markers: 2 | Within range: 3

Osteoclast Coefficient

Within range

Bone Mineral Density

Reference range: 86.73 - 180.97 | **Value:** 151.19

What this means: Reflects activity of osteoclasts responsible for bone resorption in bone remodelling.

Why it matters: Balance between osteoclast and osteoblast activity determines bone density.

Functional significance: Linked to PTH signalling, vitamin D status, inflammatory cytokines, and acid-base balance.

Common patterns: Inflammation + hormonal imbalance -> increased osteoclast activity.

Your result in this scan: Osteoclast Coefficient appears within the analyser's expected range in this scan.

Amount of Calcium Loss

Within range

Bone Mineral Density

Reference range: 0.209 - 0.751 | **Value:** 0.469

What this means: Represents net calcium loss from bone stores, reflecting mineral depletion.

Why it matters: Calcium is fundamental for bone strength and neuromuscular function.

Functional significance: Linked to vitamin D, parathyroid function, magnesium balance, and diet.

Common patterns: Low vitamin D + high PTH -> calcium leaching from bone.

Your result in this scan: Amount of Calcium Loss appears within the analyser's expected range in this scan.

Degree of Bone Hyperplasia

Mildly elevated

Bone Mineral Density

Reference range: 0.046 - 0.167 | **Value:** 0.389

What this means: Reflects the extent of reactive or excessive bone overgrowth within skeletal tissue.

Why it matters: Bone hyperplasia may indicate chronic irritation, altered remodelling, degenerative adaptation, or prolonged mechanical stress.

Functional significance: Interacts with mineral turnover, inflammatory signalling, structural load, and the balance between degeneration and repair.

Common patterns: Chronic joint stress and inflammatory degeneration can promote bone overgrowth; mineral imbalance may further distort remodelling quality.

Your result in this scan: Indicates stronger hyperplastic remodelling pressure, often linked to chronic strain, inflammation, or compensatory skeletal adaptation.

Degree of Osteoporosis

Moderately elevated

Bone Mineral Density

Reference range: 0.124 - 0.453 | **Value:** 0.528

What this means: Reflects extent of bone mineral density reduction and structural fragility.

Why it matters: Osteoporosis increases fracture risk and reflects systemic ageing or imbalance.

Functional significance: Linked to hormonal decline, nutrient deficiency, inflammation, and inactivity.

Common patterns: Oestrogen decline + inflammation + calcium loss -> osteoporosis.

Your result in this scan: Indicates reduced bone density and increased fragility.

Bone Mineral Density

Within range

Bone Mineral Density

Reference range: 0.796 - 0.433 | **Value:** 0.435

What this means: Reflects the mineral density and load-bearing strength of bone tissue within the structural system.

Why it matters: Bone mineral density is central to fracture resistance, posture, skeletal reserve, and long-term integrity of the frame.

Functional significance: Depends on calcium-phosphorus balance, vitamin D and K status, parathyroid signalling, collagen matrix quality, and mechanical loading.

Common patterns: Low vitamin D, calcium imbalance, and poor collagen support can reduce bone density; inflammation and endocrine imbalance may worsen quality over time.

Your result in this scan: Bone Mineral Density appears within the analyser's expected range in this scan.

Rheumatoid Bone Disease

Medium priority

Complete system marker table — all markers and all severity grades are shown below.

Flagged markers: 3 | Within range: 2

Degree of Cervical Calcification

Mildly elevated

Rheumatoid Bone Disease

Reference range: 421 - 490 | **Value:** 502.068

What this means: Represents calcification deposits in cervical spine tissues, indicating chronic degeneration.

Why it matters: Calcification reduces flexibility and may compress nerves.

Functional significance: Linked to chronic inflammation, mineral imbalance, and poor tissue turnover.

Common patterns: Inflammation + calcium dysregulation -> calcification.

Your result in this scan: Indicates stiffness, degeneration, and restricted mobility.

Degree of Lumbar Calcification

Within range

Rheumatoid Bone Disease

Reference range: 4.326 - 7.531 | **Value:** 7.352

What this means: Reflects calcific degeneration within lumbar structures affecting mobility and nerve function.

Why it matters: Calcification contributes to stiffness and chronic back pain.

Functional significance: Linked to inflammation, mineral imbalance, and degenerative processes.

Common patterns: Chronic inflammation -> calcium deposition in tissues.

Your result in this scan: Degree of Lumbar Calcification appears within the analyser's expected range in this scan.

Bone Hyperplasia Coefficient

Mildly elevated

Rheumatoid Bone Disease

Reference range: 2.954 - 5.543 | **Value:** 6

What this means: Reflects a quantified tendency toward reactive bone proliferation and compensatory overgrowth within skeletal structures.

Why it matters: A raised bone hyperplasia tendency can signal chronic mechanical loading, degenerative change, inflammatory irritation, or altered remodelling chemistry.

Functional significance: Closely linked to mineral regulation, local tissue stress, endocrine influences on bone turnover, and connective support around joints.

Common patterns: Mechanical wear plus inflammatory signalling can drive hyperplastic bone responses; low connective support may worsen compensatory skeletal change.

Your result in this scan: Indicates heightened hyperplastic remodelling, often associated with chronic strain or inflammatory-degenerative bone adaptation.

Osteoporosis Coefficient

Within range

Rheumatoid Bone Disease

Reference range: 2.019 - 4.721 | **Value:** 4.705

What this means: Composite marker indicating overall bone fragility and density decline.

Why it matters: Provides integrated assessment of skeletal vulnerability.

Functional significance: Linked to endocrine status, mineral balance, and inflammation.

Common patterns: Hormonal imbalance + mineral loss -> bone fragility.

Your result in this scan: Osteoporosis Coefficient appears within the analyser's expected range in this scan.

Rheumatism Coefficient

Moderately elevated

Rheumatoid Bone Disease

Reference range: 4.023 - 11.627 | **Value:** 16.843

What this means: Reflects connective tissue and joint inflammatory burden associated with rheumatic-type patterns affecting movement and periarticular tissues.

Why it matters: Rheumatic patterns can impair mobility, increase pain sensitivity, and indicate wider immune-inflammatory strain on musculoskeletal connective tissues.

Functional significance: Interacts with immune activation, inflammatory cytokines, gut-immune cross-talk, mineral balance, and oxidative stress in joints and fascia.

Common patterns: Gut dysbiosis and immune activation can amplify rheumatic patterns; oxidative stress and low collagen support may worsen joint and fascia symptoms.

Your result in this scan: Indicates stronger joint and connective inflammation, immune reactivity, or chronic musculoskeletal inflammatory stress.

Bone Growth Index

Medium priority

Complete system marker table — all markers and all severity grades are shown below.

Flagged markers: 3 | Within range: 2

Bone alkaline phosphatase

Mildly reduced

Bone Growth Index

Reference range: 0.433 - 0.796 | **Value:** 0.399

What this means: Reflects osteoblastic activity and bone formation dynamics, indicating the degree of active skeletal matrix building and mineralisation support.

Why it matters: Bone alkaline phosphatase is a key marker of bone formation and helps show whether remodelling is actively supporting skeletal repair and maintenance.

Functional significance: Interacts with vitamin D status, calcium-phosphorus handling, osteoblast activity, endocrine regulation, and overall remodelling balance.

Common patterns: Low vitamin D or mineral imbalance may limit effective osteoblastic mineralisation; inflammatory or repair states can increase bone alkaline phosphatase demand.

Your result in this scan: Suggests reduced bone-building drive or slower osteoblastic activity within skeletal maintenance.

Osteocalcin

Mildly reduced

Bone Growth Index

Reference range: 0.525 - 0.817 | **Value:** 0.508

What this means: Reflects osteoblast-derived bone matrix activity and mineralisation signalling, with additional relevance to glucose and metabolic cross-talk.

Why it matters: Osteocalcin helps indicate active bone turnover and links skeletal metabolism with broader endocrine and metabolic regulation.

Functional significance: Interacts with vitamin K, vitamin D, calcium handling, bone formation, and metabolic-hormonal signalling.

Common patterns: Low vitamin K or vitamin D can impair osteocalcin function; altered bone turnover may also overlap with metabolic signalling disturbances.

Your result in this scan: Suggests weaker osteoblastic matrix activity or slower supportive bone turnover.

Status of long bone healing

Mildly reduced

Bone Growth Index

Reference range: 0.713 - 0.992 | **Value:** 0.606

What this means: Reflects the efficiency of repair and recovery processes within long bone tissue after structural stress or injury.

Why it matters: Long bone healing depends on coordinated inflammation, matrix formation, mineral deposition, and vascular supply, making it a useful indicator of structural repair capacity.

Functional significance: Linked to collagen matrix quality, calcium-phosphorus balance, vitamin C, vitamin D, osteoblastic function, and inflammatory resolution.

Common patterns: Low vitamin C, protein insufficiency, mineral imbalance, and inflammatory dysregulation can slow long bone healing and reduce repair quality.

Your result in this scan: Suggests slower repair kinetics, weak matrix support, reduced mineralisation, or impaired structural recovery capacity.

Short bone cartilage healing situation

Within range

Bone Growth Index

Reference range: 0.202 - 0.991 | **Value:** 0.362

What this means: Reflects healing and recovery quality within cartilage and short-bone associated supportive structures.

Why it matters: Cartilage healing is inherently slow and depends on strong connective tissue support, making it highly sensitive to chronic inflammation, mineral imbalance, and poor matrix turnover.

Functional significance: Interacts with collagen support, sulphur-containing compounds, vitamin C, inflammatory control, and local tissue biomechanics.

Common patterns: Chronic inflammation, low collagen support, and oxidative burden can weaken cartilage repair and worsen degenerative joint symptoms.

Your result in this scan: Short bone cartilage healing situation appears within the analyser's expected range in this scan.

Epiphyseal line

Within range

Bone Growth Index

Reference range: 0.432 - 0.826 | **Value:** 0.769

What this means: Reflects growth plate activity and skeletal maturation status.

Why it matters: Important for growth regulation and developmental assessment.

Functional significance: Linked to endocrine signalling (GH, sex hormones) and nutrition.

Common patterns: Hormonal imbalance -> abnormal growth patterns.

Your result in this scan: Epiphyseal line appears within the analyser's expected range in this scan.

Blood Sugar

Within range

Complete system marker table — all markers and all severity grades are shown below.

Flagged markers: 0 | Within range: 3

Coefficient of Insulin Secretion

Within range

Blood Sugar

Reference range: 2.967 - 3.528 | **Value:** 3.043

What this means: Reflects the responsiveness of pancreatic beta cells in secreting insulin relative to glucose exposure and metabolic demand.

Why it matters: Insulin secretion is central to glucose control, energy storage, and metabolic regulation; dysfunction contributes to insulin resistance and metabolic syndrome.

Functional significance: Linked to pancreatic beta-cell integrity, glucose load, incretin signalling, inflammatory state, and insulin sensitivity at tissue level.

Common patterns: High refined carbohydrate intake + inflammation + visceral adiposity -> insulin resistance -> compensatory insulin hypersecretion -> eventual beta-cell strain.

Your result in this scan: Coefficient of Insulin Secretion appears within the analyser's expected range in this scan.

Blood Sugar Coefficient

Within range

Blood Sugar

Reference range: 2.163 - 7.321 | **Value:** 3.191

What this means: Represents the overall balance of circulating glucose levels relative to metabolic demand and regulatory control.

Why it matters: Glucose balance underpins energy availability, cognitive function, and metabolic stability; dysregulation leads to fatigue, inflammation, and chronic disease risk.

Functional significance: Linked to insulin-glucagon balance, hepatic glucose output, dietary intake, mitochondrial utilisation, and stress hormones.

Common patterns: High glycaemic load + impaired insulin signalling -> elevated blood sugar coefficient -> oxidative stress and inflammatory burden.

Your result in this scan: Blood Sugar Coefficient appears within the analyser's expected range in this scan.

Urine Sugar Coefficient

Within range

Blood Sugar

Reference range: 2.204 - 2.819 | **Value:** 2.535

What this means: Reflects the presence of glucose excretion via urine, indicating overflow beyond renal reabsorption thresholds.

Why it matters: Urinary glucose reflects significant dysregulation of blood sugar control and can signal advanced metabolic disturbance.

Functional significance: Linked to renal glucose threshold, plasma glucose levels, insulin function, and kidney filtration capacity.

Common patterns: Chronic hyperglycaemia + insulin resistance -> renal threshold exceeded -> glycosuria.

Your result in this scan: Urine Sugar Coefficient appears within the analyser's expected range in this scan.

Trace Element

Medium priority

Complete system marker table — all markers and all severity grades are shown below.

Flagged markers: 3 | Within range: 16

Calcium

Within range

Trace Element

Reference range: 1.219 - 3.021 | **Value:** 1.518

What this means: Represents calcium status within the body, reflecting its availability for structural, neuromuscular, and signalling functions.

Why it matters: Calcium is essential for bone integrity, nerve conduction, muscle contraction, and intracellular signalling pathways.

Functional significance: Linked to parathyroid regulation, vitamin D status, bone metabolism, and cellular signalling mechanisms.

Common patterns: Vitamin D imbalance + parathyroid dysregulation -> altered calcium homeostasis affecting bone and cellular function.

Your result in this scan: Calcium appears within the analyser's expected range in this scan.

Iron

Mildly reduced

Trace Element

Reference range: 1.151 - 1.847 | **Value:** 0.743

What this means: Reflects systemic iron status and availability for oxygen transport, enzymatic activity, and mitochondrial function.

Why it matters: Iron is critical for haemoglobin, oxygen delivery, and cellular energy production; imbalance affects fatigue and oxidative stress.

Functional significance: Linked to haem synthesis, absorption, inflammation, liver storage, and red blood cell turnover.

Common patterns: Inflammation + poor absorption or excess intake -> altered iron balance impacting oxygen transport and redox state.

Your result in this scan: Suggests reduced iron availability, potentially leading to impaired oxygen delivery and fatigue.

Zinc

Within range

Trace Element

Reference range: 1.143 - 1.989 | **Value:** 1.336

What this means: Represents zinc status, a critical cofactor in enzymatic activity, immune regulation, and cellular repair.

Why it matters: Zinc supports immune function, DNA synthesis, hormone regulation, and antioxidant defence.

Functional significance: Linked to digestive absorption, immune signalling, wound healing, and hormonal balance.

Common patterns: Poor diet + malabsorption -> zinc deficiency -> impaired immunity and repair.

Your result in this scan: Zinc appears within the analyser's expected range in this scan.

Selenium

Within range

Trace Element

Reference range: 0.847 - 2.045 | **Value:** 1.096

What this means: Reflects selenium availability for antioxidant systems and thyroid hormone metabolism.

Why it matters: Selenium is essential for glutathione peroxidase activity and thyroid hormone conversion.

Functional significance: Linked to thyroid function, antioxidant defence, and immune modulation.

Common patterns: Low intake + oxidative stress -> reduced selenium function -> impaired detox and thyroid activity.

Your result in this scan: Selenium appears within the analyser's expected range in this scan.

Phosphorus

Within range

Trace Element

Reference range: 1.195 - 2.134 | **Value:** 2.087

What this means: Represents phosphorus status, essential for energy metabolism and cellular structure.

Why it matters: Phosphorus is a key component of ATP, DNA, and cellular membranes.

Functional significance: Linked to kidney function, bone metabolism, and energy production.

Common patterns: Kidney dysfunction + dietary imbalance -> altered phosphorus levels.

Your result in this scan: Phosphorus appears within the analyser's expected range in this scan.

Potassium

Within range

Trace Element

Reference range: 0.689 - 0.987 | **Value:** 0.977

What this means: Reflects potassium balance influencing cellular electrical activity and fluid balance.

Why it matters: Potassium is critical for nerve conduction, muscle contraction, and cardiovascular stability.

Functional significance: Linked to kidney function, hydration, and electrolyte regulation.

Common patterns: Electrolyte imbalance + renal dysfunction -> potassium disturbance.

Your result in this scan: Potassium appears within the analyser's expected range in this scan.

Magnesium

Within range

Trace Element

Reference range: 0.568 - 0.992 | **Value:** 0.761

What this means: Represents magnesium status, essential for enzymatic activity and neuromuscular stability.

Why it matters: Magnesium supports ATP production, muscle relaxation, and nervous system balance.

Functional significance: Linked to stress, mitochondrial function, and electrolyte balance.

Common patterns: Stress + poor intake -> magnesium depletion.

Your result in this scan: Magnesium appears within the analyser's expected range in this scan.

Copper

Moderately reduced

Trace Element

Reference range: 0.474 - 0.749 | **Value:** 0.23

What this means: Reflects copper balance involved in redox activity and connective tissue formation.

Why it matters: Copper is essential for iron metabolism and antioxidant enzymes.

Functional significance: Linked to zinc balance and liver function.

Common patterns: Zinc imbalance -> copper dysregulation.

Your result in this scan: Suggests deficiency affecting iron metabolism.

Cobalt

Within range

Trace Element

Reference range: 2.326 - 5.531 | **Value:** 2.493

What this means: Represents cobalt status, linked to vitamin B12 structure and function.

Why it matters: Cobalt is essential for red blood cell production and neurological function.

Functional significance: Linked to B12 metabolism and gut absorption.

Common patterns: Poor absorption -> B12 deficiency.

Your result in this scan: Cobalt appears within the analyser's expected range in this scan.

Manganese

Within range

Trace Element

Reference range: 0.497 - 0.879 | **Value:** 0.732

What this means: Reflects manganese status for enzymatic and antioxidant functions.

Why it matters: Manganese supports bone formation and metabolic enzymes.

Functional significance: Linked to mitochondrial and antioxidant activity.

Common patterns: Environmental exposure -> imbalance.

Your result in this scan: Manganese appears within the analyser's expected range in this scan.

Iodine

Within range

Trace Element

Reference range: 1.421 - 5.490 | **Value:** 5.311

What this means: Represents iodine availability for thyroid hormone production.

Why it matters: Iodine is essential for T3 and T4 synthesis.

Functional significance: Linked to thyroid function and metabolism.

Common patterns: Low intake -> hypothyroid patterns.

Your result in this scan: Iodine appears within the analyser's expected range in this scan.

Nickel

Within range

Trace Element

Reference range: 2.462 - 5.753 | **Value:** 2.481

What this means: Reflects nickel exposure and trace biological role.

Why it matters: Nickel may influence enzyme activity at trace levels.

Functional significance: Linked to environmental exposure.

Common patterns: Environmental sources -> imbalance.

Your result in this scan: Nickel appears within the analyser's expected range in this scan.

Fluorine

Within range

Trace Element

Reference range: 1.954 - 4.543 | **Value:** 2.019

What this means: Represents fluorine exposure influencing bone and dental health.

Why it matters: Fluorine contributes to enamel strength but excess may affect tissues.

Functional significance: Linked to environmental intake.

Common patterns: Water intake -> fluorine levels.

Your result in this scan: Fluorine appears within the analyser's expected range in this scan.

Molybdenum

Within range

Trace Element

Reference range: 0.938 - 1.712 | **Value:** 1.43

What this means: Reflects molybdenum status in enzymatic detox pathways.

Why it matters: Molybdenum supports sulphite metabolism and detoxification.

Functional significance: Linked to liver enzymes.

Common patterns: Low intake -> impaired detox.

Your result in this scan: Molybdenum appears within the analyser's expected range in this scan.

Vanadium

Within range

Trace Element

Reference range: 1.019 - 3.721 | **Value:** 2.638

What this means: Represents vanadium involvement in glucose and lipid metabolism.

Why it matters: Vanadium may influence insulin signalling.

Functional significance: Linked to metabolic regulation.

Common patterns: Metabolic dysfunction -> altered signalling.

Your result in this scan: Vanadium appears within the analyser's expected range in this scan.

Tin

Within range

Trace Element

Reference range: 1.023 - 7.627 | **Value:** 7.159

What this means: Reflects exposure to tin and trace biological impact.

Why it matters: Primarily reflects environmental exposure.

Functional significance: Linked to toxicity.

Common patterns: Environmental contamination.

Your result in this scan: Tin appears within the analyser's expected range in this scan.

Silicon

Mildly reduced

Trace Element

Reference range: 1.425 - 5.872 | **Value:** 1.367

What this means: Represents silicon role in connective tissue and bone health.

Why it matters: Supports collagen and structural integrity.

Functional significance: Linked to bone metabolism.

Common patterns: Nutritional imbalance.

Your result in this scan: Suggests deficiency.

Strontium

Within range

Trace Element

Reference range: 1.142 - 5.862 | **Value:** 3.508

What this means: Represents strontium status related to bone mineralisation and structural integrity.

Why it matters: Strontium supports bone formation and may influence osteoblastic activity and mineral balance.

Functional significance: Linked to calcium metabolism, bone turnover, and skeletal health.

Common patterns: Calcium imbalance + dietary variation -> altered strontium incorporation affecting bone dynamics.

Your result in this scan: Strontium appears within the analyser's expected range in this scan.

Boron

Within range

Trace Element

Reference range: 1.124 - 3.453 | **Value:** 2.537

What this means: Reflects boron-related support for bone-mineral metabolism, steroid hormone balance, membrane signalling, and inflammatory regulation.

Why it matters: Boron may influence calcium and magnesium utilisation, vitamin D dynamics, bone support, and aspects of sex hormone metabolism and inflammatory tone.

Functional significance: Interacts with mineral retention, endocrine balance, bone matrix health, and possibly cognition and inflammatory regulation through membrane-signalling effects.

Common patterns: Low boron may align with weaker bone support, less efficient magnesium-calcium handling, and reduced hormone-mineral synergy across endocrine and inflammatory pathways.

Your result in this scan: Boron appears within the analyser's expected range in this scan.

Vitamin

High priority

Complete system marker table — all markers and all severity grades are shown below.

Flagged markers: 6 | Within range: 4

Vitamin A

Mildly reduced

Vitamin

Reference range: 0.346 - 0.401 | **Value:** 0.338

What this means: Reflects retinoid status relevant to epithelial integrity, immune signalling, vision pathways, and gene expression regulation.

Why it matters: Vitamin A is essential for mucosal barrier integrity, immune tolerance, retinal function, and regulation of cellular differentiation.

Functional significance: Interacts with gut integrity, immune response modulation, liver storage, thyroid signalling, and epithelial repair processes.

Common patterns: Low vitamin A weakens gut and respiratory barriers, increasing antigen load and immune activation; also affects skin and hormonal signalling.

Your result in this scan: Suggests impaired mucosal barrier, weakened immune tolerance, poor epithelial repair, or reduced visual adaptation.

Vitamin B1

Mildly reduced

Vitamin

Reference range: 2.124 - 4.192 | **Value:** 1.679

What this means: Reflects thiamine availability required for carbohydrate metabolism, pyruvate dehydrogenase activity, nerve conduction, and mitochondrial ATP production.

Why it matters: Vitamin B1 is essential for converting glucose into usable energy and for maintaining neurological and cardiovascular function under metabolic stress.

Functional significance: Linked to glucose handling, mitochondrial entry of glycolytic products, autonomic function, digestive motility, and high carbohydrate demand states.

Common patterns: Low B1 → impaired pyruvate entry into mitochondria → reduced ATP production and greater lactate-type fatigue; nervous system and digestive motility may also be affected.

Your result in this scan: Suggests impaired carbohydrate-to-energy conversion, neurological irritability, fatigue, poor autonomic tone, or deficiency related to intake, stress, or higher metabolic demand.

Vitamin B2

Within range

Vitamin

Reference range: 1.549 - 2.213 | **Value:** 1.549

What this means: Reflects riboflavin availability required for FAD/FMN-dependent redox reactions in mitochondrial respiration, detoxification, and fatty acid metabolism.

Why it matters: Vitamin B2 supports ATP production, antioxidant recycling, methylation support, and effective metabolism of fats and other B vitamins.

Functional significance: Interacts with mitochondrial electron transport, glutathione recycling, liver detox pathways, and activation of other nutrient cofactors.

Common patterns: Low B2 → weaker electron transport + reduced glutathione recycling → fatigue and oxidative vulnerability; detoxification and lipid metabolism may also become less efficient.

Your result in this scan: Vitamin B2 appears within the analyser's expected range in this scan.

Vitamin B3

Within range

Vitamin

Reference range: 14.477 - 21.348 | **Value:** 14.72

What this means: Reflects niacin-related status relevant to NAD and NADP-dependent redox chemistry, mitochondrial energy transfer, DNA repair, and lipid metabolism.

Why it matters: Vitamin B3 is foundational to cellular respiration, detoxification redox balance, skin integrity, nervous system function, and healthy handling of fats and carbohydrates.

Functional significance: Closely linked to mitochondrial electron transfer, liver detox demand, glucose-fat metabolism, stress resilience, and repair processes requiring redox capacity.

Common patterns: Low B3 can reduce NAD-dependent energy production and weaken detox redox capacity; this may worsen fatigue, skin sensitivity, mood instability, and poor metabolic resilience.

Your result in this scan: Vitamin B3 appears within the analyser's expected range in this scan.

Vitamin B6

Within range

Vitamin

Reference range: 0.824 - 1.942 | **Value:** 1.616

What this means: Reflects pyridoxine-dependent cofactor status required for amino acid metabolism, neurotransmitter synthesis, glycogen phosphorylase activity, and homocysteine regulation.

Why it matters: Vitamin B6 is central to neurotransmitter production, protein handling, immune modulation, glycaemic stability, and efficient methylation-transsulphuration balance.

Functional significance: Interfaces with neurotransmitter synthesis, glycogen mobilisation, histamine regulation, homocysteine metabolism, and immune-inflammatory control.

Common patterns: Low B6 → weaker serotonin/GABA synthesis + poorer glycogen handling + altered histamine regulation; this may show as mood instability, PMS-type patterns, allergy tendency, and fatigue.

Your result in this scan: Vitamin B6 appears within the analyser's expected range in this scan.

Vitamin B12

Within range

Vitamin

Reference range: 6.428 - 21.396 | **Value:** 6.962

What this means: Reflects cobalamin-related status required for methylation, red blood cell production, myelin integrity, homocysteine regulation, and mitochondrial support.

Why it matters: Vitamin B12 is essential for neurological function, cognitive clarity, methylation chemistry, healthy haematopoiesis, and sustained energy output.

Functional significance: Interacts with intrinsic factor and stomach health, folate metabolism, cobalt-related status, methyl donation, and nerve membrane resilience.

Common patterns: Low stomach acid or intrinsic factor dysfunction can reduce B12 handling; low B12 may drive fatigue, memory issues, neuropathy, elevated homocysteine, and reduced methylation resilience.

Your result in this scan: Vitamin B12 appears within the analyser's expected range in this scan.

Vitamin C

Moderately reduced

Vitamin

Reference range: 4.543 - 5.023 | **Value:** 3.55

What this means: Reflects ascorbate status required for collagen synthesis, antioxidant defence, immune activity, and adrenal support.

Why it matters: Vitamin C is essential for connective tissue repair, immune defence, adrenal resilience, and protection against oxidative stress.

Functional significance: Closely linked to adrenal cortisol dynamics, collagen formation, iron absorption, and redox balance.

Common patterns: Low vitamin C impairs collagen, increases oxidative stress, and weakens adrenal response to stress and immune demand.

Your result in this scan: Suggests reduced antioxidant protection, weaker collagen repair, adrenal strain, or increased susceptibility to infection.

Vitamin D3

Mildly reduced

Vitamin

Reference range: 5.327 - 7.109 | **Value:** 5.259

What this means: Reflects vitamin D status relevant to calcium regulation, immune modulation, gene expression, and endocrine signalling.

Why it matters: Vitamin D is essential for bone health, immune balance, hormone regulation, and anti-inflammatory signalling.

Functional significance: Interacts with calcium-magnesium balance, parathyroid hormone, immune tolerance, and endocrine stability.

Common patterns: Low vitamin D contributes to immune dysregulation, bone weakness, and inflammatory patterns.

Your result in this scan: Suggests impaired immune modulation, weaker bone-mineral balance, or increased inflammatory susceptibility.

Vitamin E

Moderately reduced

Vitamin

Reference range: 4.826 - 6.013 | **Value:** 4.11

What this means: Reflects lipid-soluble antioxidant status protecting membranes from oxidative damage.

Why it matters: Vitamin E protects cell membranes, supports cardiovascular health, and reduces oxidative stress in lipid environments.

Functional significance: Interacts with lipid metabolism, oxidative stress, mitochondrial function, and inflammatory signalling.

Common patterns: Low vitamin E increases lipid oxidation and vascular stress.

Your result in this scan: Suggests increased oxidative damage to lipids and membranes.

Vitamin K

Mildly reduced

Vitamin

Reference range: 0.717 - 1.486 | **Value:** 0.599

What this means: Reflects vitamin K status involved in blood clotting, calcium regulation, and bone metabolism.

Why it matters: Vitamin K is required for proper calcium placement, preventing vascular calcification and supporting bone health.

Functional significance: Interacts with vitamin D, calcium metabolism, liver function, and coagulation pathways.

Common patterns: Vitamin K imbalance affects bone and vascular calcium deposition.

Your result in this scan: Suggests impaired clotting or poor calcium distribution.

Amino Acid

Medium priority

Complete system marker table — all markers and all severity grades are shown below.

Flagged markers: 2 | Within range: 9

Lysine

Mildly reduced

Amino Acid

Reference range: 0.962 - 1.213 | **Value:** 0.853

What this means: Reflects lysine availability, an essential amino acid critical for protein synthesis, collagen formation, immune modulation, and calcium metabolism.

Why it matters: Lysine supports tissue repair, immune defence, and connective tissue integrity, and is particularly relevant in viral susceptibility and wound healing.

Functional significance: Linked to collagen cross-linking, antibody production, carnitine synthesis, and overall protein turnover.

Common patterns: Low lysine -> reduced collagen integrity + impaired immune response; may co-occur with gut dysfunction or malabsorption.

Your result in this scan: Suggests reduced protein quality or intake, impaired absorption, or increased demand for tissue repair and immune activity.

Tryptophan

Within range

Amino Acid

Reference range: 4.978 - 6.289 | **Value:** 6.008

What this means: Reflects tryptophan availability, a precursor for serotonin, melatonin, and kynurenine pathway metabolites.

Why it matters: Tryptophan is essential for mood regulation, sleep quality, and immune modulation.

Functional significance: Linked to neurotransmitter synthesis, circadian rhythm regulation, and immune-inflammatory balance.

Common patterns: Inflammation shifts tryptophan -> kynurenine pathway, reducing serotonin; stress and gut dysfunction can worsen depletion.

Your result in this scan: Tryptophan appears within the analyser's expected range in this scan.

Phenylalanine

Within range

Amino Acid

Reference range: 1.928 - 2.491 | **Value:** 2.057

What this means: Reflects phenylalanine availability, a precursor to tyrosine and downstream catecholamines.

Why it matters: Phenylalanine supports dopamine, noradrenaline, and adrenaline production, influencing focus, motivation, and stress response.

Functional significance: Linked to neurotransmitter synthesis, cognitive performance, and stress adaptation.

Common patterns: Low phenylalanine -> low dopamine -> fatigue and poor motivation; stress may increase demand.

Your result in this scan: Phenylalanine appears within the analyser's expected range in this scan.

Methionine

Within range

Amino Acid

Reference range: 1.245 - 1.637 | **Value:** 1.398

What this means: Reflects methionine availability, a key methyl donor involved in detoxification and epigenetic regulation.

Why it matters: Methionine is central to methylation, detoxification, and liver function.

Functional significance: Linked to SAMe production, homocysteine cycling, glutathione synthesis, and detox pathways.

Common patterns: Methionine imbalance -> homocysteine dysregulation -> cardiovascular and detox stress.

Your result in this scan: Methionine appears within the analyser's expected range in this scan.

Threonine

Within range

Amino Acid

Reference range: 1.194 - 1.685 | **Value:** 1.37

What this means: Reflects threonine status, important for protein synthesis and mucosal integrity.

Why it matters: Threonine supports gut lining integrity and immune defence.

Functional significance: Linked to mucin production, gut barrier function, and immune support.

Common patterns: Low threonine -> weakened gut lining -> increased permeability and immune activation.

Your result in this scan: Threonine appears within the analyser's expected range in this scan.

Isoleucine

Mildly reduced

Amino Acid

Reference range: 4.582 - 5.657 | **Value:** 4.175

What this means: Reflects isoleucine status, a branched-chain amino acid (BCAA) involved in energy metabolism and muscle repair.

Why it matters: Isoleucine supports glucose uptake and muscle recovery.

Functional significance: Linked to energy production, muscle metabolism, and blood sugar regulation.

Common patterns: Low BCAAs -> fatigue and muscle weakness; metabolic stress increases demand.

Your result in this scan: Suggests reduced muscle repair or energy support.

Leucine

Within range

Amino Acid

Reference range: 6.982 - 9.256 | **Value:** 7.041

What this means: Reflects leucine status, a key BCAA regulating muscle protein synthesis via mTOR signalling.

Why it matters: Leucine is critical for muscle maintenance, repair, and metabolic signalling.

Functional significance: Linked to anabolic signalling, muscle metabolism, and insulin sensitivity.

Common patterns: Low leucine -> reduced muscle maintenance; ageing and stress increase need.

Your result in this scan: Leucine appears within the analyser's expected range in this scan.

Valine

Within range

Amino Acid

Reference range: 6.982 - 9.677 | **Value:** 7.565

What this means: Reflects valine status, a BCAA involved in energy provision and muscle metabolism.

Why it matters: Valine supports endurance, muscle repair, and metabolic stability.

Functional significance: Linked to nitrogen balance, muscle metabolism, and CNS function.

Common patterns: Low valine -> fatigue and reduced endurance.

Your result in this scan: Valine appears within the analyser's expected range in this scan.

Histidine

Within range

Amino Acid

Reference range: 5.113 - 6.258 | **Value:** 5.261

What this means: Reflects histidine status, involved in haemoglobin structure and histamine production.

Why it matters: Histidine supports oxygen transport and immune responses.

Functional significance: Linked to haemoglobin, histamine signalling, and antioxidant systems.

Common patterns: Histidine imbalance -> histamine issues or oxygen transport inefficiency.

Your result in this scan: Histidine appears within the analyser's expected range in this scan.

Arginine

Within range

Amino Acid

Reference range: 1.812 - 2.337 | **Value:** 1.975

What this means: Reflects arginine status, a semi-essential amino acid involved in nitric oxide production and vascular function.

Why it matters: Arginine supports circulation, immune response, and wound healing.

Functional significance: Linked to nitric oxide synthesis, vascular dilation, and immune activity.

Common patterns: Low arginine -> poor circulation and healing.

Your result in this scan: Arginine appears within the analyser's expected range in this scan.

Homocysteine

Within range

Amino Acid

Reference range: 0.983 - 1.265 | **Value:** 1.073

What this means: Reflects homocysteine levels, a key intermediate in methylation and cardiovascular risk pathways.

Why it matters: Elevated homocysteine is associated with cardiovascular risk, impaired methylation, and oxidative stress.

Functional significance: Linked to B-vitamin status, methylation cycles, detoxification, and vascular health.

Common patterns: High homocysteine -> cardiovascular risk + oxidative stress + poor detox.

Your result in this scan: Homocysteine appears within the analyser's expected range in this scan.

Coenzyme

Medium priority

Complete system marker table — all markers and all severity grades are shown below.

Flagged markers: 3 | Within range: 3

Nicotinamide

Within range

Coenzyme

Reference range: 2.074 - 3.309 | **Value:** 2.293

What this means: Reflects NAD precursor status supporting redox reactions and energy metabolism.

Why it matters: Nicotinamide supports mitochondrial function, DNA repair, and metabolic resilience.

Functional significance: Closely linked to NAD cycles, detoxification, and cellular repair.

Common patterns: Low NAD precursors reduce cellular resilience and increase fatigue.

Your result in this scan: Nicotinamide appears within the analyser's expected range in this scan.

Biotin

Mildly reduced

Coenzyme

Reference range: 1.833 - 2.979 | **Value:** 1.241

What this means: Reflects biotin status required for fatty acid metabolism and glucose regulation.

Why it matters: Biotin supports hair, skin, and metabolic enzyme function.

Functional significance: Interacts with carbohydrate metabolism and mitochondrial enzymes.

Common patterns: Low biotin affects hair, skin, and glucose metabolism.

Your result in this scan: Suggests impaired fat metabolism and skin issues.

Pantothenic acid

Moderately reduced

Coenzyme

Reference range: 1.116 - 2.101 | **Value:** 0.604

What this means: Reflects vitamin B5 status required for CoA synthesis and energy metabolism.

Why it matters: Essential for fatty acid metabolism, adrenal function, and energy production.

Functional significance: Interacts with adrenal hormones and mitochondrial pathways.

Common patterns: Low B5 reduces CoA availability and energy production.

Your result in this scan: Suggests fatigue and adrenal strain.

Folic acid

Within range

Coenzyme

Reference range: 1.449 - 2.246 | **Value:** 2.236

What this means: Reflects folate status required for methylation and DNA synthesis.

Why it matters: Essential for cell division, red blood cell formation, and methylation balance.

Functional significance: Interacts with B12, homocysteine, and DNA repair.

Common patterns: Low folate increases homocysteine and reduces repair capacity.

Your result in this scan: Folic acid appears within the analyser's expected range in this scan.

Coenzyme Q10

Mildly reduced

Coenzyme

Reference range: 0.831 - 1.588 | **Value:** 0.643

What this means: Reflects availability of CoQ10, a key electron transport cofactor in the mitochondrial inner membrane required for oxidative phosphorylation and cellular antioxidant protection.

Why it matters: CoQ10 is essential for ATP generation in high-demand tissues such as heart, brain, muscle, and liver, and it also helps protect membranes from oxidative damage.

Functional significance: Closely linked to mitochondrial respiratory chain efficiency, redox balance, statin-related depletion, cardiac energy demand, and overall fatigue resilience.

Common patterns: Low CoQ10 → poorer mitochondrial ATP production → fatigue, low exercise tolerance, and weaker cardiac-muscular energy support; oxidative stress can further deplete reserves.

Your result in this scan: Suggests impaired mitochondrial electron transport, lower ATP output, greater oxidative vulnerability, or depletion associated with age, medication use, or chronic illness burden.

Glutathione

Within range

Coenzyme

Reference range: 0.726 - 1.281 | **Value:** 1.114

What this means: Reflects glutathione status as a master antioxidant and detox molecule.

Why it matters: Central to detoxification, oxidative stress control, and immune resilience.

Functional significance: Interacts with liver detox, selenium, and redox balance.

Common patterns: Low glutathione increases toxin sensitivity and inflammation.

Your result in this scan: Glutathione appears within the analyser's expected range in this scan.

Essential Fatty Acid

Medium priority

Complete system marker table — all markers and all severity grades are shown below.

Flagged markers: 2 | Within range: 2

Linoleic acid

Mildly reduced

Essential Fatty Acid

Reference range: 0.642 - 0.985 | **Value:** 0.447

What this means: Reflects linoleic-acid status as a key omega-6 essential fatty acid involved in membrane structure, epidermal barrier integrity, and eicosanoid precursor balance.

Why it matters: Linoleic acid is required for healthy skin-barrier function, cell-membrane composition, and downstream inflammatory signalling. Deficiency can impair barrier resilience, while excess relative to omega-3 pathways can promote inflammatory imbalance.

Functional significance: Linked to dietary fat quality, bile-dependent absorption, membrane fluidity, epidermal ceramide formation, and the broader omega-6 to omega-3 balance shaping inflammatory tone.

Common patterns: Low bile flow or fat malabsorption -> lower linoleic-acid availability and weaker skin-barrier resilience; omega-6 excess without omega-3 balance can intensify inflammatory and metabolic stress.

Your result in this scan: Suggests reduced essential fatty acid availability, impaired fat absorption, weakened skin-barrier integrity, or lower membrane-lipid support.

α-Linolenic acid

Within range

Essential Fatty Acid

Reference range: 0.814 - 1.202 | **Value:** 0.852

What this means: Reflects alpha-linolenic-acid availability as a plant-derived omega-3 essential fatty acid supporting membrane fluidity, inflammatory modulation, and precursor input into longer-chain omega-3 pathways.

Why it matters: Alpha-linolenic acid contributes to anti-inflammatory balance, cardiovascular resilience, and membrane health, especially when broader omega-3 status is suboptimal.

Functional significance: Linked to dietary fat quality, desaturase and elongase activity, bile-dependent absorption, oxidative protection, and the overall omega-6 to omega-3 ratio.

Common patterns: Low alpha-linolenic-acid intake or poor absorption -> weaker anti-inflammatory balance; excess omega-6 exposure can further blunt the functional impact of available omega-3 precursors.

Your result in this scan: α-Linolenic acid appears within the analyser's expected range in this scan.

γ-Linolenic acid

Within range

Essential Fatty Acid

Reference range: 0.921 - 1.334 | **Value:** 0.996

What this means: Reflects gamma-linolenic-acid related functional support within omega-6 metabolism, relevant to membrane signalling, skin integrity, and downstream eicosanoid balance.

Why it matters: Gamma-linolenic acid can support anti-inflammatory and barrier-regulating pathways within omega-6 metabolism, particularly where delta-6-desaturase conversion is impaired.

Functional significance: Linked to fatty-acid conversion efficiency, skin-barrier resilience, hormone-related inflammatory balance, and the body's ability to generate more regulatory lipid mediators.

Common patterns: Low conversion efficiency from linoleic acid can reduce gamma-linolenic-acid support and contribute to skin dryness, PMS-type inflammatory patterns, and barrier fragility.

Your result in this scan: γ-Linolenic acid appears within the analyser's expected range in this scan.

Arachidonic acid

Mildly reduced

Essential Fatty Acid

Reference range: 0.661 - 0.808 | **Value:** 0.575

What this means: Reflects arachidonic-acid status as a membrane-bound omega-6 fatty acid involved in cell signalling and the generation of multiple inflammatory and regulatory eicosanoids.

Why it matters: Arachidonic acid is essential to membrane function and signalling, but excessive dominance can amplify inflammatory cascades, pain pathways, and vascular reactivity.

Functional significance: Linked to membrane phospholipids, inflammatory mediator production, immune activation, pain signalling, and the balance between pro-inflammatory and resolving lipid pathways.

Common patterns: Omega-3 deficiency plus high arachidonic-acid dominance -> stronger inflammatory signalling, pain amplification, and vascular reactivity; oxidative stress can worsen downstream tissue effects.

Your result in this scan: May suggest reduced membrane availability for eicosanoid signalling, lower dietary intake, altered absorption, or context-dependent reduced inflammatory drive.

Endocrine System

Medium priority

Complete system marker table — all markers and all severity grades are shown below.

Flagged markers: 2 | Within range: 5

Thyroid secretion index

Within range

Endocrine System

Reference range: 2.954 - 5.543 | **Value:** 2.996

What this means: Reflects thyroid gland secretory output, including synthesis and release of thyroxine (T4) and triiodothyronine (T3) under TSH regulation, alongside iodine handling and glandular responsiveness.

Why it matters: Thyroid output is a primary determinant of basal metabolic rate, thermogenesis, mitochondrial respiration, cardiac drive, gastrointestinal motility, and tissue-level energy availability.

Functional significance: Interacts with hepatic T4-to-T3 conversion, selenium-dependent deiodinase activity, adrenal cortisol balance, iodine sufficiency, and cellular receptor responsiveness.

Common patterns: Chronic stress → elevated cortisol → impaired T4-to-T3 conversion and reduced thyroid signalling; liver dysfunction → weaker peripheral activation; gut dysfunction → impaired iodine, selenium, and tyrosine handling.

Your result in this scan: Thyroid secretion index appears within the analyser's expected range in this scan.

Parathyroid hormone secretion index

Mildly reduced

Endocrine System

Reference range: 2.845 - 4.017 | **Value:** 2.121

What this means: Reflects parathyroid hormone regulatory activity involved in calcium-phosphorus homeostasis, vitamin D coordination, and bone-mineral signalling.

Why it matters: Parathyroid regulation is essential for neuromuscular excitability, skeletal integrity, intracellular signalling, and the maintenance of stable calcium availability.

Functional significance: Closely linked to vitamin D status, magnesium sufficiency, renal handling of minerals, bone turnover, and acid-base buffering patterns.

Common patterns: Low vitamin D or magnesium imbalance → altered PTH signalling → disturbed calcium handling → muscle tension, bone turnover changes, and neuromuscular irritability.

Your result in this scan: Suggests reduced parathyroid signalling, altered calcium regulation, magnesium-related suppression, or reduced adaptive response to mineral demand.

Adrenal glands Index

Mildly reduced

Endocrine System

Reference range: 2.412 - 2.974 | **Value:** 2.149

What this means: Reflects adrenal gland output and reserve, including cortisol, catecholamine, and DHEA-related stress adaptation under hypothalamic-pituitary-adrenal axis control.

Why it matters: Adrenal regulation is central to circadian rhythm, blood glucose stability, inflammatory tone, cardiovascular response, and tolerance to physical or psychological stress.

Functional significance: Interfaces with thyroid signalling, autonomic tone, insulin sensitivity, immune regulation, sleep architecture, and mitochondrial demand.

Common patterns: Chronic stress → cortisol dysregulation → sleep disruption + insulin resistance + reduced thyroid conversion; prolonged overactivation may transition into lower reserve and poor resilience.

Your result in this scan: Suggests reduced adrenal output, diminished stress resilience, flattened adaptive response, or depleted reserve after prolonged stress burden.

Pituitary secretion index

Within range

Endocrine System

Reference range: 2.163 - 7.34 | **Value:** 5.512

What this means: Reflects pituitary secretory signalling across trophic hormones such as TSH, ACTH, LH, FSH, prolactin, and growth-hormone-related regulation.

Why it matters: The pituitary acts as a central endocrine relay, translating hypothalamic signals into coordinated thyroid, adrenal, reproductive, and growth-related responses.

Functional significance: Depends on hypothalamic input, circadian timing, feedback-loop integrity, nutrient sufficiency, and low neuroinflammatory disruption of neuroendocrine signalling.

Common patterns: Hypothalamic stress or circadian disruption → pituitary dysregulation → downstream thyroid/adrenal/reproductive imbalance; poor feedback integrity can amplify multi-axis endocrine instability.

Your result in this scan: Pituitary secretion index appears within the analyser's expected range in this scan.

Pineal secretion index

Within range

Endocrine System

Reference range: 3.210 - 6.854 | **Value:** 6.778

What this means: Reflects pineal gland activity and melatonin-related circadian signalling.

Why it matters: Pineal function regulates sleep-wake cycles, hormonal timing, and recovery processes.

Functional significance: Linked to light exposure, circadian rhythm, and neuroendocrine balance.

Common patterns: Disrupted light exposure -> circadian misalignment -> hormonal imbalance.

Your result in this scan: Pineal secretion index appears within the analyser's expected range in this scan.

Thymus gland secretion index

Within range

Endocrine System

Reference range: 2.967 - 3.528 | **Value:** 2.969

What this means: Reflects thymic regulatory activity associated with immune maturation, T-cell education, and endocrine-immune interaction within the neuroendocrine network.

Why it matters: Thymic function is relevant to immune tolerance, adaptive immune competence, inflammatory regulation, and resilience against chronic stress-related immune dysregulation.

Functional significance: Interacts with adrenal cortisol tone, immune signalling, micronutrient status, age-related involution patterns, and systemic inflammatory burden.

Common patterns: Chronic cortisol excess → thymic suppression → weaker immune tolerance and adaptive response; immune activation and inflammatory load can further destabilise endocrine-immune balance.

Your result in this scan: Thymus gland secretion index appears within the analyser's expected range in this scan.

Gland secretion index

Within range

Endocrine System

Reference range: 2.204 - 2.819 | **Value:** 2.492

What this means: Reflects generalised endocrine gland secretory performance across integrated glandular networks, providing a broad view of overall endocrine signalling tone.

Why it matters: Global glandular output influences metabolic drive, reproductive rhythm, stress adaptation, tissue repair, sleep physiology, and inflammatory balance.

Functional significance: Represents integrated output from thyroid, adrenal, pituitary, pineal, and related endocrine structures, and therefore should be interpreted alongside more specific glandular markers.

Common patterns: Chronic stress, circadian disruption, poor nutrient status, and inflammatory burden can suppress coordinated endocrine output; conversely, compensatory activation may appear during stress, metabolic imbalance, or hormonal dysregulation.

Your result in this scan: Gland secretion index appears within the analyser's expected range in this scan.

Immune System

Within range

Complete system marker table — all markers and all severity grades are shown below.

Flagged markers: 0 | Within range: 9

Lymph node Index

Within range

Immune System

Reference range: 133.437 - 140.47 | **Value:** 134.342

What this means: Reflects lymph node functional activity within immune surveillance, antigen presentation, lymph filtration, and adaptive immune coordination.

Why it matters: Lymph nodes are central processing sites for immune recognition, lymphatic drainage, and orchestration of immune responses to pathogens, toxins, and tissue debris.

Functional significance: Integrates lymphatic flow, antigen burden, mucosal immune traffic, inflammatory signalling, and the balance between immune activation and immune tolerance.

Common patterns: Poor lymphatic drainage → antigen and waste retention → greater inflammatory load; chronic gut or respiratory antigen exposure can increase nodal activation and broader immune stress patterns.

Your result in this scan: Lymph node Index appears within the analyser's expected range in this scan.

Tonsil immune Index

Within range

Immune System

Reference range: 0.124 - 0.453 | **Value:** 0.307

What this means: Reflects tonsillar immune activity as part of upper airway mucosal defence, immune sampling, and early response to inhaled or ingested antigens.

Why it matters: Tonsillar function contributes to first-line immune defence, oral-respiratory immune tolerance, and early recognition of environmental and microbial challenges.

Functional significance: Closely linked to respiratory mucosal integrity, microbiome exposure, allergy burden, sinus-throat inflammation, and adaptive immune priming.

Common patterns: Respiratory antigen load or recurrent infection → persistent tonsillar activation → wider mucosal immune stress; chronic upper airway inflammation can feed into allergy patterns and systemic immune load.

Your result in this scan: Tonsil immune Index appears within the analyser's expected range in this scan.

Bone marrow Index

Within range

Immune System

Reference range: 0.146 - 3.218 | **Value:** 1.068

What this means: Reflects bone marrow contribution to haematopoiesis and immune cell production, including the generation of white blood cells and immune progenitor activity.

Why it matters: Bone marrow is foundational to immune competence, tissue repair, oxygen transport support, and the replenishment of circulating defence cells.

Functional significance: Depends on nutrient sufficiency, low toxic burden, stable inflammatory signalling, and adequate marrow microenvironment support.

Common patterns: Chronic toxic exposure or nutritional depletion → impaired marrow resilience; persistent infection or inflammation → increased marrow demand and immune cell turnover.

Your result in this scan: Bone marrow Index appears within the analyser's expected range in this scan.

Spleen index

Within range

Immune System

Reference range: 34.367 - 35.642 | **Value:** 35.434

What this means: Reflects splenic immune and filtration activity, including blood surveillance, immune cell storage, removal of damaged cells, and response to circulating antigenic burden.

Why it matters: The spleen plays a major role in blood-borne immune defence, clearance of cellular debris, and coordination of systemic immune monitoring.

Functional significance: Interfaces with blood quality, circulating immune complexes, red cell turnover, inflammatory burden, and adaptive immune mobilisation.

Common patterns: Poor blood quality, chronic inflammation, or higher immune-complex burden → greater splenic filtration demand; splenic strain can coexist with liver burden and immune activation.

Your result in this scan: Spleen index appears within the analyser's expected range in this scan.

Thymus index

Within range

Immune System

Reference range: 58.425 - 61.213 | **Value:** 59.292

What this means: Reflects thymic contribution to T-cell maturation, immune tolerance, and adaptive immune education.

Why it matters: The thymus is central to immune discrimination between self and non-self, helping shape adaptive immune resilience and tolerance patterns.

Functional significance: Closely linked to adrenal-cortisol tone, age-related involution, micronutrient status, inflammatory load, and overall neuroendocrine-immune balance.

Common patterns: Chronic cortisol excess → thymic suppression → reduced adaptive tolerance; immune activation and inflammatory burden can destabilise thymic-immune balance further.

Your result in this scan: Thymus index appears within the analyser's expected range in this scan.

Immunoglobulin index

Within range

Immune System

Reference range: 3.712 - 6.981 | **Value:** 4.089

What this means: Reflects humoral immune activity through immunoglobulin production and antibody-mediated defence capacity.

Why it matters: Immunoglobulins are critical for recognition and neutralisation of pathogens, mucosal protection, and long-term adaptive immune memory.

Functional significance: Depends on adequate B-cell activity, mucosal antigen handling, protein-nutrient sufficiency, and balanced immune tolerance versus overreactivity.

Common patterns: Gut or respiratory barrier compromise → higher antigen exposure → increased immunoglobulin demand; persistent food/environmental triggers can push chronic humoral activation.

Your result in this scan: Immunoglobulin index appears within the analyser's expected range in this scan.

Respiratory Immunity

Within range

Immune System

Reference range: 3.241 - 9.814 | **Value:** 3.384

What this means: Reflects immune competence within the respiratory tract, including airway mucosal defence, pathogen response, and inflammatory tolerance in the lungs and upper airways.

Why it matters: Respiratory immune balance is essential for resistance to recurrent infections, control of allergic reactivity, and healthy tolerance to inhaled environmental exposures.

Functional significance: Linked to mucosal integrity, microbiome exposure, pollution burden, histamine signalling, and the balance between immune defence and airway inflammation.

Common patterns: Pollution, allergens, or chronic sinus-throat burden → ongoing respiratory immune activation; mucosal weakness can increase infection cycles and downstream systemic immune stress.

Your result in this scan: Respiratory Immunity appears within the analyser's expected range in this scan.

GI Immunity

Within range

Immune System

Reference range: 0.638 - 1.712 | **Value:** 0.949

What this means: Reflects immune activity associated with the gut-associated lymphoid tissue (GALT), mucosal barrier defence, and immune tolerance within the gastrointestinal tract.

Why it matters: Most immune activity is intimately linked to the gut; imbalance here can drive food reactivity, chronic inflammation, autoimmunity risk, and systemic immune dysregulation.

Functional significance: Depends on barrier integrity, microbiome balance, secretory immunity, antigen load, and coordination between tolerance and inflammatory response.

Common patterns: Gut dysbiosis + permeability → antigen translocation → immune activation → inflammatory spillover across skin, mood, hormones, and systemic tolerance pathways.

Your result in this scan: GI Immunity appears within the analyser's expected range in this scan.

Mucosal Immunity

Within range

Immune System

Reference range: 4.111 - 18.741 | **Value:** 10.307

What this means: Reflects integrated mucosal immune defence across barrier tissues, including secretory immune activity, epithelial protection, and tolerance to environmental and dietary exposures.

Why it matters: Mucosal immunity is a frontline defence system and a key regulator of infection resistance, allergy patterns, and systemic inflammatory tone.

Functional significance: Interacts with secretory IgA dynamics, epithelial integrity, microbiome balance, histamine responses, and nutrient-dependent barrier repair.

Common patterns: Barrier weakness → higher antigen penetration → immune activation; dysbiosis or nutrient insufficiency can reduce mucosal resilience and increase allergy or infection cycles.

Your result in this scan: Mucosal Immunity appears within the analyser's expected range in this scan.

Thyroid

Medium priority

Complete system marker table — all markers and all severity grades are shown below.

Flagged markers: 3 | Within range: 1

FT4

Mildly elevated

Thyroid

Reference range: 0.103 - 0.316 | **Value:** 0.6

What this means: Represents circulating levels of thyroxine available for peripheral conversion into active thyroid hormone (T3).

Why it matters: FT4 is a key determinant of metabolic rate, influencing energy production, temperature regulation, and overall endocrine balance.

Functional significance: Linked to thyroid gland output, iodine status, deiodinase activity, and feedback from the hypothalamic-pituitary-thyroid axis.

Common patterns: Iodine imbalance + stress + inflammation -> altered thyroid output -> disrupted metabolic regulation.

Your result in this scan: Indicates increased thyroid hormone availability, which may reflect hyperthyroid tendencies or compensatory activation.

Thyroglobulin

Mildly elevated

Thyroid

Reference range: 0.114 - 0.202 | **Value:** 0.294

What this means: Reflects the protein precursor used in thyroid hormone synthesis within the thyroid gland.

Why it matters: Thyroglobulin is essential for the production of T3 and T4 and can indicate thyroid gland activity or dysfunction.

Functional significance: Linked to thyroid tissue integrity, iodine availability, and autoimmune activity.

Common patterns: Autoimmune activity + iodine imbalance -> altered thyroglobulin dynamics.

Your result in this scan: Indicates increased thyroid activity or possible glandular stress.

Thyroglobulin

Within range

Thyroid

Reference range: 0.421 - 0.734 | **Value:** 0.711

What this means: Reflects the protein precursor used in thyroid hormone synthesis within the thyroid gland.

Why it matters: Thyroglobulin is essential for the production of T3 and T4 and can indicate thyroid gland activity or dysfunction.

Functional significance: Linked to thyroid tissue integrity, iodine availability, and autoimmune activity.

Common patterns: Autoimmune activity + iodine imbalance -> altered thyroglobulin dynamics.

Your result in this scan: Thyroglobulin appears within the analyser's expected range in this scan.

T3

Mildly elevated

Thyroid

Reference range: 0.161 - 0.308 | **Value:** 0.518

What this means: Represents levels of active thyroid hormone responsible for metabolic activation at cellular level.

Why it matters: T3 directly regulates mitochondrial activity, energy production, and metabolic rate.

Functional significance: Linked to FT4 conversion, selenium status, liver function, and stress.

Common patterns: Stress + inflammation -> impaired T4 to T3 conversion.

Your result in this scan: Indicates elevated T3 activity, potentially reflecting hypermetabolic state.

Human Toxin

Within range

Complete system marker table — all markers and all severity grades are shown below.

Flagged markers: 0 | Within range: 4

Stimulating Beverage

Within range

Human Toxin

Reference range: 0.209 - 0.751 | **Value:** 0.475

What this means: Reflects physiological burden associated with stimulant-beverage exposure, especially caffeine-type intake that can influence adrenal output, autonomic tone, sleep quality, and vascular reactivity.

Why it matters: High stimulant load can mask fatigue, increase sympathetic drive, destabilise blood sugar, impair sleep recovery, and elevate cardiovascular or stress-related strain.

Functional significance: Closely linked to adrenal-cortisol rhythm, catecholamine tone, heart-rate variability, sleep depth, dehydration patterns, and compensatory use during low-energy states.

Common patterns: Low energy + poor sleep -> greater stimulant reliance -> higher cortisol and sympathetic activation -> worsened recovery debt, palpitations, and blood-sugar volatility.

Your result in this scan: Stimulating Beverage appears within the analyser's expected range in this scan.

Electromagnetic Radiation

Within range

Human Toxin

Reference range: 0.046 - 0.167 | **Value:** 0.109

What this means: Reflects potential physiological stress associated with electromagnetic exposure, interpreted here as an environmental-load marker rather than a standalone diagnostic measure.

Why it matters: In sensitive individuals, cumulative environmental stress may contribute to sleep disturbance, autonomic irritability, headaches, or reduced recovery resilience, even when mechanisms remain multifactorial.

Functional significance: Best interpreted alongside nervous-system tone, sleep disruption, oxidative stress burden, and total environmental sensitivity rather than in isolation.

Common patterns: High total-load sensitivity + poor sleep + autonomic overdrive can make environmental exposures feel more provocative; oxidative and stress burden often coexist.

Your result in this scan: Electromagnetic Radiation appears within the analyser's expected range in this scan.

Free Radical

Within range

Human Toxin

Reference range: 0.124 - 0.453 | **Value:** 0.178

What this means: Reflects oxidative free-radical burden and the degree to which reactive species may be exceeding antioxidant buffering capacity.

Why it matters: Excess free-radical activity can damage lipids, proteins, DNA, membranes, and mitochondria, thereby accelerating inflammation, ageing, and tissue dysfunction.

Functional significance: Closely linked to mitochondrial efficiency, detoxification load, inflammation, lipid oxidation, glutathione status, selenium-dependent antioxidant systems, and toxic exposure.

Common patterns: Toxic burden + mitochondrial strain + inflammation -> higher free-radical activity -> membrane damage, fatigue, vascular stress, and accelerated tissue ageing.

Your result in this scan: Free Radical appears within the analyser's expected range in this scan.

Toxic Pesticide Residue

Within range

Human Toxin

Reference range: 0.013 - 0.313 | **Value:** 0.305

What this means: Reflects burden associated with pesticide-type toxic exposure, suggesting environmental chemical load requiring hepatic and antioxidant clearance support.

Why it matters: Pesticide burden can increase detoxification demand, oxidative stress, endocrine disruption risk, and cumulative toxic load that affects neurological, immune, and metabolic resilience.

Functional significance: Linked to liver Phase I and Phase II detoxification, glutathione reserve, endocrine sensitivity, mitochondrial function, and cumulative environmental exposure history.

Common patterns: Pesticide load -> higher hepatic detox demand + glutathione usage -> oxidative stress and endocrine strain; gut and nervous-system symptoms may rise as tolerance falls.

Your result in this scan: Toxic Pesticide Residue appears within the analyser's expected range in this scan.

Heavy Metal

High priority

Complete system marker table — all markers and all severity grades are shown below.

Flagged markers: 4 | Within range: 4

Lead

Mildly elevated

Heavy Metal

Reference range: 0.052 - 0.643 | **Value:** 0.976

What this means: Reflects lead burden as a toxic-metal exposure marker relevant to neurological stress, haematologic interference, mitochondrial impairment, and long-term detox burden.

Why it matters: Lead can disrupt nervous-system function, impair haem-related processes, increase oxidative stress, and reduce developmental and cognitive resilience.

Functional significance: Linked to glutathione demand, mineral competition, neurological function, blood formation, detoxification capacity, and cumulative environmental exposure.

Common patterns: Lead burden -> oxidative stress + mineral displacement + neuronal strain -> fatigue, cognitive fog, irritability, and reduced detox reserve.

Your result in this scan: Indicates increased lead exposure burden or lower tolerance to lead-related toxicity, often associated with neurocognitive, haematologic, or oxidative stress patterns.

Mercury

Mildly elevated

Heavy Metal

Reference range: 0.013 - 0.336 | **Value:** 0.468

What this means: Reflects mercury burden as a toxic-metal exposure marker with relevance to neurological irritation, immune dysregulation, oxidative stress, and detoxification strain.

Why it matters: Mercury can interfere with sulfhydryl-rich enzymes, disrupt neuronal stability, increase oxidative burden, and challenge immune and mitochondrial resilience.

Functional significance: Closely linked to glutathione systems, selenium balance, nervous-system sensitivity, mitochondrial stress, and cumulative toxic exposure.

Common patterns: Mercury burden + low selenium/glutathione -> greater neuronal and oxidative stress; immune reactivity and fatigue patterns may intensify as detox reserve falls.

Your result in this scan: Indicates increased mercury burden or reduced tolerance to mercury-associated stress, potentially affecting cognition, mood, immune balance, and oxidative defence.

Cadmium

Mildly elevated

Heavy Metal

Reference range: 0.527 - 1.523 | **Value:** 1.556

What this means: Reflects cadmium burden as a toxic-metal exposure marker associated with oxidative damage, kidney stress, and competition with essential minerals such as zinc.

Why it matters: Cadmium can accumulate over time, increasing detox burden, impairing antioxidant systems, and stressing renal and metabolic regulation.

Functional significance: Linked to kidney function, zinc-dependent processes, glutathione demand, smoking or environmental exposure history, and chronic inflammatory load.

Common patterns: Cadmium burden -> zinc displacement + oxidative stress + renal strain -> weaker detox resilience and altered immune-barrier function.

Your result in this scan: Indicates increased cadmium exposure or reduced tolerance to cadmium-related toxicity, often associated with renal stress, mineral imbalance, and oxidative burden.

Chromium

Within range

Heavy Metal

Reference range: 0.176 - 1.183 | **Value:** 0.345

What this means: Reflects chromium burden in this context as an exposure-related heavy-metal marker rather than the nutritional trivalent chromium used in glucose metabolism.

Why it matters: Toxic chromium exposure can provoke oxidative stress, tissue irritation, and detoxification burden, especially when environmental or occupational exposure is relevant.

Functional significance: Best interpreted alongside exposure history, glutathione-antioxidant reserve, respiratory or skin reactivity, and broader toxic-load indicators.

Common patterns: Toxic chromium exposure can align with respiratory, skin, and oxidative burden patterns; reduced detox reserve may increase symptomatic sensitivity.

Your result in this scan: Chromium appears within the analyser's expected range in this scan.

Arsenic

Within range

Heavy Metal

Reference range: 0.153 - 0.621 | **Value:** 0.596

What this means: Reflects arsenic burden as a toxic-metal and toxicoid exposure marker relevant to detoxification stress, vascular effects, mitochondrial dysfunction, and skin or nerve irritation.

Why it matters: Arsenic burden can impair energy metabolism, increase oxidative stress, and affect vascular, neurological, and skin-related systems over time.

Functional significance: Linked to methylation capacity, glutathione support, vascular health, skin changes, mitochondrial function, and cumulative exposure burden.

Common patterns: Arsenic burden -> oxidative stress + mitochondrial inhibition + vascular strain; methylation weakness may reduce effective handling and prolong tissue impact.

Your result in this scan: Arsenic appears within the analyser's expected range in this scan.

Antimony

Within range

Heavy Metal

Reference range: 0.162 - 0.412 | **Value:** 0.323

What this means: Reflects antimony burden as a toxic-metal exposure marker associated with environmental or occupational load and increased detoxification demand.

Why it matters: Although less commonly discussed than lead or mercury, antimony burden may still contribute to toxic stress, oxidative load, and reduced resilience in sensitive individuals.

Functional significance: Linked to cumulative environmental burden, liver-detox reserve, glutathione demand, and broad oxidative-inflammatory stress patterns.

Common patterns: Higher cumulative metal burden -> greater glutathione usage and detox pressure; antimony should be interpreted within the broader heavy-metal landscape rather than alone.

Your result in this scan: Antimony appears within the analyser's expected range in this scan.

Thallium

Within range

Heavy Metal

Reference range: 0.182 - 0.542 | **Value:** 0.388

What this means: Reflects thallium burden as a highly concerning toxic-metal exposure marker relevant to neurological, gastrointestinal, and mitochondrial stress.

Why it matters: Thallium can significantly disrupt cellular metabolism and nerve function, making even moderate burden clinically important within a toxic-load context.

Functional significance: Linked to mitochondrial disruption, nervous-system irritation, detox reserve, potassium-related transport pathways, and cumulative environmental exposure.

Common patterns: Thallium burden -> mitochondrial stress + neurological irritation + systemic fatigue; weak detox reserve may amplify symptom severity and recovery delay.

Your result in this scan: Thallium appears within the analyser's expected range in this scan.

Aluminum

Mildly elevated

Heavy Metal

Reference range: 0.192 - 0.412 | **Value:** 0.57

What this means: Reflects aluminium burden as a toxic-metal exposure marker relevant to neurological stress, bone-mineral interference, and cumulative detoxification demand.

Why it matters: Aluminium burden has been associated with oxidative stress, altered neuronal function, mineral disruption, and chronic low-grade toxic load in sensitive individuals.

Functional significance: Linked to nervous-system sensitivity, bone-mineral balance, detox reserve, oxidative stress, and competition with essential minerals.

Common patterns: Aluminium burden -> oxidative stress + mineral interference + neurological strain; weaker detox reserve and poor silica support may reduce resilience.

Your result in this scan: Indicates increased aluminium-related exposure burden or reduced tolerance to its neurological and mineral-disruptive effects.

Basic Physical Quality

Low priority

Complete system marker table — all markers and all severity grades are shown below.

Flagged markers: 1 | Within range: 4

Response Ability

Within range

Basic Physical Quality

Reference range: 59.786 - 65.424 | **Value:** 61.674

What this means: Reflects the body's general capacity to perceive, adapt to, and recover from internal or external stressors, functioning as a broad resilience marker rather than a single organ-specific measure.

Why it matters: Response ability matters because many clinical patterns are shaped not only by the burden itself, but by how effectively the organism can mount an adaptive response without tipping into exhaustion, collapse, or prolonged dysregulation.

Functional significance: Linked to autonomic flexibility, adrenal reserve, mitochondrial energy availability, immune responsiveness, inflammatory control, and the overall coordination of recovery systems under stress.

Common patterns: Chronic stress + low mitochondrial output + poor sleep + inflammatory load -> reduced response ability; stronger autonomic flexibility and recovery capacity improve it.

Your result in this scan: Response Ability appears within the analyser's expected range in this scan.

Mental Power

Within range

Basic Physical Quality

Reference range: 58.715 - 63.213 | **Value:** 61.23

What this means: Reflects perceived neurocognitive drive and mental stamina, including sustained attention, executive energy, and resistance to cognitive fatigue.

Why it matters: Mental stamina depends on stable glucose delivery, mitochondrial output, neurotransmitter availability, sleep quality, and low inflammatory interference with brain function.

Functional significance: Integrates cerebral perfusion, catecholamine balance, thyroid-adrenal support, and phospholipid membrane integrity.

Common patterns: Low thyroid or poor sleep → reduced ATP output and mental endurance; chronic stress → temporary overdrive followed by reduced resilience; gut inflammation and glycaemic instability can further erode clarity and focus.

Your result in this scan: Mental Power appears within the analyser's expected range in this scan.

Water Shortage

Within range

Basic Physical Quality

Reference range: 33.967 - 37.642 | **Value:** 34.512

What this means: Reflects dehydration burden or reduced effective body-water availability, indicating that tissue hydration and fluid distribution may be insufficient for optimal physiological function.

Why it matters: Adequate hydration is essential for circulation, kidney clearance, electrolyte balance, thermoregulation, mucosal integrity, and efficient metabolic reactions at the cellular level.

Functional significance: Linked to intracellular and extracellular fluid balance, electrolyte status, kidney handling, adrenal-sodium regulation, perspiration loss, and the distinction between true hydration and fluid retention.

Common patterns: Low intake + electrolyte imbalance + high loss through stress, heat, or diuresis -> water shortage; dehydration can worsen circulation, kidney strain, headaches, fatigue, and mucosal dryness.

Your result in this scan: Water Shortage appears within the analyser's expected range in this scan.

Hypoxia

Mildly reduced

Basic Physical Quality

Reference range: 133.642 - 141.476 | **Value:** 130.994

What this means: Reflects reduced oxygen availability at the tissue level, indicating a functional mismatch between oxygen demand and delivery or utilisation.

Why it matters: Hypoxia matters because inadequate oxygenation compromises ATP production, impairs tissue repair, increases fatigue, and amplifies vulnerability in high-demand systems such as brain, heart, and muscle.

Functional significance: Linked to lung function, haemoglobin and iron status, microcirculation, cardiac output, mitochondrial efficiency, inflammatory tissue congestion, and the distinction between low delivery and poor utilisation.

Common patterns: Poor lung mechanics + low perfusion + reduced haemoglobin or mitochondrial dysfunction -> hypoxic patterns; inflammation and oedema can further impair local oxygen diffusion.

Your result in this scan: Suggests lower evidence of meaningful tissue oxygen deficit and more favourable oxygen delivery-utilisation balance.

PH

Within range

Basic Physical Quality

Reference range: 3.156 - 3.694 | **Value:** 3.248

What this means: Reflects overall acid-base tendency within the source framework, functioning as a broad indicator of systemic buffering balance rather than a standalone compartment-specific laboratory pH value.

Why it matters: Acid-base balance matters because enzyme activity, mitochondrial output, mineral stability, respiratory compensation, and renal buffering all depend on an adequately regulated internal pH environment.

Functional significance: Linked to renal acid excretion, respiratory CO2 handling, dietary acid load, mineral buffering, mitochondrial metabolism, and inflammatory or catabolic states that shift the internal acid-base

terrain.

Common patterns: High metabolic stress + poor renal buffering + altered ventilation -> pH imbalance; chronic inflammatory load, tissue hypoxia, and catabolic states can all destabilise acid-base resilience.

Your result in this scan: PH appears within the analyser's expected range in this scan.

Allergy

High priority

Complete system marker table — all markers and all severity grades are shown below.

Flagged markers: 6 | Within range: 7

Drug Sensitivity

Within range

Allergy

Reference range: 0.431 - 1.329 | **Value:** 0.563

What this means: Reflects hypersensitivity tendency to pharmaceutical compounds, indicating how strongly immune and detoxification systems may react to drug-related antigenic or chemical exposure.

Why it matters: Drug reactivity can signal altered immune tolerance, histamine burden, impaired biotransformation capacity, or heightened sensitivity to xenobiotic load.

Functional significance: Closely linked to liver Phase I and Phase II detoxification, mast-cell and histamine activity, barrier integrity, and prior antigen sensitisation patterns.

Common patterns: Poor liver clearance plus high histamine tone can increase drug reactivity; gut barrier dysfunction may also heighten immune sensitisation and reduce tolerance to chemical compounds.

Your result in this scan: Drug Sensitivity appears within the analyser's expected range in this scan.

Alcohol allergy index

Within range

Allergy

Reference range: 0.432 - 1.246 | **Value:** 1.207

What this means: Reflects hypersensitivity or low tolerance tendency toward alcohol exposure, including immune reactivity, aldehyde-handling burden, and histamine-related response patterns.

Why it matters: Alcohol reactivity can indicate poor aldehyde clearance, histamine intolerance, liver stress, and reduced resilience to inflammatory or toxic load.

Functional significance: Interacts with liver detoxification, aldehyde metabolism, sulphur pathways, histamine balance, gut permeability, and inflammatory signalling.

Common patterns: Low aldehyde clearance plus liver burden can intensify alcohol sensitivity; alcohol can also worsen gut permeability and histamine symptoms, amplifying immune reactivity.

Your result in this scan: Alcohol allergy index appears within the analyser's expected range in this scan.

Pollen allergy index

Within range

Allergy

Reference range: 0.143 - 1.989 | **Value:** 1.646

What this means: Reflects hypersensitivity tendency to airborne pollen antigens, indicating mucosal immune overreactivity and reduced tolerance to seasonal environmental exposures.

Why it matters: Pollen reactivity often reflects histamine excess, barrier hyperreactivity, and imbalanced immune tolerance at the respiratory mucosal surface.

Functional significance: Closely linked to Th2-skewed immune patterns, mast-cell activation, histamine metabolism, respiratory mucosal integrity, and total antigenic burden.

Common patterns: Respiratory mucosal weakness plus high histamine tone can drive pollen symptoms; liver overload and gut dysbiosis may reduce tolerance and prolong immune reactivity.

Your result in this scan: Pollen allergy index appears within the analyser's expected range in this scan.

Injection allergy index

Mildly elevated

Allergy

Reference range: 0.847 - 1.045 | **Value:** 1.184

What this means: Reflects hypersensitivity tendency to injected substances, suggesting reactivity to direct antigenic or chemical exposure bypassing digestive barriers.

Why it matters: Injection sensitivity may indicate heightened immune vigilance, mast-cell instability, or low tolerance to introduced compounds and preservatives.

Functional significance: Interacts with histamine signalling, mast-cell reactivity, detox reserve, prior sensitisation history, and the broader immune tolerance threshold.

Common patterns: High histamine tone and reduced detox resilience can increase injection-related sensitivity; systemic immune priming may amplify reactions to introduced compounds.

Your result in this scan: Indicates increased reactivity to injected exposures, often associated with immune sensitisation, mast-cell activation, or lower chemical tolerance.

Chemical Sensitivity

Mildly elevated

Allergy

Reference range: 0.842 - 1.643 | **Value:** 1.896

What this means: Reflects hypersensitivity tendency to chemical products, indicating low tolerance to environmental chemical exposure and potential overload of detoxification and barrier systems.

Why it matters: Chemical sensitivity often points to reduced biotransformation reserve, increased histamine activity, impaired barrier resilience, or chronic cumulative toxic burden.

Functional significance: Strongly linked to liver detox pathways, glutathione status, histamine regulation, mucosal and skin barrier integrity, and cumulative environmental exposure patterns.

Common patterns: Low glutathione plus liver overload can reduce tolerance to chemical products; skin and respiratory barriers may become more reactive when histamine and toxic burden are high.

Your result in this scan: Indicates increased chemical sensitivity, poor tolerance to fragrances or compounds, or a detox-histamine-barrier pattern prone to environmental reactivity.

Paint allergy index

Within range

Allergy

Reference range: 0.346 - 1.401 | **Value:** 0.489

What this means: Reflects sensitivity to paint-related compounds and volatile chemical exposures, suggesting heightened reactivity to inhaled or contact-based environmental irritants.

Why it matters: Paint sensitivity can reveal low tolerance to solvents, volatile compounds, and chemically induced mucosal or skin inflammation.

Functional significance: Closely linked to liver detox reserve, respiratory mucosal integrity, histamine activity, glutathione-dependent clearance, and cumulative environmental burden.

Common patterns: Volatile chemical exposure plus low detox reserve can provoke respiratory and skin reactivity; glutathione insufficiency may reduce resilience to solvent-type burden.

Your result in this scan: Paint allergy index appears within the analyser's expected range in this scan.

Dust allergy index

Mildly elevated

Allergy

Reference range: 0.543 - 1.023 | **Value:** 1.263

What this means: Reflects hypersensitivity tendency to dust and particulate exposure, indicating respiratory mucosal reactivity and immune overresponse to airborne antigenic burden.

Why it matters: Dust sensitivity is commonly associated with histamine excess, respiratory barrier weakness, and chronic antigenic stimulation of the upper airway immune system.

Functional significance: Interacts with respiratory mucosal immunity, mast-cell activity, histamine metabolism, sinus reactivity, and the broader total-load of inhaled irritants.

Common patterns: Dust burden can amplify histamine symptoms when respiratory barriers are weak; gut-immune imbalance and liver congestion may reduce tolerance and prolong reactivity.

Your result in this scan: Indicates heightened reactivity to dust particulates, respiratory immune activation, or chronic mucosal sensitivity.

Smoke allergy index

Mildly elevated

Allergy

Reference range: 0.826 - 1.013 | **Value:** 2.532

What this means: Reflects hypersensitivity tendency to smoke exposure, indicating respiratory tract irritation, oxidative burden, and mucosal immune overreactivity.

Why it matters: Smoke sensitivity points to reduced respiratory resilience, inflammatory reactivity, and often a lower threshold for oxidative or chemical irritation.

Functional significance: Closely linked to respiratory mucosal integrity, antioxidant capacity, glutathione reserve, histamine activity, and detoxification tolerance.

Common patterns: Low glutathione and high histamine tone can intensify smoke reactions; repeated exposure may amplify respiratory inflammation and wider immune burden.

Your result in this scan: Indicates increased smoke sensitivity, often associated with oxidative stress, mucosal irritation, histamine response, or poor respiratory-barrier protection.

Hair dye allergy index

Moderately elevated

Allergy

Reference range: 0.717 - 1.486 | **Value:** 2.856

What this means: Reflects cutaneous and chemical hypersensitivity tendency to hair-dye compounds, suggesting lower tolerance to topical chemical exposure and sensitisation of skin immune pathways.

Why it matters: Hair-dye reactivity often signals barrier fragility, chemical sensitivity, histamine activation, or immune sensitisation to repeated topical exposure.

Functional significance: Interacts with skin barrier integrity, liver detox reserve, histamine response, glutathione status, and cumulative exposure to aromatic chemical compounds.

Common patterns: Low glutathione and impaired skin barrier function can worsen topical chemical reactions; liver overload and histamine excess may intensify visible skin responses.

Your result in this scan: Indicates stronger sensitivity to hair-dye compounds, often associated with barrier compromise, histamine response, or reduced chemical tolerance.

Animal fur allergy index

Within range

Allergy

Reference range: 0.124 - 1.192 | **Value:** 0.571

What this means: Reflects hypersensitivity tendency to animal fur or dander exposure, indicating respiratory and mucosal immune reactivity to environmental protein antigens.

Why it matters: Animal-dander sensitivity commonly reflects histamine-driven immune response, mucosal overreactivity, and reduced tolerance to persistent airborne antigen exposure.

Functional significance: Closely linked to mast-cell activity, respiratory mucosal integrity, histamine clearance, and total environmental antigen burden.

Common patterns: Respiratory barrier weakness plus high histamine tone can intensify dander reactions; liver and gut burden may lower overall tolerance to chronic antigen exposure.

Your result in this scan: Animal fur allergy index appears within the analyser's expected range in this scan.

Metal jewelry allergy index

Within range

Allergy

Reference range: 0.549 - 1.213 | **Value:** 0.743

What this means: Reflects cutaneous hypersensitivity tendency to metal contact, indicating local immune sensitisation and lower tolerance to metal-triggered inflammatory skin reactions.

Why it matters: Metal reactivity may point to skin barrier weakness, metal sensitivity, and heightened local immune activation upon repeated contact exposure.

Functional significance: Interacts with skin barrier integrity, histamine response, metal burden tolerance, and the broader detoxification landscape.

Common patterns: Barrier weakness and poor detox tolerance can heighten contact-metal reactions; systemic histamine and immune reactivity may make local responses more visible or persistent.

Your result in this scan: Metal jewelry allergy index appears within the analyser's expected range in this scan.

Seafood allergy index

Mildly elevated

Allergy

Reference range: 0.449 - 1.246 | **Value:** 2.379

What this means: Reflects hypersensitivity tendency to seafood antigens, indicating altered oral-gut immune tolerance and increased risk of food-triggered immune reactivity.

Why it matters: Seafood sensitivity may signal impaired digestive tolerance, histamine overload, or heightened antigen response across the gut-immune axis.

Functional significance: Strongly linked to gut permeability, digestive adequacy, histamine metabolism, mucosal immunity, and prior sensitisation patterns.

Common patterns: Gut barrier compromise can increase exposure to food antigens and amplify seafood sensitivity; histamine excess may worsen symptom intensity and duration.

Your result in this scan: Indicates greater food-triggered hypersensitivity, often associated with histamine symptoms, gut permeability, or reduced oral tolerance.

Milk allergy index

Within range

Allergy

Reference range: 0.477 - 1.348 | **Value:** 0.739

What this means: Reflects hypersensitivity tendency to milk-related antigens, indicating altered digestive-immune tolerance and increased food-reactive potential at the gut barrier.

Why it matters: Milk sensitivity may arise from immune reactivity to milk proteins, poor digestive handling, mucosal inflammation, or broader gut permeability patterns.

Functional significance: Closely linked to stomach and pancreatic digestive adequacy, gut permeability, mucosal immunity, histamine activity, and immune tolerance mechanisms.

Common patterns: Low digestive capacity plus increased gut permeability can heighten milk reactivity; immune activation and histamine patterns may extend symptoms beyond the digestive tract.

Your result in this scan: Milk allergy index appears within the analyser's expected range in this scan.

Obesity

High priority

Complete system marker table — all markers and all severity grades are shown below.

Flagged markers: 4 | Within range: 1

Abnormal lipid metabolism coefficient

Moderately reduced

Obesity

Reference range: 1.992 - 3.713 | **Value:** 0.89

What this means: Reflects baseline mitochondrial energy expenditure under resting conditions.

Why it matters: Determines energy balance, weight regulation, and metabolic efficiency.

Functional significance: Linked to thyroid hormones, mitochondrial density, and lean mass.

Common patterns: Hypothyroid + low muscle mass -> low BMR.

Your result in this scan: Suggests metabolic slowdown and fatigue.

Brown adipose tissue abnormalities coefficient

Moderately reduced

Obesity

Reference range: 2.791 - 4.202 | **Value:** 1.858

What this means: Reflects functional irregularity in brown adipose tissue, the thermogenic fat compartment involved in heat production, metabolic flexibility, and energy expenditure.

Why it matters: Brown adipose tissue helps dissipate energy through thermogenesis rather than storage, so altered function may contribute to weight gain tendency, cold intolerance, reduced metabolic responsiveness, and impaired glucose-lipid handling.

Functional significance: Linked to mitochondrial uncoupling activity, thyroid signalling, sympathetic tone, insulin sensitivity, cold-adaptation capacity, and overall metabolic rate regulation.

Common patterns: Low thyroid activity + low sympathetic thermogenic drive + mitochondrial inefficiency -> weaker brown-fat activation -> lower energy expenditure and greater obesity susceptibility.

Your result in this scan: Suggests more favourable brown-fat thermogenic activity and less evidence of impaired adaptive energy expenditure.

Hyperinsulinemia coefficient

Within range

Obesity

Reference range: 0.097 - 0.215 | **Value:** 0.206

What this means: Represents elevated insulin activity within reproductive-metabolic interactions, often linked to hormonal imbalance.

Why it matters: Hyperinsulinemia can disrupt reproductive hormones and metabolic stability.

Functional significance: Linked to insulin resistance, ovarian function, and endocrine signalling.

Common patterns: Insulin resistance -> reproductive dysfunction.

Your result in this scan: Hyperinsulinemia coefficient appears within the analyser's expected range in this scan.

Nucleus of the hypothalamus abnormal coefficient

Mildly elevated

Obesity

Reference range: 0.332 - 0.626 | **Value:** 0.719

What this means: Reflects dysregulation within hypothalamic centres that coordinate appetite, satiety, autonomic tone, circadian signalling, and energy-balance regulation.

Why it matters: The hypothalamus is central to body-weight regulation because it integrates hunger, leptin and insulin signals, stress cues, temperature regulation, and circadian rhythm into behavioural and metabolic output.

Functional significance: Linked to leptin sensitivity, insulin signalling, cortisol rhythm, circadian stability, reward-driven eating, inflammatory signalling, and autonomic regulation of metabolism.

Common patterns: Sleep disruption + leptin resistance + chronic stress + inflammation -> hypothalamic dysregulation -> appetite instability, weight gain tendency, and poorer metabolic adaptation.

Your result in this scan: Indicates disrupted appetite-satiety regulation, circadian-metabolic dysregulation, or hypothalamic stress affecting body-weight control and energy homeostasis.

Triglyceride content of abnormal coefficient

Moderately elevated

Obesity

Reference range: 1.341 - 1.991 | **Value:** 5.222

What this means: Reflects mitochondrial coupling efficiency in ATP production versus heat generation.

Why it matters: Determines fatigue, performance, and metabolic flexibility.

Functional significance: Linked to mitochondrial health, oxidative phosphorylation, and nutrient utilisation.

Common patterns: Mitochondrial dysfunction -> low efficiency.

Your result in this scan: Indicates efficient ATP generation.

Skin

High priority

Complete system marker table — all markers and all severity grades are shown below.

Flagged markers: 6 | Within range: 4

Skin Free Radical Index

Within range

Skin

Reference range: 0.124 - 3.453 | **Value:** 2.972

What this means: Reflects the oxidative stress burden at the level of skin tissue, indicating the balance between free radical generation and antioxidant defence within the dermal and epidermal layers.

Why it matters: Skin is highly exposed to environmental oxidative stress (UV, pollution), and elevated free radical activity accelerates ageing, inflammation, pigmentation, and barrier dysfunction.

Functional significance: Linked to systemic antioxidant status, mitochondrial function, detoxification capacity, and environmental exposure load.

Common patterns: UV exposure + toxin load + low antioxidants -> elevated skin oxidative stress -> premature ageing, pigmentation, inflammation.

Your result in this scan: Skin Free Radical Index appears within the analyser's expected range in this scan.

Skin Collagen Index

Mildly reduced

Skin

Reference range: 4.471 - 6.079 | **Value:** 2.899

What this means: Reflects collagen support within skin tissue, indicating the quality of dermal structural proteins and visible connective tissue resilience.

Why it matters: Skin collagen influences firmness, elasticity, repair capacity, and the visible pace of connective tissue ageing.

Functional significance: Depends on vitamin C, protein availability, copper and silicon balance, hormone status, and oxidative stress control.

Common patterns: Oxidative stress, glycation, hormone decline, and low vitamin C can reduce skin collagen integrity and worsen visible ageing or slower healing.

Your result in this scan: Suggests reduced dermal matrix support, weaker repair, accelerated skin ageing, or lower structural resilience within the skin.

Skin Grease Index

Mildly elevated

Skin

Reference range: 14.477 - 21.348 | **Value:** 25.821

What this means: Reflects sebum production and lipid balance on the skin surface, indicating sebaceous gland activity and skin barrier lubrication.

Why it matters: Sebum plays a protective role in maintaining skin hydration and antimicrobial defence, but imbalance can lead to dryness or acne-related conditions.

Functional significance: Linked to androgen activity, lipid metabolism, gut-skin axis, and inflammatory signalling.

Common patterns: Hormonal imbalance + inflammation + gut dysbiosis -> increased sebum -> acne; low lipids -> dry skin and barrier weakness.

Your result in this scan: Indicates increased sebum production, potentially linked to hormonal imbalance, inflammation, or acne-prone patterns.

Skin Immunity Index

Mildly elevated

Skin

Reference range: 1.035 - 3.230 | **Value:** 4.02

What this means: Reflects immune defence activity at the skin level, indicating the ability of the skin barrier to respond to pathogens, irritation, and environmental stress.

Why it matters: Skin immunity is the first line of defence against infection and environmental insults, and dysfunction can lead to sensitivity, infections, or inflammatory skin conditions.

Functional significance: Linked to gut health, microbiome balance, systemic immunity, and inflammatory regulation.

Common patterns: Gut dysbiosis + immune imbalance -> skin inflammation (eczema, acne, psoriasis patterns).

Your result in this scan: May indicate heightened immune activation, sensitivity, or inflammatory skin reactivity.

Skin Moisture Index

Moderately elevated

Skin

Reference range: 0.218 - 0.953 | **Value:** 2.145

What this means: Reflects hydration retention and barrier moisture quality within skin tissue.

Why it matters: Adequate skin moisture is important for barrier defence, comfort, wound healing, and resistance to irritation.

Functional significance: Linked to essential fatty acids, vitamin A status, collagen integrity, mineral balance, thyroid function, and inflammatory-barrier stress.

Common patterns: Low essential fatty acids, low vitamin A, and reduced collagen support can weaken moisture retention; thyroid and inflammatory patterns may further impair barrier quality.

Your result in this scan: May indicate preserved hydration or transient fluid retention requiring broader skin context.

Skin Moisture Loss

Within range

Skin

Reference range: 2.214 - 4.158 | **Value:** 3.125

What this means: Reflects transepidermal water loss (TEWL), indicating the integrity of the skin barrier and its ability to retain moisture.

Why it matters: Excessive moisture loss compromises barrier function, increases sensitivity, and accelerates ageing and inflammation.

Functional significance: Linked to lipid barrier integrity, ceramide levels, hydration status, and environmental exposure.

Common patterns: Low lipids + dehydration + oxidative stress -> increased TEWL -> dry, sensitive skin.

Your result in this scan: Skin Moisture Loss appears within the analyser's expected range in this scan.

Skin Red Blood Trace Index

Within range

Skin

Reference range: 0.824 - 1.942 | **Value:** 1.055

What this means: Reflects microcirculation and capillary activity within skin tissue, indicating blood supply and oxygen delivery to the dermis.

Why it matters: Adequate circulation supports nutrient delivery, repair, and healthy skin tone, while imbalance may indicate inflammation or vascular issues.

Functional significance: Linked to vascular health, oxygenation, inflammation, and autonomic tone.

Common patterns: Poor circulation -> dull skin; inflammation -> redness; vascular instability -> sensitivity.

Your result in this scan: Skin Red Blood Trace Index appears within the analyser's expected range in this scan.

Skin Elasticity Index

Mildly reduced

Skin

Reference range: 2.717 - 3.512 | **Value:** 1.571

What this means: Reflects the recoil capacity and structural flexibility of skin tissue, indicating the condition of collagen-elastin architecture and hydration-related matrix support.

Why it matters: Skin elasticity is a visible marker of connective tissue quality, ageing pace, oxidative stress burden, and extracellular matrix resilience.

Functional significance: Depends on collagen turnover, elastin preservation, vitamin C status, copper support, hydration, hormone balance, and oxidative protection.

Common patterns: Oxidative stress, glycation, low collagen support, and hormone decline can reduce skin elasticity; poor hydration and inflammation worsen the pattern.

Your result in this scan: Suggests reduced connective tissue resilience, accelerated matrix ageing, oxidative degradation, or weaker structural support within the dermis.

Skin Melanin Index

Mildly elevated

Skin

Reference range: 0.346 - 0.501 | **Value:** 0.612

What this means: Reflects skin pigment expression and melanin-related tone, with relevance to oxidative exposure, inflammatory change, and hormonal influence on skin tissue.

Why it matters: Altered melanin patterns can provide clues about inflammatory repair demand, endocrine shifts, UV burden, and local tissue reactivity within the skin barrier.

Functional significance: Interacts with oxidative stress, hormone patterns, liver function, inflammatory signalling, and skin barrier integrity.

Common patterns: Hormonal imbalance, liver burden, or inflammatory skin stress can alter melanin patterns; oxidative exposure may further amplify pigment signalling.

Your result in this scan: Suggests increased pigment signalling, often associated with oxidative exposure, irritation, hormonal shifts, or post-inflammatory change.

Tin

Within range

Skin

Reference range: 0.842 - 1.858 | **Value:** 1.419

What this means: Reflects exposure to tin and trace biological impact.

Why it matters: Primarily reflects environmental exposure.

Functional significance: Linked to toxicity.

Common patterns: Environmental contamination.

Your result in this scan: Tin appears within the analyser's expected range in this scan.

Eye

High priority

Complete system marker table — all markers and all severity grades are shown below.

Flagged markers: 6 | Within range: 2

Bags under the eyes

Within range

Eye

Reference range: 0.510 - 3.109 | **Value:** 3.004

What this means: Reflects periorbital fluid retention and tissue congestion, often indicating lymphatic stagnation, altered renal-fluid handling, or poor overnight recovery.

Why it matters: Periorbital puffiness can signal systemic fluid-regulation stress, impaired lymphatic clearance, sleep disruption, or local microcirculatory sluggishness.

Functional significance: Linked to kidney-fluid balance, lymphatic drainage, sodium-water handling, sleep quality, and local vascular permeability.

Common patterns: Poor sleep + lymphatic congestion + kidney-fluid imbalance -> visible periorbital puffiness; inflammatory salt-water retention may intensify the pattern.

Your result in this scan: Bags under the eyes appears within the analyser's expected range in this scan.

Collagen eye wrinkle

Mildly reduced

Eye

Reference range: 2.031 - 3.107 | **Value:** 1.31

What this means: Reflects collagen degradation and reduced extracellular-matrix integrity in the periorbital region, where connective tissue is thin and highly sensitive to oxidative and hormonal stress.

Why it matters: Periorbital wrinkling is a visible marker of collagen turnover quality, hydration status, oxidative burden, and ageing-related connective-tissue decline.

Functional significance: Depends on vitamin C status, amino-acid availability, collagen synthesis, antioxidant defence, hormone balance, and local hydration.

Common patterns: Low vitamin C + oxidative stress + reduced collagen support -> greater periorbital wrinkling; hormonal decline and dehydration can deepen the pattern.

Your result in this scan: Suggests relatively preserved matrix integrity and slower collagen breakdown in the periorbital tissues.

Dark circles

Mildly elevated

Eye

Reference range: 0.831 - 3.188 | **Value:** 3.883

What this means: Reflects microcirculatory and pigmentation changes in the periorbital region, often linked to vascular congestion or fatigue.

Why it matters: Dark circles can indicate poor sleep, circulatory stagnation, or systemic fatigue affecting tissue oxygenation.

Functional significance: Linked to microcirculation, lymphatic drainage, sleep quality, and stress.

Common patterns: Sleep disruption + stress -> periorbital stagnation.

Your result in this scan: Indicates increased vascular congestion or pigmentation changes.

Lymphatic obstruction

Within range

Eye

Reference range: 1.116 - 4.101 | **Value:** 3.52

What this means: Represents impaired lymphatic drainage affecting fluid balance and tissue detoxification.

Why it matters: Lymphatic flow is essential for removing waste and maintaining tissue fluid balance.

Functional significance: Linked to immune function, inflammation, and interstitial fluid dynamics.

Common patterns: Inflammation + stagnation -> lymphatic obstruction.

Your result in this scan: Lymphatic obstruction appears within the analyser's expected range in this scan.

Sagging

Mildly elevated

Eye

Reference range: 0.233 - 0.559 | **Value:** 0.751

What this means: Reflects loss of tissue elasticity and structural support in skin and connective tissue.

Why it matters: Sagging indicates reduced collagen integrity, ageing processes, or nutritional deficiency.

Functional significance: Linked to collagen synthesis, hormonal balance, and oxidative stress.

Common patterns: Ageing + oxidative stress -> tissue laxity.

Your result in this scan: Indicates reduced elasticity and structural support.

Edema

Moderately elevated

Eye

Reference range: 0.332 - 0.726 | **Value:** 1.234

What this means: Represents fluid accumulation in tissues due to imbalance in vascular and lymphatic dynamics.

Why it matters: Edema indicates impaired fluid regulation and possible inflammation or circulatory issues.

Functional significance: Linked to kidney function, vascular permeability, and lymphatic drainage.

Common patterns: Inflammation + poor circulation -> fluid accumulation.

Your result in this scan: Indicates fluid retention or impaired drainage.

Eye cell activity

Mildly elevated

Eye

Reference range: 0.118 - 0.892 | **Value:** 0.895

What this means: Reflects metabolic and functional activity of ocular cells involved in vision.

Why it matters: Cellular activity determines visual acuity and eye health.

Functional significance: Linked to mitochondrial function, circulation, and nutrient supply.

Common patterns: Oxidative stress -> reduced eye cell function.

Your result in this scan: Indicates increased activity or stress response.

Visual fatigue

Mildly elevated

Eye

Reference range: 2.017 - 5.157 | **Value:** 8.04

What this means: Represents strain on visual processing systems due to overuse or metabolic fatigue.

Why it matters: Visual fatigue affects focus, clarity, and cognitive performance.

Functional significance: Linked to screen exposure, neural processing, and ocular metabolism.

Common patterns: Screen time + stress -> visual fatigue.

Your result in this scan: Indicates excessive strain and overuse.

Collagen

High priority

Complete system marker table — all markers and all severity grades are shown below.

Flagged markers: 6 | Within range: 8

Overall Eye Function

Within range

Collagen

Reference range: 6.352 - 8.325 | **Value:** 8.31

What this means: Represents the overall functional status of the visual system, integrating ocular tissue vitality, circulation, and visual performance.

Why it matters: The eye reflects neurovascular, metabolic, and oxidative balance, and can show strain when circulation, energy supply, or tissue support is suboptimal.

Functional significance: Linked to retinal metabolism, optic signalling, ocular microcirculation, oxidative stress, screen strain, and broader nervous-system load.

Common patterns: Visual strain + poor sleep + oxidative stress + microcirculatory weakness -> reduced visual efficiency and greater ocular fatigue.

Your result in this scan: Overall Eye Function appears within the analyser's expected range in this scan.

Endocrine system

Within range

Collagen

Reference range: 7.245 - 8.562 | **Value:** 7.491

What this means: Reflects integrity of collagen matrix within the endocrine system, influencing structural stability and functional resilience.

Why it matters: Collagen is the primary structural protein underpinning tissue integrity and repair across the body.

Functional significance: Linked to vitamin C, glycine, proline, copper, oxidative stress, and glycation damage.

Common patterns: Oxidative stress + glycation -> collagen breakdown in endocrine system leading to dysfunction.

Your result in this scan: Endocrine system appears within the analyser's expected range in this scan.

Hair and skin

Within range

Collagen

Reference range: 4.533 - 6.179 | **Value:** 4.806

What this means: Reflects integrity of collagen matrix within the hair and skin, influencing structural stability and functional resilience.

Why it matters: Collagen is the primary structural protein underpinning tissue integrity and repair across the body.

Functional significance: Linked to vitamin C, glycine, proline, copper, oxidative stress, and glycation damage.

Common patterns: Oxidative stress + glycation -> collagen breakdown in hair and skin leading to dysfunction.

Your result in this scan: Hair and skin appears within the analyser's expected range in this scan.

Endocrine system

Within range

Collagen

Reference range: 6.178 - 8.651 | **Value:** 8.253

What this means: Reflects integrity of collagen matrix within the endocrine system, influencing structural stability and functional resilience.

Why it matters: Collagen is the primary structural protein underpinning tissue integrity and repair across the body.

Functional significance: Linked to vitamin C, glycine, proline, copper, oxidative stress, and glycation damage.

Common patterns: Oxidative stress + glycation -> collagen breakdown in endocrine system leading to dysfunction.

Your result in this scan: Endocrine system appears within the analyser's expected range in this scan.

Circulatory system

Within range

Collagen

Reference range: 3.586 - 4.337 | **Value:** 4.153

What this means: Reflects integrity of collagen matrix within the circulatory system, influencing structural stability and functional resilience.

Why it matters: Collagen is the primary structural protein underpinning tissue integrity and repair across the body.

Functional significance: Linked to vitamin C, glycine, proline, copper, oxidative stress, and glycation damage.

Common patterns: Oxidative stress + glycation -> collagen breakdown in circulatory system leading to dysfunction.

Your result in this scan: Circulatory system appears within the analyser's expected range in this scan.

Digestive system

Mildly reduced

Collagen

Reference range: 3.492 - 4.723 | **Value:** 3.168

What this means: Reflects integrity of collagen matrix within the digestive system, influencing structural stability and functional resilience.

Why it matters: Collagen is the primary structural protein underpinning tissue integrity and repair across the body.

Functional significance: Linked to vitamin C, glycine, proline, copper, oxidative stress, and glycation damage.

Common patterns: Oxidative stress + glycation -> collagen breakdown in digestive system leading to dysfunction.

Your result in this scan: Suggests weakened connective tissue, poor repair, or structural fragility.

Immune system

Mildly reduced

Collagen

Reference range: 3.376 - 4.582 | **Value:** 2.938

What this means: Reflects integrity of collagen matrix within the immune system, influencing structural stability and functional resilience.

Why it matters: Collagen is the primary structural protein underpinning tissue integrity and repair across the body.

Functional significance: Linked to vitamin C, glycine, proline, copper, oxidative stress, and glycation damage.

Common patterns: Oxidative stress + glycation -> collagen breakdown in immune system leading to dysfunction.

Your result in this scan: Suggests weakened connective tissue, poor repair, or structural fragility.

Motion systems

Mildly reduced

Collagen

Reference range: 6.458 - 8.133 | **Value:** 5.573

What this means: Reflects integrity of collagen matrix within the motion systems, influencing structural stability and functional resilience.

Why it matters: Collagen is the primary structural protein underpinning tissue integrity and repair across the body.

Functional significance: Linked to vitamin C, glycine, proline, copper, oxidative stress, and glycation damage.

Common patterns: Oxidative stress + glycation -> collagen breakdown in motion systems leading to dysfunction.

Your result in this scan: Suggests weakened connective tissue, poor repair, or structural fragility.

Muscle Tissue

Mildly reduced

Collagen

Reference range: 6.552 - 8.268 | **Value:** 5.774

What this means: Reflects integrity of collagen matrix within the muscle tissue, influencing structural stability and functional resilience.

Why it matters: Collagen is the primary structural protein underpinning tissue integrity and repair across the body.

Functional significance: Linked to vitamin C, glycine, proline, copper, oxidative stress, and glycation damage.

Common patterns: Oxidative stress + glycation -> collagen breakdown in muscle tissue leading to dysfunction.

Your result in this scan: Suggests weakened connective tissue, poor repair, or structural fragility.

Fat Metabolism

Mildly reduced

Collagen

Reference range: 6.338 - 8.368 | **Value:** 4.551

What this means: Reflects integrity of collagen matrix within the fat metabolism, influencing structural stability and functional resilience.

Why it matters: Collagen is the primary structural protein underpinning tissue integrity and repair across the body.

Functional significance: Linked to vitamin C, glycine, proline, copper, oxidative stress, and glycation damage.

Common patterns: Oxidative stress + glycation -> collagen breakdown in fat metabolism leading to dysfunction.

Your result in this scan: Suggests weakened connective tissue, poor repair, or structural fragility.

Detoxification and metabolism

Within range

Collagen

Reference range: 6.187 - 8.466 | **Value:** 6.519

What this means: Reflects integrity of collagen matrix within the detoxification and metabolism, influencing structural stability and functional resilience.

Why it matters: Collagen is the primary structural protein underpinning tissue integrity and repair across the body.

Functional significance: Linked to vitamin C, glycine, proline, copper, oxidative stress, and glycation damage.

Common patterns: Oxidative stress + glycation -> collagen breakdown in detoxification and metabolism leading to dysfunction.

Your result in this scan: Detoxification and metabolism appears within the analyser's expected range in this scan.

Reproductive system

Within range

Collagen

Reference range: 3.778 - 4.985 | **Value:** 3.997

What this means: Reflects integrity of collagen matrix within the reproductive system, influencing structural stability and functional resilience.

Why it matters: Collagen is the primary structural protein underpinning tissue integrity and repair across the body.

Functional significance: Linked to vitamin C, glycine, proline, copper, oxidative stress, and glycation damage.

Common patterns: Oxidative stress + glycation -> collagen breakdown in reproductive system leading to dysfunction.

Your result in this scan: Reproductive system appears within the analyser's expected range in this scan.

Nervous system

Mildly reduced

Collagen

Reference range: 3.357 - 4.239 | **Value:** 2.643

What this means: Reflects integrity of collagen matrix within the nervous system, influencing structural stability and functional resilience.

Why it matters: Collagen is the primary structural protein underpinning tissue integrity and repair across the body.

Functional significance: Linked to vitamin C, glycine, proline, copper, oxidative stress, and glycation damage.

Common patterns: Oxidative stress + glycation -> collagen breakdown in nervous system leading to dysfunction.

Your result in this scan: Suggests weakened connective tissue, poor repair, or structural fragility.

Skeleton

Within range

Collagen

Reference range: 6.256 - 8.682 | **Value:** 7.975

What this means: Reflects integrity of collagen matrix within the skeleton, influencing structural stability and functional resilience.

Why it matters: Collagen is the primary structural protein underpinning tissue integrity and repair across the body.

Functional significance: Linked to vitamin C, glycine, proline, copper, oxidative stress, and glycation damage.

Common patterns: Oxidative stress + glycation -> collagen breakdown in skeleton leading to dysfunction.

Your result in this scan: Skeleton appears within the analyser's expected range in this scan.

Channels and collaterals

High priority

Complete system marker table — all markers and all severity grades are shown below.

Flagged markers: 4 | Within range: 12

Hand Tai Yin Lung Meridian

Within range

Channels and collaterals

Reference range: 48.264 - 65.371 | **Value:** 63.778

What this means: Reflects the functional state of the Lung meridian within the channels and collaterals model, associated with respiratory vitality, boundary regulation, grief-stress patterns, and surface defensive function.

Why it matters: In an East Asian functional framework, the Lung channel is linked to respiration, skin and boundary integrity, immune-defensive responsiveness, and the body's capacity to distribute protective energy to the exterior.

Functional significance: Bridges respiratory function, skin barrier resilience, grief-related autonomic patterns, mucosal immunity, and circulation of protective regulatory tone through the upper body.

Common patterns: Respiratory weakness plus skin-barrier and allergy tendencies often cluster with Lung-channel imbalance; unresolved grief, shallow breathing, and recurrent mucosal irritation may sustain the pattern.

Your result in this scan: Hand Tai Yin Lung Meridian appears within the analyser's expected range in this scan.

Hand Yangming Large Intestine Meridian

Within range

Channels and collaterals

Reference range: 56.749 - 67.522 | **Value:** 61.756

What this means: Reflects the functional state of the Large Intestine meridian, associated with elimination, release, mucosal boundary handling, and paired regulation with the Lung channel.

Why it matters: This channel is linked to bowel clearance, the ability to let go of waste burden, and the relationship between mucosal detoxification and external defensive health.

Functional significance: Connects bowel transit, mucosal inflammation, skin expression, paired Lung-Large Intestine dynamics, and detoxification load through the elimination pathway.

Common patterns: Constipation, skin reactivity, and respiratory-allergy patterns often coexist with Large Intestine meridian imbalance; poor clearance increases toxic and inflammatory recirculation.

Your result in this scan: Hand Yangming Large Intestine Meridian appears within the analyser's expected range in this scan.

Foot Yangming Stomach Meridian

Within range

Channels and collaterals

Reference range: 0.481 - 1.043 | **Value:** 0.485

What this means: Reflects the functional state of the Stomach meridian, associated with digestive receiving, nourishment processing, appetite regulation, and tissue support derived from food transformation.

Why it matters: Within this framework, the Stomach channel governs how nourishment is received and processed, influencing energy, muscle tissue, and the body's tolerance to dietary load.

Functional significance: Links appetite, gastric function, damp-heat or irritation patterns, upper digestive congestion, and the body's capacity to derive stable nourishment from intake.

Common patterns: Reflux, bloating, appetite disturbance, and facial or oral inflammatory signs often cluster with Stomach meridian stress; chronic irritation can impair nutrient assimilation and energy quality.

Your result in this scan: Foot Yangming Stomach Meridian appears within the analyser's expected range in this scan.

Foot Tai Yin Lung Meridian

Within range

Channels and collaterals

Reference range: 0.327 - 0.937 | **Value:** 0.465

What this means: Reflects the functional state of the Spleen meridian in practical TCM usage despite the source label discrepancy, associated with transformation of food and fluid, tissue nourishment, and dampness handling.

Why it matters: This channel is central to digestive assimilation, fluid distribution, muscle tone, and the prevention of stagnating damp patterns that undermine energy and tissue resilience.

Functional significance: Links digestion, fluid metabolism, oedema tendency, fatigue after eating, bruising or tissue weakness, and the generation of usable post-digestive energy.

Common patterns: Bloating, loose stools, oedema, fatigue, and sugar craving often cluster with Spleen-type channel weakness; poor transformation can also worsen damp skin and hormonal patterns.

Your result in this scan: Foot Tai Yin Lung Meridian appears within the analyser's expected range in this scan.

Hand Shao Yin Heart Sutra

Mildly reduced

Channels and collaterals

Reference range: 1.672 - 1.978 | **Value:** 1.477

What this means: Reflects the functional state of the Heart channel, associated with circulation, emotional coherence, sleep quality, and the stability of mental-spiritual regulation in traditional pattern terms.

Why it matters: The Heart channel is linked to blood movement, calmness of mind, restful sleep, and the coherence of emotional and cognitive function under stress.

Functional significance: Bridges cardiac rhythm, sleep disruption, anxiety-agitation patterns, microcirculatory tone, and the interface between emotional load and physical regulation.

Common patterns: Sleep disruption, palpitations, anxiety, and mental restlessness often cluster with Heart-channel imbalance; inflammation and stress overdrive may deepen the pattern.

Your result in this scan: Suggests depletion of heart-blood or calming capacity, poor sleep quality, emotional fragility, or reduced circulatory vitality in this regulatory axis.

Hand the small intestine by the sun

Within range

Channels and collaterals

Reference range: 0.192 - 0.412 | **Value:** 0.199

What this means: Reflects the functional state of the Small Intestine channel, associated with separation of clear from turbid, discernment, and downstream handling of digestive and inflammatory burden.

Why it matters: This channel is traditionally linked to sorting and separating function, both physically in digestion and metaphorically in clarity, making it relevant to absorption quality and inflammatory spillover.

Functional significance: Connects small-bowel assimilation, discernment of usable nutrients versus waste, inflammatory gut burden, and paired Heart-Small Intestine regulation.

Common patterns: Malabsorption, bloating, inflammatory gut patterns, and cognitive irritability may cluster with Small Intestine channel imbalance; this can also affect the Heart-paired axis of calm versus agitation.

Your result in this scan: Hand the small intestine by the sun appears within the analyser's expected range in this scan.

Foot Tai Yang Bladder Meridian

Mildly reduced

Channels and collaterals

Reference range: 4.832 - 5.147 | **Value:** 3.773

What this means: Reflects the functional state of the Bladder meridian, associated with fluid regulation, posterior chain tension, nervous-system vigilance, and stress storage along the back body.

Why it matters: The Bladder channel is often linked to urinary function, autonomic vigilance, musculoskeletal tension, and the body's tendency to hold stress through the back and sympathetic system.

Functional significance: Connects urinary balance, back tension, fear-stress patterns, sleep disruption, and paired Kidney-Bladder regulation in fluid and autonomic control.

Common patterns: Back tension, urinary irritation, poor stress recovery, and sleep disruption often coexist with Bladder-channel imbalance; sympathetic dominance can maintain the pattern.

Your result in this scan: Suggests depleted fluid-regulatory resilience, lower stress tolerance, posterior-chain weakness, or reduced channel vitality in elimination and autonomic support.

Foot Shao Yin Kidney

Within range

Channels and collaterals

Reference range: 3.321 - 4.244 | **Value:** 3.937

What this means: Reflects the functional state of the Kidney channel, associated with foundational reserves, endocrine-adrenal resilience, reproductive vitality, bone support, and fear-stress regulation in traditional pattern terms.

Why it matters: The Kidney channel is considered central to constitutional reserve, growth and repair, fertility, stress endurance, and the deeper energetic support of brain, bones, and hormones.

Functional significance: Links adrenal-endocrine resilience, bone density, reproductive function, fluid metabolism, fatigue patterns, and the ability to recover from chronic depletion.

Common patterns: Chronic stress, poor sleep, hormonal decline, fertility issues, and bone weakness often cluster with Kidney-channel depletion; adrenal burden may further destabilise the pattern.

Your result in this scan: Foot Shao Yin Kidney appears within the analyser's expected range in this scan.

Hand Jue Yin Pericardium Meridian

Within range

Channels and collaterals

Reference range: 1.338 - 1.672 | **Value:** 1.34

What this means: Reflects the functional state of the Pericardium channel, associated with emotional buffering, chest tension, circulation, and protection against stress-induced internal agitation.

Why it matters: This channel is traditionally linked to protecting the Heart from emotional overload, making it relevant to chest tightness, anxiety, circulation, and relational stress patterns.

Functional significance: Connects mood regulation, chest pressure, circulation, sleep, and the buffering of emotional stress within the cardiovascular-nervous interface.

Common patterns: Stress, chest tightness, palpitatory sensations, and emotional overload often cluster with Pericardium-channel imbalance; sleep and mood may deteriorate alongside the pattern.

Your result in this scan: Hand Jue Yin Pericardium Meridian appears within the analyser's expected range in this scan.

Hand Shao Yang Triple Burner Meridian

Mildly reduced

Channels and collaterals

Reference range: 0.669 - 1.544 | **Value:** 0.548

What this means: Reflects the functional state of the Triple Burner channel, associated with coordination across upper, middle, and lower body regulatory compartments, especially fluid movement and stress adaptation.

Why it matters: This channel represents systemic communication and distribution, making it relevant to fluid movement, endocrine stress signalling, and the integration of multiple body regions.

Functional significance: Bridges adrenal-stress patterns, temperature regulation, fluid circulation, lymphatic movement, and whole-body coordination across compartments.

Common patterns: Fluid retention, endocrine stress, temperature instability, and lymphatic congestion may cluster with Triple Burner imbalance; poor cross-system coordination can sustain chronic complexity.

Your result in this scan: Suggests poor coordination of systemic regulation, sluggish fluid communication, lower adaptive responsiveness, or depleted integrative signalling across body regions.

Foot Shao Yang Gall Bladder Meridian

Within range

Channels and collaterals

Reference range: 1.554 - 1.988 | **Value:** 1.868

What this means: Reflects the functional state of the Gall Bladder channel, associated with bile flow, decisiveness, lateral tension patterns, and the handling of damp-heat or stagnant digestive stress.

Why it matters: In functional terms, this channel links digestive fat handling, liver-gallbladder flow, irritability, and the body's ability to process pressure without stagnation.

Functional significance: Connects bile secretion, lateral body tension, headaches, irritability, decision fatigue, and paired Liver-Gall Bladder dynamics.

Common patterns: Poor bile flow, headaches, irritability, and side-body tension often cluster with Gall Bladder meridian stress; liver congestion can intensify the pattern.

Your result in this scan: Foot Shao Yang Gall Bladder Meridian appears within the analyser's expected range in this scan.

Foot Jue Yin Liver Meridian

Within range

Channels and collaterals

Reference range: 1.553 - 2.187 | **Value:** 1.652

What this means: Reflects the functional state of the Liver channel, associated with smooth flow, emotional pressure regulation, blood storage, hormonal rhythm, and tension-stagnation patterns.

Why it matters: This channel is central to stress processing, menstrual-hormonal flow, tendon flexibility, mood stability, and the movement of energy without congestion.

Functional significance: Bridges liver detoxification, bile flow, menstrual and hormonal regulation, irritability, muscular tension, and blood-vascular adaptability.

Common patterns: Stress, PMS, headaches, muscular tension, and poor bile flow commonly cluster with Liver-meridian imbalance; detox congestion and hormonal recirculation may sustain it.

Your result in this scan: Foot Jue Yin Liver Meridian appears within the analyser's expected range in this scan.

Ren channel

Within range

Channels and collaterals

Reference range: 11.719 - 18.418 | **Value:** 12.572

What this means: Reflects the functional state of the Ren vessel, associated with front-body integration, reproductive and hormonal regulation, fluid-nutrient support, and yin-type restorative processes.

Why it matters: This channel is traditionally important for reproductive, hormonal, and restorative regulation, especially across the anterior body and pelvic-lower abdominal axis.

Functional significance: Connects fertility, menstrual regularity, pelvic nourishment, endocrine balance, and restorative reserve through front-body regulation.

Common patterns: Hormonal irregularity, fertility stress, lower-abdominal weakness, and pelvic congestion often cluster with Ren-channel imbalance; endocrine depletion can deepen the pattern.

Your result in this scan: Ren channel appears within the analyser's expected range in this scan.

Governor meridian

Within range

Channels and collaterals

Reference range: 0.708 - 1.942 | **Value:** 1.536

What this means: Reflects the functional state of the Governor vessel, associated with back-body integration, spinal support, mental drive, and yang-type activating regulation.

Why it matters: This channel is traditionally linked to motivation, structural support, spinal vitality, and the activation side of energy and nervous-system tone.

Functional significance: Connects spinal tension, mental drive, nervous-system activation, temperature support, and the capacity to mobilise energy through the back body.

Common patterns: Back tension, low motivation, poor temperature resilience, or mental overdrive can cluster with Governor-vessel imbalance; sleep and stress patterns often interact strongly.

Your result in this scan: Governor meridian appears within the analyser's expected range in this scan.

Vital meridian

Within range

Channels and collaterals

Reference range: 6.138 - 21.396 | **Value:** 6.331

What this means: Reflects generalised vital-channel reserve in the source framework, pointing to overall constitutional vitality, regulatory coherence, and resilience of the channel network.

Why it matters: A broad vitality meridian concept can be clinically useful as an overview of reserve, recovery capacity, and whole-system energetic coherence when interpreted alongside specific channels.

Functional significance: Bridges fatigue, constitutional resilience, multisystem recovery, stress tolerance, and the integration of channel-level regulation across the body.

Common patterns: Chronic stress, sleep loss, endocrine depletion, and multisystem burden can all lower vital-meridian resilience; broad fatigue patterns often align here.

Your result in this scan: Vital meridian appears within the analyser's expected range in this scan.

Tai mai

Mildly reduced

Channels and collaterals

Reference range: 5.733 - 7.109 | **Value:** 5.146

What this means: Reflects the functional state of the Dai Mai concept in traditional pattern language, associated with belt-like regulation around the waist, structural containment, and lower-body fluid or stagnation patterns.

Why it matters: This channel is often considered relevant where there is pelvic congestion, abdominal heaviness, lower-body stagnation, or poor containment of structural and fluid dynamics.

Functional significance: Connects waist tension, pelvic heaviness, lower-body circulation, gynaecological congestion, and the structural holding pattern of the core-lower trunk interface.

Common patterns: Pelvic congestion, abdominal bloating, lower-body fluid retention, and menstrual stagnation may cluster with Tai Mai imbalance; connective-tissue laxity or tension can modify the pattern.

Your result in this scan: Suggests weak containment, lower-body heaviness, pelvic instability, or reduced structural-integrative support through the waist and lower abdomen.

Pulse of heart and brain

Medium priority

Complete system marker table — all markers and all severity grades are shown below.

Flagged markers: 2 | Within range: 5

Stroke index

Within range

Pulse of heart and brain

Reference range: 60.735 - 65.396 | **Value:** 62.076

What this means: Reflects stroke volume relative to body size or circulatory demand, offering a scaled view of cardiac output efficiency.

Why it matters: This marker helps assess whether forward flow is adequate for the individual's physiological needs rather than considering raw output alone.

Functional significance: Linked to ventricular pumping performance, preload, afterload, body size, and tissue perfusion demand.

Common patterns: Low stroke index + high resistance -> reduced tissue delivery efficiency; fatigue and reduced exercise tolerance commonly align with this pattern.

Your result in this scan: Stroke index appears within the analyser's expected range in this scan.

Stroke volume(SV)

Mildly reduced

Pulse of heart and brain

Reference range: 63.012 - 67.892 | **Value:** 61.901

What this means: Reflects beat-by-beat blood ejection from the heart, representing the immediate mechanical contribution of each contraction to circulation.

Why it matters: Stroke volume is a core determinant of cardiac output and directly influences systemic perfusion, oxygen delivery, and exercise tolerance.

Functional significance: Linked to ventricular filling, myocardial contractility, afterload, fluid status, and oxygen-energetic support of the heart.

Common patterns: Reduced contractility or elevated afterload -> lower SV; low SV often coexists with fatigue, poor circulation, and reduced cardiovascular resilience.

Your result in this scan: Suggests reduced pumping effectiveness, lower filling or contractility, or resistance-related impairment of forward flow.

Heart peripheral resistance (TRR)

Mildly elevated

Pulse of heart and brain

Reference range: 0.983 - 1.265 | **Value:** 1.638

What this means: Reflects total peripheral resistance encountered by blood as it moves through the systemic vascular network.

Why it matters: Higher peripheral resistance increases cardiac strain, reduces downstream perfusion efficiency, and often reflects vascular tone imbalance or endothelial stress.

Functional significance: Linked to arterial stiffness, autonomic tone, endothelial health, inflammation, and blood viscosity.

Common patterns: Sympathetic overdrive + endothelial dysfunction + thickened blood -> raised peripheral resistance and greater cardiac workload.

Your result in this scan: Indicates increased resistance within the vascular bed, often associated with vasoconstriction, arterial rigidity, stress overdrive, or inflammatory circulation burden.

Pulse wave coefficient K

Within range

Pulse of heart and brain

Reference range: 0.316 - 0.401 | **Value:** 0.391

What this means: Reflects properties of the pulse-wave transmission through arteries, providing insight into vascular elasticity, compliance, and wave reflection dynamics.

Why it matters: Pulse-wave characteristics help indicate whether vessels are supple and adaptive or whether stiffness and pressure-wave reflection are increasing cardiovascular strain.

Functional significance: Linked to arterial elasticity, vascular ageing, endothelial tone, blood pressure behaviour, and the efficiency of pulse transmission through the arterial tree.

Common patterns: Arterial stiffness + vascular ageing + pressure stress -> altered pulse-wave coefficient; this often aligns with endothelial dysfunction and reduced perfusion quality.

Your result in this scan: Pulse wave coefficient K appears within the analyser's expected range in this scan.

Cerebrovascular blood oxygen saturation(Sa)

Within range

Pulse of heart and brain

Reference range: 0.710 - 1.109 | **Value:** 0.894

What this means: Reflects the saturation of oxygen carried in blood supplying the cerebrovascular system, indicating how effectively oxygen is available for brain delivery.

Why it matters: The brain has a high oxygen demand, so reduced saturation can impair cognition, concentration, autonomic stability, and resilience under stress.

Functional significance: Linked to pulmonary oxygenation, haemoglobin function, blood-flow quality, vascular delivery, and neural metabolic demand.

Common patterns: Reduced oxygen saturation + poor cerebral perfusion -> brain fog, headaches, and fatigue; lung function and blood quality are key upstream drivers.

Your result in this scan: Cerebrovascular blood oxygen saturation(Sa) appears within the analyser's expected range in this scan.

Cerebrovascular blood oxygen volume(CaCO2)

Within range

Pulse of heart and brain

Reference range: 7.880 - 10.090 | **Value:** 8.081

What this means: Reflects the oxygen content available within blood supplying the cerebrovascular system, indicating total oxygen-carrying capacity rather than saturation alone.

Why it matters: Even with normal saturation, low oxygen content can impair cerebral metabolism if haemoglobin or circulatory support is inadequate.

Functional significance: Linked to haemoglobin status, oxygen transport capacity, pulmonary function, perfusion quality, and the metabolic needs of brain tissue.

Common patterns: Low haemoglobin or reduced oxygen transport + poor cerebral perfusion -> diminished oxygen volume to brain tissues and reduced cognitive stamina.

Your result in this scan: Cerebrovascular blood oxygen volume(CaCO2) appears within the analyser's expected range in this scan.

Cerebrovascular blood oxygen presure(PaO2)

Within range

Pulse of heart and brain

Reference range: 5.017 - 5.597 | **Value:** 5.169

What this means: Reflects the partial pressure of oxygen in blood supplying the cerebrovascular system, indicating oxygen availability for diffusion into brain tissue.

Why it matters: Oxygen pressure influences how readily oxygen can move into tissues; reduced PaO2 can compromise neural metabolism even when other circulation markers appear acceptable.

Functional significance: Linked to lung gas exchange, blood oxygenation, vascular delivery, microcirculation, and tissue oxygen diffusion gradients.

Common patterns: Poor lung function + reduced oxygenation + impaired cerebral perfusion -> lower PaO2 support to the brain; this may worsen fatigue and cognitive slowing.

Your result in this scan: Cerebrovascular blood oxygen presure(PaO2) appears within the analyser's expected range in this scan.

Blood lipids

High priority

Complete system marker table — all markers and all severity grades are shown below.

Flagged markers: 5 | Within range: 2

Blood viscosity

Moderately elevated

Blood lipids

Reference range: 4.131 - 4.562 | **Value:** 5.233

What this means: Reflects thickness of blood affecting circulation and oxygen delivery.

Why it matters: High viscosity impairs circulation and increases cardiovascular risk.

Functional significance: Linked to hydration, lipids, inflammation.

Common patterns: High viscosity increases cardiovascular burden.

Your result in this scan: Indicates sluggish circulation and vascular strain.

Total cholesterol(TC)

Within range

Blood lipids

Reference range: 1.833 - 2.979 | **Value:** 2.75

What this means: Reflects total circulating cholesterol burden across lipoprotein fractions, giving a broad view of cholesterol availability, transport, and potential accumulation.

Why it matters: Cholesterol is required for steroid hormone synthesis, membrane integrity, bile acid production, and repair, but excessive or poorly handled cholesterol can increase vascular and inflammatory strain.

Functional significance: Interacts with liver cholesterol synthesis and clearance, thyroid status, bile flow, oxidative stress, and the balance between protective and atherogenic lipoprotein patterns.

Common patterns: Low thyroid or poor bile flow can increase cholesterol retention; oxidative stress and inflammation determine whether higher cholesterol becomes more damaging to vessels and tissues.

Your result in this scan: Total cholesterol(TC) appears within the analyser's expected range in this scan.

Triglyceride(TG)

Mildly elevated

Blood lipids

Reference range: 1.116 - 2.101 | **Value:** 2.218

What this means: Reflects circulating triglyceride levels as a marker of hepatic lipid export, carbohydrate handling, and the balance between fat storage and oxidation.

Why it matters: Triglycerides often rise with insulin resistance, excess hepatic lipogenesis, and poor lipid clearance, making them a strong functional marker of cardiometabolic strain.

Functional significance: Closely linked to insulin status, liver fat burden, mitochondrial beta oxidation, thyroid rate, and the efficiency of lipoprotein processing and clearance.

Common patterns: High carbohydrate load plus hyperinsulinaemia drives triglyceride synthesis and hepatic export; low thyroid and mitochondrial inefficiency can worsen persistence of elevated triglycerides.

Your result in this scan: Indicates excess triglyceride production or retention, often associated with insulin resistance, high glycaemic load, fatty liver, or poor fat oxidation.

High-density lipoprotein (HDL-C)

Mildly elevated

Blood lipids

Reference range: 1.449 - 2.246 | **Value:** 2.955

What this means: Reflects high-density lipoprotein status involved in reverse cholesterol transport, antioxidant protection, and aspects of vascular repair signalling.

Why it matters: HDL participates in returning excess cholesterol to the liver and is often associated with healthier lipid trafficking and lower oxidative vascular stress when functionally robust.

Functional significance: Interacts with liver function, triglyceride balance, exercise status, inflammatory tone, and the overall quality of lipoprotein metabolism rather than quantity alone.

Common patterns: Low HDL commonly tracks with hypertriglyceridaemia, insulin resistance, fatty liver, and inflammatory burden; improving metabolic flexibility often improves HDL-related patterns.

Your result in this scan: Often reflects stronger lipid transport protection, though quality and function still matter alongside total metabolic and inflammatory context.

Low-density lipoprotein (LDL-C)

Within range

Blood lipids

Reference range: 0.831 - 1.588 | **Value:** 1.015

What this means: Reflects low-density lipoprotein cholesterol burden, indicating transport of cholesterol to peripheral tissues and potential exposure to oxidative modification.

Why it matters: LDL carries essential cholesterol for cell use, but when elevated or oxidatively damaged it can contribute to endothelial stress, vascular inflammation, and plaque-related risk.

Functional significance: Depends on liver receptor-mediated clearance, thyroid activity, dietary-metabolic load, oxidative stress, and the inflammatory environment affecting particle behaviour.

Common patterns: Low thyroid, poor hepatic clearance, and oxidative stress can worsen LDL-related vascular burden; inflammation makes LDL more damaging than quantity alone suggests.

Your result in this scan: Low-density lipoprotein (LDL-C) appears within the analyser's expected range in this scan.

Neutral fat(MB)

Mildly elevated

Blood lipids

Reference range: 0.726 - 1.281 | **Value:** 3.096

What this means: Reflects neutral fat burden within circulation, generally corresponding to stored or transported triglyceride-rich lipid material.

Why it matters: Higher neutral fat tends to signal inefficient fat handling, excess storage drive, insulin resistance, and a greater burden on vascular and hepatic lipid processing.

Functional significance: Closely linked to triglyceride metabolism, hepatic fat export, insulin signalling, adipose storage patterns, and mitochondrial utilisation of fats as fuel.

Common patterns: Neutral fat often rises alongside high triglycerides, hyperinsulinaemia, and liver congestion; persistent elevation can track with obesity and cardiometabolic strain.

Your result in this scan: Indicates elevated lipid storage pressure, altered transport of triglyceride-rich particles, or broader metabolic inefficiency in fat utilisation and clearance.

Circulating immune complex(CIC)

Mildly elevated

Blood lipids

Reference range: 13.012 - 17.291 | **Value:** 18.601

What this means: Reflects immune complex burden indicating antigen-antibody interaction.

Why it matters: Elevated CIC indicates immune activation and possible autoimmune burden.

Functional significance: Linked to immune clearance, liver function, and inflammation.

Common patterns: High CIC linked to chronic inflammation or autoimmunity.

Your result in this scan: Indicates immune activation or poor clearance.

Prostate

High priority

Complete system marker table — all markers and all severity grades are shown below.

Flagged markers: 3 | Within range: 0

Degree of Prostatic Hyperplasia

Moderately elevated

Prostate

Reference range: 1.023 - 3.230 | **Value:** 5.03

What this means: Reflects the extent of enlargement or proliferative change in prostate tissue, suggesting altered growth signalling and reduced ease of urinary outflow.

Why it matters: Prostatic enlargement matters because it can impair urinary flow, increase nocturia, create incomplete bladder emptying, and signal longer-term androgen-inflammatory dysregulation.

Functional significance: Linked to age-related androgen balance, DHT signalling, inflammatory burden, pelvic congestion, insulin-metabolic factors, and local tissue-remodelling tendencies.

Common patterns: Ageing + inflammatory load + androgen-metabolic signalling can drive prostatic hyperplasia; pelvic congestion and poor circulation may worsen urinary symptoms.

Your result in this scan: Indicates increased prostatic enlargement, tissue proliferation, or growth-related urinary obstruction risk.

Degree of Prostatic Calcification

Mildly elevated

Prostate

Reference range: 1.471 - 6.079 | **Value:** 8.907

What this means: Reflects calcific change within prostate tissue, suggesting chronic inflammation, stagnation, tissue-remodelling burden, or longer-standing glandular congestion.

Why it matters: Prostatic calcification can point to chronic irritative history, unresolved inflammation, pelvic stagnation, and altered prostate tissue quality affecting comfort and function.

Functional significance: Linked to chronic prostatitis patterns, pelvic congestion, mineral deposition in inflamed tissue, impaired drainage, and long-term glandular remodelling.

Common patterns: Chronic inflammation + pelvic stagnation + impaired local drainage can promote calcific change; this may coexist with prostatitis symptoms or hyperplastic patterns.

Your result in this scan: Indicates increased calcification burden, chronic inflammatory history, tissue stagnation, or altered glandular architecture.

Prostatitis Syndrome

Moderately elevated

Prostate

Reference range: 2.213 - 2.717 | **Value:** 6.11

What this means: Reflects an inflammatory-irritative pattern involving the prostate, suggesting pain, congestion, urinary discomfort, or immune-reactive stress in the gland and surrounding pelvic tissues.

Why it matters: Prostatitis-type patterns matter because they can affect urinary ease, pelvic comfort, sexual function, quality of life, and the chronic inflammatory burden of the male pelvis.

Functional significance: Linked to glandular inflammation, pelvic-floor tension, microbial or sterile inflammatory triggers, urinary irritation, congestion, and autonomic stress patterns.

Common patterns: Pelvic tension + inflammation + poor drainage can maintain prostatitis-type patterns even without clear infection; stress and autonomic guarding often amplify symptoms.

Your result in this scan: Indicates increased prostatic inflammation, pelvic discomfort, urinary irritation, congestion, or chronic pain-type burden in the male pelvic axis.

Male Sexual Function

Medium priority

Complete system marker table — all markers and all severity grades are shown below.

Flagged markers: 3 | Within range: 0

Testosterone

Moderately reduced

Male Sexual Function

Reference range: 3.342 - 9.461 | **Value:** 2.492

What this means: Reflects testosterone status influencing energy, libido, muscle, and metabolic health.

Why it matters: Key hormone for vitality, strength, and reproductive function.

Functional significance: Linked to adrenal, metabolic, and reproductive systems.

Common patterns: Metabolic dysfunction affects testosterone.

Your result in this scan: Fatigue, low libido.

Gonadotropin

Mildly reduced

Male Sexual Function

Reference range: 4.111 - 18.741 | **Value:** 3.558

What this means: Reflects general gonadotropin activity controlling reproductive hormones.

Why it matters: Central to fertility and endocrine signalling.

Functional significance: Linked to pituitary-gonadal axis.

Common patterns: Pituitary imbalance -> reproductive issues.

Your result in this scan: Low reproductive drive.

Erection Transmitter

Mildly reduced

Male Sexual Function

Reference range: 3.241 - 9.814 | **Value:** 3.03

What this means: Reflects neurotransmitter activity related to erectile function.

Why it matters: Indicates vascular and neurological integrity in sexual function.

Functional significance: Linked to nitric oxide, circulation, and nervous system.

Common patterns: Circulation + NO pathways -> erection.

Your result in this scan: Reduced erectile signalling.

Sperm and semen

Within range

Complete system marker table — all markers and all severity grades are shown below.

Flagged markers: 0 | Within range: 4

Semen volume

Within range

Sperm and semen

Reference range: 1.502 - 6.028 | **Value:** 5.263

What this means: Reflects semen production and reproductive capacity.

Why it matters: Important for fertility and reproductive health.

Functional significance: Linked to prostate, hydration, and hormone status.

Common patterns: Hydration + hormones affect semen.

Your result in this scan: Semen volume appears within the analyser's expected range in this scan.

Semen liquefying time

Within range

Sperm and semen

Reference range: 10.283 - 30.282 | **Value:** 26.982

What this means: Reflects the time required for semen to liquefy after ejaculation, indicating seminal enzymatic activity and the functional quality of seminal fluid.

Why it matters: Normal liquefaction is important because excessively delayed liquefaction can impair sperm release and motility, affecting fertility potential.

Functional significance: Linked to prostatic secretions, seminal enzyme activity, inflammatory status, mineral balance, and the quality of accessory-gland contributions to semen.

Common patterns: Prostatic inflammation + altered seminal enzymes + mineral imbalance can prolong liquefaction time and reduce effective sperm mobility.

Your result in this scan: Semen liquefying time appears within the analyser's expected range in this scan.

Number of sperms

Within range

Sperm and semen

Reference range: 2.483 - 3.932 | **Value:** 3

What this means: Reflects sperm count within semen, indicating the output capacity of spermatogenesis and the overall quantitative component of male fertility potential.

Why it matters: Sperm number matters because adequate count is one of the core determinants of conception potential, although quality and motility remain equally important.

Functional significance: Linked to testicular function, gonadotropin support, testosterone balance, heat exposure, oxidative stress, nutrient status, toxic burden, and spermatogenic efficiency.

Common patterns: Heat stress + oxidative burden + toxin exposure + endocrine imbalance can all reduce sperm count; sleep, metabolic health, and testicular environment strongly influence output.

Your result in this scan: Number of sperms appears within the analyser's expected range in this scan.

Sperm motility rate

Within range

Sperm and semen

Reference range: 0.637 - 0.877 | **Value:** 0.689

What this means: Reflects the proportion of sperm demonstrating effective movement, indicating their functional ability to travel and support fertilisation potential.

Why it matters: Motility is a critical fertility determinant because sperm must move efficiently through the reproductive tract to reach the ovum.

Functional significance: Linked to mitochondrial energy production, membrane integrity, oxidative stress, seminal fluid quality, mineral status, and inflammatory burden in the male reproductive tract.

Common patterns: Oxidative stress + mitochondrial dysfunction + poor membrane lipid quality can all reduce motility; inflammation and heat exposure may worsen the decline.

Your result in this scan: Sperm motility rate appears within the analyser's expected range in this scan.

Male Hormone

Medium priority

Complete system marker table — all markers and all severity grades are shown below.

Flagged markers: 2 | Within range: 4

Androgen

Within range

Male Hormone

Reference range: 1.372 - 1.878 | **Value:** 1.492

What this means: Reflects androgen hormone activity influencing muscle mass, libido, energy, and metabolic function.

Why it matters: Androgens regulate strength, drive, metabolic rate, and reproductive function.

Functional significance: Linked to adrenal function, insulin sensitivity, and reproductive axis.

Common patterns: Insulin resistance can elevate androgen; stress can suppress it.

Your result in this scan: Androgen appears within the analyser's expected range in this scan.

Estrogen

Within range

Male Hormone

Reference range: 0.521 - 1.036 | **Value:** 0.672

What this means: Reflects oestrogen status within reproductive-endocrine signalling, including production, peripheral conversion, receptor activity, hepatic conjugation, and enterohepatic recirculation.

Why it matters: Oestrogen influences menstrual physiology, adipose distribution, bone turnover, mood, vascular tone, thyroid-binding dynamics, and inflammatory signalling.

Functional significance: Closely linked to hepatic detoxification, microbiome beta-glucuronidase activity, ovarian/adipose production, progesterone balance, and insulin-metabolic status.

Common patterns: Poor liver conjugation + gut dysbiosis → oestrogen recirculation; insulin resistance and excess adiposity → higher aromatisation; chronic constipation can further increase reabsorption of hormone metabolites.

Your result in this scan: Estrogen appears within the analyser's expected range in this scan.

Progesterone(P)

Mildly reduced

Male Hormone

Reference range: 0.686 - 1.432 | **Value:** 0.651

What this means: Reflects progesterone levels important for reproductive balance and nervous system calming.

Why it matters: Progesterone balances oestrogen and supports fertility and mood stability.

Functional significance: Linked to ovulation, adrenal function, and GABA pathways.

Common patterns: Low progesterone + high oestrogen -> PMS.

Your result in this scan: Oestrogen dominance, anxiety, PMS.

Luteinizing Hormone (LH)

Mildly reduced

Male Hormone

Reference range: 0.679 - 1.324 | **Value:** 0.593

What this means: Reflects LH signalling involved in ovulation and testosterone production.

Why it matters: Critical for reproductive timing and endocrine balance.

Functional significance: Linked to pituitary signalling and gonadal function.

Common patterns: LH imbalance -> ovulatory dysfunction.

Your result in this scan: Reduced reproductive signalling.

Prolactin(PRL)

Within range

Male Hormone

Reference range: 0.738 - 1.742 | **Value:** 1.386

What this means: Reflects prolactin hormone involved in lactation and reproductive regulation.

Why it matters: Elevated prolactin can suppress fertility and alter cycles.

Functional significance: Linked to stress, pituitary function, and dopamine balance.

Common patterns: Stress -> elevated prolactin -> cycle disruption.

Your result in this scan: Prolactin(PRL) appears within the analyser's expected range in this scan.

Follicle Stimulating Hormone(FSH)

Within range

Male Hormone

Reference range: 0.387 - 0.956 | **Value:** 0.54

What this means: Reflects FSH levels involved in follicle development and reproductive function.

Why it matters: Essential for fertility and ovarian/testicular function.

Functional significance: Linked to pituitary and gonadal axis.

Common patterns: High FSH -> reduced ovarian reserve.

Your result in this scan: Follicle Stimulating Hormone(FSH) appears within the analyser's expected range in this scan.

Human Immunity

High priority

Complete system marker table — all markers and all severity grades are shown below.

Flagged markers: 4 | Within range: 1

Tissue barrier immune index

Mildly elevated

Human Immunity

Reference range: 7.435 - 8.847 | **Value:** 9.626

What this means: Reflects immune competence at tissue barrier interfaces, including epithelial surveillance, local immune tolerance, and defence against antigen penetration across skin and mucosal surfaces.

Why it matters: Barrier immunity determines how effectively the body prevents inappropriate antigen entry while maintaining tolerance to normal environmental exposures.

Functional significance: Closely linked to epithelial integrity, tight junction regulation, microbiome balance, histamine response, and inflammatory repair signalling.

Common patterns: Compromised skin/gut/airway barriers → higher antigen entry → stronger innate and adaptive immune activation; chronic barrier stress can sustain allergy, eczema, gut symptoms, and inflammatory overload.

Your result in this scan: Indicates heightened barrier immune activation, ongoing irritant exposure, inflammatory repair demand, or excessive reactivity to environmental or dietary antigens.

Innate immune cell index

Mildly elevated

Human Immunity

Reference range: 7.424 - 9.052 | **Value:** 10.784

What this means: Reflects functional activity of innate immune cells involved in first-response defence, including phagocytic surveillance, early inflammatory signalling, and pathogen recognition.

Why it matters: Innate immunity provides rapid protection against microbial challenge and shapes the intensity and direction of downstream adaptive immune responses.

Functional significance: Linked to inflammatory cytokine tone, nutrient status, microbial exposure, oxidative balance, and resolution capacity after acute immune activation.

Common patterns: Gut endotoxin translocation or recurrent infection → persistent innate activation → cytokine burden → fatigue, brain fog, and wider inflammatory spillover into endocrine and metabolic systems.

Your result in this scan: Indicates active innate immune stimulation, inflammatory signalling, endotoxin exposure, infection burden, or unresolved first-response activation.

Innate immune molecule index

Within range

Human Immunity

Reference range: 7.143 - 9.214 | **Value:** 7.585

What this means: Reflects soluble innate immune mediators involved in early defence, inflammatory communication, and coordination of antimicrobial responses.

Why it matters: These molecules shape the speed, intensity, and resolution of early immune events and strongly influence whether inflammation becomes protective or excessive.

Functional significance: Connected to cytokine signalling, complement-type responses, barrier challenge, endotoxin exposure, oxidative stress, and tissue repair signalling.

Common patterns: Persistent microbial products or tissue damage → sustained innate mediator release → chronic inflammatory tone; unresolved activation can impair recovery, sleep quality, and metabolic resilience.

Your result in this scan: Innate immune molecule index appears within the analyser's expected range in this scan.

Cellular immune index

Mildly elevated

Human Immunity

Reference range: 5.769 - 7.643 | **Value:** 8.547

What this means: Reflects cell-mediated immune activity, particularly T-cell-driven defence, immune coordination, and targeted response to infected or abnormal cells.

Why it matters: Cellular immunity is essential for antiviral defence, immune surveillance, and maintaining balance between effective immune response and excess tissue-reactive inflammation.

Functional significance: Depends on thymic support, micronutrient sufficiency, antigen exposure patterns, cortisol tone, and healthy communication between innate and adaptive immunity.

Common patterns: High cortisol or chronic nutrient depletion → weaker cellular response; persistent viral or antigenic challenge → increased cell-mediated activation and inflammatory strain.

Your result in this scan: Indicates active cell-mediated immune stimulation, chronic antigenic challenge, autoimmune-type drive, or heightened inflammatory reactivity.

Humoral immunity index

Mildly elevated

Human Immunity

Reference range: 6.424 - 8.219 | **Value:** 8.708

What this means: Reflects antibody-mediated immune activity and the functional strength of humoral defence across antigen recognition, immune memory, and adaptive immune responsiveness.

Why it matters: Humoral immunity is central to the body's ability to neutralise pathogens, maintain mucosal defence, and manage repeated antigen exposure without excessive chronic reactivity.

Functional significance: Interacts with immunoglobulin production, mucosal barrier integrity, protein status, gut antigen load, and the balance between immune defence and hypersensitivity.

Common patterns: High chronic antigen exposure from gut or respiratory barriers can push humoral activation; persistent stimulation may contribute to allergy, food reactivity, and circulating immune-complex burden.

Your result in this scan: Indicates stronger antibody-driven immune activity, which may reflect robust defence or excessive reactivity depending on antigen load and inflammatory context.

Human Consciousness Level

High priority

Complete system marker table — all markers and all severity grades are shown below.

Flagged markers: 6 | Within range: 11

Shame

Within range

Human Consciousness Level

Reference range: 2.214 - 3.925 | **Value:** 3.056

What this means: Reflects a low-resource self-evaluative emotional state characterised by contraction, withdrawal, and reduced psychological resilience within the source framework.

Why it matters: Persistent shame states can be associated with chronic stress activation, social withdrawal, reduced motivation, and impaired recovery capacity.

Functional significance: Best interpreted as a psycho-emotional pattern marker linked to autonomic tone, trauma load, stress physiology, and reduced self-regulatory capacity rather than a standalone diagnosis.

Common patterns: Shame -> withdrawal + low motivation + autonomic contraction; persistent stress, trauma history, and low self-worth patterns may sustain the state.

Your result in this scan: Shame appears within the analyser's expected range in this scan.

Guilt

Within range

Human Consciousness Level

Reference range: 2.154 - 3.478 | **Value:** 2.543

What this means: Reflects a self-critical emotional state centred on remorse, self-blame, or unresolved internal conflict within the source framework.

Why it matters: Persistent guilt can maintain stress activation, rumination, low mood, and impaired emotional recovery.

Functional significance: Linked to cognitive rumination, moral-emotional conflict, stress physiology, and reduced mental recovery rather than a discrete psychiatric construct.

Common patterns: Guilt -> rumination + stress activation + reduced sleep quality; chronic internal conflict can feed anxiety, low mood, and autonomic tension.

Your result in this scan: Guilt appears within the analyser's expected range in this scan.

Apathy

Mildly reduced

Human Consciousness Level

Reference range: 2.343 - 4.089 | **Value:** 2.263

What this means: Reflects reduced motivational drive, emotional engagement, and activation capacity within the source framework.

Why it matters: Apathy-like patterns can accompany burnout, depression, chronic fatigue, and nervous-system depletion, reducing follow-through and resilience.

Functional significance: Linked to low dopaminergic drive, energy depletion, chronic stress collapse, inflammatory fatigue, or reduced reward-system engagement.

Common patterns: Chronic fatigue + low dopamine + burnout -> apathy patterns; inflammatory load and low mitochondrial output may deepen disengagement.

Your result in this scan: Suggests lower prominence of motivational collapse or disengagement.

Grief

Within range

Human Consciousness Level

Reference range: 1.845 - 3.241 | **Value:** 1.997

What this means: Reflects an emotional-processing state associated with loss, sadness, and unresolved bereavement-related contraction within the source framework.

Why it matters: Grief can influence respiration, sleep, immunity, appetite, and autonomic tone, especially when unresolved or chronic.

Functional significance: Linked to Lung-channel symbolic patterns, stress physiology, emotional processing capacity, social attachment systems, and reduced recovery reserve.

Common patterns: Grief -> chest tightness + sleep disruption + reduced vitality; unresolved bereavement may sustain autonomic dysregulation and immune vulnerability.

Your result in this scan: Grief appears within the analyser's expected range in this scan.

Fear

Mildly reduced

Human Consciousness Level

Reference range: 2.155 - 4.031 | **Value:** 1.681

What this means: Reflects heightened threat perception and protective vigilance within the source framework, associated with autonomic activation and anticipatory stress.

Why it matters: Fear states can amplify sympathetic drive, worsen sleep, tighten muscles, disrupt digestion, and reduce reflective decision-making.

Functional significance: Linked to HPA-axis activation, amygdala-driven vigilance, adrenal stress, trauma load, and reduced parasympathetic recovery.

Common patterns: Fear -> sympathetic overdrive + sleep disruption + muscle tension + digestive suppression; chronic uncertainty can sustain the pattern.

Your result in this scan: Suggests lower dominance of fear-based vigilance within the current psycho-emotional state.

Desire

Within range

Human Consciousness Level

Reference range: 1.484 - 2.485 | **Value:** 1.711

What this means: Reflects motivational pull toward acquisition, reward, or craving-driven engagement within the source framework.

Why it matters: When disproportionate, desire patterns can show up as craving, impulsivity, overconsumption, or unstable reward regulation.

Functional significance: Linked to dopaminergic reward pathways, impulse regulation, blood-sugar instability, addictive coping, and unmet emotional needs.

Common patterns: Blood-sugar volatility + dopamine dysregulation + stress coping -> stronger desire/craving patterns; this may affect eating, stimulants, or compulsive behaviours.

Your result in this scan: Desire appears within the analyser's expected range in this scan.

Anger

Within range

Human Consciousness Level

Reference range: 1.256 - 2.529 | **Value:** 1.466

What this means: Reflects a state of irritability, frustration, or forceful activation within the source framework.

Why it matters: Anger patterns can increase cardiovascular load, muscle tension, inflammatory signalling, and relational stress.

Functional significance: Linked to sympathetic activation, liver-tension symbolism, catecholamine surge, poor frustration tolerance, and sleep disruption.

Common patterns: Stress + sleep debt + blood-sugar instability -> irritability and anger; tension headaches, PMS, and liver-gallbladder patterns may cluster here.

Your result in this scan: Anger appears within the analyser's expected range in this scan.

Pride

Mildly reduced

Human Consciousness Level

Reference range: 1.378 - 2.489 | **Value:** 1.025

What this means: Reflects an elevated self-protective or self-affirming emotional posture within the source framework.

Why it matters: Depending on context, pride can reflect healthy self-regard or compensatory defensiveness masking vulnerability and poor regulation.

Functional significance: Best interpreted as a relational-emotional stance marker tied to self-concept, coping style, and adaptive versus defensive regulation.

Common patterns: Stress, insecurity, or overcompensation may elevate pride-like presentation; social threat can shift this marker in either direction.

Your result in this scan: Suggests less prominence of pride-based posturing or defensive elevation in the current pattern.

Courage

Within range

Human Consciousness Level

Reference range: 1.727 - 3.315 | **Value:** 1.8

What this means: Reflects the capacity to remain engaged and act despite discomfort, uncertainty, or stress within the source framework.

Why it matters: Courage is associated with adaptive resilience, behavioural activation, and willingness to engage with challenge without complete collapse or avoidance.

Functional significance: Linked to autonomic flexibility, motivation, emotional regulation, and the capacity to maintain function under pressure.

Common patterns: Better sleep + autonomic flexibility + stronger self-regulation often support courage-like engagement; fear and depletion tend to suppress it.

Your result in this scan: Courage appears within the analyser's expected range in this scan.

Neutrality

Mildly reduced

Human Consciousness Level

Reference range: 2.471 - 3.892 | **Value:** 2.16

What this means: Reflects a balanced, less-reactive emotional stance within the source framework, associated with reduced polarisation and steadier regulation.

Why it matters: Greater neutrality can signal improved stress tolerance, reduced emotional volatility, and better capacity for reflective decision-making.

Functional significance: Linked to autonomic balance, lower reactivity, emotional flexibility, and more stable cognitive-affective regulation.

Common patterns: Sleep stability + lower sympathetic load + better emotional regulation often support neutrality; chronic stress tends to erode it.

Your result in this scan: Suggests more reactive or polarised emotional processing and lower regulation steadiness.

Willingness

Within range

Human Consciousness Level

Reference range: 2.216 - 4.094 | **Value:** 2.598

What this means: Reflects an engaged, cooperative motivational stance within the source framework, associated with readiness to participate and adapt.

Why it matters: Willingness supports behavioural follow-through, openness to change, and resilience in treatment or recovery processes.

Functional significance: Linked to motivation, executive engagement, emotional readiness, and reduced avoidance under stress.

Common patterns: Better motivation, reduced fear, and stronger energy often support willingness; chronic depletion or distrust can reduce it.

Your result in this scan: Willingness appears within the analyser's expected range in this scan.

Acceptance

Within range

Human Consciousness Level

Reference range: 1.668 - 4.053 | **Value:** 2.239

What this means: Reflects a state of reduced resistance and greater adaptive acknowledgment of reality within the source framework.

Why it matters: Acceptance can reduce internal stress friction, improve treatment engagement, and support more stable emotional processing.

Functional significance: Linked to emotional regulation, reduced rumination, stress adaptation, and improved coping flexibility.

Common patterns: Reduced rumination + better nervous-system regulation often support acceptance; chronic threat and shame states tend to oppose it.

Your result in this scan: Acceptance appears within the analyser's expected range in this scan.

Reason

Within range

Human Consciousness Level

Reference range: 1.352 - 3.436 | **Value:** 2.693

What this means: Reflects the relative dominance of reflective, analytical, and cognitively organised processing within the source framework.

Why it matters: Reasoned processing supports planning, judgment, emotional modulation, and the ability to step back from impulsive reactivity.

Functional significance: Linked to executive function, prefrontal regulation, cognitive clarity, sleep quality, and reduced stress-noise interference.

Common patterns: Sleep deprivation + inflammatory fatigue + stress overdrive can impair reasoned processing; good metabolic and cognitive support improves it.

Your result in this scan: Reason appears within the analyser's expected range in this scan.

Love

Mildly reduced

Human Consciousness Level

Reference range: 2.138 - 3.754 | **Value:** 1.871

What this means: Reflects an affiliative, connected, and emotionally integrated state within the source framework.

Why it matters: Connected and prosocial emotional states are often associated with better regulation, relational safety, and reduced chronic stress physiology.

Functional significance: Linked to vagal tone, social safety signalling, oxytocin-associated states, attachment security, and emotional integration.

Common patterns: Higher vagal tone + felt safety + reduced threat physiology can support love/connection states; trauma and chronic stress often suppress them.

Your result in this scan: Suggests lower access to affiliative regulation or reduced sense of connection and emotional safety.

Joy

Within range

Human Consciousness Level

Reference range: 2.129 - 4.021 | **Value:** 2.654

What this means: Reflects access to positive affect, vitality, and unforced wellbeing within the source framework.

Why it matters: Sustained inability to access joy can signal fatigue, depressive load, burnout, anhedonia, or nervous-system dysregulation.

Functional significance: Linked to reward physiology, dopamine balance, social engagement, energy availability, and reduced inflammatory-depressive burden.

Common patterns: Burnout + low dopamine + poor sleep + inflammatory load can reduce joy; restored energy and social engagement often improve it.

Your result in this scan: Joy appears within the analyser's expected range in this scan.

Peace

Mildly reduced

Human Consciousness Level

Reference range: 2.243 - 4.022 | **Value:** 2.099

What this means: Reflects a deeply settled, non-reactive internal state within the source framework, associated with low internal conflict and high autonomic ease.

Why it matters: Peace-like states indicate reduced threat physiology, greater emotional integration, and improved resilience to stress reactivity.

Functional significance: Linked to parasympathetic dominance, reduced rumination, lower inflammatory stress signalling, and improved recovery tone.

Common patterns: Good sleep + parasympathetic tone + reduced chronic stress support peace; trauma activation and burnout tend to oppose it.

Your result in this scan: Suggests internal agitation, unresolved stress, or poor access to settled regulation.

Enlightenment

Within range

Human Consciousness Level

Reference range: 2.225 - 3.822 | **Value:** 3.556

What this means: Reflects the highest conceptual level in the source framework and should be interpreted as a symbolic marker of integrative awareness rather than a clinical state.

Why it matters: This marker is not a conventional medical construct and should not be over-medicalised; at most, it may function as an abstract resilience or integration indicator in the source system.

Functional significance: Best treated as a non-diagnostic, interpretive construct linked loosely to self-awareness, regulation, and integrative functioning.

Common patterns: Use cautiously and avoid over-interpretation; this marker is more philosophical than clinical and should always remain secondary to concrete physiological findings.

Your result in this scan: Enlightenment appears within the analyser's expected range in this scan.

Respiratory Function

Low priority

Complete system marker table — all markers and all severity grades are shown below.

Flagged markers: 1 | Within range: 3

Volume Tidal(VT)

Mildly reduced

Respiratory Function

Reference range: 5.431 - 7.479 | **Value:** 4.051

What this means: Reflects the volume of air moved in and out of the lungs during a normal resting breath, indicating baseline breathing efficiency and ventilatory patterning.

Why it matters: Tidal volume is a useful marker of whether breathing is shallow and inefficient or adequately supports gas exchange without excessive work.

Functional significance: Linked to diaphragmatic recruitment, autonomic tone, chest-wall movement, respiratory drive, and the balance between calm nasal breathing and stress-pattern upper-chest breathing.

Common patterns: Stress breathing -> lower VT with higher rate; diaphragmatic weakness, pain, or restriction can also reduce effective tidal volume.

Your result in this scan: Suggests shallow breathing, reduced diaphragmatic engagement, low ventilatory efficiency, or chronic under-breathing patterns.

Inspiratory Volume(TI)

Within range

Respiratory Function

Reference range: 4.126 - 6.045 | **Value:** 4.442

What this means: Reflects inspiratory volume capacity and the efficiency of drawing air into the lungs, indicating how effectively inhalation mechanics are functioning.

Why it matters: Inspiratory performance is central to oxygen uptake, diaphragmatic conditioning, and the ability to expand beyond shallow stress-driven breathing.

Functional significance: Linked to diaphragmatic strength, rib-cage mobility, inspiratory-muscle function, thoracic posture, and restrictive or pain-limited breathing mechanics.

Common patterns: Poor posture + anxiety + diaphragmatic weakness -> lower inspiratory volume; restrictive lung or chest-wall patterns can intensify the limitation.

Your result in this scan: Inspiratory Volume(TI) appears within the analyser's expected range in this scan.

Functional Residual Capacity(FRC)

Within range

Respiratory Function

Reference range: 5.147 - 6.219 | **Value:** 5.503

What this means: Reflects the volume of air remaining in the lungs after a normal exhalation, indicating resting lung-volume set point and the balance between elastic recoil and airway closure.

Why it matters: FRC is important because it stabilises gas exchange between breaths. Abnormal values can suggest restriction, air trapping, altered recoil, or poor breathing mechanics.

Functional significance: Linked to lung elasticity, small-airway patency, chest-wall mechanics, obesity-related restriction, and the degree of hyperinflation or airway collapse.

Common patterns: Restriction lowers FRC; obstructive air trapping may increase it. Postural compression and abdominal pressure can also reduce effective resting lung volume.

Your result in this scan: Functional Residual Capacity(FRC) appears within the analyser's expected range in this scan.

Residual Volume(RV)

Within range

Respiratory Function

Reference range: 4.363 - 5.647 | **Value:** 5.322

What this means: Reflects the volume of air remaining in the lungs after maximal exhalation, indicating how completely the lungs can empty and whether air trapping is present.

Why it matters: Residual volume is highly relevant to obstructive patterns, hyperinflation, and the efficiency of expiration. Excess air trapping increases work of breathing and impairs gas exchange.

Functional significance: Linked to airway resistance, expiratory flow limitation, lung elasticity, bronchial narrowing, and small-airway collapse during exhalation.

Common patterns: Higher airway resistance + loss of elastic recoil -> raised RV; chronic inflammation, mucus burden, and bronchial narrowing can all contribute.

Your result in this scan: Residual Volume(RV) appears within the analyser's expected range in this scan.

Lecithin

Medium priority

Complete system marker table — all markers and all severity grades are shown below.

Flagged markers: 2 | Within range: 4

Phospholipid index

Within range

Lecithin

Reference range: 3.654 - 4.478 | **Value:** 4.471

What this means: Represents the status of phospholipids as key structural components of cell membranes and lipoproteins.

Why it matters: Phospholipids are essential for membrane integrity, cellular signalling, and lipid transport.

Functional significance: Linked to liver function, lipid metabolism, and membrane fluidity.

Common patterns: Liver dysfunction + oxidative stress -> membrane instability.

Your result in this scan: Phospholipid index appears within the analyser's expected range in this scan.

Sphingolipid index

Mildly reduced

Lecithin

Reference range: 3.121 - 3.853 | **Value:** 2.842

What this means: Reflects sphingolipid balance involved in membrane structure and cellular signalling.

Why it matters: Sphingolipids regulate cell signalling, apoptosis, and membrane stability.

Functional significance: Linked to lipid metabolism and inflammatory signalling.

Common patterns: Inflammation -> sphingolipid dysregulation.

Your result in this scan: Suggests reduced signalling capacity or membrane dysfunction.

PH

Within range

Lecithin

Reference range: 3.341 - 4.214 | **Value:** 3.894

What this means: Reflects overall acid-base tendency within the source framework, functioning as a broad indicator of systemic buffering balance rather than a standalone compartment-specific laboratory pH value.

Why it matters: Acid-base balance matters because enzyme activity, mitochondrial output, mineral stability, respiratory compensation, and renal buffering all depend on an adequately regulated internal pH environment.

Functional significance: Linked to renal acid excretion, respiratory CO2 handling, dietary acid load, mineral buffering, mitochondrial metabolism, and inflammatory or catabolic states that shift the internal acid-base

terrain.

Common patterns: High metabolic stress + poor renal buffering + altered ventilation -> pH imbalance; chronic inflammatory load, tissue hypoxia, and catabolic states can all destabilise acid-base resilience.

Your result in this scan: PH appears within the analyser's expected range in this scan.

Lecithin index

Within range

Lecithin

Reference range: 4.365 - 5.142 | **Value:** 4.165

What this means: Represents lecithin levels contributing to lipid transport and membrane structure.

Why it matters: Lecithin supports lipid metabolism, liver function, and bile production.

Functional significance: Linked to phosphatidylcholine, liver health, and fat digestion.

Common patterns: Liver stress -> lecithin depletion.

Your result in this scan: Lecithin index appears within the analyser's expected range in this scan.

Brain phospholipid index

Mildly reduced

Lecithin

Reference range: 4.225 - 5.713 | **Value:** 4.038

What this means: Reflects phospholipid availability and membrane integrity within brain tissue, influencing synaptic signalling, myelin stability, receptor function, and neuronal resilience.

Why it matters: Phospholipids are fundamental to cell membrane fluidity, neurotransmission, myelin support, and effective communication across neural networks.

Functional significance: Depends on choline status, essential fatty acid balance, hepatic phospholipid metabolism, oxidative protection, and efficient membrane turnover.

Common patterns: Low phospholipid support + oxidative stress → poorer membrane fluidity → reduced neurotransmission and memory performance; liver dysfunction can also impair phospholipid synthesis and trafficking.

Your result in this scan: Suggests reduced membrane integrity, impaired synaptic signalling, weaker myelin support, or suboptimal availability of choline and structural lipids.

Liposomal index

Within range

Lecithin

Reference range: 3.112 - 4.081 | **Value:** 4.071

What this means: Reflects liposomal or phospholipid-vesicle style transport capacity, indicating how effectively lipid-based carriers may support membrane delivery, absorption, and intracellular nutrient trafficking.

Why it matters: Liposomal transport concepts are relevant to membrane interaction, nutrient-delivery efficiency, and the broader integrity of phospholipid-based transport environments.

Functional significance: Linked to phospholipid sufficiency, bile-dependent fat handling, membrane fluidity, intracellular transport efficiency, and liver-lipid processing.

Common patterns: Low phospholipid sufficiency plus poor bile flow can reduce lipid-carrier efficiency and membrane-delivery support; this may coexist with broader membrane and hepatic lipid dysfunction.

Your result in this scan: Liposomal index appears within the analyser's expected range in this scan.

Fatty acid

Medium priority

Complete system marker table — all markers and all severity grades are shown below.

Flagged markers: 2 | Within range: 2

Saturated fatty acid index

Within range

Fatty acid

Reference range: 1.334 - 2.407 | **Value:** 1.437

What this means: Represents the proportion of saturated fats within overall lipid balance.

Why it matters: Excess saturated fats can influence inflammation and metabolic health.

Functional significance: Linked to diet, lipid metabolism, and insulin sensitivity.

Common patterns: High processed fat intake -> metabolic dysfunction.

Your result in this scan: Saturated fatty acid index appears within the analyser's expected range in this scan.

Unsaturated fatty acid index

Within range

Fatty acid

Reference range: 2.224 - 3.153 | **Value:** 2.834

What this means: Reflects levels of unsaturated fats supporting membrane fluidity and metabolic function.

Why it matters: Unsaturated fats are essential for cardiovascular and metabolic health.

Functional significance: Linked to omega balance and lipid metabolism.

Common patterns: Diet influences fatty acid balance.

Your result in this scan: Unsaturated fatty acid index appears within the analyser's expected range in this scan.

Essential fatty acid index

Mildly reduced

Fatty acid

Reference range: 2.144 - 3.238 | **Value:** 2.087

What this means: Represents availability of essential fatty acids required for cellular function.

Why it matters: Essential fatty acids support inflammation control and membrane health.

Functional significance: Linked to omega-3 and omega-6 balance.

Common patterns: Poor diet -> deficiency.

Your result in this scan: Suggests deficiency affecting cellular health.

Triglyceride index

Mildly reduced

Fatty acid

Reference range: 3.361 - 4.644 | **Value:** 2.377

What this means: Reflects circulating triglyceride levels and lipid storage status.

Why it matters: Triglycerides are a key marker of metabolic health and energy storage.

Functional significance: Linked to diet, insulin, and liver function.

Common patterns: Insulin resistance -> high triglycerides.

Your result in this scan: Suggests low lipid storage.

TECHNICAL REFERENCE

Appendix

This section contains supporting technical material and summary reference tables used in the report.

Section	Marker	Range	Value	Status
Cardiovascular and Cerebrovascular	Blood Viscosity	48.264 - 65.371	53.723	Within range
Cardiovascular and Cerebrovascular	Cholesterol Crystal	56.749 - 67.522	66.455	Within range
Cardiovascular and Cerebrovascular	Blood Fat	0.481 - 1.043	0.902	Within range
Cardiovascular and Cerebrovascular	Vascular Resistance	0.327 - 0.937	1.226	Mildly elevated
Cardiovascular and Cerebrovascular	Vascular Elasticity	1.672 - 1.978	1.934	Within range
Cardiovascular and Cerebrovascular	Myocardial Blood Demand	0.192 - 0.412	0.407	Within range
Cardiovascular and Cerebrovascular	Myocardial Perfusion	4.832 - 5.147	5.132	Within range
Cardiovascular and Cerebrovascular	Myocardial Oxygen Consumption	3.321 - 4.244	3.382	Within range
Cardiovascular and Cerebrovascular	Stroke Volume	1.338 - 1.672	1.29	Mildly reduced
Cardiovascular and Cerebrovascular	LV Ejection Impedance	0.669 - 1.544	1.456	Within range
Cardiovascular and Cerebrovascular	LV Pump Power	1.554 - 1.988	0.991	Moderately reduced
Cardiovascular and Cerebrovascular	Coronary Artery Elasticity	1.553 - 2.187	1.66	Within range
Cardiovascular and Cerebrovascular	Coronary Perfusion Pressure	11.719 - 18.418	19.312	Moderately elevated
Cardiovascular and Cerebrovascular	Cerebral Blood Vessel Elasticity	0.708 - 1.942	1.021	Within range
Cardiovascular and Cerebrovascular	Brain Tissue Blood Supply Status	6.138 - 21.396	20.196	Within range
Gastrointestinal Function	Pepsin Secretion Coefficient	59.847 - 65.234	59.915	Within range
Gastrointestinal Function	Gastric Peristalsis Function Coefficient	58.425 - 61.213	58.71	Within range
Gastrointestinal Function	Gastric Absorption Function Coefficient	34.367 - 35.642	29.734	Moderately reduced

Section	Marker	Range	Value	Status
Gastrointestinal Function	Small Intestine Motility	133.437 - 140.476	126.057	Moderately reduced
Gastrointestinal Function	Small Intestine Absorption	3.572 - 6.483	2.82	Moderately reduced
Large Intestine Function	Colon Motility	4.572 - 6.483	3.088	Moderately reduced
Large Intestine Function	Colon Absorption	2.946 - 3.815	2.952	Within range
Large Intestine Function	Gut Microbiome	1.734 - 2.621	0.807	Moderately reduced
Large Intestine Function	Intraluminal pressure coefficient	1.173 - 2.297	2.639	Mildly elevated
Liver Function	Protein Metabolism	116.34 - 220.621	116.569	Within range
Liver Function	Energy Production Function	0.713 - 0.992	0.881	Within range
Liver Function	Detoxification Function	0.202 - 0.991	0.977	Within range
Liver Function	Bile Secretion Function	0.432 - 0.826	0.717	Within range
Liver Function	Liver Fat Content	0.097 - 0.419	0.636	Moderately elevated
Gallbladder Function	Serum Globulin (A/G)	126 - 159	126.767	Within range
Gallbladder Function	Total Bilirubin (TBIL)	0.232 - 0.686	0.542	Within range
Gallbladder Function	Alkaline Phosphatase (ALP)	0.082 - 0.342	0.316	Within range
Gallbladder Function	Serum Total Bile Acid (TBA	0.317 - 0.695	0.704	Mildly elevated
Gallbladder Function	Bilirubin (DBIL)	0.218 - 0.549	0.56	Mildly elevated
Pancreatic Function	Insulin	2.845 - 4.017	3.036	Within range
Pancreatic Function	Pancreatic Polypeptide (PP)	3.210 - 6.854	3.26	Within range
Pancreatic Function	Glucagon	2.412 - 2.974	2.666	Within range
Kidney Function	Urobilinogen Index	2.762 - 5.424	6.215	Mildly elevated
Kidney Function	Uric acid Index	1.435 - 1.987	2.689	Moderately elevated
Kidney Function	BUN	4.725 - 8.631	4.766	Within range
Kidney Function	Proteinuria Index	1.571 - 4.079	4.761	Mildly elevated
Lung Function	Vital Capacity	3348 - 3529	3489	Within range
Lung Function	Total Lung Capacity	4301 - 4782	4286	Mildly reduced
Lung Function	Airway Resistance	1.374 - 1.709	1.621	Within range
Lung Function	Arterial Oxygen Content	17.903 - 21.012	20.78	Within range
Brain Nerve	Brain Tissue Blood Supply	143.37 - 210.81	168.036	Within range
Brain Nerve	Cerebral Arteriosclerosis	0.103 - 0.642	0.196	Within range
Brain Nerve	Cranial Nerve Function	0.253 - 0.659	0.292	Within range

Section	Marker	Range	Value	Status
Brain Nerve	Sentiment Index	0.109 - 0.351	0.262	Within range
Brain Nerve	Memory	0.442 - 0.817	0.379	Mildly reduced
Bone Disease	Adhesion Degree of Shoulder Muscle	None	< u 0.2	Unclear
Bone Disease	Age of Ligament	None	10%-40%	Unclear
Bone Mineral Density	Osteoclast Coefficient	86.73 - 180.97	151.19	Within range
Bone Mineral Density	Amount of Calcium Loss	0.209 - 0.751	0.469	Within range
Bone Mineral Density	Degree of Bone Hyperplasia	0.046 - 0.167	0.389	Mildly elevated
Bone Mineral Density	Degree of Osteoporosis	0.124 - 0.453	0.528	Moderately elevated
Bone Mineral Density	Bone Mineral Density	0.796 - 0.433	0.435	Within range
Rheumatoid Bone Disease	Degree of Cervical Calcification	421 - 490	502.068	Mildly elevated
Rheumatoid Bone Disease	Degree of Lumbar Calcification	4.326 - 7.531	7.352	Within range
Rheumatoid Bone Disease	Bone Hyperplasia Coefficient	2.954 - 5.543	6	Mildly elevated
Rheumatoid Bone Disease	Osteoporosis Coefficient	2.019 - 4.721	4.705	Within range
Rheumatoid Bone Disease	Rheumatism Coefficient	4.023 - 11.627	16.843	Moderately elevated
Bone Growth Index	Bone alkaline phosphatase	0.433 - 0.796	0.399	Mildly reduced
Bone Growth Index	Osteocalcin	0.525 - 0.817	0.508	Mildly reduced
Bone Growth Index	Status of long bone healing	0.713 - 0.992	0.606	Mildly reduced
Bone Growth Index	Short bone cartilage healing situation	0.202 - 0.991	0.362	Within range
Bone Growth Index	Epiphyseal line	0.432 - 0.826	0.769	Within range
Blood Sugar	Coefficient of Insulin Secretion	2.967 - 3.528	3.043	Within range
Blood Sugar	Blood Sugar Coefficient	2.163 - 7.321	3.191	Within range
Blood Sugar	Urine Sugar Coefficient	2.204 - 2.819	2.535	Within range
Trace Element	Calcium	1.219 - 3.021	1.518	Within range
Trace Element	Iron	1.151 - 1.847	0.743	Mildly reduced
Trace Element	Zinc	1.143 - 1.989	1.336	Within range
Trace Element	Selenium	0.847 - 2.045	1.096	Within range
Trace Element	Phosphorus	1.195 - 2.134	2.087	Within range
Trace Element	Potassium	0.689 - 0.987	0.977	Within range
Trace Element	Magnesium	0.568 - 0.992	0.761	Within range
Trace Element	Copper	0.474 - 0.749	0.23	Moderately reduced
Trace Element	Cobalt	2.326 - 5.531	2.493	Within range

Section	Marker	Range	Value	Status
Trace Element	Manganese	0.497 - 0.879	0.732	Within range
Trace Element	Iodine	1.421 - 5.490	5.311	Within range
Trace Element	Nickel	2.462 - 5.753	2.481	Within range
Trace Element	Fluorine	1.954 - 4.543	2.019	Within range
Trace Element	Molybdenum	0.938 - 1.712	1.43	Within range
Trace Element	Vanadium	1.019 - 3.721	2.638	Within range
Trace Element	Tin	1.023 - 7.627	7.159	Within range
Trace Element	Silicon	1.425 - 5.872	1.367	Mildly reduced
Trace Element	Strontium	1.142 - 5.862	3.508	Within range
Trace Element	Boron	1.124 - 3.453	2.537	Within range
Vitamin	Vitamin A	0.346 - 0.401	0.338	Mildly reduced
Vitamin	Vitamin B1	2.124 - 4.192	1.679	Mildly reduced
Vitamin	Vitamin B2	1.549 - 2.213	1.549	Within range
Vitamin	Vitamin B3	14.477 - 21.348	14.72	Within range
Vitamin	Vitamin B6	0.824 - 1.942	1.616	Within range
Vitamin	Vitamin B12	6.428 - 21.396	6.962	Within range
Vitamin	Vitamin C	4.543 - 5.023	3.55	Moderately reduced
Vitamin	Vitamin D3	5.327 - 7.109	5.259	Mildly reduced
Vitamin	Vitamin E	4.826 - 6.013	4.11	Moderately reduced
Vitamin	Vitamin K	0.717 - 1.486	0.599	Mildly reduced
Amino Acid	Lysine	0.962 - 1.213	0.853	Mildly reduced
Amino Acid	Tryptophan	4.978 - 6.289	6.008	Within range
Amino Acid	Phenylalanine	1.928 - 2.491	2.057	Within range
Amino Acid	Methionine	1.245 - 1.637	1.398	Within range
Amino Acid	Threonine	1.194 - 1.685	1.37	Within range
Amino Acid	Isoleucine	4.582 - 5.657	4.175	Mildly reduced
Amino Acid	Leucine	6.982 - 9.256	7.041	Within range
Amino Acid	Valine	6.982 - 9.677	7.565	Within range
Amino Acid	Histidine	5.113 - 6.258	5.261	Within range
Amino Acid	Arginine	1.812 - 2.337	1.975	Within range
Amino Acid	Homocysteine	0.983 - 1.265	1.073	Within range
Coenzyme	Nicotinamide	2.074 - 3.309	2.293	Within range

Section	Marker	Range	Value	Status
Coenzyme	Biotin	1.833 - 2.979	1.241	Mildly reduced
Coenzyme	Pantothenic acid	1.116 - 2.101	0.604	Moderately reduced
Coenzyme	Folic acid	1.449 - 2.246	2.236	Within range
Coenzyme	Coenzyme Q10	0.831 - 1.588	0.643	Mildly reduced
Coenzyme	Glutathione	0.726 - 1.281	1.114	Within range
Essential Fatty Acid	Linoleic acid	0.642 - 0.985	0.447	Mildly reduced
Essential Fatty Acid	α-Linolenic acid	0.814 - 1.202	0.852	Within range
Essential Fatty Acid	γ-Linolenic acid	0.921 - 1.334	0.996	Within range
Essential Fatty Acid	Arachidonic acid	0.661 - 0.808	0.575	Mildly reduced
Endocrine System	Thyroid secretion index	2.954 - 5.543	2.996	Within range
Endocrine System	Parathyroid hormone secretion index	2.845 - 4.017	2.121	Mildly reduced
Endocrine System	Adrenal glands Index	2.412 - 2.974	2.149	Mildly reduced
Endocrine System	Pituitary secretion index	2.163 - 7.34	5.512	Within range
Endocrine System	Pineal secretion index	3.210 - 6.854	6.778	Within range
Endocrine System	Thymus gland secretion index	2.967 - 3.528	2.969	Within range
Endocrine System	Gland secretion index	2.204 - 2.819	2.492	Within range
Immune System	Lymph node Index	133.437 - 140.47	134.342	Within range
Immune System	Tonsil immune Index	0.124 - 0.453	0.307	Within range
Immune System	Bone marrow Index	0.146 - 3.218	1.068	Within range
Immune System	Spleen index	34.367 - 35.642	35.434	Within range
Immune System	Thymus index	58.425 - 61.213	59.292	Within range
Immune System	Immunoglobulin index	3.712 - 6.981	4.089	Within range
Immune System	Respiratory Immunity	3.241 - 9.814	3.384	Within range
Immune System	GI Immunity	0.638 - 1.712	0.949	Within range
Immune System	Mucosal Immunity	4.111 - 18.741	10.307	Within range
Thyroid	FT4	0.103 - 0.316	0.6	Mildly elevated
Thyroid	Thyroglobulin	0.114 - 0.202	0.294	Mildly elevated
Thyroid	Thyroglobulin	0.421 - 0.734	0.711	Within range
Thyroid	T3	0.161 - 0.308	0.518	Mildly elevated
Human Toxin	Stimulating Beverage	0.209 - 0.751	0.475	Within range
Human Toxin	Electromagnetic Radiation	0.046 - 0.167	0.109	Within range
Human Toxin	Free Radical	0.124 - 0.453	0.178	Within range
Human Toxin	Toxic Pesticide Residue	0.013 - 0.313	0.305	Within range

Section	Marker	Range	Value	Status
Heavy Metal	Lead	0.052 - 0.643	0.976	Mildly elevated
Heavy Metal	Mercury	0.013 - 0.336	0.468	Mildly elevated
Heavy Metal	Cadmium	0.527 - 1.523	1.556	Mildly elevated
Heavy Metal	Chromium	0.176 - 1.183	0.345	Within range
Heavy Metal	Arsenic	0.153 - 0.621	0.596	Within range
Heavy Metal	Antimony	0.162 - 0.412	0.323	Within range
Heavy Metal	Thallium	0.182 - 0.542	0.388	Within range
Heavy Metal	Aluminum	0.192 - 0.412	0.57	Mildly elevated
Basic Physical Quality	Response Ability	59.786 - 65.424	61.674	Within range
Basic Physical Quality	Mental Power	58.715 - 63.213	61.23	Within range
Basic Physical Quality	Water Shortage	33.967 - 37.642	34.512	Within range
Basic Physical Quality	Hypoxia	133.642 - 141.476	130.994	Mildly reduced
Basic Physical Quality	PH	3.156 - 3.694	3.248	Within range
Allergy	Drug Sensitivity	0.431 - 1.329	0.563	Within range
Allergy	Alcohol allergy index	0.432 - 1.246	1.207	Within range
Allergy	Pollen allergy index	0.143 - 1.989	1.646	Within range
Allergy	Injection allergy index	0.847 - 1.045	1.184	Mildly elevated
Allergy	Chemical Sensitivity	0.842 - 1.643	1.896	Mildly elevated
Allergy	Paint allergy index	0.346 - 1.401	0.489	Within range
Allergy	Dust allergy index	0.543 - 1.023	1.263	Mildly elevated
Allergy	Smoke allergy index	0.826 - 1.013	2.532	Mildly elevated
Allergy	Hair dye allergy index	0.717 - 1.486	2.856	Moderately elevated
Allergy	Animal fur allergy index	0.124 - 1.192	0.571	Within range
Allergy	Metal jewelry allergy index	0.549 - 1.213	0.743	Within range
Allergy	Seafood allergy index	0.449 - 1.246	2.379	Mildly elevated
Allergy	Milk allergy index	0.477 - 1.348	0.739	Within range
Obesity	Abnormal lipid metabolism coefficient	1.992 - 3.713	0.89	Moderately reduced
Obesity	Brown adipose tissue abnormalities coefficient	2.791 - 4.202	1.858	Moderately reduced
Obesity	Hyperinsulinemia coefficient	0.097 - 0.215	0.206	Within range
Obesity	Nucleus of the hypothalamus abnormal coefficient	0.332 - 0.626	0.719	Mildly elevated

Section	Marker	Range	Value	Status
Obesity	Triglyceride content of abnormal coefficient	1.341 - 1.991	5.222	Moderately elevated
Skin	Skin Free Radical Index	0.124 - 3.453	2.972	Within range
Skin	Skin Collagen Index	4.471 - 6.079	2.899	Mildly reduced
Skin	Skin Grease Index	14.477 - 21.348	25.821	Mildly elevated
Skin	Skin Immunity Index	1.035 - 3.230	4.02	Mildly elevated
Skin	Skin Moisture Index	0.218 - 0.953	2.145	Moderately elevated
Skin	Skin Moisture Loss	2.214 - 4.158	3.125	Within range
Skin	Skin Red Blood Trace Index	0.824 - 1.942	1.055	Within range
Skin	Skin Elasticity Index	2.717 - 3.512	1.571	Mildly reduced
Skin	Skin Melanin Index	0.346 - 0.501	0.612	Mildly elevated
Skin	Tin	0.842 - 1.858	1.419	Within range
Eye	Bags under the eyes	0.510 - 3.109	3.004	Within range
Eye	Collagen eye wrinkle	2.031 - 3.107	1.31	Mildly reduced
Eye	Dark circles	0.831 - 3.188	3.883	Mildly elevated
Eye	Lymphatic obstruction	1.116 - 4.101	3.52	Within range
Eye	Sagging	0.233 - 0.559	0.751	Mildly elevated
Eye	Edema	0.332 - 0.726	1.234	Moderately elevated
Eye	Eye cell activity	0.118 - 0.892	0.895	Mildly elevated
Eye	Visual fatigue	2.017 - 5.157	8.04	Mildly elevated
Collagen	Overall Eye Function	6.352 - 8.325	8.31	Within range
Collagen	Endocrine system	7.245 - 8.562	7.491	Within range
Collagen	Hair and skin	4.533 - 6.179	4.806	Within range
Collagen	Endocrine system	6.178 - 8.651	8.253	Within range
Collagen	Circulatory system	3.586 - 4.337	4.153	Within range
Collagen	Digestive system	3.492 - 4.723	3.168	Mildly reduced
Collagen	Immune system	3.376 - 4.582	2.938	Mildly reduced
Collagen	Motion systems	6.458 - 8.133	5.573	Mildly reduced
Collagen	Muscle Tissue	6.552 - 8.268	5.774	Mildly reduced
Collagen	Fat Metabolism	6.338 - 8.368	4.551	Mildly reduced
Collagen	Detoxification and metabolism	6.187 - 8.466	6.519	Within range
Collagen	Reproductive system	3.778 - 4.985	3.997	Within range

Section	Marker	Range	Value	Status
Collagen	Nervous system	3.357 - 4.239	2.643	Mildly reduced
Collagen	Skeleton	6.256 - 8.682	7.975	Within range
Channels and collaterals	Hand Tai Yin Lung Meridian	48.264 - 65.371	63.778	Within range
Channels and collaterals	Hand Yangming Large Intestine Meridian	56.749 - 67.522	61.756	Within range
Channels and collaterals	Foot Yangming Stomach Meridian	0.481 - 1.043	0.485	Within range
Channels and collaterals	Foot Tai Yin Lung Meridian	0.327 - 0.937	0.465	Within range
Channels and collaterals	Hand Shao Yin Heart Sutra	1.672 - 1.978	1.477	Mildly reduced
Channels and collaterals	Hand the small intestine by the sun	0.192 - 0.412	0.199	Within range
Channels and collaterals	Foot Tai Yang Bladder Meridian	4.832 - 5.147	3.773	Mildly reduced
Channels and collaterals	Foot Shao Yin Kidney	3.321 - 4.244	3.937	Within range
Channels and collaterals	Hand Jue Yin Pericardium Meridian	1.338 - 1.672	1.34	Within range
Channels and collaterals	Hand Shao Yang Triple Burner Meridian	0.669 - 1.544	0.548	Mildly reduced
Channels and collaterals	Foot Shao Yang Gall Bladder Meridian	1.554 - 1.988	1.868	Within range
Channels and collaterals	Foot Jue Yin Liver Meridian	1.553 - 2.187	1.652	Within range
Channels and collaterals	Ren channel	11.719 - 18.418	12.572	Within range
Channels and collaterals	Governor meridian	0.708 - 1.942	1.536	Within range
Channels and collaterals	Vital meridian	6.138 - 21.396	6.331	Within range
Channels and collaterals	Tai mai	5.733 - 7.109	5.146	Mildly reduced
Pulse of heart and brain	Stroke index	60.735 - 65.396	62.076	Within range
Pulse of heart and brain	Stroke volume(SV)	63.012 - 67.892	61.901	Mildly reduced
Pulse of heart and brain	Heart peripheral resistance (TRR)	0.983 - 1.265	1.638	Mildly elevated
Pulse of heart and brain	Pulse wave coefficient K	0.316 - 0.401	0.391	Within range
Pulse of heart and brain	Cerebrovascular blood oxygen saturation(Sa)	0.710 - 1.109	0.894	Within range
Pulse of heart and brain	Cerebrovascular blood oxygen volume(CaCO2)	7.880 - 10.090	8.081	Within range
Pulse of heart and brain	Cerebrovascular blood oxygen presure(PaO2)	5.017 - 5.597	5.169	Within range
Blood lipids	Blood viscosity	4.131 - 4.562	5.233	Moderately elevated
Blood lipids	Total cholesterol(TC)	1.833 - 2.979	2.75	Within range
Blood lipids	Triglyceride(TG)	1.116 - 2.101	2.218	Mildly elevated
Blood lipids	High-density lipoprotein (HDL-C)	1.449 - 2.246	2.955	Mildly elevated
Blood lipids	Low-density lipoprotein (LDL-C)	0.831 - 1.588	1.015	Within range
Blood lipids	Neutral fat(MB)	0.726 - 1.281	3.096	Mildly elevated
Blood lipids	Circulating immune complex(CIC)	13.012 - 17.291	18.601	Mildly elevated

Section	Marker	Range	Value	Status
Prostate	Degree of Prostatic Hyperplasia	1.023 - 3.230	5.03	Moderately elevated
Prostate	Degree of Prostatic Calcification	1.471 - 6.079	8.907	Mildly elevated
Prostate	Prostatitis Syndrome	2.213 - 2.717	6.11	Moderately elevated
Male Sexual Function	Testosterone	3.342 - 9.461	2.492	Moderately reduced
Male Sexual Function	Gonadotropin	4.111 - 18.741	3.558	Mildly reduced
Male Sexual Function	Erection Transmitter	3.241 - 9.814	3.03	Mildly reduced
Sperm and semen	Semen volume	1.502 - 6.028	5.263	Within range
Sperm and semen	Semen liquefying time	10.283 - 30.282	26.982	Within range
Sperm and semen	Number of sperms	2.483 - 3.932	3	Within range
Sperm and semen	Sperm motility rate	0.637 - 0.877	0.689	Within range
Male Hormone	Androgen	1.372 - 1.878	1.492	Within range
Male Hormone	Estrogen	0.521 - 1.036	0.672	Within range
Male Hormone	Progesterone(P)	0.686 - 1.432	0.651	Mildly reduced
Male Hormone	Luteinizing Hormone (LH)	0.679 - 1.324	0.593	Mildly reduced
Male Hormone	Prolactin(PRL)	0.738 - 1.742	1.386	Within range
Male Hormone	Follicle Stimulating Hormone(FSH)	0.387 - 0.956	0.54	Within range
Human Immunity	Tissue barrier immune index	7.435 - 8.847	9.626	Mildly elevated
Human Immunity	Innate immune cell index	7.424 - 9.052	10.784	Mildly elevated
Human Immunity	Innate immune molecule index	7.143 - 9.214	7.585	Within range
Human Immunity	Cellular immune index	5.769 - 7.643	8.547	Mildly elevated
Human Immunity	Humoral immunity index	6.424 - 8.219	8.708	Mildly elevated
Human Consciousness Level	Shame	2.214 - 3.925	3.056	Within range
Human Consciousness Level	Guilt	2.154 - 3.478	2.543	Within range
Human Consciousness Level	Apathy	2.343 - 4.089	2.263	Mildly reduced
Human Consciousness Level	Grief	1.845 - 3.241	1.997	Within range
Human Consciousness Level	Fear	2.155 - 4.031	1.681	Mildly reduced
Human Consciousness Level	Desire	1.484 - 2.485	1.711	Within range
Human Consciousness Level	Anger	1.256 - 2.529	1.466	Within range
Human Consciousness Level	Pride	1.378 - 2.489	1.025	Mildly reduced
Human Consciousness Level	Courage	1.727 - 3.315	1.8	Within range
Human Consciousness Level	Neutrality	2.471 - 3.892	2.16	Mildly reduced

Section	Marker	Range	Value	Status
Human Consciousness Level	Willingness	2.216 - 4.094	2.598	Within range
Human Consciousness Level	Acceptance	1.668 - 4.053	2.239	Within range
Human Consciousness Level	Reason	1.352 - 3.436	2.693	Within range
Human Consciousness Level	Love	2.138 - 3.754	1.871	Mildly reduced
Human Consciousness Level	Joy	2.129 - 4.021	2.654	Within range
Human Consciousness Level	Peace	2.243 - 4.022	2.099	Mildly reduced
Human Consciousness Level	Enlightenment	2.225 - 3.822	3.556	Within range
Respiratory Function	Volume Tidal(VT)	5.431 - 7.479	4.051	Mildly reduced
Respiratory Function	Inspiratory Volume(TI)	4.126 - 6.045	4.442	Within range
Respiratory Function	Functional Residual Capacity(FRC)	5.147 - 6.219	5.503	Within range
Respiratory Function	Residual Volume(RV)	4.363 - 5.647	5.322	Within range
Lecithin	Phospholipid index	3.654 - 4.478	4.471	Within range
Lecithin	Sphingolipid index	3.121 - 3.853	2.842	Mildly reduced
Lecithin	PH	3.341 - 4.214	3.894	Within range
Lecithin	Lecithin index	4.365 - 5.142	4.165	Within range
Lecithin	Brain phospholipid index	4.225 - 5.713	4.038	Mildly reduced
Lecithin	Liposomal index	3.112 - 4.081	4.071	Within range
Fatty acid	Saturated fatty acid index	1.334 - 2.407	1.437	Within range
Fatty acid	Unsaturated fatty acid index	2.224 - 3.153	2.834	Within range
Fatty acid	Essential fatty acid index	2.144 - 3.238	2.087	Mildly reduced
Fatty acid	Triglyceride index	3.361 - 4.644	2.377	Mildly reduced

These scan findings are best interpreted alongside symptoms, history, and appropriate clinical assessment.