

NaturalApproaches.com

Personalised Wellness Scan Report

Wellness screening summary

Patient details

Name: Child Example

Sex: Male

Age: 5

Height: 100.0 cm

Weight: 16.0 kg

Profile: Child

Scan time: 21/03/2026 23:13

This report is for wellness education and discussion only. It is not a medical diagnosis.

Report generated: 17/04/2026 16:24 BST

Eze Chiana
care@naturalapproaches.com
https://www.naturalapproaches.com

NAVIGATION

Table of contents

Cover

At a glance

Scan score overview

Priority category review

Clinical recommendations

Recommended support plan

Complete product recommendation mapping

Complete system marker tables

Appendix

OVERVIEW

At a glance

This section summarises the dominant patterns in the scan, the wider downstream themes they may be influencing, and the immediate priorities for follow-up.

Clinical snapshot

The scan is most suggestive of neurocognitive support needs, with likely downstream effects on repair, resilience, and wider function. There is also a secondary picture of reduced digestive and absorptive efficiency and reduced energy-production efficiency.

Practitioner overview

Taken together, the findings point most strongly to neurocognitive support needs. Priority should be given to neurocognitive nutrient support, membrane and phospholipid support, and trace element repletion.

Why this matters

This matters because reduced neurocognitive support can affect regulation, stress tolerance, mental clarity, and day-to-day functional capacity. It may also help explain the overlap with mitochondrial and energy-production strain and connective-tissue and repair pattern.

Primary pattern

Neurocognitive support pattern

This pattern points to neurocognitive support needs, most evident across neurocognitive nutrient support, membrane and phospholipid support, trace element repletion, and developmental support review.

Follow-up focus: neurocognitive nutrient support, membrane and phospholipid support, trace element repletion, developmental support review

Contributing patterns

- **Mitochondrial and energy-production strain** — Reduced energy-production efficiency, most evident across mitochondrial support, cofactor repletion, and energy-production support.
- **Connective-tissue and repair pattern** — Connective-tissue and repair pattern, most evident across repair support, connective-tissue support, and collagen support.

- **Toxic burden and detoxification strain** — Toxic-load pressure, most evident across exposure review, detoxification support, and antioxidant reserve.

Immediate priorities for follow-up

- From a clinical perspective, the most relevant next steps are to focus on neurocognitive nutrient support, membrane and phospholipid support, trace element repletion, developmental support review, and mitochondrial support. Addressing these areas is likely to produce the greatest overall shift in the current pattern.

OVERVIEW

Health Index overview

This section provides a weighted overview of the scan, helping organise attention across major body systems before moving into more detailed interpretation.



Overall Scan Score

Generally Balanced

- Robust functional balance (85-100)
- Generally balanced (70-84)
- Needs focused support (55-69)
- Priority review recommended (0-54)

How to read this dashboard

This dashboard provides a weighted clinical overview of the scan. The Health Index and body-system cards are not based on a simple count of flagged markers. They are derived from the severity, distribution, and clustering of findings across related sections, with greater weighting given to patterns that may carry broader functional significance.

Each body-system card may include more than one underlying scan section, which is why the totals shown here may differ from the marker counts shown later on individual section pages. In general, lower scores suggest a more active pattern worth reviewing in follow-up, while higher scores suggest a more stable pattern on this dashboard view.

These scores are designed to help organise clinical attention. They are not a diagnosis and should always be interpreted alongside symptoms, history, and appropriate clinical assessment.

Endocrine & Hormonal



4 flagged · 2 within range

Includes: Growth and development

Metabolic



23 flagged · 37 within range

Includes: Trace Element, Vitamin, Amino Acid, Coenzyme, Essential Fatty Acid, Lecithin, Fatty acid

Neurocognitive



5 flagged · 7 within range

Includes: ADHD, Cognitive function

Priority overview

Trace Element

Priority: Low priority

Flagged markers: 7

Section Index: 85

Vitamin

Priority: Low priority

Flagged markers: 5

Section Index: 85

Growth and development

Priority: Low priority

Flagged markers: 4

Section Index: 85

Cognitive function

Priority: Within range

Flagged markers: 4

Section Index: 89

Fatty acid

Priority: Within range

Flagged markers: 2

Section Index: 89

Amino Acid

Priority: Within range

Flagged markers: 5

Section Index: 90

RECOMMENDATIONS

Clinical recommendations

This section translates the scan findings into high-level clinical priorities, showing which functional areas appear most relevant for follow-up support.

Nutrient repletion

Given the dominant pattern, priority should be given to restoring nutrient sufficiency and absorptive capacity, particularly in the context of neurocognitive support needs.

Why this was suggested: Trace Element contains 7 flagged markers, led by Iron, Zinc, Potassium.

Cognitive function

Support should focus on cognitive function and the broader drivers influencing it.

Why this was suggested: Cognitive function contains 4 flagged markers, led by Observation, Memory, Thinking Ability.

Growth and development

Support should focus on growth and developmental support in the context of the wider scan picture.

Why this was suggested: Growth and development contains 4 flagged markers, led by Height and body mass index, Height sitting height index, BMI index.

RECOMMENDATIONS

Recommended support plan

This section organises suggested support options into a staged plan, combining foundational and targeted measures based on the dominant scan patterns.

Neurocognitive support pattern Support Plan

This plan focuses on addressing the primary functional pattern, starting with foundational support before moving into targeted optimisation.

This plan is organised around neurocognitive support needs, with additional support for reduced digestive and absorptive efficiency and reduced energy-production efficiency.

Phase 1 — Foundation

This phase focuses on core digestive, nutritional, and repair support.

Neuro Support Blend

Natural Approaches

Core

Included to strengthen cognitive support within the broader neurocognitive support pattern picture.

Omega-3 Complex

Natural Approaches

Core

Included to strengthen membrane support within the broader neurocognitive support pattern picture.

Digestive Bitters

Natural Approaches

Included to strengthen digestive support within the broader absorption and assimilation strain picture.

Phase 2 — Targeted Support

This phase adds more specific support aligned with the strongest secondary patterns.

Gut Repair Complex

Natural Approaches

Core

Included to strengthen mucosal repair within the broader absorption and assimilation strain picture.

Mineral Repletion Blend

Natural Approaches

Core

Included to strengthen nutrient repletion within the broader absorption and assimilation strain picture.

REFERENCE MAPPING

Complete product recommendation mapping

This section lists all products identified through the scan analysis. It includes full mapping across systems and markers to support practitioner review, product selection, and fulfilment planning.

Product	Source	Systems	Key markers
Digestive Bitters	Natural Approaches	Absorption and assimilation strain, digestive support	
Gut Repair Complex	Natural Approaches	Absorption and assimilation strain, mucosal repair	
Mineral Repletion Blend	Natural Approaches	Absorption and assimilation strain, nutrient repletion	
Neuro Support Blend	Natural Approaches	Neurocognitive support pattern, cognitive support	
Omega-3 Complex	Natural Approaches	Neurocognitive support pattern, membrane support	

Trace Element

Low priority

Priority interpretation

Trace Element is one of the more active sections in this scan, with multiple flagged markers combining into a broader functional pattern.

Moderately and severely abnormal markers are prioritised here for faster clinical reading.

Flagged markers: 7 | Within range: 12 | Section Index: 85

Top findings

- Zinc is moderately reduced
- Potassium is moderately reduced
- Iron is moderately reduced

Priority marker explanations

Zinc

Result: 0.888 | **Reference range:** 1.143 - 1.989 | Moderately reduced

See Appendix for full clinical interpretation of this marker.

Potassium

Result: 0.441 | **Reference range:** 0.689 - 0.987 | Moderately reduced

See Appendix for full clinical interpretation of this marker.

Iron

Result: 0.584 | **Reference range:** 1.151 - 1.847 | Moderately reduced

See Appendix for full clinical interpretation of this marker.

Vitamin

Low priority

Priority interpretation

Vitamin is one of the more active sections in this scan, with multiple flagged markers combining into a broader functional pattern.

Moderately and severely abnormal markers are prioritised here for faster clinical reading.

Flagged markers: 5 | Within range: 5 | Section Index: 85

Top findings

- Vitamin C is moderately reduced
- Vitamin E is mildly reduced
- Vitamin B6 is mildly reduced

Priority marker explanations

Vitamin C

Result: 3.567 | **Reference range:** 4.543 - 5.023 | Moderately reduced

See Appendix for full clinical interpretation of this marker.

TECHNICAL REFERENCE

Complete system marker tables

This section includes every parsed system and every marker across the report, including those within range, as the full explained reference layer.

Trace Element

High priority

Complete system marker table — all markers and all severity grades are shown below.

Flagged markers: 7 | Within range: 12

Calcium

Within range

Trace Element

Reference range: 1.219 - 3.021 | **Value:** 1.948

What this means: Represents calcium status within the body, reflecting its availability for structural, neuromuscular, and signalling functions.

Why it matters: Calcium is essential for bone integrity, nerve conduction, muscle contraction, and intracellular signalling pathways.

Functional significance: Linked to parathyroid regulation, vitamin D status, bone metabolism, and cellular signalling mechanisms.

Common patterns: Vitamin D imbalance + parathyroid dysregulation -> altered calcium homeostasis affecting bone and cellular function.

Your result in this scan: Calcium appears within the analyser's expected range in this scan.

Iron

Moderately reduced

Trace Element

Reference range: 1.151 - 1.847 | **Value:** 0.584

What this means: Reflects systemic iron status and availability for oxygen transport, enzymatic activity, and mitochondrial function.

Why it matters: Iron is critical for haemoglobin, oxygen delivery, and cellular energy production; imbalance affects fatigue and oxidative stress.

Functional significance: Linked to haem synthesis, absorption, inflammation, liver storage, and red blood cell turnover.

Common patterns: Inflammation + poor absorption or excess intake -> altered iron balance impacting oxygen transport and redox state.

Your result in this scan: Suggests reduced iron availability, potentially leading to impaired oxygen delivery and fatigue.

Zinc

Moderately reduced

Trace Element

Reference range: 1.143 - 1.989 | **Value:** 0.888

What this means: Represents zinc status, a critical cofactor in enzymatic activity, immune regulation, and cellular repair.

Why it matters: Zinc supports immune function, DNA synthesis, hormone regulation, and antioxidant defence.

Functional significance: Linked to digestive absorption, immune signalling, wound healing, and hormonal balance.

Common patterns: Poor diet + malabsorption -> zinc deficiency -> impaired immunity and repair.

Your result in this scan: Suggests reduced zinc availability, potentially impairing immune function and repair processes.

Selenium

Within range

Trace Element

Reference range: 0.847 - 2.045 | **Value:** 0.943

What this means: Reflects selenium availability for antioxidant systems and thyroid hormone metabolism.

Why it matters: Selenium is essential for glutathione peroxidase activity and thyroid hormone conversion.

Functional significance: Linked to thyroid function, antioxidant defence, and immune modulation.

Common patterns: Low intake + oxidative stress -> reduced selenium function -> impaired detox and thyroid activity.

Your result in this scan: Selenium appears within the analyser's expected range in this scan.

Phosphorus

Mildly reduced

Trace Element

Reference range: 1.195 - 2.134 | **Value:** 0.886

What this means: Represents phosphorus status, essential for energy metabolism and cellular structure.

Why it matters: Phosphorus is a key component of ATP, DNA, and cellular membranes.

Functional significance: Linked to kidney function, bone metabolism, and energy production.

Common patterns: Kidney dysfunction + dietary imbalance -> altered phosphorus levels.

Your result in this scan: Suggests reduced phosphorus affecting energy production.

Potassium

Moderately reduced

Trace Element

Reference range: 0.689 - 0.987 | **Value:** 0.441

What this means: Reflects potassium balance influencing cellular electrical activity and fluid balance.

Why it matters: Potassium is critical for nerve conduction, muscle contraction, and cardiovascular stability.

Functional significance: Linked to kidney function, hydration, and electrolyte regulation.

Common patterns: Electrolyte imbalance + renal dysfunction -> potassium disturbance.

Your result in this scan: Suggests deficiency affecting cardiac and neuromuscular function.

Magnesium

Within range

Trace Element

Reference range: 0.568 - 0.992 | **Value:** 0.912

What this means: Represents magnesium status, essential for enzymatic activity and neuromuscular stability.

Why it matters: Magnesium supports ATP production, muscle relaxation, and nervous system balance.

Functional significance: Linked to stress, mitochondrial function, and electrolyte balance.

Common patterns: Stress + poor intake -> magnesium depletion.

Your result in this scan: Magnesium appears within the analyser's expected range in this scan.

Copper

Mildly reduced

Trace Element

Reference range: 0.474 - 0.749 | **Value:** 0.32

What this means: Reflects copper balance involved in redox activity and connective tissue formation.

Why it matters: Copper is essential for iron metabolism and antioxidant enzymes.

Functional significance: Linked to zinc balance and liver function.

Common patterns: Zinc imbalance -> copper dysregulation.

Your result in this scan: Suggests deficiency affecting iron metabolism.

Cobalt

Mildly reduced

Trace Element

Reference range: 2.326 - 5.531 | **Value:** 1.777

What this means: Represents cobalt status, linked to vitamin B12 structure and function.

Why it matters: Cobalt is essential for red blood cell production and neurological function.

Functional significance: Linked to B12 metabolism and gut absorption.

Common patterns: Poor absorption -> B12 deficiency.

Your result in this scan: Suggests deficiency affecting energy and nerves.

Manganese

Within range

Trace Element

Reference range: 0.497 - 0.879 | **Value:** 0.793

What this means: Reflects manganese status for enzymatic and antioxidant functions.

Why it matters: Manganese supports bone formation and metabolic enzymes.

Functional significance: Linked to mitochondrial and antioxidant activity.

Common patterns: Environmental exposure -> imbalance.

Your result in this scan: Manganese appears within the analyser's expected range in this scan.

Iodine

Within range

Trace Element

Reference range: 1.421 - 5.490 | **Value:** 3.709

What this means: Represents iodine availability for thyroid hormone production.

Why it matters: Iodine is essential for T3 and T4 synthesis.

Functional significance: Linked to thyroid function and metabolism.

Common patterns: Low intake -> hypothyroid patterns.

Your result in this scan: Iodine appears within the analyser's expected range in this scan.

Nickel

Within range

Trace Element

Reference range: 2.462 - 5.753 | **Value:** 2.628

What this means: Reflects nickel exposure and trace biological role.

Why it matters: Nickel may influence enzyme activity at trace levels.

Functional significance: Linked to environmental exposure.

Common patterns: Environmental sources -> imbalance.

Your result in this scan: Nickel appears within the analyser's expected range in this scan.

Fluorine

Mildly reduced

Trace Element

Reference range: 1.954 - 4.543 | **Value:** 1.692

What this means: Represents fluorine exposure influencing bone and dental health.

Why it matters: Fluorine contributes to enamel strength but excess may affect tissues.

Functional significance: Linked to environmental intake.

Common patterns: Water intake -> fluorine levels.

Your result in this scan: Suggests low exposure.

Molybdenum

Within range

Trace Element

Reference range: 0.938 - 1.712 | **Value:** 0.999

What this means: Reflects molybdenum status in enzymatic detox pathways.

Why it matters: Molybdenum supports sulphite metabolism and detoxification.

Functional significance: Linked to liver enzymes.

Common patterns: Low intake -> impaired detox.

Your result in this scan: Molybdenum appears within the analyser's expected range in this scan.

Vanadium

Within range

Trace Element

Reference range: 1.019 - 3.721 | **Value:** 3.226

What this means: Represents vanadium involvement in glucose and lipid metabolism.

Why it matters: Vanadium may influence insulin signalling.

Functional significance: Linked to metabolic regulation.

Common patterns: Metabolic dysfunction -> altered signalling.

Your result in this scan: Vanadium appears within the analyser's expected range in this scan.

Tin

Within range

Trace Element

Reference range: 1.023 - 7.627 | **Value:** 1.761

What this means: Reflects exposure to tin and trace biological impact.

Why it matters: Primarily reflects environmental exposure.

Functional significance: Linked to toxicity.

Common patterns: Environmental contamination.

Your result in this scan: Tin appears within the analyser's expected range in this scan.

Silicon

Within range

Trace Element

Reference range: 1.425 - 5.872 | **Value:** 2.5

What this means: Represents silicon role in connective tissue and bone health.

Why it matters: Supports collagen and structural integrity.

Functional significance: Linked to bone metabolism.

Common patterns: Nutritional imbalance.

Your result in this scan: Silicon appears within the analyser's expected range in this scan.

Strontium

Within range

Trace Element

Reference range: 1.142 - 5.862 | **Value:** 4.538

What this means: Represents strontium status related to bone mineralisation and structural integrity.

Why it matters: Strontium supports bone formation and may influence osteoblastic activity and mineral balance.

Functional significance: Linked to calcium metabolism, bone turnover, and skeletal health.

Common patterns: Calcium imbalance + dietary variation -> altered strontium incorporation affecting bone dynamics.

Your result in this scan: Strontium appears within the analyser's expected range in this scan.

Boron

Within range

Trace Element

Reference range: 1.124 - 3.453 | **Value:** 3.09

What this means: Reflects boron-related support for bone-mineral metabolism, steroid hormone balance, membrane signalling, and inflammatory regulation.

Why it matters: Boron may influence calcium and magnesium utilisation, vitamin D dynamics, bone support, and aspects of sex hormone metabolism and inflammatory tone.

Functional significance: Interacts with mineral retention, endocrine balance, bone matrix health, and possibly cognition and inflammatory regulation through membrane-signalling effects.

Common patterns: Low boron may align with weaker bone support, less efficient magnesium-calcium handling, and reduced hormone-mineral synergy across endocrine and inflammatory pathways.

Your result in this scan: Boron appears within the analyser's expected range in this scan.

Vitamin

High priority

Complete system marker table — all markers and all severity grades are shown below.

Flagged markers: 5 | Within range: 5

Vitamin A

Within range

Vitamin

Reference range: 0.346 - 0.401 | **Value:** 0.369

What this means: Reflects retinoid status relevant to epithelial integrity, immune signalling, vision pathways, and gene expression regulation.

Why it matters: Vitamin A is essential for mucosal barrier integrity, immune tolerance, retinal function, and regulation of cellular differentiation.

Functional significance: Interacts with gut integrity, immune response modulation, liver storage, thyroid signalling, and epithelial repair processes.

Common patterns: Low vitamin A weakens gut and respiratory barriers, increasing antigen load and immune activation; also affects skin and hormonal signalling.

Your result in this scan: Vitamin A appears within the analyser's expected range in this scan.

Vitamin B1

Within range

Vitamin

Reference range: 2.124 - 4.192 | **Value:** 2.255

What this means: Reflects thiamine availability required for carbohydrate metabolism, pyruvate dehydrogenase activity, nerve conduction, and mitochondrial ATP production.

Why it matters: Vitamin B1 is essential for converting glucose into usable energy and for maintaining neurological and cardiovascular function under metabolic stress.

Functional significance: Linked to glucose handling, mitochondrial entry of glycolytic products, autonomic function, digestive motility, and high carbohydrate demand states.

Common patterns: Low B1 → impaired pyruvate entry into mitochondria → reduced ATP production and greater lactate-type fatigue; nervous system and digestive motility may also be affected.

Your result in this scan: Vitamin B1 appears within the analyser's expected range in this scan.

Vitamin B2

Mildly reduced

Vitamin

Reference range: 1.549 - 2.213 | **Value:** 1.311

What this means: Reflects riboflavin availability required for FAD/FMN-dependent redox reactions in mitochondrial respiration, detoxification, and fatty acid metabolism.

Why it matters: Vitamin B2 supports ATP production, antioxidant recycling, methylation support, and effective metabolism of fats and other B vitamins.

Functional significance: Interacts with mitochondrial electron transport, glutathione recycling, liver detox pathways, and activation of other nutrient cofactors.

Common patterns: Low B2 → weaker electron transport + reduced glutathione recycling → fatigue and oxidative vulnerability; detoxification and lipid metabolism may also become less efficient.

Your result in this scan: Suggests reduced redox cofactor support, poorer mitochondrial energy flow, weaker antioxidant recycling, or higher nutritional demand.

Vitamin B3

Within range

Vitamin

Reference range: 14.477 - 21.348 | **Value:** 19.999

What this means: Reflects niacin-related status relevant to NAD and NADP-dependent redox chemistry, mitochondrial energy transfer, DNA repair, and lipid metabolism.

Why it matters: Vitamin B3 is foundational to cellular respiration, detoxification redox balance, skin integrity, nervous system function, and healthy handling of fats and carbohydrates.

Functional significance: Closely linked to mitochondrial electron transfer, liver detox demand, glucose-fat metabolism, stress resilience, and repair processes requiring redox capacity.

Common patterns: Low B3 can reduce NAD-dependent energy production and weaken detox redox capacity; this may worsen fatigue, skin sensitivity, mood instability, and poor metabolic resilience.

Your result in this scan: Vitamin B3 appears within the analyser's expected range in this scan.

Vitamin B6

Mildly reduced

Vitamin

Reference range: 0.824 - 1.942 | **Value:** 0.678

What this means: Reflects pyridoxine-dependent cofactor status required for amino acid metabolism, neurotransmitter synthesis, glycogen phosphorylase activity, and homocysteine regulation.

Why it matters: Vitamin B6 is central to neurotransmitter production, protein handling, immune modulation, glycaemic stability, and efficient methylation-transsulphuration balance.

Functional significance: Interfaces with neurotransmitter synthesis, glycogen mobilisation, histamine regulation, homocysteine metabolism, and immune-inflammatory control.

Common patterns: Low B6 → weaker serotonin/GABA synthesis + poorer glycogen handling + altered histamine regulation; this may show as mood instability, PMS-type patterns, allergy tendency, and fatigue.

Your result in this scan: Suggests reduced neurotransmitter and amino acid metabolism support, poorer stress resilience, histamine handling issues, or impaired glycaemic and methylation balance.

Vitamin B12

Mildly reduced

Vitamin

Reference range: 6.428 - 21.396 | **Value:** 4.76

What this means: Reflects cobalamin-related status required for methylation, red blood cell production, myelin integrity, homocysteine regulation, and mitochondrial support.

Why it matters: Vitamin B12 is essential for neurological function, cognitive clarity, methylation chemistry, healthy haematopoiesis, and sustained energy output.

Functional significance: Interacts with intrinsic factor and stomach health, folate metabolism, cobalt-related status, methyl donation, and nerve membrane resilience.

Common patterns: Low stomach acid or intrinsic factor dysfunction can reduce B12 handling; low B12 may drive fatigue, memory issues, neuropathy, elevated homocysteine, and reduced methylation resilience.

Your result in this scan: Suggests impaired methylation support, weaker myelin maintenance, reduced red blood cell formation, gastric-intrinsic-factor issues, or fatigue and neuropathy risk.

Vitamin C

Moderately reduced

Vitamin

Reference range: 4.543 - 5.023 | **Value:** 3.567

What this means: Reflects ascorbate status required for collagen synthesis, antioxidant defence, immune activity, and adrenal support.

Why it matters: Vitamin C is essential for connective tissue repair, immune defence, adrenal resilience, and protection against oxidative stress.

Functional significance: Closely linked to adrenal cortisol dynamics, collagen formation, iron absorption, and redox balance.

Common patterns: Low vitamin C impairs collagen, increases oxidative stress, and weakens adrenal response to stress and immune demand.

Your result in this scan: Suggests reduced antioxidant protection, weaker collagen repair, adrenal strain, or increased susceptibility to infection.

Vitamin D3

Within range

Vitamin

Reference range: 5.327 - 7.109 | **Value:** 6.572

What this means: Reflects vitamin D status relevant to calcium regulation, immune modulation, gene expression, and endocrine signalling.

Why it matters: Vitamin D is essential for bone health, immune balance, hormone regulation, and anti-inflammatory signalling.

Functional significance: Interacts with calcium-magnesium balance, parathyroid hormone, immune tolerance, and endocrine stability.

Common patterns: Low vitamin D contributes to immune dysregulation, bone weakness, and inflammatory patterns.

Your result in this scan: Vitamin D3 appears within the analyser's expected range in this scan.

Vitamin E

Mildly reduced

Vitamin

Reference range: 4.826 - 6.013 | **Value:** 4.405

What this means: Reflects lipid-soluble antioxidant status protecting membranes from oxidative damage.

Why it matters: Vitamin E protects cell membranes, supports cardiovascular health, and reduces oxidative stress in lipid environments.

Functional significance: Interacts with lipid metabolism, oxidative stress, mitochondrial function, and inflammatory signalling.

Common patterns: Low vitamin E increases lipid oxidation and vascular stress.

Your result in this scan: Suggests increased oxidative damage to lipids and membranes.

Vitamin K

Within range

Vitamin

Reference range: 0.717 - 1.486 | **Value:** 1.07

What this means: Reflects vitamin K status involved in blood clotting, calcium regulation, and bone metabolism.

Why it matters: Vitamin K is required for proper calcium placement, preventing vascular calcification and supporting bone health.

Functional significance: Interacts with vitamin D, calcium metabolism, liver function, and coagulation pathways.

Common patterns: Vitamin K imbalance affects bone and vascular calcium deposition.

Your result in this scan: Vitamin K appears within the analyser's expected range in this scan.

Amino Acid

High priority

Complete system marker table — all markers and all severity grades are shown below.

Flagged markers: 5 | Within range: 6

Lysine

Mildly reduced

Amino Acid

Reference range: 0.962 - 1.213 | **Value:** 0.872

What this means: Reflects lysine availability, an essential amino acid critical for protein synthesis, collagen formation, immune modulation, and calcium metabolism.

Why it matters: Lysine supports tissue repair, immune defence, and connective tissue integrity, and is particularly relevant in viral susceptibility and wound healing.

Functional significance: Linked to collagen cross-linking, antibody production, carnitine synthesis, and overall protein turnover.

Common patterns: Low lysine -> reduced collagen integrity + impaired immune response; may co-occur with gut dysfunction or malabsorption.

Your result in this scan: Suggests reduced protein quality or intake, impaired absorption, or increased demand for tissue repair and immune activity.

Tryptophan

Within range

Amino Acid

Reference range: 4.978 - 6.289 | **Value:** 5.071

What this means: Reflects tryptophan availability, a precursor for serotonin, melatonin, and kynurenine pathway metabolites.

Why it matters: Tryptophan is essential for mood regulation, sleep quality, and immune modulation.

Functional significance: Linked to neurotransmitter synthesis, circadian rhythm regulation, and immune-inflammatory balance.

Common patterns: Inflammation shifts tryptophan -> kynurenine pathway, reducing serotonin; stress and gut dysfunction can worsen depletion.

Your result in this scan: Tryptophan appears within the analyser's expected range in this scan.

Phenylalanine

Mildly reduced

Amino Acid

Reference range: 1.928 - 2.491 | **Value:** 1.845

What this means: Reflects phenylalanine availability, a precursor to tyrosine and downstream catecholamines.

Why it matters: Phenylalanine supports dopamine, noradrenaline, and adrenaline production, influencing focus, motivation, and stress response.

Functional significance: Linked to neurotransmitter synthesis, cognitive performance, and stress adaptation.

Common patterns: Low phenylalanine -> low dopamine -> fatigue and poor motivation; stress may increase demand.

Your result in this scan: Suggests reduced catecholamine production, often associated with low motivation, fatigue, or poor focus.

Methionine

Within range

Amino Acid

Reference range: 1.245 - 1.637 | **Value:** 1.337

What this means: Reflects methionine availability, a key methyl donor involved in detoxification and epigenetic regulation.

Why it matters: Methionine is central to methylation, detoxification, and liver function.

Functional significance: Linked to SAMe production, homocysteine cycling, glutathione synthesis, and detox pathways.

Common patterns: Methionine imbalance -> homocysteine dysregulation -> cardiovascular and detox stress.

Your result in this scan: Methionine appears within the analyser's expected range in this scan.

Threonine

Mildly reduced

Amino Acid

Reference range: 1.194 - 1.685 | **Value:** 0.929

What this means: Reflects threonine status, important for protein synthesis and mucosal integrity.

Why it matters: Threonine supports gut lining integrity and immune defence.

Functional significance: Linked to mucin production, gut barrier function, and immune support.

Common patterns: Low threonine -> weakened gut lining -> increased permeability and immune activation.

Your result in this scan: Suggests compromised gut barrier or protein deficiency.

Isoleucine

Within range

Amino Acid

Reference range: 4.582 - 5.657 | **Value:** 5.593

What this means: Reflects isoleucine status, a branched-chain amino acid (BCAA) involved in energy metabolism and muscle repair.

Why it matters: Isoleucine supports glucose uptake and muscle recovery.

Functional significance: Linked to energy production, muscle metabolism, and blood sugar regulation.

Common patterns: Low BCAAs -> fatigue and muscle weakness; metabolic stress increases demand.

Your result in this scan: Isoleucine appears within the analyser's expected range in this scan.

Leucine

Mildly reduced

Amino Acid

Reference range: 6.982 - 9.256 | **Value:** 6.597

What this means: Reflects leucine status, a key BCAA regulating muscle protein synthesis via mTOR signalling.

Why it matters: Leucine is critical for muscle maintenance, repair, and metabolic signalling.

Functional significance: Linked to anabolic signalling, muscle metabolism, and insulin sensitivity.

Common patterns: Low leucine -> reduced muscle maintenance; ageing and stress increase need.

Your result in this scan: Suggests reduced anabolic signalling or muscle repair capacity.

Valine

Mildly reduced

Amino Acid

Reference range: 6.982 - 9.677 | **Value:** 5.696

What this means: Reflects valine status, a BCAA involved in energy provision and muscle metabolism.

Why it matters: Valine supports endurance, muscle repair, and metabolic stability.

Functional significance: Linked to nitrogen balance, muscle metabolism, and CNS function.

Common patterns: Low valine -> fatigue and reduced endurance.

Your result in this scan: Suggests reduced energy availability or muscle support.

Histidine

Within range

Amino Acid

Reference range: 5.113 - 6.258 | **Value:** 5.725

What this means: Reflects histidine status, involved in haemoglobin structure and histamine production.

Why it matters: Histidine supports oxygen transport and immune responses.

Functional significance: Linked to haemoglobin, histamine signalling, and antioxidant systems.

Common patterns: Histidine imbalance -> histamine issues or oxygen transport inefficiency.

Your result in this scan: Histidine appears within the analyser's expected range in this scan.

Arginine

Within range

Amino Acid

Reference range: 1.812 - 2.337 | **Value:** 2.17

What this means: Reflects arginine status, a semi-essential amino acid involved in nitric oxide production and vascular function.

Why it matters: Arginine supports circulation, immune response, and wound healing.

Functional significance: Linked to nitric oxide synthesis, vascular dilation, and immune activity.

Common patterns: Low arginine -> poor circulation and healing.

Your result in this scan: Arginine appears within the analyser's expected range in this scan.

Homocysteine

Within range

Amino Acid

Reference range: 0.983 - 1.265 | **Value:** 1.032

What this means: Reflects homocysteine levels, a key intermediate in methylation and cardiovascular risk pathways.

Why it matters: Elevated homocysteine is associated with cardiovascular risk, impaired methylation, and oxidative stress.

Functional significance: Linked to B-vitamin status, methylation cycles, detoxification, and vascular health.

Common patterns: High homocysteine -> cardiovascular risk + oxidative stress + poor detox.

Your result in this scan: Homocysteine appears within the analyser's expected range in this scan.

Coenzyme

Medium priority

Complete system marker table — all markers and all severity grades are shown below.

Flagged markers: 1 | Within range: 5

Nicotinamide

Moderately reduced

Coenzyme

Reference range: 2.074 - 3.309 | **Value:** 1.218

What this means: Reflects NAD precursor status supporting redox reactions and energy metabolism.

Why it matters: Nicotinamide supports mitochondrial function, DNA repair, and metabolic resilience.

Functional significance: Closely linked to NAD cycles, detoxification, and cellular repair.

Common patterns: Low NAD precursors reduce cellular resilience and increase fatigue.

Your result in this scan: Suggests reduced energy production and repair capacity.

Biotin

Within range

Coenzyme

Reference range: 1.833 - 2.979 | **Value:** 2.356

What this means: Reflects biotin status required for fatty acid metabolism and glucose regulation.

Why it matters: Biotin supports hair, skin, and metabolic enzyme function.

Functional significance: Interacts with carbohydrate metabolism and mitochondrial enzymes.

Common patterns: Low biotin affects hair, skin, and glucose metabolism.

Your result in this scan: Biotin appears within the analyser's expected range in this scan.

Pantothenic acid

Within range

Coenzyme

Reference range: 1.116 - 2.101 | **Value:** 1.5

What this means: Reflects vitamin B5 status required for CoA synthesis and energy metabolism.

Why it matters: Essential for fatty acid metabolism, adrenal function, and energy production.

Functional significance: Interacts with adrenal hormones and mitochondrial pathways.

Common patterns: Low B5 reduces CoA availability and energy production.

Your result in this scan: Pantothenic acid appears within the analyser's expected range in this scan.

Folic acid

Within range

Coenzyme

Reference range: 1.449 - 2.246 | **Value:** 1.641

What this means: Reflects folate status required for methylation and DNA synthesis.

Why it matters: Essential for cell division, red blood cell formation, and methylation balance.

Functional significance: Interacts with B12, homocysteine, and DNA repair.

Common patterns: Low folate increases homocysteine and reduces repair capacity.

Your result in this scan: Folic acid appears within the analyser's expected range in this scan.

Coenzyme Q10

Within range

Coenzyme

Reference range: 0.831 - 1.588 | **Value:** 1.584

What this means: Reflects availability of CoQ10, a key electron transport cofactor in the mitochondrial inner membrane required for oxidative phosphorylation and cellular antioxidant protection.

Why it matters: CoQ10 is essential for ATP generation in high-demand tissues such as heart, brain, muscle, and liver, and it also helps protect membranes from oxidative damage.

Functional significance: Closely linked to mitochondrial respiratory chain efficiency, redox balance, statin-related depletion, cardiac energy demand, and overall fatigue resilience.

Common patterns: Low CoQ10 → poorer mitochondrial ATP production → fatigue, low exercise tolerance, and weaker cardiac-muscular energy support; oxidative stress can further deplete reserves.

Your result in this scan: Coenzyme Q10 appears within the analyser's expected range in this scan.

Glutathione

Within range

Coenzyme

Reference range: 0.726 - 1.281 | **Value:** 0.877

What this means: Reflects glutathione status as a master antioxidant and detox molecule.

Why it matters: Central to detoxification, oxidative stress control, and immune resilience.

Functional significance: Interacts with liver detox, selenium, and redox balance.

Common patterns: Low glutathione increases toxin sensitivity and inflammation.

Your result in this scan: Glutathione appears within the analyser's expected range in this scan.

Essential Fatty Acid

Low priority

Complete system marker table — all markers and all severity grades are shown below.

Flagged markers: 1 | Within range: 3

Linoleic acid

Within range

Essential Fatty Acid

Reference range: 0.642 - 0.985 | **Value:** 0.653

What this means: Reflects linoleic-acid status as a key omega-6 essential fatty acid involved in membrane structure, epidermal barrier integrity, and eicosanoid precursor balance.

Why it matters: Linoleic acid is required for healthy skin-barrier function, cell-membrane composition, and downstream inflammatory signalling. Deficiency can impair barrier resilience, while excess relative to omega-3 pathways can promote inflammatory imbalance.

Functional significance: Linked to dietary fat quality, bile-dependent absorption, membrane fluidity, epidermal ceramide formation, and the broader omega-6 to omega-3 balance shaping inflammatory tone.

Common patterns: Low bile flow or fat malabsorption -> lower linoleic-acid availability and weaker skin-barrier resilience; omega-6 excess without omega-3 balance can intensify inflammatory and metabolic stress.

Your result in this scan: Linoleic acid appears within the analyser's expected range in this scan.

α-Linolenic acid

Within range

Essential Fatty Acid

Reference range: 0.814 - 1.202 | **Value:** 1.133

What this means: Reflects alpha-linolenic-acid availability as a plant-derived omega-3 essential fatty acid supporting membrane fluidity, inflammatory modulation, and precursor input into longer-chain omega-3 pathways.

Why it matters: Alpha-linolenic acid contributes to anti-inflammatory balance, cardiovascular resilience, and membrane health, especially when broader omega-3 status is suboptimal.

Functional significance: Linked to dietary fat quality, desaturase and elongase activity, bile-dependent absorption, oxidative protection, and the overall omega-6 to omega-3 ratio.

Common patterns: Low alpha-linolenic-acid intake or poor absorption -> weaker anti-inflammatory balance; excess omega-6 exposure can further blunt the functional impact of available omega-3 precursors.

Your result in this scan: α-Linolenic acid appears within the analyser's expected range in this scan.

γ-Linolenic acid

Within range

Essential Fatty Acid

Reference range: 0.921 - 1.334 | **Value:** 1.154

What this means: Reflects gamma-linolenic-acid related functional support within omega-6 metabolism, relevant to membrane signalling, skin integrity, and downstream eicosanoid balance.

Why it matters: Gamma-linolenic acid can support anti-inflammatory and barrier-regulating pathways within omega-6 metabolism, particularly where delta-6-desaturase conversion is impaired.

Functional significance: Linked to fatty-acid conversion efficiency, skin-barrier resilience, hormone-related inflammatory balance, and the body's ability to generate more regulatory lipid mediators.

Common patterns: Low conversion efficiency from linoleic acid can reduce gamma-linolenic-acid support and contribute to skin dryness, PMS-type inflammatory patterns, and barrier fragility.

Your result in this scan: γ-Linolenic acid appears within the analyser's expected range in this scan.

Arachidonic acid

Mildly reduced

Essential Fatty Acid

Reference range: 0.661 - 0.808 | **Value:** 0.546

What this means: Reflects arachidonic-acid status as a membrane-bound omega-6 fatty acid involved in cell signalling and the generation of multiple inflammatory and regulatory eicosanoids.

Why it matters: Arachidonic acid is essential to membrane function and signalling, but excessive dominance can amplify inflammatory cascades, pain pathways, and vascular reactivity.

Functional significance: Linked to membrane phospholipids, inflammatory mediator production, immune activation, pain signalling, and the balance between pro-inflammatory and resolving lipid pathways.

Common patterns: Omega-3 deficiency plus high arachidonic-acid dominance -> stronger inflammatory signalling, pain amplification, and vascular reactivity; oxidative stress can worsen downstream tissue effects.

Your result in this scan: May suggest reduced membrane availability for eicosanoid signalling, lower dietary intake, altered absorption, or context-dependent reduced inflammatory drive.

ADHD

Low priority

Complete system marker table — all markers and all severity grades are shown below.

Flagged markers: 1 | Within range: 3

An oxygen-hydroxy phenyl ethanol

Within range

ADHD

Reference range: 1.163 - 2.206 | **Value:** 1.45

What this means: Reflects a source-framework neurochemical proxy that most plausibly points toward catecholamine-related signalling relevant to attention, arousal, and behavioural regulation.

Why it matters: ADHD-like patterns often involve altered catecholamine balance, affecting focus, motivation, impulse regulation, and task persistence.

Functional significance: Best interpreted as a heuristic marker for dopaminergic-noradrenergic tone, not as a validated standalone metabolite diagnosis.

Common patterns: Low dopamine-noradrenaline tone -> inattentive patterns; excessive arousal + poor inhibition -> hyperactive or dysregulated attention patterns.

Your result in this scan: An oxygen-hydroxy phenyl ethanol appears within the analyser's expected range in this scan.

GE neurotransmitters

Mildly reduced

ADHD

Reference range: 0.753 - 0.972 | **Value:** 0.583

What this means: Reflects functional balance of excitatory and inhibitory neurotransmitter systems, particularly the regulation between stimulating drive and calming inhibitory control within neural networks.

Why it matters: Balanced neurotransmission is required for attention, calm focus, behavioural regulation, sleep quality, sensory processing, and stable autonomic tone.

Functional significance: Influenced by amino acid availability, B6 and magnesium status, gut-derived neurotransmitter precursors, methylation capacity, and inflammatory effects on neuronal signalling.

Common patterns: Gut dysbiosis or poor protein digestion → reduced precursor availability; magnesium/B6 deficiency → impaired GABAergic support; chronic stress → excitatory dominance, poor sleep, and reduced attentional stability.

Your result in this scan: Suggests reduced neurotransmitter output, poor precursor availability, nutrient insufficiency, or low neural drive contributing to inattention, fatigue, or low mood.

Vanilloid

Within range

ADHD

Reference range: 0.232 - 0.981 | **Value:** 0.528

What this means: Reflects a source-framework signal potentially related to sensory-reactivity and neuromodulatory processing, relevant to arousal, discomfort sensitivity, and attentional tolerance.

Why it matters: Sensory overreactivity can worsen distractibility, irritability, and difficulty sustaining focus in ADHD-like or neurodevelopmental patterns.

Functional significance: Best interpreted as a proxy for sensory-threshold regulation, stress reactivity, and neural excitability rather than a validated diagnostic analyte.

Common patterns: Sensory overload + poor inhibition + stress -> worsening attention and irritability; sleep loss can magnify the pattern.

Your result in this scan: Vanilloid appears within the analyser's expected range in this scan.

Creatine hormone

Within range

ADHD

Reference range: 0.150 - 0.240 | **Value:** 0.213

What this means: Reflects a source-framework marker most plausibly related to creatine-energy support and cognitive performance rather than a literal hormone construct.

Why it matters: Creatine-related energy buffering is relevant to fast ATP recycling in brain and muscle, potentially influencing working memory, cognitive stamina, and behavioural persistence.

Functional significance: Best interpreted as an energy-support proxy linked to mitochondrial reserve, neural ATP buffering, and cognitive endurance.

Common patterns: Low cellular energy buffering -> reduced mental stamina and task persistence; sleep debt and mitochondrial strain can worsen the presentation.

Your result in this scan: Creatine hormone appears within the analyser's expected range in this scan.

Cognitive function

High priority

Complete system marker table — all markers and all severity grades are shown below.

Flagged markers: 4 | Within range: 4

Observation

Mildly reduced

Cognitive function

Reference range: 7.432 - 9.476 | **Value:** 7.396

What this means: Reflects the capacity to notice, register, and accurately take in external information within the source framework.

Why it matters: Observation underpins attention, learning readiness, environmental awareness, and the quality of subsequent memory and reasoning.

Functional significance: Linked to sustained attention, sensory integration, cognitive alertness, and reduced distraction load.

Common patterns: Sleep loss + inattentiveness + sensory overload can reduce observation; stable attention and lower stress support it.

Your result in this scan: Suggests reduced observational focus, distractibility, or weaker intake of external information.

Memory

Mildly reduced

Cognitive function

Reference range: 6.128 - 7.455 | **Value:** 6.118

What this means: Reflects the capacity for encoding, consolidation, and recall of information within the source cognitive-performance framework.

Why it matters: Memory is central to learning, academic performance, adaptive behaviour, and the ability to integrate past experience into current functioning.

Functional significance: Linked to sleep quality, hippocampal support, attention, methylation status, energy availability, and neuroinflammatory burden.

Common patterns: Poor sleep + inattentive intake + inflammatory fatigue -> weaker memory; energy and attentional support often improve this domain.

Your result in this scan: Suggests weaker encoding or recall, often associated with stress, fatigue, sleep disruption, or poor attentional intake.

Imagination

Within range

Cognitive function

Reference range: 7.143 - 9.215 | **Value:** 8.353

What this means: Reflects the capacity for internal imagery, mental simulation, and flexible generative thinking within the source framework.

Why it matters: Imagination supports creativity, planning, learning, emotional processing, and adaptive problem-solving beyond linear reasoning alone.

Functional significance: Linked to cognitive flexibility, associative thinking, emotional-symbolic processing, and adequate mental energy.

Common patterns: Burnout, anxiety, or overcontrolled cognition can suppress imagination; safety, rest, and curiosity tend to support it.

Your result in this scan: Imagination appears within the analyser's expected range in this scan.

Vanilloid

Within range

Cognitive function

Reference range: 5.361 - 7.644 | **Value:** 6.734

What this means: Reflects a source-framework signal potentially related to sensory-reactivity and neuromodulatory processing, relevant to arousal, discomfort sensitivity, and attentional tolerance.

Why it matters: Sensory overreactivity can worsen distractibility, irritability, and difficulty sustaining focus in ADHD-like or neurodevelopmental patterns.

Functional significance: Best interpreted as a proxy for sensory-threshold regulation, stress reactivity, and neural excitability rather than a validated diagnostic analyte.

Common patterns: Sensory overload + poor inhibition + stress -> worsening attention and irritability; sleep loss can magnify the pattern.

Your result in this scan: Vanilloid appears within the analyser's expected range in this scan.

Analysis and Judgment

Within range

Cognitive function

Reference range: 8.423 - 10.217 | **Value:** 9.627

What this means: Reflects the capacity to evaluate information, compare options, and make reasoned judgments within the source cognitive-performance model.

Why it matters: Analytical judgment is central to decision-making, learning, risk assessment, and adaptive planning.

Functional significance: Linked to executive function, working memory, inhibitory control, attentional stability, and reduced emotional noise during cognition.

Common patterns: Sleep disruption + distractibility + emotional reactivity can impair analysis and judgment; stronger executive function supports it.

Your result in this scan: Analysis and Judgment appears within the analyser's expected range in this scan.

Thinking Ability

Mildly reduced

Cognitive function

Reference range: 6.714 - 8.989 | **Value:** 4.665

What this means: Reflects general cognitive processing capacity, including mental organisation, clarity, and the ability to sustain thought through complex tasks.

Why it matters: Thinking ability underpins learning, comprehension, problem-solving, and functional performance across school and life demands.

Functional significance: Linked to energy availability, attention, sleep, executive organisation, and freedom from inflammatory or stress-related cognitive drag.

Common patterns: Fatigue, sleep loss, and poor attention degrade thinking ability; stable energy and reduced stress improve it.

Your result in this scan: Suggests reduced mental clarity, slowed processing, or lower sustained cognitive capacity.

Resilience

Mildly reduced

Cognitive function

Reference range: 8.248 - 11.818 | **Value:** 6.951

What this means: Reflects the ability to recover, adapt, and remain functional after stress or setback within the source developmental-performance framework.

Why it matters: Resilience influences school functioning, stress recovery, emotional regulation, and long-term adaptive capacity.

Functional significance: Linked to nervous-system flexibility, social support, sleep quality, coping style, and energy reserve.

Common patterns: Sleep debt + chronic stress + low social safety can reduce resilience; restored regulation and support tend to strengthen it.

Your result in this scan: Suggests greater vulnerability to stress, slower recovery, or reduced capacity to bounce back after challenge.

Reasoning Ability

Within range

Cognitive function

Reference range: 5.637 - 7.714 | **Value:** 7.648

What this means: Reflects the capacity to identify relationships, infer meaning, and solve problems using logic and conceptual integration within the source framework.

Why it matters: Reasoning ability is central to academic learning, pattern recognition, planning, and adaptive response to novel situations.

Functional significance: Linked to executive function, working memory, attentional control, cognitive clarity, and reduced interference from stress and fatigue.

Common patterns: Poor sleep + inattentiveness + mental fatigue can suppress reasoning ability; stronger working memory and executive control enhance it.

Your result in this scan: Reasoning Ability appears within the analyser's expected range in this scan.

Growth and development

High priority

Complete system marker table — all markers and all severity grades are shown below.

Flagged markers: 4 | Within range: 2

Height and body mass index

Mildly reduced

Growth and development

Reference range: 4.439 - 5.847 | **Value:** 4.089

What this means: Represents the relationship between height and body mass index (BMI) as a proportional indicator of overall body composition and weight status.

Why it matters: This index provides a general assessment of whether body weight is appropriate relative to height, forming a baseline indicator of metabolic and nutritional status.

Functional significance: Linked to BMI classification, adiposity, energy balance, and long-term metabolic risk.

Common patterns: Chronic caloric imbalance + low activity or excess intake -> altered BMI relationship -> metabolic implications.

Your result in this scan: Suggests lower body mass relative to height, potentially reflecting underweight status, reduced energy reserves, or low muscle/fat mass.

Height bust index

Within range

Growth and development

Reference range: 4.126 - 5.457 | **Value:** 4.76

What this means: Reflects the proportional relationship between height and chest (thoracic) circumference, indicating upper body structural development.

Why it matters: This index provides insight into thoracic development, respiratory capacity potential, and upper-body musculoskeletal proportion.

Functional significance: Linked to skeletal growth, lung capacity, posture, and developmental patterns.

Common patterns: Growth patterns + posture + physical activity influence thoracic proportion.

Your result in this scan: Height bust index appears within the analyser's expected range in this scan.

Height sitting height index

Mildly reduced

Growth and development

Reference range: 3.346 - 4.215 | **Value:** 2.679

What this means: Represents the ratio between total height and sitting height, reflecting the proportional distribution between torso and lower limbs.

Why it matters: This index provides insight into skeletal growth patterns and developmental proportionality.

Functional significance: Linked to genetic growth patterns, spinal development, and lower limb length.

Common patterns: Growth distribution patterns -> altered body proportions.

Your result in this scan: Suggests disproportion between torso and limb length, potentially reflecting developmental variation.

BMI index

Mildly reduced

Growth and development

Reference range: 4.679 - 5.648 | **Value:** 3.743

What this means: This marker is one of the scan indicators used by the analyser to build a functional picture of this section.

Why it matters: It is usually more useful when interpreted alongside the wider pattern than in isolation.

Functional significance: This marker may help highlight relative weakness, excess, burden, or imbalance in the system being assessed.

Common patterns: Its value is usually strongest when interpreted as part of a clustered pattern rather than in isolation.

Your result in this scan: A mildly reduced result may suggest softer functional support or reserve in this area.

Grip and back muscle index

Within range

Growth and development

Reference range: 5.421 - 6.714 | **Value:** 6.32

What this means: Reflects functional muscular strength, particularly in grip and back musculature, as an indicator of physical capacity.

Why it matters: Muscle strength correlates with metabolic health, functional independence, and overall resilience.

Functional significance: Linked to muscle mass, neuromuscular coordination, activity level, and protein status.

Common patterns: Sedentary lifestyle + low protein intake -> reduced muscle strength.

Your result in this scan: Grip and back muscle index appears within the analyser's expected range in this scan.

Vital capacity index

Mildly reduced

Growth and development

Reference range: 4.512 - 6.382 | **Value:** 2.924

What this means: Represents lung capacity relative to body size, reflecting respiratory efficiency and physical fitness.

Why it matters: Vital capacity is a key indicator of pulmonary function, oxygen delivery, and endurance capacity.

Functional significance: Linked to lung function, cardiovascular fitness, posture, and muscular conditioning.

Common patterns: Low activity + poor posture -> reduced lung capacity; training improves it.

Your result in this scan: Suggests reduced respiratory capacity and potential limitation in oxygen delivery.

Lecithin

Medium priority

Complete system marker table — all markers and all severity grades are shown below.

Flagged markers: 2 | Within range: 4

Phospholipid index

Mildly reduced

Lecithin

Reference range: 3.654 - 4.478 | **Value:** 2.526

What this means: Represents the status of phospholipids as key structural components of cell membranes and lipoproteins.

Why it matters: Phospholipids are essential for membrane integrity, cellular signalling, and lipid transport.

Functional significance: Linked to liver function, lipid metabolism, and membrane fluidity.

Common patterns: Liver dysfunction + oxidative stress -> membrane instability.

Your result in this scan: Suggests reduced membrane integrity and impaired lipid transport.

Sphingolipid index

Within range

Lecithin

Reference range: 3.121 - 3.853 | **Value:** 3.419

What this means: Reflects sphingolipid balance involved in membrane structure and cellular signalling.

Why it matters: Sphingolipids regulate cell signalling, apoptosis, and membrane stability.

Functional significance: Linked to lipid metabolism and inflammatory signalling.

Common patterns: Inflammation -> sphingolipid dysregulation.

Your result in this scan: Sphingolipid index appears within the analyser's expected range in this scan.

PH

Within range

Lecithin

Reference range: 3.341 - 4.214 | **Value:** 3.457

What this means: Reflects overall acid-base tendency within the source framework, functioning as a broad indicator of systemic buffering balance rather than a standalone compartment-specific laboratory pH value.

Why it matters: Acid-base balance matters because enzyme activity, mitochondrial output, mineral stability, respiratory compensation, and renal buffering all depend on an adequately regulated internal pH environment.

Functional significance: Linked to renal acid excretion, respiratory CO2 handling, dietary acid load, mineral buffering, mitochondrial metabolism, and inflammatory or catabolic states that shift the internal acid-base

terrain.

Common patterns: High metabolic stress + poor renal buffering + altered ventilation -> pH imbalance; chronic inflammatory load, tissue hypoxia, and catabolic states can all destabilise acid-base resilience.

Your result in this scan: PH appears within the analyser's expected range in this scan.

Lecithin index

Within range

Lecithin

Reference range: 4.365 - 5.142 | **Value:** 4.743

What this means: Represents lecithin levels contributing to lipid transport and membrane structure.

Why it matters: Lecithin supports lipid metabolism, liver function, and bile production.

Functional significance: Linked to phosphatidylcholine, liver health, and fat digestion.

Common patterns: Liver stress -> lecithin depletion.

Your result in this scan: Lecithin index appears within the analyser's expected range in this scan.

Brain phospholipid index

Mildly reduced

Lecithin

Reference range: 4.225 - 5.713 | **Value:** 3.839

What this means: Reflects phospholipid availability and membrane integrity within brain tissue, influencing synaptic signalling, myelin stability, receptor function, and neuronal resilience.

Why it matters: Phospholipids are fundamental to cell membrane fluidity, neurotransmission, myelin support, and effective communication across neural networks.

Functional significance: Depends on choline status, essential fatty acid balance, hepatic phospholipid metabolism, oxidative protection, and efficient membrane turnover.

Common patterns: Low phospholipid support + oxidative stress → poorer membrane fluidity → reduced neurotransmission and memory performance; liver dysfunction can also impair phospholipid synthesis and trafficking.

Your result in this scan: Suggests reduced membrane integrity, impaired synaptic signalling, weaker myelin support, or suboptimal availability of choline and structural lipids.

Liposomal index

Within range

Lecithin

Reference range: 3.112 - 4.081 | **Value:** 4.021

What this means: Reflects liposomal or phospholipid-vesicle style transport capacity, indicating how effectively lipid-based carriers may support membrane delivery, absorption, and intracellular nutrient trafficking.

Why it matters: Liposomal transport concepts are relevant to membrane interaction, nutrient-delivery efficiency, and the broader integrity of phospholipid-based transport environments.

Functional significance: Linked to phospholipid sufficiency, bile-dependent fat handling, membrane fluidity, intracellular transport efficiency, and liver-lipid processing.

Common patterns: Low phospholipid sufficiency plus poor bile flow can reduce lipid-carrier efficiency and membrane-delivery support; this may coexist with broader membrane and hepatic lipid dysfunction.

Your result in this scan: Liposomal index appears within the analyser's expected range in this scan.

Fatty acid

Medium priority

Complete system marker table — all markers and all severity grades are shown below.

Flagged markers: 2 | Within range: 2

Saturated fatty acid index

Within range

Fatty acid

Reference range: 1.334 - 2.407 | **Value:** 2.113

What this means: Represents the proportion of saturated fats within overall lipid balance.

Why it matters: Excess saturated fats can influence inflammation and metabolic health.

Functional significance: Linked to diet, lipid metabolism, and insulin sensitivity.

Common patterns: High processed fat intake -> metabolic dysfunction.

Your result in this scan: Saturated fatty acid index appears within the analyser's expected range in this scan.

Unsaturated fatty acid index

Mildly reduced

Fatty acid

Reference range: 2.224 - 3.153 | **Value:** 1.85

What this means: Reflects levels of unsaturated fats supporting membrane fluidity and metabolic function.

Why it matters: Unsaturated fats are essential for cardiovascular and metabolic health.

Functional significance: Linked to omega balance and lipid metabolism.

Common patterns: Diet influences fatty acid balance.

Your result in this scan: Suggests reduced beneficial fats.

Essential fatty acid index

Within range

Fatty acid

Reference range: 2.144 - 3.238 | **Value:** 3.053

What this means: Represents availability of essential fatty acids required for cellular function.

Why it matters: Essential fatty acids support inflammation control and membrane health.

Functional significance: Linked to omega-3 and omega-6 balance.

Common patterns: Poor diet -> deficiency.

Your result in this scan: Essential fatty acid index appears within the analyser's expected range in this scan.

Triglyceride index

Mildly reduced

Fatty acid

Reference range: 3.361 - 4.644 | **Value:** 2.493

What this means: Reflects circulating triglyceride levels and lipid storage status.

Why it matters: Triglycerides are a key marker of metabolic health and energy storage.

Functional significance: Linked to diet, insulin, and liver function.

Common patterns: Insulin resistance -> high triglycerides.

Your result in this scan: Suggests low lipid storage.

TECHNICAL REFERENCE

Appendix

This section contains supporting technical material and summary reference tables used in the report.

Section	Marker	Range	Value	Status
Trace Element	Calcium	1.219 - 3.021	1.948	Within range
Trace Element	Iron	1.151 - 1.847	0.584	Moderately reduced
Trace Element	Zinc	1.143 - 1.989	0.888	Moderately reduced
Trace Element	Selenium	0.847 - 2.045	0.943	Within range
Trace Element	Phosphorus	1.195 - 2.134	0.886	Mildly reduced
Trace Element	Potassium	0.689 - 0.987	0.441	Moderately reduced
Trace Element	Magnesium	0.568 - 0.992	0.912	Within range
Trace Element	Copper	0.474 - 0.749	0.32	Mildly reduced
Trace Element	Cobalt	2.326 - 5.531	1.777	Mildly reduced
Trace Element	Manganese	0.497 - 0.879	0.793	Within range
Trace Element	Iodine	1.421 - 5.490	3.709	Within range
Trace Element	Nickel	2.462 - 5.753	2.628	Within range
Trace Element	Fluorine	1.954 - 4.543	1.692	Mildly reduced
Trace Element	Molybdenum	0.938 - 1.712	0.999	Within range
Trace Element	Vanadium	1.019 - 3.721	3.226	Within range
Trace Element	Tin	1.023 - 7.627	1.761	Within range
Trace Element	Silicon	1.425 - 5.872	2.5	Within range
Trace Element	Strontium	1.142 - 5.862	4.538	Within range
Trace Element	Boron	1.124 - 3.453	3.09	Within range
Vitamin	Vitamin A	0.346 - 0.401	0.369	Within range
Vitamin	Vitamin B1	2.124 - 4.192	2.255	Within range
Vitamin	Vitamin B2	1.549 - 2.213	1.311	Mildly reduced
Vitamin	Vitamin B3	14.477 - 21.348	19.999	Within range
Vitamin	Vitamin B6	0.824 - 1.942	0.678	Mildly reduced
Vitamin	Vitamin B12	6.428 - 21.396	4.76	Mildly reduced
Vitamin	Vitamin C	4.543 - 5.023	3.567	Moderately reduced
Vitamin	Vitamin D3	5.327 - 7.109	6.572	Within range

Section	Marker	Range	Value	Status
Vitamin	Vitamin E	4.826 - 6.013	4.405	Mildly reduced
Vitamin	Vitamin K	0.717 - 1.486	1.07	Within range
Amino Acid	Lysine	0.962 - 1.213	0.872	Mildly reduced
Amino Acid	Tryptophan	4.978 - 6.289	5.071	Within range
Amino Acid	Phenylalanine	1.928 - 2.491	1.845	Mildly reduced
Amino Acid	Methionine	1.245 - 1.637	1.337	Within range
Amino Acid	Threonine	1.194 - 1.685	0.929	Mildly reduced
Amino Acid	Isoleucine	4.582 - 5.657	5.593	Within range
Amino Acid	Leucine	6.982 - 9.256	6.597	Mildly reduced
Amino Acid	Valine	6.982 - 9.677	5.696	Mildly reduced
Amino Acid	Histidine	5.113 - 6.258	5.725	Within range
Amino Acid	Arginine	1.812 - 2.337	2.17	Within range
Amino Acid	Homocysteine	0.983 - 1.265	1.032	Within range
Coenzyme	Nicotinamide	2.074 - 3.309	1.218	Moderately reduced
Coenzyme	Biotin	1.833 - 2.979	2.356	Within range
Coenzyme	Pantothenic acid	1.116 - 2.101	1.5	Within range
Coenzyme	Folic acid	1.449 - 2.246	1.641	Within range
Coenzyme	Coenzyme Q10	0.831 - 1.588	1.584	Within range
Coenzyme	Glutathione	0.726 - 1.281	0.877	Within range
Essential Fatty Acid	Linoleic acid	0.642 - 0.985	0.653	Within range
Essential Fatty Acid	α-Linolenic acid	0.814 - 1.202	1.133	Within range
Essential Fatty Acid	γ-Linolenic acid	0.921 - 1.334	1.154	Within range
Essential Fatty Acid	Arachidonic acid	0.661 - 0.808	0.546	Mildly reduced
ADHD	An oxygen-hydroxy phenyl ethanol	1.163 - 2.206	1.45	Within range
ADHD	GE neurotransmitters	0.753 - 0.972	0.583	Mildly reduced
ADHD	Vanilloid	0.232 - 0.981	0.528	Within range
ADHD	Creatine hormone	0.150 - 0.240	0.213	Within range
Cognitive function	Observation	7.432 - 9.476	7.396	Mildly reduced
Cognitive function	Memory	6.128 - 7.455	6.118	Mildly reduced
Cognitive function	Imagination	7.143 - 9.215	8.353	Within range
Cognitive function	Vanilloid	5.361 - 7.644	6.734	Within range
Cognitive function	Analysis and Judgment	8.423 - 10.217	9.627	Within range
Cognitive function	Thinking Ability	6.714 - 8.989	4.665	Mildly reduced

Section	Marker	Range	Value	Status
Cognitive function	Resilience	8.248 - 11.818	6.951	Mildly reduced
Cognitive function	Reasoning Ability	5.637 - 7.714	7.648	Within range
Growth and development	Height and body mass index	4.439 - 5.847	4.089	Mildly reduced
Growth and development	Height bust index	4.126 - 5.457	4.76	Within range
Growth and development	Height sitting height index	3.346 - 4.215	2.679	Mildly reduced
Growth and development	BMI index	4.679 - 5.648	3.743	Mildly reduced
Growth and development	Grip and back muscle index	5.421 - 6.714	6.32	Within range
Growth and development	Vital capacity index	4.512 - 6.382	2.924	Mildly reduced
Lecithin	Phospholipid index	3.654 - 4.478	2.526	Mildly reduced
Lecithin	Sphingolipid index	3.121 - 3.853	3.419	Within range
Lecithin	PH	3.341 - 4.214	3.457	Within range
Lecithin	Lecithin index	4.365 - 5.142	4.743	Within range
Lecithin	Brain phospholipid index	4.225 - 5.713	3.839	Mildly reduced
Lecithin	Liposomal index	3.112 - 4.081	4.021	Within range
Fatty acid	Saturated fatty acid index	1.334 - 2.407	2.113	Within range
Fatty acid	Unsaturated fatty acid index	2.224 - 3.153	1.85	Mildly reduced
Fatty acid	Essential fatty acid index	2.144 - 3.238	3.053	Within range
Fatty acid	Triglyceride index	3.361 - 4.644	2.493	Mildly reduced

These scan findings are best interpreted alongside symptoms, history, and appropriate clinical assessment.