

NaturalApproaches.com

# Personalised Wellness Scan Report

Wellness screening summary

## Patient details

**Name:** Test Test

**Sex:** Male

**Age:** 58

**Height:** 184.0 cm

**Weight:** 90.0 kg

**Profile:** Male

**Scan time:** 21/03/2026 21:03

This report is for wellness education and discussion only. It is not a medical diagnosis.

**Report generated:** 12/04/2026 11:21 BST

Eze Chiana
care@naturalapproaches.com
https://www.naturalapproaches.com

NAVIGATION

# Table of contents

Cover

At a glance

Scan score overview

Priority category review

Body composition analysis

Clinical recommendations

Recommended support plan

Complete product recommendation mapping

Complete system marker tables

Appendix

OVERVIEW

# At a glance

This section summarises the dominant patterns in the scan, the wider downstream themes they may be influencing, and the immediate priorities for follow-up.

## Clinical snapshot

The scan is most suggestive of reduced digestive and absorptive efficiency, with likely downstream effects on repair, resilience, and wider function. There is also a secondary picture of toxic-load pressure and barrier-related inflammatory stress.

## Practitioner overview

Taken together, the findings point most strongly to reduced digestive and absorptive efficiency. Priority should be given to digestive support, absorptive support, and mucosal repair.

## Primary pattern

**Absorption and assimilation strain**

This pattern points to reduced digestive and absorptive efficiency, most evident across digestive support, absorptive support, mucosal repair, and foundational nutrient repletion.

**Follow-up focus:** digestive support, absorptive support, mucosal repair, foundational nutrient repletion

## Contributing patterns

- **Toxic burden and detoxification strain** — Toxic-load pressure, most evident across exposure review, detoxification support, and antioxidant reserve.
- **Inflammatory and barrier dysfunction pattern** — Barrier-related inflammatory stress, most evident across barrier repair, immune-modulatory support, and trigger reduction.
- **Neurocognitive support pattern** — Neurocognitive support needs, most evident across neurocognitive nutrient support, membrane and phospholipid support, and trace element repletion.

## Immediate priorities for follow-up

- Strengthen digestive function to improve breakdown, tolerance, and nutrient availability.
- Support intestinal absorption and assimilation of key nutrients.
- Prioritise mucosal repair and gut-lining resilience.
- Replete foundational nutrients to support repair, energy, and resilience.

- Review environmental and lifestyle exposures contributing to toxic load.

OVERVIEW

# Scan score overview

This section provides a weighted overview of the scan, helping organise attention across major body systems before moving into more detailed interpretation.




## Overall Scan Score

Relatively Poor

- Very Good (85-100)
- General (70-84)
- Relatively Poor (55-69)
- Very Bad (0-54)

## How to read this dashboard

This dashboard provides a weighted clinical overview of the scan. The overall score and body-system cards are not based on a simple count of flagged markers. They are derived from the severity, distribution, and clustering of findings across related sections, with greater weighting given to patterns that may carry broader functional significance.

Each body-system card may include more than one underlying scan section, which is why the totals shown here may differ from the marker counts shown later on individual section pages. In general, lower scores suggest a more active pattern worth reviewing in follow-up, while higher scores suggest a more stable pattern on this dashboard view.

These scores are designed to help organise clinical attention. They are not a diagnosis and should always be interpreted alongside symptoms, history, and appropriate clinical assessment.

### Renal & Fluid Balance




3 flagged · 1 within range

Includes: Kidney function

### Digestive




9 flagged · 10 within range

Includes: Gastrointestinal function, Large intestine function, Liver function, Gallbladder function

### Cardiovascular




11 flagged · 18 within range

Includes: Cardiovascular function, Circulation (pulse dynamics), Blood lipids

### Endocrine & Hormonal




13 flagged · 14 within range

Includes: Endocrine regulation, Thyroid function, Prostate health, Male sexual health, Male fertility, Male hormonal balance

### Skin & Barrier




12 flagged · 6 within range

Includes: Skin integrity, Ocular surface

### Musculoskeletal




25 flagged · 27 within range

Includes: Bone health, Bone mineral density, Joint inflammation, Bone remodelling, Connective tissue, Structural circulation

### Metabolic




24 flagged · 47 within range

Includes: Pancreatic function, Blood sugar regulation, Trace mineral status, Vitamin status, Amino acid status, Coenzyme status, Essential fatty acid status, Metabolic function, Lipid transport support, Fat metabolism

### Immune & Inflammatory




14 flagged · 25 within range

Includes: Immune function, Toxin load, Heavy metal burden, Allergy response, Immune resilience

### Respiratory




2 flagged · 6 within range

Includes: Lung function, Respiratory function

### Neurocognitive




7 flagged · 15 within range

Includes: Brain function, Cognitive function

### Other




1 flagged · 4 within range

## Priority overview

**Prostate**

**Priority:** High priority

**Flagged markers:** 3

**Section score:** 37

**Gastrointestinal Function**

**Priority:** High priority

**Flagged markers:** 3

**Section score:** 47

**Obesity**

**Priority:** High priority

**Flagged markers:** 4

**Section score:** 49

**Large Intestine Function**

**Priority:** High priority

**Flagged markers:** 3

**Section score:** 50

**Kidney Function**

**Priority:** Medium priority

**Flagged markers:** 3

**Section score:** 56

**Blood lipids**

**Priority:** Medium priority

**Flagged markers:** 5

**Section score:** 60

## Top sections for follow-up

- Prostate
- Gastrointestinal Function
- Obesity
- Large Intestine Function
- Kidney Function
- Blood lipids

RECOMMENDATIONS

# Clinical recommendations

This section translates the scan findings into high-level clinical priorities, showing which functional areas appear most relevant for follow-up support.

## Nutrient repletion

Priority should be given to restoring nutrient sufficiency and absorptive capacity.

**Why this was suggested:** Vitamin contains 6 flagged markers, led by Vitamin C, Vitamin E, Vitamin A.

## Inflammation and barrier support

Priority should be given to reducing inflammatory load while strengthening barrier integrity.

**Why this was suggested:** Allergy contains 6 flagged markers, led by Hair dye allergy index, Injection allergy index, Chemical Sensitivity.

## Visual and microvascular support

Support should focus on visual and microvascular function.

**Why this was suggested:** Visual and microvascular support contains 6 flagged markers, led by Edema, Collagen eye wrinkle, Dark circles.

## Connective tissue and structural support

Support should focus on connective-tissue integrity and structural resilience.

**Why this was suggested:** Connective tissue and structural support contains 6 flagged markers, led by Digestive system, Immune system, Motion systems.

## Metabolic regulation and weight balance

Support should focus on metabolic balance and weight regulation.

**Why this was suggested:** Metabolic regulation and weight balance contains 4 flagged markers, led by Abnormal lipid metabolism coefficient, Brown adipose tissue abnormalities coefficient, Triglyceride content of abnormal coefficient.

RECOMMENDATIONS

# Recommended support plan

This section organises suggested support options into a staged plan, combining foundational and targeted measures based on the dominant scan patterns.

## Absorption and assimilation strain Support Plan

This plan focuses on addressing the primary functional pattern, starting with foundational support before moving into targeted optimisation.

This plan is organised around reduced digestive and absorptive efficiency, with additional support for toxic-load pressure and barrier-related inflammatory stress.

## Phase 1 — Foundation

This phase focuses on core digestive, nutritional, and repair support.

### D-FENZ

VitalHealth

Core

Suggested because related findings appear across Cardiovascular and Cerebrovascular, Gastrointestinal Function, Large Intestine Function, and Liver Function.

### GLUTATION PLUS+

VitalHealth

Suggested because related findings appear across Heavy Metal, Liver Function, Skin, and Thyroid.

### V-CURCUMAX

VitalHealth

Suggested because related findings appear across Allergy, Bone Disease, Gallbladder Function, and Liver Function.

### V-DAILY

VitalHealth

Suggested because related findings appear across Amino Acid, Basic Physical Quality, Bone Growth Index, and Bone Mineral Density.

### V-OMEGA 3

VitalHealth

Suggested because related findings appear across Bone Growth Index, Brain Nerve, Cardiovascular and Cerebrovascular, and Essential Fatty Acid.

## Phase 2 — Targeted Support

This phase adds more specific support aligned with the strongest secondary patterns.

### V-ORGANEX

VitalHealth

Core

Suggested because related findings appear across Allergy, Basic Physical Quality, Gallbladder Function, and Liver Function.

REFERENCE MAPPING

# Complete product recommendation mapping

This section lists all products identified through the scan analysis. It includes full mapping across systems and markers to support practitioner review, product selection, and fulfilment planning.

| Product | Source | Systems | Key markers |
|---|---|---|---|
| **D-FENZ** | VitalHealth | Cardiovascular and Cerebrovascular, Gastrointestinal Function, Large Intestine Function, Liver Function | Vascular Resistance, Stroke Volume, LV Pump Power, Coronary Perfusion Pressure, Gastric Absorption Function Coefficient |
| **GLUTATION PLUS+** | VitalHealth | Heavy Metal, Liver Function, Skin, Thyroid | Aluminum, Cadmium, FT4, Lead, Liver Fat Content |
| **V-CURCUMAX** | VitalHealth | Allergy, Bone Disease, Gallbladder Function, Liver Function | Adhesion Degree of Shoulder Muscle, Bilirubin (DBIL), Bone Hyperplasia Coefficient, Chemical Sensitivity, Degree of Cervical Calcification |
| **V-DAILY** | VitalHealth | Amino Acid, Basic Physical Quality, Bone Growth Index, Bone Mineral Density | Adrenal glands Index, Biotin, Bone alkaline phosphatase, Coenzyme Q10, Collagen eye wrinkle |
| **V-OMEGA 3** | VitalHealth | Bone Growth Index, Brain Nerve, Cardiovascular and Cerebrovascular, Essential Fatty Acid | Arachidonic acid, Bone Hyperplasia Coefficient, Bone alkaline phosphatase, Collagen eye wrinkle, Copper |
| **V-ORGANEX** | VitalHealth | Allergy, Basic Physical Quality, Gallbladder Function, Liver Function | Bilirubin (DBIL), Chemical Sensitivity, Dust allergy index, Hair dye allergy index, Hypoxia |

SECTION OVERVIEW

# Priority category review

This section presents the most relevant functional categories identified in the scan. These categories have been prioritised based on the number, severity, and clustering of non-optimal markers.

Each category page summarises the strongest findings in that category and highlights the most clinically relevant marker explanations to support efficient interpretation and follow-up planning.

## Categories included in this section

- Prostate
- Gastrointestinal Function
- Obesity
- Large Intestine Function
- Kidney Function
- Blood lipids

# Prostate

High priority

## Priority interpretation

Prostate contains a meaningful cluster of non-optimal markers, enough to warrant interpretation alongside symptoms, history, and related sections.

Moderately and severely abnormal markers are prioritised here for faster clinical reading.

Flagged markers: 3 | Within range: 0 | Section score: 37

## Top findings

- Prostatitis Syndrome is moderately elevated
- Degree of Prostatic Hyperplasia is moderately elevated
- Degree of Prostatic Calcification is mildly elevated

## Priority marker explanations

### Prostatitis Syndrome

**Result:** 6.11 | **Reference range:** 2.213 - 2.717 | Moderately elevated

**What this marker means:** Reflects an inflammatory-irritative pattern involving the prostate, suggesting pain, congestion, urinary discomfort, or immune-reactive stress in the gland and surrounding pelvic tissues.

**Why it matters:** Prostatitis-type patterns matter because they can affect urinary ease, pelvic comfort, sexual function, quality of life, and the chronic inflammatory burden of the male pelvis.

**Functional significance:** Linked to glandular inflammation, pelvic-floor tension, microbial or sterile inflammatory triggers, urinary irritation, congestion, and autonomic stress patterns.

**Common patterns:** Pelvic tension + inflammation + poor drainage can maintain prostatitis-type patterns even without clear infection; stress and autonomic guarding often amplify symptoms.

**Your result in this scan:** Indicates increased prostatic inflammation, pelvic discomfort, urinary irritation, congestion, or chronic pain-type burden in the male pelvic axis.

### Degree of Prostatic Hyperplasia

**Result:** 5.03 | **Reference range:** 1.023 - 3.230 | **Moderately elevated**

**What this marker means:** Reflects the extent of enlargement or proliferative change in prostate tissue, suggesting altered growth signalling and reduced ease of urinary outflow.

**Why it matters:** Prostatic enlargement matters because it can impair urinary flow, increase nocturia, create incomplete bladder emptying, and signal longer-term androgen-inflammatory dysregulation.

**Functional significance:** Linked to age-related androgen balance, DHT signalling, inflammatory burden, pelvic congestion, insulin-metabolic factors, and local tissue-remodelling tendencies.

**Common patterns:** Ageing + inflammatory load + androgen-metabolic signalling can drive prostatic hyperplasia; pelvic congestion and poor circulation may worsen urinary symptoms.

**Your result in this scan:** Indicates increased prostatic enlargement, tissue proliferation, or growth-related urinary obstruction risk.

# Gastrointestinal Function

High priority

## Priority interpretation

Gastrointestinal Function is one of the more active sections in this scan, with multiple flagged markers combining into a broader functional pattern.

Moderately and severely abnormal markers are prioritised here for faster clinical reading.

Flagged markers: 3 | Within range: 2 | Section score: 47

## Top findings

- Small Intestine Peristalsis Function Coefficient is moderately reduced
- Small Intestine Absorption Function Coefficient is moderately reduced
- Gastric Absorption Function Coefficient is moderately reduced

## Priority marker explanations

### Small Intestine Motility

Software marker: Small Intestine Peristalsis Function Coefficient

**Result:** 126.057 | **Reference range:** 133.437 - 140.476 | Moderately reduced

**What this marker means:** Reflects coordinated small bowel motility, including segmental mixing and propulsive movement required for digestion, nutrient contact, and microbial regulation.

**Why it matters:** Adequate small intestinal motility prevents stasis, supports enzyme-substrate contact, and helps protect against bacterial overgrowth. Dysfunction is strongly linked to bloating, malabsorption, and SIBO-type presentations.

**Functional significance:** Depends on enteric nervous system signalling, vagal modulation, electrolyte balance, thyroid status, and smooth muscle responsiveness. It strongly influences transit time and absorptive efficiency.

**Common patterns:** Reduced small bowel motility → stasis → bacterial overgrowth → fermentation gases, mucosal irritation, and malabsorption; hypermotility can shorten nutrient contact time and contribute to loose stools and depletion.

**Your result in this scan:** Suggests stagnation, impaired migrating motor complex activity, autonomic dysfunction, hypothyroid tendency, or post-inflammatory motility suppression.

## Small Intestine Absorption

Software marker: Small Intestine Absorption Function Coefficient

**Result:** 2.82 | **Reference range:** 3.572 - 6.483 | **Moderately reduced**

**What this marker means:** Reflects absorptive efficiency of the small intestine across macronutrients, micronutrients, electrolytes, and water-soluble compounds, indicating the functional quality of villous and mucosal transport processes.

**Why it matters:** The small intestine is the main site of nutrient assimilation. Impaired function can produce broad downstream effects including fatigue, deficiency states, immune dysregulation, poor repair, and endocrine compromise.

**Functional significance:** Depends on villous integrity, brush border enzyme activity, bile and pancreatic support, mucosal permeability control, and healthy interaction with the microbiome.

**Common patterns:** Low bile or pancreatic enzyme output → incomplete breakdown → poor absorption; intestinal inflammation or permeability → nutrient deficiency + immune activation; malabsorption can secondarily impair thyroid, adrenal, mitochondrial, and detox pathways.

**Your result in this scan:** Suggests malabsorption, villous or brush-border compromise, inflammatory mucosal injury, pancreatic-biliary insufficiency, or dysbiosis-related impairment.

## Gastric Absorption Function Coefficient

**Result:** 29.734 | **Reference range:** 34.367 - 35.642 | **Moderately reduced**

**What this marker means:** Reflects the stomach's limited but important absorptive and transport role, including handling of water, alcohol, certain medications, and support functions linked to intrinsic factor-dependent B12 assimilation downstream.

**Why it matters:** Although the stomach is not the primary absorptive organ, impaired gastric mucosal integrity can signal wider dysfunction in secretory and barrier roles that compromise nutrient handling and increase inflammatory sensitivity.

**Functional significance:** Closely linked to parietal cell function, intrinsic factor production, mucosal integrity, acid secretion, and the quality of transition into duodenal digestion and absorption.

**Common patterns:** Parietal cell dysfunction → low acid and low intrinsic factor → impaired B12 status → fatigue, neuropathy, and methylation compromise; mucosal injury can also increase sensitivity to reflux, gastritis, and upper GI inflammation.

**Your result in this scan:** Suggests gastric mucosal compromise, reduced intrinsic factor/parietal cell support, poor secretory integrity, or inflammatory impairment of the gastric lining.

# Obesity

High priority

## Priority interpretation

Obesity is one of the more active sections in this scan, with multiple flagged markers combining into a broader functional pattern.

Moderately and severely abnormal markers are prioritised here for faster clinical reading.

Flagged markers: 4 | Within range: 1 | Section score: 49

## Top findings

- Triglyceride content of abnormal coefficient is moderately elevated
- Brown adipose tissue abnormalities coefficient is moderately reduced
- Abnormal lipid metabolism coefficient is moderately reduced

## Priority marker explanations

### Triglyceride content of abnormal coefficient

**Result:** 5.222 | **Reference range:** 1.341 - 1.991 | Moderately elevated

**What this marker means:** Reflects mitochondrial coupling efficiency in ATP production versus heat generation.

**Why it matters:** Determines fatigue, performance, and metabolic flexibility.

**Functional significance:** Linked to mitochondrial health, oxidative phosphorylation, and nutrient utilisation.

**Common patterns:** Mitochondrial dysfunction -> low efficiency.

**Your result in this scan:** Indicates efficient ATP generation.

### Brown adipose tissue abnormalities coefficient

**Result:** 1.858 | **Reference range:** 2.791 - 4.202 | **Moderately reduced**

**What this marker means:** Reflects functional irregularity in brown adipose tissue, the thermogenic fat compartment involved in heat production, metabolic flexibility, and energy expenditure.

**Why it matters:** Brown adipose tissue helps dissipate energy through thermogenesis rather than storage, so altered function may contribute to weight gain tendency, cold intolerance, reduced metabolic responsiveness, and impaired glucose-lipid handling.

**Functional significance:** Linked to mitochondrial uncoupling activity, thyroid signalling, sympathetic tone, insulin sensitivity, cold-adaptation capacity, and overall metabolic rate regulation.

**Common patterns:** Low thyroid activity + low sympathetic thermogenic drive + mitochondrial inefficiency -> weaker brown-fat activation -> lower energy expenditure and greater obesity susceptibility.

**Your result in this scan:** Suggests more favourable brown-fat thermogenic activity and less evidence of impaired adaptive energy expenditure.

### Abnormal lipid metabolism coefficient

**Result:** 0.89 | **Reference range:** 1.992 - 3.713 | **Moderately reduced**

**What this marker means:** Reflects baseline mitochondrial energy expenditure under resting conditions.

**Why it matters:** Determines energy balance, weight regulation, and metabolic efficiency.

**Functional significance:** Linked to thyroid hormones, mitochondrial density, and lean mass.

**Common patterns:** Hypothyroid + low muscle mass -> low BMR.

**Your result in this scan:** Suggests metabolic slowdown and fatigue.

# Large Intestine Function

High priority

## Priority interpretation

Large Intestine Function contains a meaningful cluster of non-optimal markers, enough to warrant interpretation alongside symptoms, history, and related sections.

Moderately and severely abnormal markers are prioritised here for faster clinical reading.

Flagged markers: 3 | Within range: 1 | Section score: 50

## Top findings

- Large intestine peristalsis function coefficient is moderately reduced
- Intestinal bacteria coefficient is moderately reduced
- Intraluminal pressure coefficient is mildly elevated

## Priority marker explanations

### Colon Motility

Software marker: Large intestine peristalsis function coefficient

**Result:** 3.088 | **Reference range:** 4.572 - 6.483 | Moderately reduced

**What this marker means:** Reflects colonic motility and propulsive function, including transit dynamics required for stool formation, waste movement, and timely elimination.

**Why it matters:** Healthy colonic transit reduces toxin reabsorption, supports microbiome balance, and prevents stagnation-associated constipation, bloating, and inflammatory burden.

**Functional significance:** Influenced by hydration status, magnesium and electrolyte balance, autonomic tone, fibre-fermentation dynamics, thyroid function, and pelvic floor or smooth muscle coordination.

**Common patterns:** Slow colonic transit → prolonged toxin and oestrogen metabolite contact → reabsorption and hepatic burden; constipation also shifts microbial ecology toward dysbiosis and inflammatory metabolite production.

**Your result in this scan:** Suggests sluggish transit, constipation tendency, autonomic suppression, dehydration, low magnesium status, or hypothyroid-associated bowel slowing.

## Gut Microbiome

Software marker: Intestinal bacteria coefficient

**Result:** 0.807 | **Reference range:** 1.734 - 2.621 | **Moderately reduced**

**What this marker means:** Reflects the functional state and relative balance of intestinal microbial ecology, including commensal support, fermentative activity, and microbial pressure within the gut lumen.

**Why it matters:** The intestinal microbiome shapes digestion, vitamin production, immune education, barrier integrity, oestrogen metabolism, and inflammatory signalling throughout the body.

**Functional significance:** Interacts directly with bile acids, fibre fermentation, mucosal immunity, short-chain fatty acid generation, endotoxin production, and enterohepatic circulation.

**Common patterns:** Dysbiosis → impaired barrier function → endotoxin translocation → hepatic and immune activation; altered beta-glucuronidase activity → oestrogen recirculation; microbial imbalance also influences mood, skin, and metabolic signalling.

**Your result in this scan:** Suggests reduced microbial diversity or beneficial flora support, often associated with antimicrobial exposure, low fibre intake, poor bile flow, or chronic stress.

# Kidney Function

Medium priority

## Priority interpretation

Kidney Function contains a meaningful cluster of non-optimal markers, enough to warrant interpretation alongside symptoms, history, and related sections.

Moderately and severely abnormal markers are prioritised here for faster clinical reading.

Flagged markers: 3 | Within range: 1 | Section score: 56

## Top findings

- Uric acid Index is moderately elevated
- Urobilinogen Index is mildly elevated
- Proteinuria Index is mildly elevated

## Priority marker explanations

### Uric acid Index

**Result:** 2.689 | **Reference range:** 1.435 - 1.987 | Moderately elevated

**What this marker means:** Reflects uric-acid burden within renal handling, indicating purine metabolism load and the kidney's role in excreting this metabolic end product.

**Why it matters:** Uric acid is clinically relevant to oxidative stress, metabolic syndrome, renal burden, vascular risk, and crystal-related inflammatory tendencies when clearance is impaired or production is excessive.

**Functional significance:** Linked to purine turnover, fructose burden, renal excretion efficiency, insulin resistance, dehydration status, and inflammatory-metabolic terrain.

**Common patterns:** Fructose excess + insulin resistance + dehydration + renal under-clearance -> higher uric-acid burden; this may overlap with metabolic syndrome and inflammatory joint patterns.

**Your result in this scan:** Indicates increased uric-acid burden, potentially associated with higher purine load, reduced renal clearance, insulin resistance, dehydration, or crystal-forming inflammatory risk.

# Blood lipids

Medium priority

## Priority interpretation

Blood lipids is one of the more active sections in this scan, with multiple flagged markers combining into a broader functional pattern.

Moderately and severely abnormal markers are prioritised here for faster clinical reading.

Flagged markers: 5 | Within range: 2 | Section score: 60

## Top findings

- Blood viscosity is moderately elevated
- Triglyceride(TG) is mildly elevated
- Neutral fat(MB) is mildly elevated

## Priority marker explanations

### Blood viscosity

**Result:** 5.233 | **Reference range:** 4.131 - 4.562 | Moderately elevated

**What this marker means:** Reflects thickness of blood affecting circulation and oxygen delivery.

**Why it matters:** High viscosity impairs circulation and increases cardiovascular risk.

**Functional significance:** Linked to hydration, lipids, inflammation.

**Common patterns:** High viscosity increases cardiovascular burden.

**Your result in this scan:** Indicates sluggish circulation and vascular strain.

DEDICATED SUMMARY

# Body Composition & Metabolic Profile

This section summarises QRMA-derived body composition and metabolic profile indicators, including hydration, lean tissue, adiposity, and weight-related measures.

Within range

QRMA category: Element of Human

## Section interpretation

This section summarises QRMA-derived body composition and metabolic profile indicators, including hydration, lean tissue, adiposity, and weight-related measures. Body fat percentage appears broadly reasonable for the profile shown. Where QRMA terminology is unconventional, clinically clearer labels are shown while preserving the original category reference.

- Hydration markers appear broadly balanced, without an obvious fluid-distribution concern on this page.
- Body fat percentage sits in a broadly reasonable range for a male profile.
- Body mass index is above the conventional range and may reflect increased metabolic load.
- Current weight sits materially above the QRMA target-weight estimate, so weight-related metrics should be interpreted alongside metabolic and inflammatory findings.
- Abdominal fat ratio is relatively high, which may align with greater central metabolic burden.

This section presents body-composition findings in a dedicated summary layout so the scan is easier to review clinically.

## Fluid & Hydration

| | |
|---|---|
| **Intracellular fluid (QRMA)** | 21.6 |
| **Extracellular fluid (QRMA)** | 11.1 |

## Lean Tissue & Structural Mass

| | |
|---|---|
| **Protein mass (QRMA)** | 8.58 |
| **Mineral mass (QRMA)** | 27 |
| **Muscle mass / muscle volume** | 41.3Kg |
| **Body cell mass (BCM)** | 30.26 Kg |
| **Muscle mass change target** | 0 Kg |

## Adiposity & Fat Distribution

| | |
|---|---|
| **Body fat mass (QRMA)** | 21.7 |
| **Body fat mass** | 21.7Kg |
| **Body fat percentage** | 24.1% |
| **Abdominal fat ratio** | 1.04 |
| **Fat mass change target** | 17.2 Kg |

## Weight & Metabolic Indicators

| | |
|---|---|
| **Height** | 184cm |
| **Body weight** | 90kg |
| **Obesity degree of body (QRMA)** | 124% |
| **Body mass index (BMI)** | 26.6 kg/m² |
| **Basal metabolic rate (BMR)** | 1825 kcal/day |
| **Target weight** | 72.8 Kg |
| **Weight change target** | 72.8 Kg |

## Other QRMA Body Composition Indices

| | |
|---|---|
| **Body shape / form assessment** | 63.3 Standard |

TECHNICAL REFERENCE

# Complete system marker tables

This section includes every parsed system and every marker across the report, including those within range, as the full explained reference layer.

# Cardiovascular and Cerebrovascular

High priority

Complete system marker table — all markers and all severity grades are shown below.

Flagged markers: 4 | Within range: 11

| Marker | Reference range | Value | Status | Visual | Meaning |
|---|---|---|---|---|---|
| Blood Viscosity | 48.264 - 65.371 | 53.723 | Within range | | Reflects the thickness and flow resistance of blood, indicating how efficiently it can move through vessels and deliver oxygen and nutrients to tissues. |
| Cholesterol Crystal | 56.749 - 67.522 | 66.455 | Within range | | Reflects altered cholesterol handling and a tendency toward less soluble or more atherogenic lipid states within circulation. |
| Blood Fat | 0.481 - 1.043 | 0.902 | Within range | | Reflects the circulating fat burden within blood, indicating how effectively lipids are being transported, cleared, and metabolically handled. |
| Vascular Resistance | 0.327 - 0.937 | 1.226 | Mildly elevated | | Reflects the resistance encountered by blood as it moves through the vascular tree, indicating vessel tone, flexibility, and microvascular openness. |
| Vascular Elasticity | 1.672 - 1.978 | 1.934 | Within range | | Reflects the compliance and recoil capacity of blood vessels, indicating how well vascular connective tissue can expand and return without excessive stiffness. |
| Myocardial Blood Demand | 0.192 - 0.412 | 0.407 | Within range | | Reflects the oxygen and nutrient demand placed on cardiac muscle tissue during its workload. |
| Myocardial Perfusion | 4.832 - 5.147 | 5.132 | Within range | | Reflects the adequacy of blood flow reaching cardiac muscle tissue, indicating how well the myocardium is being nourished and oxygenated. |

| Marker | Reference range | Value | Status | Visual | Meaning |
|---|---|---|---|---|---|
| Myocardial Oxygen Consumption | 3.321 - 4.244 | 3.382 | **Within range** | | Reflects the oxygen utilisation requirement of heart muscle tissue during its pumping activity. |
| Stroke Volume | 1.338 - 1.672 | 1.29 | **Mildly reduced** | | Reflects the volume of blood ejected by the heart with each beat, indicating pumping efficiency and circulatory support capacity. |
| LV Ejection Impedance | 0.669 - 1.544 | 1.456 | **Within range** | | Reflects the resistance faced by the left ventricle when ejecting blood into the arterial system, indicating afterload-related strain. |
| LV Pump Power | 1.554 - 1.988 | 0.991 | **Moderately reduced** | | Reflects the effective mechanical power generated by the left ventricle to sustain forward blood flow into systemic circulation. |
| Coronary Artery Elasticity | 1.553 - 2.187 | 1.66 | **Within range** | | Reflects the structural flexibility and recoil quality of coronary arterial tissue supplying the heart muscle. |
| Coronary Perfusion Pressure | 11.719 - 18.418 | 19.312 | **Moderately elevated** |  | Reflects the pressure gradient driving blood flow through the coronary circulation to nourish heart muscle tissue. |
| Cerebral Blood Vessel Elasticity | 0.708 - 1.942 | 1.021 | **Within range** | | Reflects the compliance and recoil of cerebral vessels, indicating how well connective tissue within brain circulation can adapt to flow and pressure changes. |
| Brain Tissue Blood Supply Status | 6.138 - 21.396 | 20.196 | **Within range** | | Reflects the adequacy of cerebral perfusion reaching brain tissue, indicating how well oxygen and nutrients are being delivered to support neural metabolism. |

# Gastrointestinal Function

High priority

Complete system marker table — all markers and all severity grades are shown below.

Flagged markers: 3 | Within range: 2

| Marker | Reference range | Value | Status | Visual | Meaning |
|---|---|---|---|---|---|
| Pepsin Secretion Coefficient | 59.847 - 65.234 | 59.915 | Within range | | Reflects gastric proteolytic capacity through pepsin secretion and activation, indicating how effectively the stomach can initiate protein digestion after adequate hydrochloric acid exposure. |
| Gastric Peristalsis Function Coefficient | 58.425 - 61.213 | 58.71 | Within range | | Reflects the motor efficiency of the stomach, including churning, mixing, and regulated emptying of gastric contents into the duodenum. |
| Gastric Absorption Function Coefficient | 34.367 - 35.642 | 29.734 | Moderately reduced | | Reflects the stomach's limited but important absorptive and transport role, including handling of water, alcohol, certain medications, and support functions linked to intrinsic factor-dependent B12 assimilation downstream. |
| Small Intestine Motility | 133.437 - 140.476 | 126.057 | Moderately reduced |  | Reflects coordinated small bowel motility, including segmental mixing and propulsive movement required for digestion, nutrient contact, and microbial regulation. |
| Small Intestine Absorption | 3.572 - 6.483 | 2.82 | Moderately reduced |  | Reflects absorptive efficiency of the small intestine across macronutrients, micronutrients, electrolytes, and water-soluble compounds, indicating the functional quality of villous and mucosal transport processes. |

# Large Intestine Function

**Medium priority**

Complete system marker table — all markers and all severity grades are shown below.

Flagged markers: 3 | Within range: 1

| Marker | Reference range | Value | Status | Visual | Meaning |
|---|---|---|---|---|---|
| Colon Motility | 4.572 - 6.483 | 3.088 | **Moderately reduced** | | Reflects colonic motility and propulsive function, including transit dynamics required for stool formation, waste movement, and timely elimination. |
| Colon Absorption | 2.946 - 3.815 | 2.952 | **Within range** | | Reflects absorptive activity of the colon, particularly water, electrolytes, and microbiome-derived metabolites such as short-chain fatty acids. |
| Gut Microbiome | 1.734 - 2.621 | 0.807 | **Moderately reduced** | | Reflects the functional state and relative balance of intestinal microbial ecology, including commensal support, fermentative activity, and microbial pressure within the gut lumen. |
| Intraluminal pressure coefficient | 1.173 - 2.297 | 2.639 | **Mildly elevated** | | Reflects pressure dynamics within the intestinal lumen, influenced by gas production, motility coordination, stool transit, muscular tone, and bowel distension patterns. |

# Liver Function

Low priority

Complete system marker table — all markers and all severity grades are shown below.

Flagged markers: 1 | Within range: 4

| Marker | Reference range | Value | Status | Visual | Meaning |
|---|---|---|---|---|---|
| Protein Metabolism | 116.34 - 220.621 | 116.569 | Within range | | Represents the liver's role in processing, transforming, and regulating amino acid metabolism and nitrogen balance. |
| Energy Production Function | 0.713 - 0.992 | 0.881 | Within range | | Reflects hepatic contribution to systemic energy production through glucose regulation, glycogen storage, and metabolic processing. |
| Detoxification Function | 0.202 - 0.991 | 0.977 | Within range | | Represents the liver's integrated capacity to process and eliminate endogenous and exogenous toxins. |
| Bile Secretion Function | 0.432 - 0.826 | 0.717 | Within range | | Reflects the liver's ability to produce and regulate bile flow for digestion and elimination. |
| Liver Fat Content | 0.097 - 0.419 | 0.636 | Moderately elevated | | Represents the degree of lipid accumulation within hepatic tissue. |

# Gallbladder Function

Medium priority

Complete system marker table — all markers and all severity grades are shown below.

Flagged markers: 2 | Within range: 3

| Marker | Reference range | Value | Status | Visual | Meaning |
|---|---|---|---|---|---|
| Serum Globulin (A/G) | 126 - 159 | 126.767 | Within range | | Reflects the albumin-to-globulin relationship within the source gallbladder-bile-flow cluster, functioning as a broad marker of protein balance, hepatic synthetic context, immune burden, and chronic inflammatory load rather than a bile-specific analyte alone. |
| Total Bilirubin (TBIL) | 0.232 - 0.686 | 0.542 | Within range | | Reflects total bilirubin burden across conjugated and unconjugated fractions, indicating the liver's handling of haem breakdown products and the effectiveness of bile-dependent excretion pathways. |
| Alkaline Phosphatase (ALP) | 0.082 - 0.342 | 0.316 | Within range | | Reflects alkaline phosphatase activity, a marker especially relevant to cholestatic stress, bile-duct irritation, and hepatobiliary drainage resistance, while also requiring distinction from bone-derived ALP sources. |
| Serum Total Bile Acid (TBA | 0.317 - 0.695 | 0.704 | Mildly elevated | | Reflects the circulating burden of bile acids, indicating how effectively the liver synthesises, secretes, recycles, and clears bile within the enterohepatic system. |
| Bilirubin (DBIL) | 0.218 - 0.549 | 0.56 | Mildly elevated | | Reflects direct (conjugated) bilirubin, indicating the fraction that has already been processed by the liver and is awaiting effective excretion through bile pathways. |

# Pancreatic Function

**Within range**

Complete system marker table — all markers and all severity grades are shown below.

Flagged markers: 0 | Within range: 3

| Marker | Reference range | Value | Status | Visual | Meaning |
|---|---|---|---|---|---|
| Insulin | 2.845 - 4.017 | 3.036 | **Within range** | | Reflects insulin output within the pancreatic endocrine axis, indicating beta-cell signalling involved in glucose uptake, blood-sugar regulation, and anabolic storage physiology. |
| Pancreatic Polypeptide (PP) | 3.210 - 6.854 | 3.26 | **Within range** | | Reflects pancreatic polypeptide signalling, a regulatory peptide involved in pancreatic secretory balance, gastrointestinal signalling, appetite modulation, and autonomic-pancreatic coordination. |
| Glucagon | 2.412 - 2.974 | 2.666 | **Within range** | | Reflects glucagon output within the pancreatic alpha-cell axis, indicating counter-regulatory signalling that raises blood glucose during fasting or increased energy demand. |

# Kidney Function

**Medium priority**

Complete system marker table — all markers and all severity grades are shown below.

Flagged markers: 3 | Within range: 1

| Marker | Reference range | Value | Status | Visual | Meaning |
|---|---|---|---|---|---|
| Urobilinogen Index | 2.762 - 5.424 | 6.215 | **Mildly elevated** | | Reflects the burden of urobilinogen within the renal-urinary context, indicating downstream handling of bilirubin metabolism and possible interaction between liver processing, gut conversion, and urinary excretion. |
| Uric acid Index | 1.435 - 1.987 | 2.689 | **Moderately elevated** | | Reflects uric-acid burden within renal handling, indicating purine metabolism load and the kidney's role in excreting this metabolic end product. |
| BUN | 4.725 - 8.631 | 4.766 | **Within range** | | Reflects the blood burden of urea nitrogen as a marker of protein-waste handling across the liver-kidney axis, especially renal excretion of nitrogenous end products. |
| Proteinuria Index | 1.571 - 4.079 | 4.761 | **Mildly elevated** | | Reflects urinary protein leakage, indicating compromise of the renal filtration barrier or altered glomerular permeability within the kidney axis. |

# Lung Function

Low priority

Complete system marker table — all markers and all severity grades are shown below.

Flagged markers: 1 | Within range: 3

| Marker | Reference range | Value | Status | Visual | Meaning |
|---|---|---|---|---|---|
| Vital Capacity | 3348 - 3529 | 3489 | Within range | | Reflects the maximal volume of air that can be exhaled after a full inhalation, providing a functional measure of lung expansion, ventilatory reserve, and chest-wall-respiratory muscle capacity. |
| Total Lung Capacity | 4301 - 4782 | 4286 | Mildly reduced |  | Reflects the total volume of air the lungs can contain at maximal inflation, indicating the overall size and expansion capacity of the pulmonary-thoracic system. |
| Airway Resistance | 1.374 - 1.709 | 1.621 | Within range | | Reflects the degree of opposition to airflow through the bronchial tree, indicating how easily air can move through the conducting airways. |
| Arterial Oxygen Content | 17.903 - 21.012 | 20.78 | Within range | | Reflects a source-label gas-exchange marker best interpreted cautiously as arterial blood-gas balance, with the PaCO2 label pointing more directly to carbon-dioxide handling and ventilatory adequacy than to oxygen content. |

# Brain Nerve

Low priority

Complete system marker table — all markers and all severity grades are shown below.

Flagged markers: 1 | Within range: 4

| Marker | Reference range | Value | Status | Visual | Meaning |
|---|---|---|---|---|---|
| Brain Tissue Blood Supply | 143.37 - 210.81 | 168.036 | Within range | | Reflects adequacy of cerebral perfusion to brain tissue, including microvascular blood flow, oxygen delivery, and nutrient availability required for neuronal metabolism. |
| Cerebral Arteriosclerosis | 0.103 - 0.642 | 0.196 | Within range | | Reflects structural and functional hardening of cerebral vessels, indicating reduced arterial compliance, impaired endothelial regulation, and higher cerebrovascular resistance. |
| Cranial Nerve Function | 0.253 - 0.659 | 0.292 | Within range | | Reflects the functional integrity of cranial nerve signalling involved in sensory processing, autonomic regulation, facial motor function, swallowing, vestibular balance, and vagal communication. |
| Sentiment Index | 0.109 - 0.351 | 0.262 | Within range | | Reflects functional emotional regulation capacity, incorporating neurochemical balance, limbic system tone, and stress-related affective responsiveness. |
| Memory | 0.442 - 0.817 | 0.379 | Mildly reduced | | Reflects the capacity for encoding, consolidation, and recall of information within the source cognitive-performance framework. |

# Bone Disease

Within range

Complete system marker table — all markers and all severity grades are shown below.

Flagged markers: 0 | Within range: 7

| Marker | Reference range | Value | Status | Visual | Meaning |
|---|---|---|---|---|---|
| Name: Test Test | | Sex: Male | Unclear | | This marker is one of the scan indicators used by the analyser to build a functional picture of this section. |
| Tin | | Figure: 184cm, 90kg | Unclear | | Reflects exposure to tin and trace biological impact. |
| Figure: 184cm, 90kg | | Testing Time: 21/03/2026 21:03 | Unclear | | This marker is one of the scan indicators used by the analyser to build a functional picture of this section. |
| Lumbar Fiber Protruding Dimension | | No Direction | Unclear | | Indicates structural deformation of intervertebral disc fibres within the lumbar spine, reflecting loss of disc integrity and altered load distribution. |
| Adhesion Degree of Shoulder Muscle | | < u 0.2 | Unclear | | Reflects fibrotic adhesion and reduced glide between muscle and fascial layers in the shoulder complex. |
| Limbs Circulation Limit | | + | Unclear | | Represents reduced peripheral blood flow affecting limbs, impacting oxygen and nutrient delivery. |
| Age of Ligament | | 10%-40% | Unclear | | Reflects the functional ageing state of ligament tissue, indicating the degree of connective tissue resilience, elasticity, and structural wear within supportive joint tissues. |

# Bone Mineral Density

**Medium priority**

Complete system marker table — all markers and all severity grades are shown below.

Flagged markers: 2 | Within range: 3

| Marker | Reference range | Value | Status | Visual | Meaning |
|---|---|---|---|---|---|
| Osteoclast Coefficient | 86.73 - 180.97 | 151.19 | **Within range** | | Reflects activity of osteoclasts responsible for bone resorption in bone remodelling. |
| Amount of Calcium Loss | 0.209 - 0.751 | 0.469 | **Within range** | | Represents net calcium loss from bone stores, reflecting mineral depletion. |
| Degree of Bone Hyperplasia | 0.046 - 0.167 | 0.389 | **Mildly elevated** | | Reflects the extent of reactive or excessive bone overgrowth within skeletal tissue. |
| Degree of Osteoporosis | 0.124 - 0.453 | 0.528 | **Moderately elevated** |  | Reflects extent of bone mineral density reduction and structural fragility. |
| Bone Mineral Density | 0.796 - 0.433 | 0.435 | **Within range** | | Reflects the mineral density and load-bearing strength of bone tissue within the structural system. |

# Rheumatoid Bone Disease

**Medium priority**

Complete system marker table — all markers and all severity grades are shown below.

Flagged markers: 3 | Within range: 2

| Marker | Reference range | Value | Status | Visual | Meaning |
|---|---|---|---|---|---|
| Degree of Cervical Calcification | 421 - 490 | 502.068 | **Mildly elevated** | | Represents calcification deposits in cervical spine tissues, indicating chronic degeneration. |
| Degree of Lumbar Calcification | 4.326 - 7.531 | 7.352 | **Within range** | | Reflects calcific degeneration within lumbar structures affecting mobility and nerve function. |
| Bone Hyperplasia Coefficient | 2.954 - 5.543 | 6 | **Mildly elevated** | | Reflects a quantified tendency toward reactive bone proliferation and compensatory overgrowth within skeletal structures. |
| Osteoporosis Coefficient | 2.019 - 4.721 | 4.705 | **Within range** | | Composite marker indicating overall bone fragility and density decline. |
| Rheumatism Coefficient | 4.023 - 11.627 | 16.843 | **Moderately elevated** | | Reflects connective tissue and joint inflammatory burden associated with rheumatic-type patterns affecting movement and periarticular tissues. |

# Bone Growth Index

**Medium priority**

Complete system marker table — all markers and all severity grades are shown below.

Flagged markers: 3 | Within range: 2

| Marker | Reference range | Value | Status | Visual | Meaning |
|---|---|---|---|---|---|
| Bone alkaline phosphatase | 0.433 - 0.796 | 0.399 | **Mildly reduced** | | Reflects osteoblastic activity and bone formation dynamics, indicating the degree of active skeletal matrix building and mineralisation support. |
| Osteocalcin | 0.525 - 0.817 | 0.508 | **Mildly reduced** | | Reflects osteoblast-derived bone matrix activity and mineralisation signalling, with additional relevance to glucose and metabolic cross-talk. |
| Status of long bone healing | 0.713 - 0.992 | 0.606 | **Mildly reduced** | | Reflects the efficiency of repair and recovery processes within long bone tissue after structural stress or injury. |
| Short bone cartilage healing situation | 0.202 - 0.991 | 0.362 | **Within range** | | Reflects healing and recovery quality within cartilage and short-bone associated supportive structures. |
| Epiphyseal line | 0.432 - 0.826 | 0.769 | **Within range** | | Reflects growth plate activity and skeletal maturation status. |

# Blood Sugar

**Within range**

Complete system marker table — all markers and all severity grades are shown below.

Flagged markers: 0 | Within range: 3

| Marker | Reference range | Value | Status | Visual | Meaning |
|---|---|---|---|---|---|
| Coefficient of Insulin Secretion | 2.967 - 3.528 | 3.043 | **Within range** | | Reflects the responsiveness of pancreatic beta cells in secreting insulin relative to glucose exposure and metabolic demand. |
| Blood Sugar Coefficient | 2.163 - 7.321 | 3.191 | **Within range** | | Represents the overall balance of circulating glucose levels relative to metabolic demand and regulatory control. |
| Urine Sugar Coefficient | 2.204 - 2.819 | 2.535 | **Within range** | | Reflects the presence of glucose excretion via urine, indicating overflow beyond renal reabsorption thresholds. |

# Trace Element

Medium priority

Complete system marker table — all markers and all severity grades are shown below.

Flagged markers: 3 | Within range: 16

| Marker | Reference range | Value | Status | Visual | Meaning |
|---|---|---|---|---|---|
| Calcium | 1.219 - 3.021 | 1.518 | Within range | | Represents calcium status within the body, reflecting its availability for structural, neuromuscular, and signalling functions. |
| Iron | 1.151 - 1.847 | 0.743 | Mildly reduced | | Reflects systemic iron status and availability for oxygen transport, enzymatic activity, and mitochondrial function. |
| Zinc | 1.143 - 1.989 | 1.336 | Within range | | Represents zinc status, a critical cofactor in enzymatic activity, immune regulation, and cellular repair. |
| Selenium | 0.847 - 2.045 | 1.096 | Within range | | Reflects selenium availability for antioxidant systems and thyroid hormone metabolism. |
| Phosphorus | 1.195 - 2.134 | 2.087 | Within range | | Represents phosphorus status, essential for energy metabolism and cellular structure. |
| Potassium | 0.689 - 0.987 | 0.977 | Within range | | Reflects potassium balance influencing cellular electrical activity and fluid balance. |
| Magnesium | 0.568 - 0.992 | 0.761 | Within range | | Represents magnesium status, essential for enzymatic activity and neuromuscular stability. |
| Copper | 0.474 - 0.749 | 0.23 | Moderately reduced | | Reflects copper balance involved in redox activity and connective tissue formation. |
| Cobalt | 2.326 - 5.531 | 2.493 | Within range | | Represents cobalt status, linked to vitamin B12 structure and function. |
| Manganese | 0.497 - 0.879 | 0.732 | Within range | | Reflects manganese status for enzymatic and antioxidant functions. |
| Iodine | 1.421 - 5.490 | 5.311 | Within range | | Represents iodine availability for thyroid hormone production. |

| Marker | Reference range | Value | Status | Visual | Meaning |
|---|---|---|---|---|---|
| Nickel | 2.462 - 5.753 | 2.481 | Within range | | Reflects nickel exposure and trace biological role. |
| Fluorine | 1.954 - 4.543 | 2.019 | Within range | | Represents fluorine exposure influencing bone and dental health. |
| Molybdenum | 0.938 - 1.712 | 1.43 | Within range | | Reflects molybdenum status in enzymatic detox pathways. |
| Vanadium | 1.019 - 3.721 | 2.638 | Within range | | Represents vanadium involvement in glucose and lipid metabolism. |
| Tin | 1.023 - 7.627 | 7.159 | Within range | | Reflects exposure to tin and trace biological impact. |
| Silicon | 1.425 - 5.872 | 1.367 | Mildly reduced | | Represents silicon role in connective tissue and bone health. |
| Strontium | 1.142 - 5.862 | 3.508 | Within range | | Represents strontium status related to bone mineralisation and structural integrity. |
| Boron | 1.124 - 3.453 | 2.537 | Within range | | Reflects boron-related support for bone-mineral metabolism, steroid hormone balance, membrane signalling, and inflammatory regulation. |

# Vitamin

High priority

Complete system marker table — all markers and all severity grades are shown below.

Flagged markers: 6 | Within range: 4

| Marker | Reference range | Value | Status | Visual | Meaning |
|---|---|---|---|---|---|
| Vitamin A | 0.346 - 0.401 | 0.338 | Mildly reduced | | Reflects retinoid status relevant to epithelial integrity, immune signalling, vision pathways, and gene expression regulation. |
| Vitamin B1 | 2.124 - 4.192 | 1.679 | Mildly reduced | | Reflects thiamine availability required for carbohydrate metabolism, pyruvate dehydrogenase activity, nerve conduction, and mitochondrial ATP production. |
| Vitamin B2 | 1.549 - 2.213 | 1.549 | Within range | | Reflects riboflavin availability required for FAD/FMN-dependent redox reactions in mitochondrial respiration, detoxification, and fatty acid metabolism. |
| Vitamin B3 | 14.477 - 21.348 | 14.72 | Within range | | Reflects niacin-related status relevant to NAD and NADP-dependent redox chemistry, mitochondrial energy transfer, DNA repair, and lipid metabolism. |
| Vitamin B6 | 0.824 - 1.942 | 1.616 | Within range | | Reflects pyridoxine-dependent cofactor status required for amino acid metabolism, neurotransmitter synthesis, glycogen phosphorylase activity, and homocysteine regulation. |
| Vitamin B12 | 6.428 - 21.396 | 6.962 | Within range | | Reflects cobalamin-related status required for methylation, red blood cell production, myelin integrity, homocysteine regulation, and mitochondrial support. |
| Vitamin C | 4.543 - 5.023 | 3.55 | Moderately reduced | | Reflects ascorbate status required for collagen synthesis, antioxidant defence, immune activity, and adrenal support. |

| Marker | Reference range | Value | Status | Visual | Meaning |
|---|---|---|---|---|---|
| Vitamin D3 | 5.327 - 7.109 | 5.259 | **Mildly reduced** | | Reflects vitamin D status relevant to calcium regulation, immune modulation, gene expression, and endocrine signalling. |
| Vitamin E | 4.826 - 6.013 | 4.11 | **Moderately reduced** | | Reflects lipid-soluble antioxidant status protecting membranes from oxidative damage. |
| Vitamin K | 0.717 - 1.486 | 0.599 | **Mildly reduced** | | Reflects vitamin K status involved in blood clotting, calcium regulation, and bone metabolism. |

# Amino Acid

Medium priority

Complete system marker table — all markers and all severity grades are shown below.

Flagged markers: 2 | Within range: 9

| Marker | Reference range | Value | Status | Visual | Meaning |
|---|---|---|---|---|---|
| Lysine | 0.962 - 1.213 | 0.853 | Mildly reduced | | Reflects lysine availability, an essential amino acid critical for protein synthesis, collagen formation, immune modulation, and calcium metabolism. |
| Tryptophan | 4.978 - 6.289 | 6.008 | Within range | | Reflects tryptophan availability, a precursor for serotonin, melatonin, and kynurenine pathway metabolites. |
| Phenylalanine | 1.928 - 2.491 | 2.057 | Within range | | Reflects phenylalanine availability, a precursor to tyrosine and downstream catecholamines. |
| Methionine | 1.245 - 1.637 | 1.398 | Within range | | Reflects methionine availability, a key methyl donor involved in detoxification and epigenetic regulation. |
| Threonine | 1.194 - 1.685 | 1.37 | Within range | | Reflects threonine status, important for protein synthesis and mucosal integrity. |
| Isoleucine | 4.582 - 5.657 | 4.175 | Mildly reduced | | Reflects isoleucine status, a branched-chain amino acid (BCAA) involved in energy metabolism and muscle repair. |
| Leucine | 6.982 - 9.256 | 7.041 | Within range | | Reflects leucine status, a key BCAA regulating muscle protein synthesis via mTOR signalling. |
| Valine | 6.982 - 9.677 | 7.565 | Within range | | Reflects valine status, a BCAA involved in energy provision and muscle metabolism. |
| Histidine | 5.113 - 6.258 | 5.261 | Within range | | Reflects histidine status, involved in haemoglobin structure and histamine production. |

| Marker | Reference range | Value | Status | Visual | Meaning |
|---|---|---|---|---|---|
| Arginine | 1.812 - 2.337 | 1.975 | **Within range** | | Reflects arginine status, a semi-essential amino acid involved in nitric oxide production and vascular function. |
| Homocysteine | 0.983 - 1.265 | 1.073 | **Within range** | | Reflects homocysteine levels, a key intermediate in methylation and cardiovascular risk pathways. |

# Coenzyme

**Medium priority**

Complete system marker table — all markers and all severity grades are shown below.

Flagged markers: 3 | Within range: 3

| Marker | Reference range | Value | Status | Visual | Meaning |
|---|---|---|---|---|---|
| Nicotinamide | 2.074 - 3.309 | 2.293 | Within range | | Reflects NAD precursor status supporting redox reactions and energy metabolism. |
| Biotin | 1.833 - 2.979 | 1.241 | Mildly reduced | | Reflects biotin status required for fatty acid metabolism and glucose regulation. |
| Pantothenic acid | 1.116 - 2.101 | 0.604 | Moderately reduced | | Reflects vitamin B5 status required for CoA synthesis and energy metabolism. |
| Folic acid | 1.449 - 2.246 | 2.236 | Within range | | Reflects folate status required for methylation and DNA synthesis. |
| Coenzyme Q10 | 0.831 - 1.588 | 0.643 | Mildly reduced | | Reflects availability of CoQ10, a key electron transport cofactor in the mitochondrial inner membrane required for oxidative phosphorylation and cellular antioxidant protection. |
| Glutathione | 0.726 - 1.281 | 1.114 | Within range | | Reflects glutathione status as a master antioxidant and detox molecule. |

# Essential Fatty Acid

**Medium priority**

Complete system marker table — all markers and all severity grades are shown below.

Flagged markers: 2 | Within range: 2

| Marker | Reference range | Value | Status | Visual | Meaning |
|---|---|---|---|---|---|
| Linoleic acid | 0.642 - 0.985 | 0.447 | Mildly reduced | | Reflects linoleic-acid status as a key omega-6 essential fatty acid involved in membrane structure, epidermal barrier integrity, and eicosanoid precursor balance. |
| α-Linolenic acid | 0.814 - 1.202 | 0.852 | Within range | | Reflects alpha-linolenic-acid availability as a plant-derived omega-3 essential fatty acid supporting membrane fluidity, inflammatory modulation, and precursor input into longer-chain omega-3 pathways. |
| γ-Linolenic acid | 0.921 - 1.334 | 0.996 | Within range | | Reflects gamma-linolenic-acid related functional support within omega-6 metabolism, relevant to membrane signalling, skin integrity, and downstream eicosanoid balance. |
| Arachidonic acid | 0.661 - 0.808 | 0.575 | Mildly reduced | | Reflects arachidonic-acid status as a membrane-bound omega-6 fatty acid involved in cell signalling and the generation of multiple inflammatory and regulatory eicosanoids. |

# Endocrine System

Medium priority

Complete system marker table — all markers and all severity grades are shown below.

Flagged markers: 2 | Within range: 5

| Marker | Reference range | Value | Status | Visual | Meaning |
|---|---|---|---|---|---|
| Thyroid secretion index | 2.954 - 5.543 | 2.996 | Within range | | Reflects thyroid gland secretory output, including synthesis and release of thyroxine (T4) and triiodothyronine (T3) under TSH regulation, alongside iodine handling and glandular responsiveness. |
| Parathyroid hormone secretion index | 2.845 - 4.017 | 2.121 | Mildly reduced | | Reflects parathyroid hormone regulatory activity involved in calcium-phosphorus homeostasis, vitamin D coordination, and bone-mineral signalling. |
| Adrenal glands Index | 2.412 - 2.974 | 2.149 | Mildly reduced | | Reflects adrenal gland output and reserve, including cortisol, catecholamine, and DHEA-related stress adaptation under hypothalamic-pituitary-adrenal axis control. |
| Pituitary secretion index | 2.163 - 7.34 | 5.512 | Within range | | Reflects pituitary secretory signalling across trophic hormones such as TSH, ACTH, LH, FSH, prolactin, and growth-hormone-related regulation. |
| Pineal secretion index | 3.210 - 6.854 | 6.778 | Within range | | Reflects pineal gland activity and melatonin-related circadian signalling. |
| Thymus gland secretion index | 2.967 - 3.528 | 2.969 | Within range | | Reflects thymic regulatory activity associated with immune maturation, T-cell education, and endocrine-immune interaction within the neuroendocrine network. |
| Gland secretion index | 2.204 - 2.819 | 2.492 | Within range | | Reflects generalised endocrine gland secretory performance across integrated glandular networks, providing a broad view of overall endocrine signalling tone. |

# Immune System

**Within range**

Complete system marker table — all markers and all severity grades are shown below.

Flagged markers: 0 | Within range: 9

| Marker | Reference range | Value | Status | Visual | Meaning |
|---|---|---|---|---|---|
| Lymph node Index | 133.437 - 140.47 | 134.342 | Within range | | Reflects lymph node functional activity within immune surveillance, antigen presentation, lymph filtration, and adaptive immune coordination. |
| Tonsil immune Index | 0.124 - 0.453 | 0.307 | Within range | | Reflects tonsillar immune activity as part of upper airway mucosal defence, immune sampling, and early response to inhaled or ingested antigens. |
| Bone marrow Index | 0.146 - 3.218 | 1.068 | Within range | | Reflects bone marrow contribution to haematopoiesis and immune cell production, including the generation of white blood cells and immune progenitor activity. |
| Spleen index | 34.367 - 35.642 | 35.434 | Within range | | Reflects splenic immune and filtration activity, including blood surveillance, immune cell storage, removal of damaged cells, and response to circulating antigenic burden. |
| Thymus index | 58.425 - 61.213 | 59.292 | Within range | | Reflects thymic contribution to T-cell maturation, immune tolerance, and adaptive immune education. |
| Immunoglobulin index | 3.712 - 6.981 | 4.089 | Within range | | Reflects humoral immune activity through immunoglobulin production and antibody-mediated defence capacity. |
| Respiratory Immunity | 3.241 - 9.814 | 3.384 | Within range | | Reflects immune competence within the respiratory tract, including airway mucosal defence, pathogen response, and inflammatory tolerance in the lungs and upper airways. |

| Marker | Reference range | Value | Status | Visual | Meaning |
|---|---|---|---|---|---|
| GI Immunity | 0.638 - 1.712 | 0.949 | **Within range** | | Reflects immune activity associated with the gut-associated lymphoid tissue (GALT), mucosal barrier defence, and immune tolerance within the gastrointestinal tract. |
| Mucosal Immunity | 4.111 - 18.741 | 10.307 | **Within range** | | Reflects integrated mucosal immune defence across barrier tissues, including secretory immune activity, epithelial protection, and tolerance to environmental and dietary exposures. |

# Thyroid

**Medium priority**

Complete system marker table — all markers and all severity grades are shown below.

Flagged markers: 3 | Within range: 1

| Marker | Reference range | Value | Status | Visual | Meaning |
|---|---|---|---|---|---|
| FT4 | 0.103 - 0.316 | 0.6 | **Mildly elevated** | | Represents circulating levels of thyroxine available for peripheral conversion into active thyroid hormone (T3). |
| Thyroglobulin | 0.114 - 0.202 | 0.294 | **Mildly elevated** | | Reflects the protein precursor used in thyroid hormone synthesis within the thyroid gland. |
| Thyroglobulin | 0.421 - 0.734 | 0.711 | **Within range** | | Reflects the protein precursor used in thyroid hormone synthesis within the thyroid gland. |
| T3 | 0.161 - 0.308 | 0.518 | **Mildly elevated** | | Represents levels of active thyroid hormone responsible for metabolic activation at cellular level. |

# Human Toxin

Within range

Complete system marker table — all markers and all severity grades are shown below.

Flagged markers: 0 | Within range: 4

| Marker | Reference range | Value | Status | Visual | Meaning |
|---|---|---|---|---|---|
| Stimulating Beverage | 0.209 - 0.751 | 0.475 | Within range | | Reflects physiological burden associated with stimulant-beverage exposure, especially caffeine-type intake that can influence adrenal output, autonomic tone, sleep quality, and vascular reactivity. |
| Electromagnetic Radiation | 0.046 - 0.167 | 0.109 | Within range | | Reflects potential physiological stress associated with electromagnetic exposure, interpreted here as an environmental-load marker rather than a standalone diagnostic measure. |
| Free Radical | 0.124 - 0.453 | 0.178 | Within range | | Reflects oxidative free-radical burden and the degree to which reactive species may be exceeding antioxidant buffering capacity. |
| Toxic Pesticide Residue | 0.013 - 0.313 | 0.305 | Within range | | Reflects burden associated with pesticide-type toxic exposure, suggesting environmental chemical load requiring hepatic and antioxidant clearance support. |

# Heavy Metal

High priority

Complete system marker table — all markers and all severity grades are shown below.

Flagged markers: 4 | Within range: 4

| Marker | Reference range | Value | Status | Visual | Meaning |
|---|---|---|---|---|---|
| Lead | 0.052 - 0.643 | 0.976 | Mildly elevated | | Reflects lead burden as a toxic-metal exposure marker relevant to neurological stress, haematologic interference, mitochondrial impairment, and long-term detox burden. |
| Mercury | 0.013 - 0.336 | 0.468 | Mildly elevated | | Reflects mercury burden as a toxic-metal exposure marker with relevance to neurological irritation, immune dysregulation, oxidative stress, and detoxification strain. |
| Cadmium | 0.527 - 1.523 | 1.556 | Mildly elevated | | Reflects cadmium burden as a toxic-metal exposure marker associated with oxidative damage, kidney stress, and competition with essential minerals such as zinc. |
| Chromium | 0.176 - 1.183 | 0.345 | Within range | | Reflects chromium burden in this context as an exposure-related heavy-metal marker rather than the nutritional trivalent chromium used in glucose metabolism. |
| Arsenic | 0.153 - 0.621 | 0.596 | Within range | | Reflects arsenic burden as a toxic-metal and toxicoid exposure marker relevant to detoxification stress, vascular effects, mitochondrial dysfunction, and skin or nerve irritation. |
| Antimony | 0.162 - 0.412 | 0.323 | Within range | | Reflects antimony burden as a toxic-metal exposure marker associated with environmental or occupational load and increased detoxification demand. |
| Thallium | 0.182 - 0.542 | 0.388 | Within range | | Reflects thallium burden as a highly concerning toxic-metal exposure marker relevant to |

| Marker | Reference range | Value | Status | Visual | Meaning |
|---|---|---|---|---|---|
| | | | | | neurological, gastrointestinal, and mitochondrial stress. |
| Aluminum | 0.192 - 0.412 | 0.57 | **Mildly elevated** | | Reflects aluminium burden as a toxic-metal exposure marker relevant to neurological stress, bone-mineral interference, and cumulative detoxification demand. |

# Basic Physical Quality

Low priority

Complete system marker table — all markers and all severity grades are shown below.

Flagged markers: 1 | Within range: 4

| Marker | Reference range | Value | Status | Visual | Meaning |
|---|---|---|---|---|---|
| Response Ability | 59.786 - 65.424 | 61.674 | Within range | | Reflects the body's general capacity to perceive, adapt to, and recover from internal or external stressors, functioning as a broad resilience marker rather than a single organ-specific measure. |
| Mental Power | 58.715 - 63.213 | 61.23 | Within range | | Reflects perceived neurocognitive drive and mental stamina, including sustained attention, executive energy, and resistance to cognitive fatigue. |
| Water Shortage | 33.967 - 37.642 | 34.512 | Within range | | Reflects dehydration burden or reduced effective body-water availability, indicating that tissue hydration and fluid distribution may be insufficient for optimal physiological function. |
| Hypoxia | 133.642 - 141.476 | 130.994 | Mildly reduced | | Reflects reduced oxygen availability at the tissue level, indicating a functional mismatch between oxygen demand and delivery or utilisation. |
| PH | 3.156 - 3.694 | 3.248 | Within range | | Reflects overall acid-base tendency within the source framework, functioning as a broad indicator of systemic buffering balance rather than a standalone compartment-specific laboratory pH value. |

# Allergy

High priority

Complete system marker table — all markers and all severity grades are shown below.

Flagged markers: 6 | Within range: 7

| Marker | Reference range | Value | Status | Visual | Meaning |
|---|---|---|---|---|---|
| Drug Sensitivity | 0.431 - 1.329 | 0.563 | Within range | | Reflects hypersensitivity tendency to pharmaceutical compounds, indicating how strongly immune and detoxification systems may react to drug-related antigenic or chemical exposure. |
| Alcohol allergy index | 0.432 - 1.246 | 1.207 | Within range | | Reflects hypersensitivity or low tolerance tendency toward alcohol exposure, including immune reactivity, aldehyde-handling burden, and histamine-related response patterns. |
| Pollen allergy index | 0.143 - 1.989 | 1.646 | Within range | | Reflects hypersensitivity tendency to airborne pollen antigens, indicating mucosal immune overreactivity and reduced tolerance to seasonal environmental exposures. |
| Injection allergy index | 0.847 - 1.045 | 1.184 | Mildly elevated | | Reflects hypersensitivity tendency to injected substances, suggesting reactivity to direct antigenic or chemical exposure bypassing digestive barriers. |
| Chemical Sensitivity | 0.842 - 1.643 | 1.896 | Mildly elevated | | Reflects hypersensitivity tendency to chemical products, indicating low tolerance to environmental chemical exposure and potential overload of detoxification and barrier systems. |
| Paint allergy index | 0.346 - 1.401 | 0.489 | Within range | | Reflects sensitivity to paint-related compounds and volatile chemical exposures, suggesting heightened reactivity to inhaled or contact-based environmental irritants. |

| Marker | Reference range | Value | Status | Visual | Meaning |
|---|---|---|---|---|---|
| Dust allergy index | 0.543 - 1.023 | 1.263 | Mildly elevated | | Reflects hypersensitivity tendency to dust and particulate exposure, indicating respiratory mucosal reactivity and immune overresponse to airborne antigenic burden. |
| Smoke allergy index | 0.826 - 1.013 | 2.532 | Mildly elevated | | Reflects hypersensitivity tendency to smoke exposure, indicating respiratory tract irritation, oxidative burden, and mucosal immune overreactivity. |
| Hair dye allergy index | 0.717 - 1.486 | 2.856 | Moderately elevated | | Reflects cutaneous and chemical hypersensitivity tendency to hair-dye compounds, suggesting lower tolerance to topical chemical exposure and sensitisation of skin immune pathways. |
| Animal fur allergy index | 0.124 - 1.192 | 0.571 | Within range | | Reflects hypersensitivity tendency to animal fur or dander exposure, indicating respiratory and mucosal immune reactivity to environmental protein antigens. |
| Metal jewelry allergy index | 0.549 - 1.213 | 0.743 | Within range | | Reflects cutaneous hypersensitivity tendency to metal contact, indicating local immune sensitisation and lower tolerance to metal-triggered inflammatory skin reactions. |
| Seafood allergy index | 0.449 - 1.246 | 2.379 | Mildly elevated | | Reflects hypersensitivity tendency to seafood antigens, indicating altered oral-gut immune tolerance and increased risk of food-triggered immune reactivity. |
| Milk allergy index | 0.477 - 1.348 | 0.739 | Within range | | Reflects hypersensitivity tendency to milk-related antigens, indicating altered digestive-immune tolerance and increased food-reactive potential at the gut barrier. |

# Obesity

High priority

Complete system marker table — all markers and all severity grades are shown below.

Flagged markers: 4 | Within range: 1

| Marker | Reference range | Value | Status | Visual | Meaning |
|---|---|---|---|---|---|
| Abnormal lipid metabolism coefficient | 1.992 - 3.713 | 0.89 | Moderately reduced | | Reflects baseline mitochondrial energy expenditure under resting conditions. |
| Brown adipose tissue abnormalities coefficient | 2.791 - 4.202 | 1.858 | Moderately reduced | | Reflects functional irregularity in brown adipose tissue, the thermogenic fat compartment involved in heat production, metabolic flexibility, and energy expenditure. |
| Hyperinsulinemia coefficient | 0.097 - 0.215 | 0.206 | Within range | | Represents elevated insulin activity within reproductive-metabolic interactions, often linked to hormonal imbalance. |
| Nucleus of the hypothalamus abnormal coefficient | 0.332 - 0.626 | 0.719 | Mildly elevated | | Reflects dysregulation within hypothalamic centres that coordinate appetite, satiety, autonomic tone, circadian signalling, and energy-balance regulation. |
| Triglyceride content of abnormal coefficient | 1.341 - 1.991 | 5.222 | Moderately elevated | | Reflects mitochondrial coupling efficiency in ATP production versus heat generation. |

# Skin

High priority

Complete system marker table — all markers and all severity grades are shown below.

Flagged markers: 6 | Within range: 4

| Marker | Reference range | Value | Status | Visual | Meaning |
|---|---|---|---|---|---|
| Skin Free Radical Index | 0.124 - 3.453 | 2.972 | Within range | | Reflects the oxidative stress burden at the level of skin tissue, indicating the balance between free radical generation and antioxidant defence within the dermal and epidermal layers. |
| Skin Collagen Index | 4.471 - 6.079 | 2.899 | Mildly reduced | | Reflects collagen support within skin tissue, indicating the quality of dermal structural proteins and visible connective tissue resilience. |
| Skin Grease Index | 14.477 - 21.348 | 25.821 | Mildly elevated | | Reflects sebum production and lipid balance on the skin surface, indicating sebaceous gland activity and skin barrier lubrication. |
| Skin Immunity Index | 1.035 - 3.230 | 4.02 | Mildly elevated | | Reflects immune defence activity at the skin level, indicating the ability of the skin barrier to respond to pathogens, irritation, and environmental stress. |
| Skin Moisture Index | 0.218 - 0.953 | 2.145 | Moderately elevated | | Reflects hydration retention and barrier moisture quality within skin tissue. |
| Skin Moisture Loss | 2.214 - 4.158 | 3.125 | Within range | | Reflects transepidermal water loss (TEWL), indicating the integrity of the skin barrier and its ability to retain moisture. |
| Skin Red Blood Trace Index | 0.824 - 1.942 | 1.055 | Within range | | Reflects microcirculation and capillary activity within skin tissue, indicating blood supply and oxygen delivery to the dermis. |
| Skin Elasticity Index | 2.717 - 3.512 | 1.571 | Mildly reduced | | Reflects the recoil capacity and structural flexibility of skin tissue, indicating the condition of collagen-elastin architecture and hydration-related matrix support. |

| Marker | Reference range | Value | Status | Visual | Meaning |
|---|---|---|---|---|---|
| Skin Melanin Index | 0.346 - 0.501 | 0.612 | **Mildly elevated** | | Reflects skin pigment expression and melanin-related tone, with relevance to oxidative exposure, inflammatory change, and hormonal influence on skin tissue. |
| Tin | 0.842 - 1.858 | 1.419 | **Within range** | | Reflects exposure to tin and trace biological impact. |

# Eye

High priority

Complete system marker table — all markers and all severity grades are shown below.

Flagged markers: 6 | Within range: 2

| Marker | Reference range | Value | Status | Visual | Meaning |
|---|---|---|---|---|---|
| Bags under the eyes | 0.510 - 3.109 | 3.004 | Within range | | Reflects periorbital fluid retention and tissue congestion, often indicating lymphatic stagnation, altered renal-fluid handling, or poor overnight recovery. |
| Collagen eye wrinkle | 2.031 - 3.107 | 1.31 | Mildly reduced | | Reflects collagen degradation and reduced extracellular-matrix integrity in the periorbital region, where connective tissue is thin and highly sensitive to oxidative and hormonal stress. |
| Dark circles | 0.831 - 3.188 | 3.883 | Mildly elevated | | Reflects microcirculatory and pigmentation changes in the periorbital region, often linked to vascular congestion or fatigue. |
| Lymphatic obstruction | 1.116 - 4.101 | 3.52 | Within range | | Represents impaired lymphatic drainage affecting fluid balance and tissue detoxification. |
| Sagging | 0.233 - 0.559 | 0.751 | Mildly elevated | | Reflects loss of tissue elasticity and structural support in skin and connective tissue. |
| Edema | 0.332 - 0.726 | 1.234 | Moderately elevated | | Represents fluid accumulation in tissues due to imbalance in vascular and lymphatic dynamics. |
| Eye cell activity | 0.118 - 0.892 | 0.895 | Mildly elevated | | Reflects metabolic and functional activity of ocular cells involved in vision. |
| Visual fatigue | 2.017 - 5.157 | 8.04 | Mildly elevated | | Represents strain on visual processing systems due to overuse or metabolic fatigue. |

# Collagen

High priority

Complete system marker table — all markers and all severity grades are shown below.

Flagged markers: 6 | Within range: 8

| Marker | Reference range | Value | Status | Visual | Meaning |
|---|---|---|---|---|---|
| Overall Eye Function | 6.352 - 8.325 | 8.31 | Within range | | Represents the overall functional status of the visual system, integrating ocular tissue vitality, circulation, and visual performance. |
| Endocrine system | 7.245 - 8.562 | 7.491 | Within range | | Reflects integrity of collagen matrix within the endocrine system, influencing structural stability and functional resilience. |
| Hair and skin | 4.533 - 6.179 | 4.806 | Within range | | Reflects integrity of collagen matrix within the hair and skin, influencing structural stability and functional resilience. |
| Endocrine system | 6.178 - 8.651 | 8.253 | Within range | | Reflects integrity of collagen matrix within the endocrine system, influencing structural stability and functional resilience. |
| Circulatory system | 3.586 - 4.337 | 4.153 | Within range | | Reflects integrity of collagen matrix within the circulatory system, influencing structural stability and functional resilience. |
| Digestive system | 3.492 - 4.723 | 3.168 | Mildly reduced | | Reflects integrity of collagen matrix within the digestive system, influencing structural stability and functional resilience. |
| Immune system | 3.376 - 4.582 | 2.938 | Mildly reduced | | Reflects integrity of collagen matrix within the immune system, influencing structural stability and functional resilience. |
| Motion systems | 6.458 - 8.133 | 5.573 | Mildly reduced | | Reflects integrity of collagen matrix within the motion systems, influencing structural stability and functional resilience. |

| Marker | Reference range | Value | Status | Visual | Meaning |
|---|---|---|---|---|---|
| Muscle Tissue | 6.552 - 8.268 | 5.774 | **Mildly reduced** | | Reflects integrity of collagen matrix within the muscle tissue, influencing structural stability and functional resilience. |
| Fat Metabolism | 6.338 - 8.368 | 4.551 | **Mildly reduced** | | Reflects integrity of collagen matrix within the fat metabolism, influencing structural stability and functional resilience. |
| Detoxification and metabolism | 6.187 - 8.466 | 6.519 | **Within range** | | Reflects integrity of collagen matrix within the detoxification and metabolism, influencing structural stability and functional resilience. |
| Reproductive system | 3.778 - 4.985 | 3.997 | **Within range** | | Reflects integrity of collagen matrix within the reproductive system, influencing structural stability and functional resilience. |
| Nervous system | 3.357 - 4.239 | 2.643 | **Mildly reduced** | | Reflects integrity of collagen matrix within the nervous system, influencing structural stability and functional resilience. |
| Skeleton | 6.256 - 8.682 | 7.975 | **Within range** | | Reflects integrity of collagen matrix within the skeleton, influencing structural stability and functional resilience. |

# Channels and collaterals

High priority

Complete system marker table — all markers and all severity grades are shown below.

Flagged markers: 4 | Within range: 12

| Marker | Reference range | Value | Status | Visual | Meaning |
|---|---|---|---|---|---|
| Hand Tai Yin Lung Meridian | 48.264 - 65.371 | 63.778 | Within range | | Reflects the functional state of the Lung meridian within the channels and collaterals model, associated with respiratory vitality, boundary regulation, grief-stress patterns, and surface defensive function. |
| Hand Yangming Large Intestine Meridian | 56.749 - 67.522 | 61.756 | Within range | | Reflects the functional state of the Large Intestine meridian, associated with elimination, release, mucosal boundary handling, and paired regulation with the Lung channel. |
| Foot Yangming Stomach Meridian | 0.481 - 1.043 | 0.485 | Within range | | Reflects the functional state of the Stomach meridian, associated with digestive receiving, nourishment processing, appetite regulation, and tissue support derived from food transformation. |
| Foot Tai Yin Lung Meridian | 0.327 - 0.937 | 0.465 | Within range | | Reflects the functional state of the Spleen meridian in practical TCM usage despite the source label discrepancy, associated with transformation of food and fluid, tissue nourishment, and dampness handling. |
| Hand Shao Yin Heart Sutra | 1.672 - 1.978 | 1.477 | Mildly reduced | | Reflects the functional state of the Heart channel, associated with circulation, emotional coherence, sleep quality, and the stability of mental-spiritual regulation in traditional pattern terms. |
| Hand the small intestine by the sun | 0.192 - 0.412 | 0.199 | Within range | | Reflects the functional state of the Small Intestine channel, associated with separation of clear from turbid, discernment, and downstream handling of |

| Marker | Reference range | Value | Status | Visual | Meaning |
|---|---|---|---|---|---|
| | | | | | digestive and inflammatory burden. |
| Foot Tai Yang Bladder Meridian | 4.832 - 5.147 | 3.773 | **Mildly reduced** | | Reflects the functional state of the Bladder meridian, associated with fluid regulation, posterior chain tension, nervous-system vigilance, and stress storage along the back body. |
| Foot Shao Yin Kidney | 3.321 - 4.244 | 3.937 | **Within range** | | Reflects the functional state of the Kidney channel, associated with foundational reserves, endocrine-adrenal resilience, reproductive vitality, bone support, and fear-stress regulation in traditional pattern terms. |
| Hand Jue Yin Pericardium Meridian | 1.338 - 1.672 | 1.34 | **Within range** | | Reflects the functional state of the Pericardium channel, associated with emotional buffering, chest tension, circulation, and protection against stress-induced internal agitation. |
| Hand Shao Yang Triple Burner Meridian | 0.669 - 1.544 | 0.548 | **Mildly reduced** | | Reflects the functional state of the Triple Burner channel, associated with coordination across upper, middle, and lower body regulatory compartments, especially fluid movement and stress adaptation. |
| Foot Shao Yang Gall Bladder Meridian | 1.554 - 1.988 | 1.868 | **Within range** | | Reflects the functional state of the Gall Bladder channel, associated with bile flow, decisiveness, lateral tension patterns, and the handling of damp-heat or stagnant digestive stress. |
| Foot Jue Yin Liver Meridian | 1.553 - 2.187 | 1.652 | **Within range** | | Reflects the functional state of the Liver channel, associated with smooth flow, emotional pressure regulation, blood storage, hormonal rhythm, and tension-stagnation patterns. |
| Ren channel | 11.719 - 18.418 | 12.572 | **Within range** | | Reflects the functional state of the Ren vessel, associated with front-body integration, reproductive and hormonal regulation, fluid-nutrient support, and yin-type restorative processes. |

| Marker | Reference range | Value | Status | Visual | Meaning |
|---|---|---|---|---|---|
| Governor meridian | 0.708 - 1.942 | 1.536 | **Within range** | | Reflects the functional state of the Governor vessel, associated with back-body integration, spinal support, mental drive, and yang-type activating regulation. |
| Vital meridian | 6.138 - 21.396 | 6.331 | **Within range** | | Reflects generalised vital-channel reserve in the source framework, pointing to overall constitutional vitality, regulatory coherence, and resilience of the channel network. |
| Tai mai | 5.733 - 7.109 | 5.146 | **Mildly reduced** | | Reflects the functional state of the Dai Mai concept in traditional pattern language, associated with belt-like regulation around the waist, structural containment, and lower-body fluid or stagnation patterns. |

# Pulse of heart and brain

**Medium priority**

Complete system marker table — all markers and all severity grades are shown below.

Flagged markers: 2 | Within range: 5

| Marker | Reference range | Value | Status | Visual | Meaning |
|---|---|---|---|---|---|
| Stroke index | 60.735 - 65.396 | 62.076 | Within range | | Reflects stroke volume relative to body size or circulatory demand, offering a scaled view of cardiac output efficiency. |
| Stroke volume(SV) | 63.012 - 67.892 | 61.901 | Mildly reduced | | Reflects beat-by-beat blood ejection from the heart, representing the immediate mechanical contribution of each contraction to circulation. |
| Heart peripheral resistance (TRR) | 0.983 - 1.265 | 1.638 | Mildly elevated | | Reflects total peripheral resistance encountered by blood as it moves through the systemic vascular network. |
| Pulse wave coefficient K | 0.316 - 0.401 | 0.391 | Within range | | Reflects properties of the pulse-wave transmission through arteries, providing insight into vascular elasticity, compliance, and wave reflection dynamics. |
| Cerebrovascular blood oxygen saturation(Sa) | 0.710 - 1.109 | 0.894 | Within range | | Reflects the saturation of oxygen carried in blood supplying the cerebrovascular system, indicating how effectively oxygen is available for brain delivery. |
| Cerebrovascular blood oxygen volume(CaCO2) | 7.880 - 10.090 | 8.081 | Within range | | Reflects the oxygen content available within blood supplying the cerebrovascular system, indicating total oxygen-carrying capacity rather than saturation alone. |
| Cerebrovascular blood oxygen presure(PaO2) | 5.017 - 5.597 | 5.169 | Within range | | Reflects the partial pressure of oxygen in blood supplying the cerebrovascular system, indicating oxygen availability for diffusion into brain tissue. |

# Blood lipids

High priority

Complete system marker table — all markers and all severity grades are shown below.

Flagged markers: 5 | Within range: 2

| Marker | Reference range | Value | Status | Visual | Meaning |
|---|---|---|---|---|---|
| Blood viscosity | 4.131 - 4.562 | 5.233 | Moderately elevated | | Reflects thickness of blood affecting circulation and oxygen delivery. |
| Total cholesterol(TC) | 1.833 - 2.979 | 2.75 | Within range | | Reflects total circulating cholesterol burden across lipoprotein fractions, giving a broad view of cholesterol availability, transport, and potential accumulation. |
| Triglyceride(TG) | 1.116 - 2.101 | 2.218 | Mildly elevated | | Reflects circulating triglyceride levels as a marker of hepatic lipid export, carbohydrate handling, and the balance between fat storage and oxidation. |
| High-density lipoprotein (HDL-C) | 1.449 - 2.246 | 2.955 | Mildly elevated | | Reflects high-density lipoprotein status involved in reverse cholesterol transport, antioxidant protection, and aspects of vascular repair signalling. |
| Low-density lipoprotein (LDL-C) | 0.831 - 1.588 | 1.015 | Within range | | Reflects low-density lipoprotein cholesterol burden, indicating transport of cholesterol to peripheral tissues and potential exposure to oxidative modification. |
| Neutral fat(MB) | 0.726 - 1.281 | 3.096 | Mildly elevated | | Reflects neutral fat burden within circulation, generally corresponding to stored or transported triglyceride-rich lipid material. |
| Circulating immune complex(CIC) | 13.012 - 17.291 | 18.601 | Mildly elevated | | Reflects immune complex burden indicating antigen-antibody interaction. |

# Prostate

**Medium priority**

Complete system marker table — all markers and all severity grades are shown below.

Flagged markers: 3 | Within range: 0

| Marker | Reference range | Value | Status | Visual | Meaning |
|---|---|---|---|---|---|
| Degree of Prostatic Hyperplasia | 1.023 - 3.230 | 5.03 | **Moderately elevated** | | Reflects the extent of enlargement or proliferative change in prostate tissue, suggesting altered growth signalling and reduced ease of urinary outflow. |
| Degree of Prostatic Calcification | 1.471 - 6.079 | 8.907 | **Mildly elevated** | | Reflects calcific change within prostate tissue, suggesting chronic inflammation, stagnation, tissue-remodelling burden, or longer-standing glandular congestion. |
| Prostatitis Syndrome | 2.213 - 2.717 | 6.11 | **Moderately elevated** |  | Reflects an inflammatory-irritative pattern involving the prostate, suggesting pain, congestion, urinary discomfort, or immune-reactive stress in the gland and surrounding pelvic tissues. |

# Male Sexual Function

**Medium priority**

Complete system marker table — all markers and all severity grades are shown below.

Flagged markers: 3 | Within range: 0

| Marker | Reference range | Value | Status | Visual | Meaning |
|---|---|---|---|---|---|
| Testosterone | 3.342 - 9.461 | 2.492 | **Moderately reduced** | | Reflects testosterone status influencing energy, libido, muscle, and metabolic health. |
| Gonadotropin | 4.111 - 18.741 | 3.558 | **Mildly reduced** | | Reflects general gonadotropin activity controlling reproductive hormones. |
| Erection Transmitter | 3.241 - 9.814 | 3.03 | **Mildly reduced** | | Reflects neurotransmitter activity related to erectile function. |

# Sperm and semen

**Within range**

Complete system marker table — all markers and all severity grades are shown below.

Flagged markers: 0 | Within range: 4

| Marker | Reference range | Value | Status | Visual | Meaning |
|---|---|---|---|---|---|
| Semen volume | 1.502 - 6.028 | 5.263 | **Within range** | | Reflects semen production and reproductive capacity. |
| Semen liquefying time | 10.283 - 30.282 | 26.982 | **Within range** | | Reflects the time required for semen to liquefy after ejaculation, indicating seminal enzymatic activity and the functional quality of seminal fluid. |
| Number of sperms | 2.483 - 3.932 | 3 | **Within range** | | Reflects sperm count within semen, indicating the output capacity of spermatogenesis and the overall quantitative component of male fertility potential. |
| Sperm motility rate | 0.637 - 0.877 | 0.689 | **Within range** | | Reflects the proportion of sperm demonstrating effective movement, indicating their functional ability to travel and support fertilisation potential. |

# Male Hormone

Medium priority

Complete system marker table — all markers and all severity grades are shown below.

Flagged markers: 2 | Within range: 4

| Marker | Reference range | Value | Status | Visual | Meaning |
|---|---|---|---|---|---|
| Androgen | 1.372 - 1.878 | 1.492 | Within range | | Reflects androgen hormone activity influencing muscle mass, libido, energy, and metabolic function. |
| Estrogen | 0.521 - 1.036 | 0.672 | Within range | | Reflects oestrogen status within reproductive-endocrine signalling, including production, peripheral conversion, receptor activity, hepatic conjugation, and enterohepatic recirculation. |
| Progesterone(P) | 0.686 - 1.432 | 0.651 | Mildly reduced | | Reflects progesterone levels important for reproductive balance and nervous system calming. |
| Luteinizing Hormone (LH) | 0.679 - 1.324 | 0.593 | Mildly reduced | | Reflects LH signalling involved in ovulation and testosterone production. |
| Prolactin(PRL) | 0.738 - 1.742 | 1.386 | Within range | | Reflects prolactin hormone involved in lactation and reproductive regulation. |
| Follicle Stimulating Hormone(FSH) | 0.387 - 0.956 | 0.54 | Within range | | Reflects FSH levels involved in follicle development and reproductive function. |

# Human Immunity

High priority

Complete system marker table — all markers and all severity grades are shown below.

Flagged markers: 4 | Within range: 1

| Marker | Reference range | Value | Status | Visual | Meaning |
|---|---|---|---|---|---|
| Tissue barrier immune index | 7.435 - 8.847 | 9.626 | Mildly elevated | | Reflects immune competence at tissue barrier interfaces, including epithelial surveillance, local immune tolerance, and defence against antigen penetration across skin and mucosal surfaces. |
| Innate immune cell index | 7.424 - 9.052 | 10.784 | Mildly elevated | | Reflects functional activity of innate immune cells involved in first-response defence, including phagocytic surveillance, early inflammatory signalling, and pathogen recognition. |
| Innate immune molecule index | 7.143 - 9.214 | 7.585 | Within range | | Reflects soluble innate immune mediators involved in early defence, inflammatory communication, and coordination of antimicrobial responses. |
| Cellular immune index | 5.769 - 7.643 | 8.547 | Mildly elevated | | Reflects cell-mediated immune activity, particularly T-cell-driven defence, immune coordination, and targeted response to infected or abnormal cells. |
| Humoral immunity index | 6.424 - 8.219 | 8.708 | Mildly elevated | | Reflects antibody-mediated immune activity and the functional strength of humoral defence across antigen recognition, immune memory, and adaptive immune responsiveness. |

# Human Consciousness Level

High priority

Complete system marker table — all markers and all severity grades are shown below.

Flagged markers: 6 | Within range: 11

| Marker | Reference range | Value | Status | Visual | Meaning |
|---|---|---|---|---|---|
| Shame | 2.214 - 3.925 | 3.056 | Within range | | Reflects a low-resource self-evaluative emotional state characterised by contraction, withdrawal, and reduced psychological resilience within the source framework. |
| Guilt | 2.154 - 3.478 | 2.543 | Within range | | Reflects a self-critical emotional state centred on remorse, self-blame, or unresolved internal conflict within the source framework. |
| Apathy | 2.343 - 4.089 | 2.263 | Mildly reduced | | Reflects reduced motivational drive, emotional engagement, and activation capacity within the source framework. |
| Grief | 1.845 - 3.241 | 1.997 | Within range | | Reflects an emotional-processing state associated with loss, sadness, and unresolved bereavement-related contraction within the source framework. |
| Fear | 2.155 - 4.031 | 1.681 | Mildly reduced | | Reflects heightened threat perception and protective vigilance within the source framework, associated with autonomic activation and anticipatory stress. |
| Desire | 1.484 - 2.485 | 1.711 | Within range | | Reflects motivational pull toward acquisition, reward, or craving-driven engagement within the source framework. |
| Anger | 1.256 - 2.529 | 1.466 | Within range | | Reflects a state of irritability, frustration, or forceful activation within the source framework. |
| Pride | 1.378 - 2.489 | 1.025 | Mildly reduced | | Reflects an elevated self-protective or self-affirming emotional posture within the source framework. |
| Courage | 1.727 - 3.315 | 1.8 | Within range | | Reflects the capacity to remain engaged and act despite discomfort, |

| Marker | Reference range | Value | Status | Visual | Meaning |
|---|---|---|---|---|---|
| | | | | | uncertainty, or stress within the source framework. |
| Neutrality | 2.471 - 3.892 | 2.16 | **Mildly reduced** | | Reflects a balanced, less-reactive emotional stance within the source framework, associated with reduced polarisation and steadier regulation. |
| Willingness | 2.216 - 4.094 | 2.598 | **Within range** | | Reflects an engaged, cooperative motivational stance within the source framework, associated with readiness to participate and adapt. |
| Acceptance | 1.668 - 4.053 | 2.239 | **Within range** | | Reflects a state of reduced resistance and greater adaptive acknowledgment of reality within the source framework. |
| Reason | 1.352 - 3.436 | 2.693 | **Within range** | | Reflects the relative dominance of reflective, analytical, and cognitively organised processing within the source framework. |
| Love | 2.138 - 3.754 | 1.871 | **Mildly reduced** | | Reflects an affiliative, connected, and emotionally integrated state within the source framework. |
| Joy | 2.129 - 4.021 | 2.654 | **Within range** | | Reflects access to positive affect, vitality, and unforced wellbeing within the source framework. |
| Peace | 2.243 - 4.022 | 2.099 | **Mildly reduced** | | Reflects a deeply settled, non-reactive internal state within the source framework, associated with low internal conflict and high autonomic ease. |
| Enlightenment | 2.225 - 3.822 | 3.556 | **Within range** | | Reflects the highest conceptual level in the source framework and should be interpreted as a symbolic marker of integrative awareness rather than a clinical state. |

# Respiratory Function

Low priority

Complete system marker table — all markers and all severity grades are shown below.

Flagged markers: 1 | Within range: 3

| Marker | Reference range | Value | Status | Visual | Meaning |
|---|---|---|---|---|---|
| Volume Tidal(VT) | 5.431 - 7.479 | 4.051 | Mildly reduced | | Reflects the volume of air moved in and out of the lungs during a normal resting breath, indicating baseline breathing efficiency and ventilatory patterning. |
| Inspiratory Volume(TI) | 4.126 - 6.045 | 4.442 | Within range | | Reflects inspiratory volume capacity and the efficiency of drawing air into the lungs, indicating how effectively inhalation mechanics are functioning. |
| Functional Residual Capacity(FRC) | 5.147 - 6.219 | 5.503 | Within range | | Reflects the volume of air remaining in the lungs after a normal exhalation, indicating resting lung-volume set point and the balance between elastic recoil and airway closure. |
| Residual Volume(RV) | 4.363 - 5.647 | 5.322 | Within range | | Reflects the volume of air remaining in the lungs after maximal exhalation, indicating how completely the lungs can empty and whether air trapping is present. |

# Lecithin

Medium priority

Complete system marker table — all markers and all severity grades are shown below.

Flagged markers: 2 | Within range: 4

| Marker | Reference range | Value | Status | Visual | Meaning |
|---|---|---|---|---|---|
| Phospholipid index | 3.654 - 4.478 | 4.471 | Within range | | Represents the status of phospholipids as key structural components of cell membranes and lipoproteins. |
| Sphingolipid index | 3.121 - 3.853 | 2.842 | Mildly reduced | | Reflects sphingolipid balance involved in membrane structure and cellular signalling. |
| PH | 3.341 - 4.214 | 3.894 | Within range | | Reflects overall acid-base tendency within the source framework, functioning as a broad indicator of systemic buffering balance rather than a standalone compartment-specific laboratory pH value. |
| Lecithin index | 4.365 - 5.142 | 4.165 | Within range | | Represents lecithin levels contributing to lipid transport and membrane structure. |
| Brain phospholipid index | 4.225 - 5.713 | 4.038 | Mildly reduced | | Reflects phospholipid availability and membrane integrity within brain tissue, influencing synaptic signalling, myelin stability, receptor function, and neuronal resilience. |
| Liposomal index | 3.112 - 4.081 | 4.071 | Within range | | Reflects liposomal or phospholipid-vesicle style transport capacity, indicating how effectively lipid-based carriers may support membrane delivery, absorption, and intracellular nutrient trafficking. |

# Fatty acid

Medium priority

Complete system marker table — all markers and all severity grades are shown below.

Flagged markers: 2 | Within range: 2

| Marker | Reference range | Value | Status | Visual | Meaning |
|---|---|---|---|---|---|
| Saturated fatty acid index | 1.334 - 2.407 | 1.437 | Within range | | Represents the proportion of saturated fats within overall lipid balance. |
| Unsaturated fatty acid index | 2.224 - 3.153 | 2.834 | Within range | | Reflects levels of unsaturated fats supporting membrane fluidity and metabolic function. |
| Essential fatty acid index | 2.144 - 3.238 | 2.087 | Mildly reduced | | Represents availability of essential fatty acids required for cellular function. |
| Triglyceride index | 3.361 - 4.644 | 2.377 | Mildly reduced | | Reflects circulating triglyceride levels and lipid storage status. |

TECHNICAL REFERENCE

# Appendix

This section contains supporting technical material and summary reference tables used in the report.

| Section | Marker | Range | Value | Status |
|---|---|---|---|---|
| Cardiovascular and Cerebrovascular | Blood Viscosity | 48.264 - 65.371 | 53.723 | Within range |
| Cardiovascular and Cerebrovascular | Cholesterol Crystal | 56.749 - 67.522 | 66.455 | Within range |
| Cardiovascular and Cerebrovascular | Blood Fat | 0.481 - 1.043 | 0.902 | Within range |
| Cardiovascular and Cerebrovascular | Vascular Resistance | 0.327 - 0.937 | 1.226 | Mildly elevated |
| Cardiovascular and Cerebrovascular | Vascular Elasticity | 1.672 - 1.978 | 1.934 | Within range |
| Cardiovascular and Cerebrovascular | Myocardial Blood Demand | 0.192 - 0.412 | 0.407 | Within range |
| Cardiovascular and Cerebrovascular | Myocardial Perfusion | 4.832 - 5.147 | 5.132 | Within range |
| Cardiovascular and Cerebrovascular | Myocardial Oxygen Consumption | 3.321 - 4.244 | 3.382 | Within range |
| Cardiovascular and Cerebrovascular | Stroke Volume | 1.338 - 1.672 | 1.29 | Mildly reduced |
| Cardiovascular and Cerebrovascular | LV Ejection Impedance | 0.669 - 1.544 | 1.456 | Within range |
| Cardiovascular and Cerebrovascular | LV Pump Power | 1.554 - 1.988 | 0.991 | Moderately reduced |
| Cardiovascular and Cerebrovascular | Coronary Artery Elasticity | 1.553 - 2.187 | 1.66 | Within range |
| Cardiovascular and Cerebrovascular | Coronary Perfusion Pressure | 11.719 - 18.418 | 19.312 | Moderately elevated |
| Cardiovascular and Cerebrovascular | Cerebral Blood Vessel Elasticity | 0.708 - 1.942 | 1.021 | Within range |
| Cardiovascular and Cerebrovascular | Brain Tissue Blood Supply Status | 6.138 - 21.396 | 20.196 | Within range |
| Gastrointestinal Function | Pepsin Secretion Coefficient | 59.847 - 65.234 | 59.915 | Within range |
| Gastrointestinal Function | Gastric Peristalsis Function Coefficient | 58.425 - 61.213 | 58.71 | Within range |

| Section | Marker | Range | Value | Status |
|---|---|---|---|---|
| Gastrointestinal Function | Gastric Absorption Function Coefficient | 34.367 - 35.642 | 29.734 | Moderately reduced |
| Gastrointestinal Function | Small Intestine Motility | 133.437 - 140.476 | 126.057 | Moderately reduced |
| Gastrointestinal Function | Small Intestine Absorption | 3.572 - 6.483 | 2.82 | Moderately reduced |
| Large Intestine Function | Colon Motility | 4.572 - 6.483 | 3.088 | Moderately reduced |
| Large Intestine Function | Colon Absorption | 2.946 - 3.815 | 2.952 | Within range |
| Large Intestine Function | Gut Microbiome | 1.734 - 2.621 | 0.807 | Moderately reduced |
| Large Intestine Function | Intraluminal pressure coefficient | 1.173 - 2.297 | 2.639 | Mildly elevated |
| Liver Function | Protein Metabolism | 116.34 - 220.621 | 116.569 | Within range |
| Liver Function | Energy Production Function | 0.713 - 0.992 | 0.881 | Within range |
| Liver Function | Detoxification Function | 0.202 - 0.991 | 0.977 | Within range |
| Liver Function | Bile Secretion Function | 0.432 - 0.826 | 0.717 | Within range |
| Liver Function | Liver Fat Content | 0.097 - 0.419 | 0.636 | Moderately elevated |
| Gallbladder Function | Serum Globulin (A/G) | 126 - 159 | 126.767 | Within range |
| Gallbladder Function | Total Bilirubin (TBIL) | 0.232 - 0.686 | 0.542 | Within range |
| Gallbladder Function | Alkaline Phosphatase (ALP) | 0.082 - 0.342 | 0.316 | Within range |
| Gallbladder Function | Serum Total Bile Acid (TBA | 0.317 - 0.695 | 0.704 | Mildly elevated |
| Gallbladder Function | Bilirubin (DBIL) | 0.218 - 0.549 | 0.56 | Mildly elevated |
| Pancreatic Function | Insulin | 2.845 - 4.017 | 3.036 | Within range |
| Pancreatic Function | Pancreatic Polypeptide (PP) | 3.210 - 6.854 | 3.26 | Within range |
| Pancreatic Function | Glucagon | 2.412 - 2.974 | 2.666 | Within range |
| Kidney Function | Urobilinogen Index | 2.762 - 5.424 | 6.215 | Mildly elevated |
| Kidney Function | Uric acid Index | 1.435 - 1.987 | 2.689 | Moderately elevated |
| Kidney Function | BUN | 4.725 - 8.631 | 4.766 | Within range |

| Section | Marker | Range | Value | Status |
|---|---|---|---|---|
| Kidney Function | Proteinuria Index | 1.571 - 4.079 | 4.761 | Mildly elevated |
| Lung Function | Vital Capacity | 3348 - 3529 | 3489 | Within range |
| Lung Function | Total Lung Capacity | 4301 - 4782 | 4286 | Mildly reduced |
| Lung Function | Airway Resistance | 1.374 - 1.709 | 1.621 | Within range |
| Lung Function | Arterial Oxygen Content | 17.903 - 21.012 | 20.78 | Within range |
| Brain Nerve | Brain Tissue Blood Supply | 143.37 - 210.81 | 168.036 | Within range |
| Brain Nerve | Cerebral Arteriosclerosis | 0.103 - 0.642 | 0.196 | Within range |
| Brain Nerve | Cranial Nerve Function | 0.253 - 0.659 | 0.292 | Within range |
| Brain Nerve | Sentiment Index | 0.109 - 0.351 | 0.262 | Within range |
| Brain Nerve | Memory | 0.442 - 0.817 | 0.379 | Mildly reduced |
| Bone Disease | Name: Test Test | None | Sex: Male | Unclear |
| Bone Disease | Tin | None | Figure: 184cm, 90kg | Unclear |
| Bone Disease | Figure: 184cm, 90kg | None | Testing Time: 21/03/2026 21:03 | Unclear |
| Bone Disease | Lumbar Fiber Protruding Dimension | None | No Direction | Unclear |
| Bone Disease | Adhesion Degree of Shoulder Muscle | None | < u 0.2 | Unclear |
| Bone Disease | Limbs Circulation Limit | None | + | Unclear |
| Bone Disease | Age of Ligament | None | 10%-40% | Unclear |
| Bone Mineral Density | Osteoclast Coefficient | 86.73 - 180.97 | 151.19 | Within range |
| Bone Mineral Density | Amount of Calcium Loss | 0.209 - 0.751 | 0.469 | Within range |
| Bone Mineral Density | Degree of Bone Hyperplasia | 0.046 - 0.167 | 0.389 | Mildly elevated |
| Bone Mineral Density | Degree of Osteoporosis | 0.124 - 0.453 | 0.528 | Moderately elevated |
| Bone Mineral Density | Bone Mineral Density | 0.796 - 0.433 | 0.435 | Within range |
| Rheumatoid Bone Disease | Degree of Cervical Calcification | 421 - 490 | 502.068 | Mildly elevated |
| Rheumatoid Bone Disease | Degree of Lumbar Calcification | 4.326 - 7.531 | 7.352 | Within range |
| Rheumatoid Bone Disease | Bone Hyperplasia Coefficient | 2.954 - 5.543 | 6 | Mildly elevated |

| Section | Marker | Range | Value | Status |
|---|---|---|---|---|
| Rheumatoid Bone Disease | Osteoporosis Coefficient | 2.019 - 4.721 | 4.705 | Within range |
| Rheumatoid Bone Disease | Rheumatism Coefficient | 4.023 - 11.627 | 16.843 | Moderately elevated |
| Bone Growth Index | Bone alkaline phosphatase | 0.433 - 0.796 | 0.399 | Mildly reduced |
| Bone Growth Index | Osteocalcin | 0.525 - 0.817 | 0.508 | Mildly reduced |
| Bone Growth Index | Status of long bone healing | 0.713 - 0.992 | 0.606 | Mildly reduced |
| Bone Growth Index | Short bone cartilage healing situation | 0.202 - 0.991 | 0.362 | Within range |
| Bone Growth Index | Epiphyseal line | 0.432 - 0.826 | 0.769 | Within range |
| Blood Sugar | Coefficient of Insulin Secretion | 2.967 - 3.528 | 3.043 | Within range |
| Blood Sugar | Blood Sugar Coefficient | 2.163 - 7.321 | 3.191 | Within range |
| Blood Sugar | Urine Sugar Coefficient | 2.204 - 2.819 | 2.535 | Within range |
| Trace Element | Calcium | 1.219 - 3.021 | 1.518 | Within range |
| Trace Element | Iron | 1.151 - 1.847 | 0.743 | Mildly reduced |
| Trace Element | Zinc | 1.143 - 1.989 | 1.336 | Within range |
| Trace Element | Selenium | 0.847 - 2.045 | 1.096 | Within range |
| Trace Element | Phosphorus | 1.195 - 2.134 | 2.087 | Within range |
| Trace Element | Potassium | 0.689 - 0.987 | 0.977 | Within range |
| Trace Element | Magnesium | 0.568 - 0.992 | 0.761 | Within range |
| Trace Element | Copper | 0.474 - 0.749 | 0.23 | Moderately reduced |
| Trace Element | Cobalt | 2.326 - 5.531 | 2.493 | Within range |
| Trace Element | Manganese | 0.497 - 0.879 | 0.732 | Within range |
| Trace Element | Iodine | 1.421 - 5.490 | 5.311 | Within range |
| Trace Element | Nickel | 2.462 - 5.753 | 2.481 | Within range |
| Trace Element | Fluorine | 1.954 - 4.543 | 2.019 | Within range |
| Trace Element | Molybdenum | 0.938 - 1.712 | 1.43 | Within range |
| Trace Element | Vanadium | 1.019 - 3.721 | 2.638 | Within range |

| Section | Marker | Range | Value | Status |
|---|---|---|---|---|
| Trace Element | Tin | 1.023 - 7.627 | 7.159 | Within range |
| Trace Element | Silicon | 1.425 - 5.872 | 1.367 | Mildly reduced |
| Trace Element | Strontium | 1.142 - 5.862 | 3.508 | Within range |
| Trace Element | Boron | 1.124 - 3.453 | 2.537 | Within range |
| Vitamin | Vitamin A | 0.346 - 0.401 | 0.338 | Mildly reduced |
| Vitamin | Vitamin B1 | 2.124 - 4.192 | 1.679 | Mildly reduced |
| Vitamin | Vitamin B2 | 1.549 - 2.213 | 1.549 | Within range |
| Vitamin | Vitamin B3 | 14.477 - 21.348 | 14.72 | Within range |
| Vitamin | Vitamin B6 | 0.824 - 1.942 | 1.616 | Within range |
| Vitamin | Vitamin B12 | 6.428 - 21.396 | 6.962 | Within range |
| Vitamin | Vitamin C | 4.543 - 5.023 | 3.55 | Moderately reduced |
| Vitamin | Vitamin D3 | 5.327 - 7.109 | 5.259 | Mildly reduced |
| Vitamin | Vitamin E | 4.826 - 6.013 | 4.11 | Moderately reduced |
| Vitamin | Vitamin K | 0.717 - 1.486 | 0.599 | Mildly reduced |
| Amino Acid | Lysine | 0.962 - 1.213 | 0.853 | Mildly reduced |
| Amino Acid | Tryptophan | 4.978 - 6.289 | 6.008 | Within range |
| Amino Acid | Phenylalanine | 1.928 - 2.491 | 2.057 | Within range |
| Amino Acid | Methionine | 1.245 - 1.637 | 1.398 | Within range |
| Amino Acid | Threonine | 1.194 - 1.685 | 1.37 | Within range |
| Amino Acid | Isoleucine | 4.582 - 5.657 | 4.175 | Mildly reduced |
| Amino Acid | Leucine | 6.982 - 9.256 | 7.041 | Within range |
| Amino Acid | Valine | 6.982 - 9.677 | 7.565 | Within range |
| Amino Acid | Histidine | 5.113 - 6.258 | 5.261 | Within range |
| Amino Acid | Arginine | 1.812 - 2.337 | 1.975 | Within range |
| Amino Acid | Homocysteine | 0.983 - 1.265 | 1.073 | Within range |

| Section | Marker | Range | Value | Status |
|---|---|---|---|---|
| Coenzyme | Nicotinamide | 2.074 - 3.309 | 2.293 | Within range |
| Coenzyme | Biotin | 1.833 - 2.979 | 1.241 | Mildly reduced |
| Coenzyme | Pantothenic acid | 1.116 - 2.101 | 0.604 | Moderately reduced |
| Coenzyme | Folic acid | 1.449 - 2.246 | 2.236 | Within range |
| Coenzyme | Coenzyme Q10 | 0.831 - 1.588 | 0.643 | Mildly reduced |
| Coenzyme | Glutathione | 0.726 - 1.281 | 1.114 | Within range |
| Essential Fatty Acid | Linoleic acid | 0.642 - 0.985 | 0.447 | Mildly reduced |
| Essential Fatty Acid | α-Linolenic acid | 0.814 - 1.202 | 0.852 | Within range |
| Essential Fatty Acid | γ-Linolenic acid | 0.921 - 1.334 | 0.996 | Within range |
| Essential Fatty Acid | Arachidonic acid | 0.661 - 0.808 | 0.575 | Mildly reduced |
| Endocrine System | Thyroid secretion index | 2.954 - 5.543 | 2.996 | Within range |
| Endocrine System | Parathyroid hormone secretion index | 2.845 - 4.017 | 2.121 | Mildly reduced |
| Endocrine System | Adrenal glands Index | 2.412 - 2.974 | 2.149 | Mildly reduced |
| Endocrine System | Pituitary secretion index | 2.163 - 7.34 | 5.512 | Within range |
| Endocrine System | Pineal secretion index | 3.210 - 6.854 | 6.778 | Within range |
| Endocrine System | Thymus gland secretion index | 2.967 - 3.528 | 2.969 | Within range |
| Endocrine System | Gland secretion index | 2.204 - 2.819 | 2.492 | Within range |
| Immune System | Lymph node Index | 133.437 - 140.47 | 134.342 | Within range |
| Immune System | Tonsil immune Index | 0.124 - 0.453 | 0.307 | Within range |
| Immune System | Bone marrow Index | 0.146 - 3.218 | 1.068 | Within range |
| Immune System | Spleen index | 34.367 - 35.642 | 35.434 | Within range |
| Immune System | Thymus index | 58.425 - 61.213 | 59.292 | Within range |
| Immune System | Immunoglobulin index | 3.712 - 6.981 | 4.089 | Within range |
| Immune System | Respiratory Immunity | 3.241 - 9.814 | 3.384 | Within range |
| Immune System | GI Immunity | 0.638 - 1.712 | 0.949 | Within range |

| Section | Marker | Range | Value | Status |
|---|---|---|---|---|
| Immune System | Mucosal Immunity | 4.111 - 18.741 | 10.307 | Within range |
| Thyroid | FT4 | 0.103 - 0.316 | 0.6 | Mildly elevated |
| Thyroid | Thyroglobulin | 0.114 - 0.202 | 0.294 | Mildly elevated |
| Thyroid | Thyroglobulin | 0.421 - 0.734 | 0.711 | Within range |
| Thyroid | T3 | 0.161 - 0.308 | 0.518 | Mildly elevated |
| Human Toxin | Stimulating Beverage | 0.209 - 0.751 | 0.475 | Within range |
| Human Toxin | Electromagnetic Radiation | 0.046 - 0.167 | 0.109 | Within range |
| Human Toxin | Free Radical | 0.124 - 0.453 | 0.178 | Within range |
| Human Toxin | Toxic Pesticide Residue | 0.013 - 0.313 | 0.305 | Within range |
| Heavy Metal | Lead | 0.052 - 0.643 | 0.976 | Mildly elevated |
| Heavy Metal | Mercury | 0.013 - 0.336 | 0.468 | Mildly elevated |
| Heavy Metal | Cadmium | 0.527 - 1.523 | 1.556 | Mildly elevated |
| Heavy Metal | Chromium | 0.176 - 1.183 | 0.345 | Within range |
| Heavy Metal | Arsenic | 0.153 - 0.621 | 0.596 | Within range |
| Heavy Metal | Antimony | 0.162 - 0.412 | 0.323 | Within range |
| Heavy Metal | Thallium | 0.182 - 0.542 | 0.388 | Within range |
| Heavy Metal | Aluminum | 0.192 - 0.412 | 0.57 | Mildly elevated |
| Basic Physical Quality | Response Ability | 59.786 - 65.424 | 61.674 | Within range |
| Basic Physical Quality | Mental Power | 58.715 - 63.213 | 61.23 | Within range |
| Basic Physical Quality | Water Shortage | 33.967 - 37.642 | 34.512 | Within range |
| Basic Physical Quality | Hypoxia | 133.642 - 141.476 | 130.994 | Mildly reduced |
| Basic Physical Quality | PH | 3.156 - 3.694 | 3.248 | Within range |
| Allergy | Drug Sensitivity | 0.431 - 1.329 | 0.563 | Within range |
| Allergy | Alcohol allergy index | 0.432 - 1.246 | 1.207 | Within range |
| Allergy | Pollen allergy index | 0.143 - 1.989 | 1.646 | Within range |

| Section | Marker | Range | Value | Status |
|---|---|---|---|---|
| Allergy | Injection allergy index | 0.847 - 1.045 | 1.184 | Mildly elevated |
| Allergy | Chemical Sensitivity | 0.842 - 1.643 | 1.896 | Mildly elevated |
| Allergy | Paint allergy index | 0.346 - 1.401 | 0.489 | Within range |
| Allergy | Dust allergy index | 0.543 - 1.023 | 1.263 | Mildly elevated |
| Allergy | Smoke allergy index | 0.826 - 1.013 | 2.532 | Mildly elevated |
| Allergy | Hair dye allergy index | 0.717 - 1.486 | 2.856 | Moderately elevated |
| Allergy | Animal fur allergy index | 0.124 - 1.192 | 0.571 | Within range |
| Allergy | Metal jewelry allergy index | 0.549 - 1.213 | 0.743 | Within range |
| Allergy | Seafood allergy index | 0.449 - 1.246 | 2.379 | Mildly elevated |
| Allergy | Milk allergy index | 0.477 - 1.348 | 0.739 | Within range |
| Obesity | Abnormal lipid metabolism coefficient | 1.992 - 3.713 | 0.89 | Moderately reduced |
| Obesity | Brown adipose tissue abnormalities coefficient | 2.791 - 4.202 | 1.858 | Moderately reduced |
| Obesity | Hyperinsulinemia coefficient | 0.097 - 0.215 | 0.206 | Within range |
| Obesity | Nucleus of the hypothalamus abnormal coefficient | 0.332 - 0.626 | 0.719 | Mildly elevated |
| Obesity | Triglyceride content of abnormal coefficient | 1.341 - 1.991 | 5.222 | Moderately elevated |
| Skin | Skin Free Radical Index | 0.124 - 3.453 | 2.972 | Within range |
| Skin | Skin Collagen Index | 4.471 - 6.079 | 2.899 | Mildly reduced |
| Skin | Skin Grease Index | 14.477 - 21.348 | 25.821 | Mildly elevated |
| Skin | Skin Immunity Index | 1.035 - 3.230 | 4.02 | Mildly elevated |
| Skin | Skin Moisture Index | 0.218 - 0.953 | 2.145 | Moderately elevated |
| Skin | Skin Moisture Loss | 2.214 - 4.158 | 3.125 | Within range |
| Skin | Skin Red Blood Trace Index | 0.824 - 1.942 | 1.055 | Within range |
| Skin | Skin Elasticity Index | 2.717 - 3.512 | 1.571 | Mildly reduced |

| Section | Marker | Range | Value | Status |
|---|---|---|---|---|
| Skin | Skin Melanin Index | 0.346 - 0.501 | 0.612 | Mildly elevated |
| Skin | Tin | 0.842 - 1.858 | 1.419 | Within range |
| Eye | Bags under the eyes | 0.510 - 3.109 | 3.004 | Within range |
| Eye | Collagen eye wrinkle | 2.031 - 3.107 | 1.31 | Mildly reduced |
| Eye | Dark circles | 0.831 - 3.188 | 3.883 | Mildly elevated |
| Eye | Lymphatic obstruction | 1.116 - 4.101 | 3.52 | Within range |
| Eye | Sagging | 0.233 - 0.559 | 0.751 | Mildly elevated |
| Eye | Edema | 0.332 - 0.726 | 1.234 | Moderately elevated |
| Eye | Eye cell activity | 0.118 - 0.892 | 0.895 | Mildly elevated |
| Eye | Visual fatigue | 2.017 - 5.157 | 8.04 | Mildly elevated |
| Collagen | Overall Eye Function | 6.352 - 8.325 | 8.31 | Within range |
| Collagen | Endocrine system | 7.245 - 8.562 | 7.491 | Within range |
| Collagen | Hair and skin | 4.533 - 6.179 | 4.806 | Within range |
| Collagen | Endocrine system | 6.178 - 8.651 | 8.253 | Within range |
| Collagen | Circulatory system | 3.586 - 4.337 | 4.153 | Within range |
| Collagen | Digestive system | 3.492 - 4.723 | 3.168 | Mildly reduced |
| Collagen | Immune system | 3.376 - 4.582 | 2.938 | Mildly reduced |
| Collagen | Motion systems | 6.458 - 8.133 | 5.573 | Mildly reduced |
| Collagen | Muscle Tissue | 6.552 - 8.268 | 5.774 | Mildly reduced |
| Collagen | Fat Metabolism | 6.338 - 8.368 | 4.551 | Mildly reduced |
| Collagen | Detoxification and metabolism | 6.187 - 8.466 | 6.519 | Within range |
| Collagen | Reproductive system | 3.778 - 4.985 | 3.997 | Within range |
| Collagen | Nervous system | 3.357 - 4.239 | 2.643 | Mildly reduced |
| Collagen | Skeleton | 6.256 - 8.682 | 7.975 | Within range |
| Channels and collaterals | Hand Tai Yin Lung Meridian | 48.264 - 65.371 | 63.778 | Within range |

| Section | Marker | Range | Value | Status |
|---|---|---|---|---|
| Channels and collaterals | Hand Yangming Large Intestine Meridian | 56.749 - 67.522 | 61.756 | Within range |
| Channels and collaterals | Foot Yangming Stomach Meridian | 0.481 - 1.043 | 0.485 | Within range |
| Channels and collaterals | Foot Tai Yin Lung Meridian | 0.327 - 0.937 | 0.465 | Within range |
| Channels and collaterals | Hand Shao Yin Heart Sutra | 1.672 - 1.978 | 1.477 | Mildly reduced |
| Channels and collaterals | Hand the small intestine by the sun | 0.192 - 0.412 | 0.199 | Within range |
| Channels and collaterals | Foot Tai Yang Bladder Meridian | 4.832 - 5.147 | 3.773 | Mildly reduced |
| Channels and collaterals | Foot Shao Yin Kidney | 3.321 - 4.244 | 3.937 | Within range |
| Channels and collaterals | Hand Jue Yin Pericardium Meridian | 1.338 - 1.672 | 1.34 | Within range |
| Channels and collaterals | Hand Shao Yang Triple Burner Meridian | 0.669 - 1.544 | 0.548 | Mildly reduced |
| Channels and collaterals | Foot Shao Yang Gall Bladder Meridian | 1.554 - 1.988 | 1.868 | Within range |
| Channels and collaterals | Foot Jue Yin Liver Meridian | 1.553 - 2.187 | 1.652 | Within range |
| Channels and collaterals | Ren channel | 11.719 - 18.418 | 12.572 | Within range |
| Channels and collaterals | Governor meridian | 0.708 - 1.942 | 1.536 | Within range |
| Channels and collaterals | Vital meridian | 6.138 - 21.396 | 6.331 | Within range |
| Channels and collaterals | Tai mai | 5.733 - 7.109 | 5.146 | Mildly reduced |
| Pulse of heart and brain | Stroke index | 60.735 - 65.396 | 62.076 | Within range |
| Pulse of heart and brain | Stroke volume(SV) | 63.012 - 67.892 | 61.901 | Mildly reduced |
| Pulse of heart and brain | Heart peripheral resistance (TRR) | 0.983 - 1.265 | 1.638 | Mildly elevated |
| Pulse of heart and brain | Pulse wave coefficient K | 0.316 - 0.401 | 0.391 | Within range |
| Pulse of heart and brain | Cerebrovascular blood oxygen saturation(Sa) | 0.710 - 1.109 | 0.894 | Within range |
| Pulse of heart and brain | Cerebrovascular blood oxygen volume(CaCO2) | 7.880 - 10.090 | 8.081 | Within range |
| Pulse of heart and brain | Cerebrovascular blood oxygen presure(PaO2) | 5.017 - 5.597 | 5.169 | Within range |
| Blood lipids | Blood viscosity | 4.131 - 4.562 | 5.233 | Moderately elevated |

| Section | Marker | Range | Value | Status |
|---|---|---|---|---|
| Blood lipids | Total cholesterol(TC) | 1.833 - 2.979 | 2.75 | Within range |
| Blood lipids | Triglyceride(TG) | 1.116 - 2.101 | 2.218 | Mildly elevated |
| Blood lipids | High-density lipoprotein (HDL-C) | 1.449 - 2.246 | 2.955 | Mildly elevated |
| Blood lipids | Low-density lipoprotein (LDL-C) | 0.831 - 1.588 | 1.015 | Within range |
| Blood lipids | Neutral fat(MB) | 0.726 - 1.281 | 3.096 | Mildly elevated |
| Blood lipids | Circulating immune complex(CIC) | 13.012 - 17.291 | 18.601 | Mildly elevated |
| Prostate | Degree of Prostatic Hyperplasia | 1.023 - 3.230 | 5.03 | Moderately elevated |
| Prostate | Degree of Prostatic Calcification | 1.471 - 6.079 | 8.907 | Mildly elevated |
| Prostate | Prostatitis Syndrome | 2.213 - 2.717 | 6.11 | Moderately elevated |
| Male Sexual Function | Testosterone | 3.342 - 9.461 | 2.492 | Moderately reduced |
| Male Sexual Function | Gonadotropin | 4.111 - 18.741 | 3.558 | Mildly reduced |
| Male Sexual Function | Erection Transmitter | 3.241 - 9.814 | 3.03 | Mildly reduced |
| Sperm and semen | Semen volume | 1.502 - 6.028 | 5.263 | Within range |
| Sperm and semen | Semen liquefying time | 10.283 - 30.282 | 26.982 | Within range |
| Sperm and semen | Number of sperms | 2.483 - 3.932 | 3 | Within range |
| Sperm and semen | Sperm motility rate | 0.637 - 0.877 | 0.689 | Within range |
| Male Hormone | Androgen | 1.372 - 1.878 | 1.492 | Within range |
| Male Hormone | Estrogen | 0.521 - 1.036 | 0.672 | Within range |
| Male Hormone | Progesterone(P) | 0.686 - 1.432 | 0.651 | Mildly reduced |
| Male Hormone | Luteinizing Hormone (LH) | 0.679 - 1.324 | 0.593 | Mildly reduced |
| Male Hormone | Prolactin(PRL) | 0.738 - 1.742 | 1.386 | Within range |
| Male Hormone | Follicle Stimulating Hormone(FSH) | 0.387 - 0.956 | 0.54 | Within range |
| Human Immunity | Tissue barrier immune index | 7.435 - 8.847 | 9.626 | Mildly elevated |
| Human Immunity | Innate immune cell index | 7.424 - 9.052 | 10.784 | Mildly elevated |

| Section | Marker | Range | Value | Status |
|---|---|---|---|---|
| Human Immunity | Innate immune molecule index | 7.143 - 9.214 | 7.585 | Within range |
| Human Immunity | Cellular immune index | 5.769 - 7.643 | 8.547 | Mildly elevated |
| Human Immunity | Humoral immunity index | 6.424 - 8.219 | 8.708 | Mildly elevated |
| Human Consciousness Level | Shame | 2.214 - 3.925 | 3.056 | Within range |
| Human Consciousness Level | Guilt | 2.154 - 3.478 | 2.543 | Within range |
| Human Consciousness Level | Apathy | 2.343 - 4.089 | 2.263 | Mildly reduced |
| Human Consciousness Level | Grief | 1.845 - 3.241 | 1.997 | Within range |
| Human Consciousness Level | Fear | 2.155 - 4.031 | 1.681 | Mildly reduced |
| Human Consciousness Level | Desire | 1.484 - 2.485 | 1.711 | Within range |
| Human Consciousness Level | Anger | 1.256 - 2.529 | 1.466 | Within range |
| Human Consciousness Level | Pride | 1.378 - 2.489 | 1.025 | Mildly reduced |
| Human Consciousness Level | Courage | 1.727 - 3.315 | 1.8 | Within range |
| Human Consciousness Level | Neutrality | 2.471 - 3.892 | 2.16 | Mildly reduced |
| Human Consciousness Level | Willingness | 2.216 - 4.094 | 2.598 | Within range |
| Human Consciousness Level | Acceptance | 1.668 - 4.053 | 2.239 | Within range |
| Human Consciousness Level | Reason | 1.352 - 3.436 | 2.693 | Within range |
| Human Consciousness Level | Love | 2.138 - 3.754 | 1.871 | Mildly reduced |
| Human Consciousness Level | Joy | 2.129 - 4.021 | 2.654 | Within range |
| Human Consciousness Level | Peace | 2.243 - 4.022 | 2.099 | Mildly reduced |
| Human Consciousness Level | Enlightenment | 2.225 - 3.822 | 3.556 | Within range |
| Respiratory Function | Volume Tidal(VT) | 5.431 - 7.479 | 4.051 | Mildly reduced |
| Respiratory Function | Inspiratory Volume(TI) | 4.126 - 6.045 | 4.442 | Within range |
| Respiratory Function | Functional Residual Capacity(FRC) | 5.147 - 6.219 | 5.503 | Within range |
| Respiratory Function | Residual Volume(RV) | 4.363 - 5.647 | 5.322 | Within range |
| Lecithin | Phospholipid index | 3.654 - 4.478 | 4.471 | Within range |
| Lecithin | Sphingolipid index | 3.121 - 3.853 | 2.842 | Mildly reduced |

| Section | Marker | Range | Value | Status |
|---|---|---|---|---|
| Lecithin | PH | 3.341 - 4.214 | 3.894 | Within range |
| Lecithin | Lecithin index | 4.365 - 5.142 | 4.165 | Within range |
| Lecithin | Brain phospholipid index | 4.225 - 5.713 | 4.038 | Mildly reduced |
| Lecithin | Liposomal index | 3.112 - 4.081 | 4.071 | Within range |
| Fatty acid | Saturated fatty acid index | 1.334 - 2.407 | 1.437 | Within range |
| Fatty acid | Unsaturated fatty acid index | 2.224 - 3.153 | 2.834 | Within range |
| Fatty acid | Essential fatty acid index | 2.144 - 3.238 | 2.087 | Mildly reduced |
| Fatty acid | Triglyceride index | 3.361 - 4.644 | 2.377 | Mildly reduced |

These scan findings are best interpreted alongside symptoms, history, and appropriate clinical assessment.