

---

# Eric Wan - [ezw23@drexel.edu](mailto:ezw23@drexel.edu) - Lab3

## Table of Contents

Problem 1 - Eulers .....	1
Problem 2 - Runge-Kutta .....	2
Problem 3 - ODE45 .....	3
Problem 4 - Analytical .....	4
Question 5 - Commenting on Solutions .....	5

## Problem 1 - Eulers

```
step of 10

dt = 10;
pI = 3/15000;
tI = 0;
tEnd = 500;
tSpan = tI:dt:tEnd;
p = zeros(size(tSpan));
p(1) = pI;
formatSpec = 'For a step of %.2f, half of population infected at %.2f
    hours with ratio of %.4f for Eulers \n';
type ebola.m
for i=2:length(tSpan)
    pprime = ebola(tSpan(i-1), p(i-1));
    p(i) = p(i-1) + dt*pprime;
end
figure
plot(tSpan,p,'b')
grid on
hold on
legend show
xlabel('Time')
ylabel('P(t)')
title('First Order DiffEq Analysis')
fprintf(formatSpec, dt, tSpan(26), p(26))

% step of 5
dt = 5;
tSpan = tI:dt:tEnd;
p = zeros(size(tSpan));
p(1) = pI;
for i=2:length(tSpan)
    pprime = ebola(tSpan(i-1), p(i-1));
    p(i) = p(i-1) + dt*pprime;
end
plot(tSpan,p,'r')
fprintf(formatSpec, dt, tSpan(47), p(47))

% step of 1
```


```
dt = 1;  
tSpan = tI:dt:tEnd;  
p = zeros(size(tSpan));  
p(1) = pI;  
for i=2:length(tSpan)  
    pprime = ebola(tSpan(i-1), p(i-1));  
    p(i) = p(i-1) + dt*pprime;  
end  
plot(tSpan,p,'g')  
fprintf(formatSpec, dt, tSpan(218), p(218))
```

```
function [ dPdt ] = ebola( t, P )  
dPdt = 0.04*P*(1-P);  
end
```

*For a step of 10.00, half of population infected at 250.00 hours with ratio of 0.5298 for Eulers*

*For a step of 5.00, half of population infected at 230.00 hours with ratio of 0.4967 for Eulers*

*For a step of 1.00, half of population infected at 217.00 hours with ratio of 0.5053 for Eulers*



## Problem 2 - Runge-Kutta

```
dt = 5;
```


```

tSpan = tI:dt:tEnd;
k = zeros(size(tSpan));
k(1) = pI;

for i=2:length(tSpan)
k1 = ebola(tSpan(i-1), k(i-1));
k2 = ebola((tSpan(i-1)+(dt/2)), (k(i-1) + (dt/2*k1)));
k(i) = k(i-1) + dt*k2;
end
plot(tSpan,k,'k')
formatSpec = 'For a step of %.2f, half of population infected at %.2f
hours with ratio of %.4f for Runge-Kutta\n';
fprintf(formatSpec, dt, tSpan(44), k(44))

```

*For a step of 5.00, half of population infected at 215.00 hours with ratio of 0.5116 for Runge-Kutta*



## Problem 3 - ODE45


```

dt = 5;
tSpan = tI:dt:tEnd;
p(1) = pI;
[tode, pode] = ode45(@ebola, [tSpan], pI);
plot(tode,pode,'m')
legend('dt = 10', 'dt = 5', 'dt = 1', 'Runge-Kutta', 'ODE45')

```

```
formatSpec = 'For a step of %.2f, half of population infected at %.2f
hours with ratio of %.4f for ODE45\n';
fprintf(formatSpec, dt, tode(44), pde(44))
```

*For a step of 5.00, half of population infected at 215.00 hours with ratio of 0.5211 for ODE45*



## Problem 4 - Analytical

rewriting eulers step of 5

```
dt = 5;
tSpan = tI:dt:tEnd;
p = zeros(size(tSpan));
p(1) = pI;
for i=2:length(tSpan)
    pprime = ebola(tSpan(i-1), p(i-1));
    p(i) = p(i-1) + dt*pprime;
end

n = numel(tSpan);
pts = zeros(1,n);
for i = 1:n
    pa = exp(0.04*tSpan(i))/(exp(0.04*tSpan(i)) + 4999);
    pts(i) = pa;
end
```

```

plot(tSpan, pts, 'y')
legend('dt = 10', 'dt = 5', 'dt = 1', 'Runge-
Kutta', 'ODE45', 'Analytical')


err1 = mean((pts - p).^2); % mse between Analytical and Euler
fprintf('The MSE between Analytical and Euler is %f\n', err1)
err2 = mean((pts - pde).^2); % mse between Analytical and ODE45
fprintf('The MSE between Analytical and ODE45 is %f\n', err2)
err3 = mean((pts - k).^2); % mse between Analytical and Runge-Kutta
fprintf('The MSE between Analytical and Runge-Kutta is %f\n', err3)

```

The MSE between Analytical and Euler is 0.003660

The MSE between Analytical and ODE45 is 0.000000

The MSE between Analytical and Runge-Kutta is 0.000011



## Question 5 - Commenting on Solutions

```

%{
ODE45 is the most accurate with very little MSE error to the
analytical
solution
Runge-Kutta is very close to the analytical solution
Eulers is the farthest off from the analytical solution
%}

```

*Published with MATLAB® R2015a*