

Introducing..... Linear Regression!

Frank Edwards


We can define a line as:

$$y = mx + b$$

Where m is the slope and b is the y-intercept.


Lines: slopes

$$y = x + 2$$




Lines: slopes

$$y = 2x + 2$$




Lines: slopes

$$y = -2x + 2$$




Lines: slopes

$$y = 0.2x + 2$$




Lines: intercepts

$$y = x + 0$$




Lines: intercepts

$$y = x + 2$$



Lines: intercepts

$$y = x - 3$$



The linear regression model

We can describe the relationship between a predictor variable X and an outcome variable Y with a line:

$$y = mx + b$$

What does m describe?

The linear regression model

We can describe the relationship between a predictor variable X and an outcome variable Y with a line:

$$y = mx + b$$

What does m describe?

- The increase in y for a one-unit increase in x

What does b describe

The linear regression model

We can describe the relationship between a predictor variable X and an outcome variable Y with a line:

$$y = mx + b$$

What does m describe?

- The increase in y for a one-unit increase in x

What does b describe

- The location of y when $x = 0$

The linear regression model: expected value

We can describe the relationship between a predictor variable X and the expected value E of an outcome variable Y with the line:

$$E[Y] = \beta_0 + \beta_1 X$$

The linear regression model: expected value

We can describe the relationship between a predictor variable X and the expected value E of an outcome variable Y with the line:

$$E[Y] = \beta_0 + \beta_1 X$$

What does β_0 describe?

The linear regression model: expected value

We can describe the relationship between a predictor variable X and the expected value E of an outcome variable Y with the line:

$$E[Y] = \beta_0 + \beta_1 X$$

What does β_0 describe?

What does β_1 describe?

The error term in linear regression

We can describe the relationship between a predictor variable X and an outcome variable Y with the line:

$$Y = \beta_0 + \beta_1 X + \varepsilon$$

Where β_0 is the y-intercept of the line, β_1 is the slope of the line, and ε is the error between the fitted line and the coordinates (X, Y)

The linear regression model

$$Y = \beta_0 + \beta_1 X + \varepsilon$$

β_0 : The value of y when x is equal to zero

The linear regression model

$$Y = \beta_0 + \beta_1 X + \varepsilon$$

β_0 : The value of y when x is equal to zero

β_1 : The average increase in y when x increases by one unit

The linear regression model

$$Y = \beta_0 + \beta_1 X + \varepsilon$$

β_0 : The value of y when x is equal to zero

β_1 : The average increase in y when x increases by one unit

ε : The distance between the line $y = \beta_0 + \beta_1 X$ and the actual observed values of y .

The linear regression model

$$Y = \beta_0 + \beta_1 X + \varepsilon$$

β_0 : The value of y when x is equal to zero

β_1 : The average increase in y when x increases by one unit

ε : The distance between the line $y = \beta_0 + \beta_1 X$ and the actual observed values of y .

The line $y = \beta_0 + \beta_1 X$ provides an expected value for y based on the values of x .

The linear regression model as a prediction engine

- If $\beta_0 = 2$ and $\beta_1 = 1.5$, what is the expected value of y when $x = 4$?

The linear regression model as a prediction engine

- If $\beta_0 = 2$ and $\beta_1 = 1.5$, what is the expected value of y when $x = 4$?
- When $x = 2$?

The linear regression model and prediction

We put a *hat* on variables to indicate that they are estimated from the data, or predicted.

A regression line predicts values Y , \hat{Y} with the equation:

$$\hat{Y} = \beta_0 + \beta_1 X$$

and the residual, or prediction error is the difference between the observed and predicted values of Y

$$\varepsilon = Y_{obs} - \hat{Y}$$

Understanding the regression line for real data

```
## # A tibble: 10 x 2
##       x       y
##   <dbl> <dbl>
## 1 -0.311 -0.237
## 2  0.932  0.883
## 3  1.04   1.62
## 4  3.55   3.41
## 5  0.507  0.254
## 6  0.655  1.11
## 7  1.39   1.29
## 8  0.868  0.477
## 9  0.859  0.561
## 10 0.509  0.626
```

$$\beta_0 = 0.05, \beta_1 = 0.95$$

- Estimate \hat{Y} . Recall that $\hat{Y} = \beta_0 + \beta_1 X$

Understanding the regression line for real data


```
## # A tibble: 10 x 2
##       x       y
##   <dbl> <dbl>
## 1 -0.311 -0.237
## 2  0.932  0.883
## 3  1.04   1.62
## 4  3.55   3.41
## 5  0.507  0.254
## 6  0.655  1.11
## 7  1.39   1.29
## 8  0.868  0.477
## 9  0.859  0.561
## 10 0.509  0.626
```

$$\beta_0 = 0.05, \beta_1 = 0.95$$

- Estimate \hat{Y} . Recall that $\hat{Y} = \beta_0 + \beta_1 X$
- Estimate ε . Recall that $\varepsilon = Y - \hat{Y}$


Understanding the regression line

$$\beta_0 = 0.05, \beta_1 = 0.95$$




Understanding the regression line: adding the fit

$$\beta_0 = 0.05, \beta_1 = 0.95$$




Understanding the regression line: adding \hat{y}

$$\beta_0 = 0.05, \beta_1 = 0.95$$



Understanding the regression line: adding ε

$$\beta_0 = 0.05, \beta_1 = 0.95$$



Ordinary least squares

We usually fit a linear regression using a method called *ordinary least squares*, or OLS.

Ordinary least squares

We usually fit a linear regression using a method called *ordinary least squares*, or OLS.

- This method seeks to minimize the distance between \hat{Y} and Y .

Ordinary least squares

We usually fit a linear regression using a method called *ordinary least squares*, or OLS.

- This method seeks to minimize the distance between \hat{Y} and Y .
- To do so, we minimize the sum of squared residuals (SSR)

Ordinary least squares

We usually fit a linear regression using a method called *ordinary least squares*, or OLS.

- This method seeks to minimize the distance between \hat{Y} and Y .
- To do so, we minimize the sum of squared residuals (SSR)

In other words, we solve for the values of β_0 and β_1 that result in the smallest possible value for SSR

$$\text{SSR} = \sum_{i=1}^n \varepsilon_i^2 = \sum_{i=1}^n (y_i - \beta_0 - \beta_1 x_i)^2$$

Ordinary least squares

We usually fit a linear regression using a method called *ordinary least squares*, or OLS.

- This method seeks to minimize the distance between \hat{Y} and Y .
- To do so, we minimize the sum of squared residuals (SSR)

In other words, we solve for the values of β_0 and β_1 that result in the smallest possible value for SSR

$$\text{SSR} = \sum_{i=1}^n \varepsilon_i^2 = \sum_{i=1}^n (y_i - \beta_0 - \beta_1 x_i)^2$$

Estimating a regression model in R, the basics

Let's use `iris` to estimate this regression model:

$$\text{Sepal.Length} = \beta_0 + \beta_1 \text{Petal.Length} + \varepsilon$$

```
my_cool_model <- lm(Sepal.Length ~ Petal.Length,  
                    data = iris)
```

```
my_cool_model
```

```
##  
## Call:  
## lm(formula = Sepal.Length ~ Petal.Length, data = iris)  
##  
## Coefficients:  
## (Intercept)  Petal.Length  
##          4.3066          0.4089
```

What does this model tell us?

$$\cdot \beta_0 = 4.3$$

What does this model tell us?


- $\beta_0 = 4.3$
- $\beta_1 = 0.4$

What does this model tell us?


- $\beta_0 = 4.3$
- $\beta_1 = 0.4$

$$E(\text{Sepal.Length}) = \beta_0 + \beta_1 \text{Petal.Length}$$


Visualized: observed data



Visualized: regression line



Visualized: expected values (\hat{y})



Visualized: residuals (epsilon)

