

Contents

| | |
|-------------------------------|----|
| 1. Topology | 5 |
| 2. Measure Theory | 8 |
| 3. Probability Theory | 9 |
| 4. Wasserstein Distance | 10 |

1. Topology

Definition 1.1 (Topology):

Let X be a set. A **topology over X** is a subset Σ of 2^X such that:

1. $A \subseteq \Sigma \implies \bigcup_{E \in A} E$. **Infinite or finite unions of sets.**
2. $A, B \in \Sigma \implies A \cap B \in \Sigma$. **Finite intersections of sets.**
3. $X \in \Sigma$

Definition 1.2 (Topological Space):

(X, Σ) is a **topological space** iff. Σ is a **topology** of X .

Definition 1.3 (Everywhere dense):

Let (X, Σ) **topological space**, and $H \subseteq X$. H is said **everywhere dense in Σ** iff. $\forall E \in \Sigma, E \neq \emptyset : H \cap E \neq \emptyset$. **We can find a bit of H in every corner of the topology Σ .**

Definition 1.4 (Separable):

Let (X, Σ) be a **topological space**. (X, Σ) is said **separable** iff $\exists H \subseteq X, H$ is countable : H is **everywhere dense in Σ** . **There is a sequence of elements $\{x_n \in X\}_{n=1}^{\infty}$ such that every set in the topology contains at least one element x_i .**

Definition 1.5 (Metric Space):

(X, d) is a **metric space** iff.

1. $X \neq \emptyset$
2. $d : X \times X \longrightarrow \mathbb{R}_{\geq 0}$ such that (**d is a distance**):
 1. $\forall x, y \in X : d(x, y) = 0 \implies x = y$. **there are no different elements at zero-distance.**
 2. $\forall x, y \in X : d(x, y) = d(y, x)$. **symmetry.**
 3. $\forall x, y, z \in X : d(x, z) \leq d(x, y) + d(y, z)$. **triangular inequality.**

Definition 1.6 (open ε -ball):

Let (X, d) be a **metric space**, $x \in X$, and $\varepsilon \in \mathbb{R}_{>0}$. We call $B_{\varepsilon}(x) = \{y \in X \mid d(x, y) < \varepsilon\}$ an **open ε -ball**. **A ball of ε radius centered at some point.**

Definition 1.7 (Neighborhood):

Let (X, d) be a **metric space**, $S \subseteq X$, $x \in S$, and $\varepsilon \in \mathbb{R}_{>0}$ such that the **open ε -ball** $B_\varepsilon(x) \subseteq S$. Then S is said a **neighborhood of x** . A neighborhood of an element is simply a set that contains an open ball containing the element.

Definition 1.8 (Open Set):

Let (X, d) be a **metric space** and $U \subseteq X$. U is an **open set** iff. $\forall u \in U : \exists \varepsilon \in \mathbb{R}_{>0} : B_\varepsilon(u) \subseteq U$. An open set is simply a set which is also neighborhood for all its points.

Definition 1.9 (Induced Topology):

Let (X, d) be a **metric space**. Σ is said an **induced topology** iff. $\Sigma = \{U \subseteq X \mid U \text{ is an open set in } (X, d)\}$

Definition 1.10 (Metrizible):

Let (X, Σ) be a **topological space**. (X, Σ) is said **metrizable** iff. $\exists (X, d)$ **metric space** : Σ is a **topology induced** by (X, d) .

Definition 1.11 (Cauchy Sequence):

Let (X, d) be a **metric space**, $[x_n \in X]$ a sequence. $[x_n]$ is said a **cauchy sequence** iff. $\forall \varepsilon \in \mathbb{R}_{>0} : \exists N \in \mathbb{N} : \forall m, n \in \mathbb{N} : d(x_n, x_m) \leq \varepsilon$. There is a point after which all pairs of elements are close to each other.

Definition 1.12 (Convergent Sequence):

Let (X, d) be a **metric space**, $l \in X$, $[x_n \in X]$ a sequence. $[x_n]$ is said a **convergent sequence to the limit l** iff. $\forall \varepsilon \in \mathbb{R}_{>0} : \exists N \in \mathbb{R}_{>0} : \forall n > N : d(x_n, l) < \varepsilon$. If such a limit exists the sequence is simply said **convergent**.

Definition 1.13 (Complete Metric Space):

Let (X, d) be a **metric space**. (X, d) is said a **complete metric space** iff. every **cauchy sequence** is **convergent**.

Definition 1.14 (Polish Space):

Let (X, Σ) be a topological space. (X, Σ) is said a **Polish Space** iff. (X, Σ) is separable, metrizable, and a complete metric space for some metric.

2. Measure Theory

Definition 2.1 (σ -algebra):

Let X be a set. $\Sigma \subseteq 2^X$ is said a **sigma algebra of X** iff.:

1. $X \in \Sigma$
2. $E \in \Sigma \implies X \setminus E \in \Sigma$. **close under complement.**
3. $\{A_n \in \Sigma\}_{n=1}^{\infty} \implies \bigcup_{i=1}^{\infty} A_i \in \Sigma$. **close under infinite unions.**

Definition 2.2 (generate σ -algebra):

Let X be a set and $G \subseteq 2^X$. The **σ -algebra generated by G** , denoted $\sigma(G)$, is the smallest **σ -algebra** such that:

1. $G \subseteq \sigma(G)$.
2. $\forall \Sigma$ σ -algebra : $G \subseteq \Sigma \implies \sigma(G) \subseteq \Sigma$. **Every other σ -algebra that contains G contains also the generated one, $\sigma(G)$.**

Definition 2.3 (σ -algebra product):

Let Σ_1 and Σ_2 be **σ -algebras** on X_1 and X_2 respectively. The **product σ -algebra** denoted $\Sigma_1 \otimes \Sigma_2$ is defined as **$\sigma(\{S_1 \times S_2 \mid S_1 \in \Sigma_1, S_2 \in \Sigma_2\})$**

Definition 2.4 (measurable space):

(X, Σ) is said **measurable** iff. Σ is a **sigma-algebra** of X .

Definition 2.5 (measure):

Given (X, Σ) **measurable space**. $\mu : \Sigma \longrightarrow \mathbb{R} \cup \{+\infty, -\infty\}$ is said a **measure** iff.

1. $E \in \Sigma \implies \mu(E) \geq 0$. **positive.**
2. $\{E_n \in \Sigma\}_{n=1}^{\infty}$ such that $E_i \cap E_j = \emptyset$ for $i \neq j \implies \mu\left(\bigcup_{i=1}^{\infty} E_i\right) = \sum_{i=1}^{\infty} \mu(E_i)$. **The measure of disjoint sets is the sum of the measures of each set.**
3. $\exists E \in \Sigma : \mu(E) < \infty$. **For at least an element μ is finite.**

Definition 2.6 (measure space):

(X, Σ, μ) is said a **measure space** iff. (X, Σ) is a **sigma algebra** and μ is a **measure** of (X, Σ) .

3. Probability Theory

Definition 3.1 (Probability Space):

(Ω, Σ, p) is said a **probability space** iff.

1. (Ω, Σ, p) is a **measure space**.
2. $p(\Omega) = 1$.

Intuitively, Ω represents the set of all possible outcomes, it is also known as **sample space**. Σ represents the set of all possible events. These are nothing more than set of outcomes. It is also known as **event space**. p is a measure on the event space, it is also known as **probability function**. It maps events to their likelihood.

Example 3.1 (Fair Die):

Consider the probability space (Ω, Σ, p) , where:

1. $\Omega = \{1, 2, 3, 4, 5, 6\}$ is the sample space, representing the possible outcomes of rolling a standard six-sided die.
2. $\Sigma = 2^\Omega$ is the event space.
3. $p : \Sigma \rightarrow [0, 1]$ is the probability measure function, defined as $P(E) = \frac{|E|}{6}$ for any event $E \in \Sigma$.

For example, consider the event $A = \{1, 2, 3\}$, which represents rolling a 1, 2, or 3. This event is an element of Σ . The probability of event A occurring is $p(A) = \frac{|A|}{6} = \frac{3}{6} = \frac{1}{2}$.

Definition 3.2 (Coupling):

Let $(\Omega_1, \Sigma_1, \mu_1)$ and $(\Omega_2, \Sigma_2, \mu_2)$ be **probability spaces**. A **coupling** is a **probability space** $(\Omega_1 \times \Omega_2, \Sigma_1 \otimes \Sigma_2, \gamma)$ such that:

1. $\forall E \in \Sigma_1 : \gamma(E \times \Omega_2) = \mu_1(E)$.
2. $\forall E \in \Sigma_2 : \gamma(\Omega_1 \times E) = \mu_2(E)$.

4. Wasserstein Distance
