

# Discrete Morse Theory and Rips' Lemma for Trees

Fabian Roll (TUM)

DGD Days 2021  
15.-17.09.21


Project: C04 - Persistence and Stability of Geometric Complexes

Advisor: Prof. Dr. Ulrich Bauer

# Background

**Definition.**  $X$  metric space. The Vietoris-Rips complex at scale  $r$  is the simplicial complex


$$\text{VR}_r(X) = \{S \subseteq X \text{ finite} \mid S \neq \emptyset, \text{diam } S \leq r\}.$$



$r = 3, 6, 9$


# Background

## Persistent Homology



# Background

## Persistent Homology



Software: U. Bauer “Ripser: [efficient](#) computation of Vietoris-Rips persistence barcodes”

# Geometry and Combinatorics of Input influences Computation Time


5000 ordered points	time for $H_0$ & $H_1$
random in $\mathbb{R}^3$	1m 17s
random on $S^2$	5m 39s
graph (covid data)	12s

# Geometry and Combinatorics of Input influences Computation Time

5000 ordered points	time for $H_0$ & $H_1$
random in $\mathbb{R}^3$	1m 17s
random on $S^2$	5m 39s
graph (covid data)	12s
graph (covid data, reversed order)	6m 19s
graph (covid data, random order)	2m 52s

# Gromov Hyperbolicity


**Definition.** A **length space** is  $\delta$ -hyperbolic if all triangles are  $\delta$ -slim.



Wikipedia

# Gromov Hyperbolicity


**Definition.** A metric space is  $\delta$ -hyperbolic if for all four points





# Gromov Hyperbolicity

**Definition.** A metric space is  $\delta$ -hyperbolic if for all four points




w.l.o.g.  $l \geq l \geq l$

we have  $\delta \geq l - l$ .

# Gromov Hyperbolicity

**Definition.** A metric space is  $\delta$ -hyperbolic if for all four points




w.l.o.g.  $\ell \geq \ell \geq \ell$

we have  $\delta \geq \ell - \ell$ .

**Example.** finite metric space, trees are 0-hyperbolic, hyperbolic plane, ...

# Gromov Hyperbolicity

**Definition.** A metric space is  $\delta$ -hyperbolic if for all four points




w.l.o.g.  $\ell \geq \ell \geq \ell$

we have  $\delta \geq \ell - \ell$ .


**Example.** finite metric space, trees are 0-hyperbolic, hyperbolic plane, ...

**Lemma (Rips).**  $X$  length space and  $\delta$ -hyperbolic. Then,  $VR_r(X)$  is contractible for  $r \geq 4\delta$ .


# Discrete Morse Theory



# Discrete Morse Theory



# Discrete Morse Theory



# Apparent Pairs

Persistence computation: choose total order on  $K$

# Apparent Pairs

Persistence computation: choose total order on  $K$

**Definition.**  $(\sigma, \tau)$  apparent pair if and only if

- ▶  $\sigma$  maximal facet of  $\tau$  and
- ▶  $\tau$  minimal cofacet of  $\sigma$

in the chosen total order.




# Apparent Pairs

Persistence computation: choose total order on  $K$

**Definition.**  $(\sigma, \tau)$  apparent pair if and only if

- ▶  $\sigma$  maximal facet of  $\tau$  and
- ▶  $\tau$  minimal cofacet of  $\sigma$

in the chosen total order.




# Apparent Pairs

Persistence computation: choose total order on  $K$

**Definition.**  $(\sigma, \tau)$  apparent pair if and only if

- ▶  $\sigma$  maximal facet of  $\tau$  and
- ▶  $\tau$  minimal cofacet of  $\sigma$

in the chosen total order.



**Lemma.** The apparent pairs form a discrete gradient.


# Apparent Pairs

Persistence computation: choose total order on  $K$

**Definition.**  $(\sigma, \tau)$  apparent pair if and only if

- ▶  $\sigma$  maximal facet of  $\tau$  and
- ▶  $\tau$  minimal cofacet of  $\sigma$


in the chosen total order.



**Lemma.** The apparent pairs form a discrete gradient. → Ripser


# Results

- ▶  $X$  vertex set of a directed rooted tree
- ▶ sort simplices of  $VR_{\infty}(X)$  by diameter, then lexicographically



# Results

- ▶  $X$  vertex set of a directed rooted tree
- ▶ sort simplices of  $VR_\infty(X)$  by diameter, then lexicographically




**Theorem.** If vertices are ordered in the direction of the tree and  $r \geq r_c = (\text{connectivity threshold})$ , then

$$VR_r(X) \searrow \{0\}$$

by the apparent pairs gradient

# Results

- ▶  $X$  vertex set of a directed rooted tree
- ▶ sort simplices of  $VR_\infty(X)$  by diameter, then lexicographically



**Theorem.** If vertices are ordered in the direction of the tree and  $r \geq r_c = (\text{connectivity threshold})$ , then


$$VR_r(X) \searrow \{0\}$$

by the apparent pairs gradient

**Remark.**  $r \geq r_c = 4\delta + r_c$  with  $\delta = 0$

# Results

- ▶  $X$  vertex set of a directed rooted tree
- ▶ sort simplices of  $VR_\infty(X)$  by diameter, then lexicographically




**Theorem.** If vertices are ordered in the direction of the tree and  $r \geq r_c = (\text{connectivity threshold})$ , then

$$VR_r(X) \searrow \{0\}$$

by the apparent pairs gradient


**Remark.**  $r \geq r_c = 4\delta + r_c$  with  $\delta = 0 \rightarrow$  similar to Rips' lemma

For general  $X$  sort points:





For general  $X$  sort points:



5000 ordered points	time for $H_0$ & $H_1$	non-apparent
random tree	7s	0
graph (covid data)	12s	4544
graph (covid data, reversed order)	6m 19s	19344
graph (covid data, random order)	2m 52s	13957

# Future Work

- ▶ find analogous statements for general  $\delta$ -hyperbolic spaces with sharp bounds
- ▶ use results/heuristics to find approximation algorithms for Vietoris-Rips persistent homology