

Connecting Morse theory and persistent homology of geometric complexes

Fabian Roll (TUM)


ÖMG Meeting 2023, Graz
Minisymposium: Computational Topology and Geometry

Joint work with Ulrich Bauer

Čech complexes

Definition. The *Čech complex* $\check{\text{C}}\text{ech}_r(X)$ of $X \subseteq \mathbb{R}^d$ is the simplicial complex

$$\check{\text{C}}\text{ech}_r(X) = \{J \subseteq X \mid |J| < \infty \text{ and } \bigcap_{y \in J} D_r(y) \neq \emptyset\}.$$



Nerves

Definition (Alexandroff 1928). Let $\mathcal{U} = (U_i)_{i \in I}$ be a cover of a space X . The *nerve* of \mathcal{U} is the simplicial complex

$$\text{Nrv}(\mathcal{U}) = \{J \subseteq I \mid |J| < \infty \text{ and } \bigcap_{i \in J} U_i \neq \emptyset\}$$

recording the intersection pattern of the cover.

Nerves

Definition (Alexandroff 1928). Let $\mathcal{U} = (U_i)_{i \in I}$ be a cover of a space X . The *nerve* of \mathcal{U} is the simplicial complex

$$\text{Nrv}(\mathcal{U}) = \{J \subseteq I \mid |J| < \infty \text{ and } \bigcap_{i \in J} U_i \neq \emptyset\}$$

recording the intersection pattern of the cover.

Theorem (Borsuk 1948, and many more). Let \mathcal{U} be a nice cover of X . Then $\text{Nrv}(\mathcal{U})$ is homotopy equivalent to X .

Nerves

Definition (Alexandroff 1928). Let $\mathcal{U} = (U_i)_{i \in I}$ be a cover of a space X . The *nerve* of \mathcal{U} is the simplicial complex

$$\text{Nrv}(\mathcal{U}) = \{J \subseteq I \mid |J| < \infty \text{ and } \bigcap_{i \in J} U_i \neq \emptyset\}$$

recording the intersection pattern of the cover.

Theorem (Borsuk 1948, and many more). Let \mathcal{U} be a nice cover of X . Then $\text{Nrv}(\mathcal{U})$ is homotopy equivalent to X .

Here *nice* can mean different things:

- open, numerable cover, contractible intersections
- finite, closed, convex cover




U. Bauer, M. Kerber, F. Roll, and A. Rolle


A unified view on the functorial nerve theorem and its variations

Expositiones Mathematicae, 2023. doi:10.1016/j.exmath.2023.04.005

Persistent homology



Persistent homology



Persistent homology

The persistent homology of a filtration L_\bullet can be computed by the *exhaustive Matrix reduction* algorithm that

Persistent homology

The persistent homology of a filtration L_\bullet can be computed by the *exhaustive Matrix reduction* algorithm that

- requires a total order on the simplices $\sigma_1 < \dots < \sigma_n$ inducing a simplexwise refinement $K_\bullet = (K_i = \{\sigma_1, \dots, \sigma_i\})_i$ of L_\bullet

Persistent homology

The persistent homology of a filtration L_\bullet can be computed by the *exhaustive Matrix reduction* algorithm that

- requires a total order on the simplices $\sigma_1 < \dots < \sigma_n$ inducing a simplexwise refinement $K_\bullet = (K_i = \{\sigma_1, \dots, \sigma_i\})_i$ of L_\bullet .
- uses a variant of Gaussian elimination:
 - ▶ $R = D$ boundary matrix (\mathbb{Z}_2 coefficients) of K_\bullet ,
 - $V = I$ identity matrix

Persistent homology

The persistent homology of a filtration L_\bullet can be computed by the *exhaustive Matrix reduction* algorithm that

- requires a total order on the simplices $\sigma_1 < \dots < \sigma_n$ inducing a simplexwise refinement $K_\bullet = (K_i = \{\sigma_1, \dots, \sigma_i\})_i$ of L_\bullet .
- uses a variant of Gaussian elimination:
 - ▶ $R = D$ boundary matrix (\mathbb{Z}_2 coefficients) of K_\bullet ,
 - ▶ $V = I$ identity matrix
 - ▶ while $\exists i < j$ with $(R_j)_k \neq 0$, where $k = \text{pivot } R_i$
 - ▶ add R_i to R_j
 - ▶ add V_i to V_j

Persistent homology

The persistent homology of a filtration L_\bullet can be computed by the *exhaustive Matrix reduction* algorithm that

- requires a total order on the simplices $\sigma_1 < \dots < \sigma_n$ inducing a simplexwise refinement $K_\bullet = (K_i = \{\sigma_1, \dots, \sigma_i\})_i$ of L_\bullet
- uses a variant of Gaussian elimination:
 - ▶ $R = D$ boundary matrix (\mathbb{Z}_2 coefficients) of K_\bullet ,
 - ▶ $V = I$ identity matrix
 - ▶ while $\exists i < j$ with $(R_j)_k \neq 0$, where $k = \text{pivot } R_i$
 - ▶ add R_i to R_j
 - ▶ add V_i to V_j
- yields a reduced matrix $R = D \cdot V$ (columns have distinct pivots) and V is full rank upper triangular


Persistent homology

The persistent homology of a filtration L_\bullet can be computed by the *exhaustive Matrix reduction* algorithm that

- requires a total order on the simplices $\sigma_1 < \dots < \sigma_n$ inducing a simplexwise refinement $K_\bullet = (K_i = \{\sigma_1, \dots, \sigma_i\})_i$ of L_\bullet
- uses a variant of Gaussian elimination:
 - ▶ $R = D$ boundary matrix (\mathbb{Z}_2 coefficients) of K_\bullet ,
 - ▶ $V = I$ identity matrix
 - ▶ while $\exists i < j$ with $(R_j)_k \neq 0$, where $k = \text{pivot } R_i$
 - ▶ add R_i to R_j
 - ▶ add V_i to V_j
- yields a reduced matrix $R = D \cdot V$ (columns have distinct pivots) and V is full rank upper triangular
- determines the barcode through $\{[\text{pivot } R_i, i) \mid R_i \neq 0\}$


Delaunay complexes


Voronoi diagram and Delaunay triangulation




Delaunay complexes

Definition. The *Delaunay complex* $\text{Del}_r(X)$ of $X \subseteq \mathbb{R}^d$ is the nerve of the cover by closed Voronoi balls of radius r centered at points in X .





Delaunay complex




Wrap complex

Discrete Morse theory

A *discrete Morse function* is a


- monotone function $f: K \rightarrow \mathbb{R}$ that
- partitions the complex into pairs and critical simplices, yielding the *discrete gradient* V




Discrete Morse theory

A *discrete Morse function* is a

- monotone function $f: K \rightarrow \mathbb{R}$ that
- partitions the complex into pairs and critical simplices, yielding the *discrete gradient* V




Discrete Morse functions - and their gradients - encode collapses:




Discrete Morse theory

A *discrete Morse function* is a

- monotone function $f: K \rightarrow \mathbb{R}$ that
- partitions the complex into pairs and critical simplices, yielding the *discrete gradient* V




Discrete Morse functions - and their gradients - encode collapses:




Discrete Morse theory

A *discrete Morse function* is a

- monotone function $f: K \rightarrow \mathbb{R}$ that
- partitions the complex into pairs and critical simplices, yielding the *discrete gradient* V




Discrete Morse functions - and their gradients - encode collapses:



Generalized discrete Morse theory

Generalized gradients consist of intervals (in the face poset) instead of just facet pairs:



Morse Theory of Čech and Delaunay complexes

Proposition (Bauer, Edelsbrunner 2014). The Čech and Delaunay complexes are sublevel sets of generalized discrete Morse functions.


Morse Theory of Čech and Delaunay complexes

Proposition (Bauer, Edelsbrunner 2014). The Čech and Delaunay complexes are sublevel sets of generalized discrete Morse functions.

Theorem (Bauer, Edelsbrunner 2017).


Čech, Delaunay, and Wrap complexes (at any scale r) are related by collapses encoded by a single discrete gradient field:

$$\text{Čech}_r(X) \searrow \text{Del}_r(X) \searrow \text{Wrap}_r(X).$$




Wrap complexes

Consider the gradient V of the Delaunay radius function $\text{Del}(X) \rightarrow \mathbb{R}$.



Wrap complexes

Consider the gradient V of the Delaunay radius function $\text{Del}(X) \rightarrow \mathbb{R}$.




Definition (Edelsbrunner 1995; Bauer, Edelsbrunner 2017).

$\text{Wrap}_r(X)$ is the *descending complex* of V on $\text{Del}_r(X)$: the smallest subcomplex of $\text{Del}_r(X)$ that

- contains all critical simplices
- is a union of intervals of V .

Wrap complexes

Consider the gradient V of the Delaunay radius function $\text{Del}(X) \rightarrow \mathbb{R}$.




Definition (Edelsbrunner 1995; Bauer, Edelsbrunner 2017).

$\text{Wrap}_r(X)$ is the *descending complex* of V on $\text{Del}_r(X)$: the smallest subcomplex of $\text{Del}_r(X)$ that

- contains all critical simplices
- is a union of intervals of V .

Wrap complexes

Consider the gradient V of the Delaunay radius function $\text{Del}(X) \rightarrow \mathbb{R}$.




Definition (Edelsbrunner 1995; Bauer, Edelsbrunner 2017).

$\text{Wrap}_r(X)$ is the *descending complex* of V on $\text{Del}_r(X)$: the smallest subcomplex of $\text{Del}_r(X)$ that

- contains all critical simplices
- is a union of intervals of V .

Wrap complexes

Consider the gradient V of the Delaunay radius function $\text{Del}(X) \rightarrow \mathbb{R}$.




Definition (Edelsbrunner 1995; Bauer, Edelsbrunner 2017).

$\text{Wrap}_r(X)$ is the *descending complex* of V on $\text{Del}_r(X)$: the smallest subcomplex of $\text{Del}_r(X)$ that

- contains all critical simplices
- is a union of intervals of V .

Wrap complexes

Consider the gradient V of the Delaunay radius function $\text{Del}(X) \rightarrow \mathbb{R}$.




Definition (Edelsbrunner 1995; Bauer, Edelsbrunner 2017).

$\text{Wrap}_r(X)$ is the *descending complex* of V on $\text{Del}_r(X)$: the smallest subcomplex of $\text{Del}_r(X)$ that

- contains all critical simplices
- is a union of intervals of V .

Wrap complexes

Consider the gradient V of the Delaunay radius function $\text{Del}(X) \rightarrow \mathbb{R}$.




Definition (Edelsbrunner 1995; Bauer, Edelsbrunner 2017).

$\text{Wrap}_r(X)$ is the *descending complex* of V on $\text{Del}_r(X)$: the smallest subcomplex of $\text{Del}_r(X)$ that

- contains all critical simplices
- is a union of intervals of V .

Wrap complexes

Consider the gradient V of the Delaunay radius function $\text{Del}(X) \rightarrow \mathbb{R}$.




Definition (Edelsbrunner 1995; Bauer, Edelsbrunner 2017).

$\text{Wrap}_r(X)$ is the *descending complex* of V on $\text{Del}_r(X)$: the smallest subcomplex of $\text{Del}_r(X)$ that

- contains all critical simplices
- is a union of intervals of V .

Wrap complexes

Consider the gradient V of the Delaunay radius function $\text{Del}(X) \rightarrow \mathbb{R}$.




Definition (Edelsbrunner 1995; Bauer, Edelsbrunner 2017).


$\text{Wrap}_r(X)$ is the *descending complex* of V on $\text{Del}_r(X)$: the smallest subcomplex of $\text{Del}_r(X)$ that

- contains all critical simplices
- is a union of intervals of V .


Wrap complexes and exhaustively reduced cycles




Wrap complexes and exhaustively reduced cycles




Wrap complexes and exhaustively reduced cycles




Wrap complexes and exhaustively reduced cycles



Wrap complexes and exhaustively reduced cycles



Wrap complexes and exhaustively reduced cycles



Algebraic gradient flows and persistent homology

Loose ends in the literature:

- Forman (1998) defines a *flow* $C_*(K) \rightarrow C_*(K)$ associated to a gradient (discrete Morse theory)

Algebraic gradient flows and persistent homology

Loose ends in the literature:

- Forman (1998) defines a *flow* $C_*(K) \rightarrow C_*(K)$ associated to a gradient (discrete Morse theory)
- Kozlov/Sköldberg/Jöllenbeck–Welker (2006/08/09) generalize discrete Morse theory to based chain complexes (*algebraic Morse theory*)

Algebraic gradient flows and persistent homology

Loose ends in the literature:

- Forman (1998) defines a *flow* $C_*(K) \rightarrow C_*(K)$ associated to a gradient (discrete Morse theory)
- Kozlov/Sköldberg/Jöllenbeck–Welker (2006/08/09) generalize discrete Morse theory to based chain complexes (*algebraic Morse theory*)
- Cohen-Steiner–Lieutier–Vuillamy (2022) consider shape reconstruction from point clouds using lexicographically minimal cycles

Algebraic gradient flows and persistent homology

Loose ends in the literature:

- Forman (1998) defines a *flow* $C_*(K) \rightarrow C_*(K)$ associated to a gradient (discrete Morse theory)
- Kozlov/Sköldberg/Jöllenbeck–Welker (2006/08/09) generalize discrete Morse theory to based chain complexes (*algebraic Morse theory*)
- Cohen-Steiner–Lieutier–Vuillamy (2022) consider shape reconstruction from point clouds using lexicographically minimal cycles

We interpret persistent homology in terms of algebraic Morse theory:

- persistence pairs correspond to algebraic gradient pairs

Algebraic gradient flows and persistent homology

Loose ends in the literature:

- Forman (1998) defines a *flow* $C_*(K) \rightarrow C_*(K)$ associated to a gradient (discrete Morse theory)
- Kozlov/Sköldberg/Jöllenbeck–Welker (2006/08/09) generalize discrete Morse theory to based chain complexes (*algebraic Morse theory*)
- Cohen-Steiner–Lieutier–Vuillamy (2022) consider shape reconstruction from point clouds using lexicographically minimal cycles

We interpret persistent homology in terms of algebraic Morse theory:

- persistence pairs correspond to algebraic gradient pairs
- exhaustive Matrix reduction corresponds to gradient flow

Algebraic gradient flows and persistent homology

Loose ends in the literature:

- Forman (1998) defines a *flow* $C_*(K) \rightarrow C_*(K)$ associated to a gradient (discrete Morse theory)
- Kozlov/Sköldberg/Jöllenbeck–Welker (2006/08/09) generalize discrete Morse theory to based chain complexes (*algebraic Morse theory*)
- Cohen-Steiner–Lieutier–Vuillamy (2022) consider shape reconstruction from point clouds using lexicographically minimal cycles

We interpret persistent homology in terms of algebraic Morse theory:

- persistence pairs correspond to algebraic gradient pairs
- exhaustive Matrix reduction corresponds to gradient flow
- the resulting representative cycles are lexicographically minimal

Lexicographic minimality

Given a total order $\sigma_1 < \dots < \sigma_m$ of the simplices of K

Lexicographic minimality

Given a total order $\sigma_1 < \dots < \sigma_m$ of the simplices of K

- the chains $C_n(K)$ can be ordered totally by

$$c_1 < c_2 : \Leftrightarrow \max\{(c_1 \setminus c_2) \cup (c_2 \setminus c_1)\} \in c_2$$

(sometimes also referred to as the *colexicographic order*)

Lexicographic minimality

Given a total order $\sigma_1 < \dots < \sigma_m$ of the simplices of K

- the chains $C_n(K)$ can be ordered totally by

$$c_1 < c_2 :\Leftrightarrow \max\{(c_1 \setminus c_2) \cup (c_2 \setminus c_1)\} \in c_2$$

(sometimes also referred to as the *colexicographic order*)

- a chain $c \in C_n(K)$ is *lexicographically minimal* if there exists no strictly smaller homologous chain $(c + \partial e) < c$

Lexicographic minimality

Given a total order $\sigma_1 < \dots < \sigma_m$ of the simplices of K

- the chains $C_n(K)$ can be ordered totally by

$$c_1 < c_2 : \Leftrightarrow \max\{(c_1 \setminus c_2) \cup (c_2 \setminus c_1)\} \in c_2$$

(sometimes also referred to as the *colexicographic order*)

- a chain $c \in C_n(K)$ is *lexicographically minimal* if there exists no strictly smaller homologous chain $(c + \partial e) < c$

Given a monotonic function $f: K \rightarrow \mathbb{R}$ and a total order on $\text{Vert } K$, the f -*lexicographic total order* on simplices is given by ordering the simplices

Lexicographic minimality

Given a total order $\sigma_1 < \dots < \sigma_m$ of the simplices of K

- the chains $C_n(K)$ can be ordered totally by

$$c_1 < c_2 :\Leftrightarrow \max\{(c_1 \setminus c_2) \cup (c_2 \setminus c_1)\} \in c_2$$

(sometimes also referred to as the *colexicographic order*)

- a chain $c \in C_n(K)$ is *lexicographically minimal* if there exists no strictly smaller homologous chain $(c + \partial e) < c$

Given a monotonic function $f: K \rightarrow \mathbb{R}$ and a total order on $\text{Vert } K$, the f -*lexicographic total order* on simplices is given by ordering the simplices

- by their value under f

Lexicographic minimality

Given a total order $\sigma_1 < \dots < \sigma_m$ of the simplices of K

- the chains $C_n(K)$ can be ordered totally by

$$c_1 < c_2 : \Leftrightarrow \max\{(c_1 \setminus c_2) \cup (c_2 \setminus c_1)\} \in c_2$$

(sometimes also referred to as the *colexicographic order*)

- a chain $c \in C_n(K)$ is *lexicographically minimal* if there exists no strictly smaller homologous chain $(c + \partial e) < c$

Given a monotonic function $f: K \rightarrow \mathbb{R}$ and a total order on $\text{Vert } K$, the f -*lexicographic total order* on simplices is given by ordering the simplices

- by their value under f
- then by dimension

Lexicographic minimality

Given a total order $\sigma_1 < \dots < \sigma_m$ of the simplices of K

- the chains $C_n(K)$ can be ordered totally by

$$c_1 < c_2 : \Leftrightarrow \max\{(c_1 \setminus c_2) \cup (c_2 \setminus c_1)\} \in c_2$$

(sometimes also referred to as the *colexicographic order*)


- a chain $c \in C_n(K)$ is *lexicographically minimal* if there exists no strictly smaller homologous chain $(c + \partial e) < c$

Given a monotonic function $f: K \rightarrow \mathbb{R}$ and a total order on $\text{Vert } K$, the f -*lexicographic total order* on simplices is given by ordering the simplices

- by their value under f
- then by dimension
- then by the lexicographic order induced by the total vertex order.

Minimal cycles and Wrap complexes

Theorem (Bauer, R 2022). Let $X \subset \mathbb{R}^d$ be a finite subset in general position and let $r \in \mathbb{R}$. Then the lexicographically minimal cycles of $\text{Del}_r(X)$ are supported on $\text{Wrap}_r(X)$.



Making discrete Morse theory algebraic

Algebraic Morse theory expresses discrete Morse theory in purely algebraic terms.

Making discrete Morse theory algebraic

Algebraic Morse theory expresses discrete Morse theory in purely algebraic terms.

Definition. Let (C_*, Σ_*) be a based chain complex. A basis element $\sigma \in \Sigma_d$ is a *facet* of $\tau \in \Sigma_{d+1}$ if $\langle \sigma, \partial\tau \rangle \neq 0$. Taking the transitive closure yields the *face poset*.

Making discrete Morse theory algebraic

Algebraic Morse theory expresses discrete Morse theory in purely algebraic terms.

Definition. Let (C_*, Σ_*) be a based chain complex. A basis element $\sigma \in \Sigma_d$ is a *facet* of $\tau \in \Sigma_{d+1}$ if $\langle \sigma, \partial\tau \rangle \neq 0$. Taking the transitive closure yields the *face poset*.

- The basis elements Σ_* take the role of the simplices in discrete Morse theory.

Making discrete Morse theory algebraic

Algebraic Morse theory expresses discrete Morse theory in purely algebraic terms.

Definition. Let (C_*, Σ_*) be a based chain complex. A basis element $\sigma \in \Sigma_d$ is a *facet* of $\tau \in \Sigma_{d+1}$ if $\langle \sigma, \partial\tau \rangle \neq 0$. Taking the transitive closure yields the *face poset*.

- The basis elements Σ_* take the role of the simplices in discrete Morse theory.
- All further definitions of discrete Morse theory apply verbatim.

Algebraic gradients from persistent homology

Let $R = D \cdot V$ be the reduced boundary matrix
(for the Delaunay filtration, breaking ties lexicographically).

- The pairs of columns (R_i, V_i) (with $R_i \neq 0$) form an algebraic gradient.

Algebraic gradients from persistent homology

Let $R = D \cdot V$ be the reduced boundary matrix
(for the Delaunay filtration, breaking ties lexicographically).

- The pairs of columns (R_i, V_i) (with $R_i \neq 0$) form an algebraic gradient.
- We consider another one, the *persistence gradient*, given by the pairs $(\text{pivot } R_i, V_i)$ (with $R_i \neq 0$).

Algebraic gradients from persistent homology

Let $R = D \cdot V$ be the reduced boundary matrix
(for the Delaunay filtration, breaking ties lexicographically).

- The pairs of columns (R_i, V_i) (with $R_i \neq 0$) form an algebraic gradient.
- We consider another one, the *persistence gradient*, given by the pairs (pivot R_i, V_i) (with $R_i \neq 0$).
- If (σ, τ) is an *apparent pair*, then $(\text{pivot } R_\tau, V_\tau) = (\sigma, \tau)$.

Algebraic gradients from persistent homology

Let $R = D \cdot V$ be the reduced boundary matrix
(for the Delaunay filtration, breaking ties lexicographically).

- The pairs of columns (R_i, V_i) (with $R_i \neq 0$) form an algebraic gradient.
- We consider another one, the *persistence gradient*, given by the pairs (pivot R_i, V_i) (with $R_i \neq 0$).
- If (σ, τ) is an *apparent pair*, then (pivot R_τ, V_τ) = (σ, τ) .
- Thus the apparent pairs gradient is a subset of the persistence gradient.

Algebraic gradients from persistent homology

Let $R = D \cdot V$ be the reduced boundary matrix
(for the Delaunay filtration, breaking ties lexicographically).

- The pairs of columns (R_i, V_i) (with $R_i \neq 0$) form an algebraic gradient.
- We consider another one, the *persistence gradient*, given by the pairs (pivot R_i, V_i) (with $R_i \neq 0$).
- If (σ, τ) is an *apparent pair*, then (pivot R_τ, V_τ) = (σ, τ) .
- Thus the apparent pairs gradient is a subset of the persistence gradient.
- The apparent pairs of persistence zero refine the gradient of the Delaunay radius function.

Key insights leading to the theorem

- The following are equivalent for any cycle $z \in Z_*(K_i)$:

Key insights leading to the theorem

- The following are equivalent for any cycle $z \in Z_*(K_i)$:
 - ▶ z is lexicographically minimal in its homology class $[z] \in H_*(K_i)$

Key insights leading to the theorem

- The following are equivalent for any cycle $z \in Z_*(K_i)$:
 - ▶ z is lexicographically minimal in its homology class $[z] \in H_*(K_i)$
 - ▶ z is invariant under the flow induced by the persistence gradient

Key insights leading to the theorem

- The following are equivalent for any cycle $z \in Z_*(K_i)$:
 - ▶ z is lexicographically minimal in its homology class $[z] \in H_*(K_i)$
 - ▶ z is invariant under the flow induced by the persistence gradient
 - ▶ z is exhaustively reduced

Key insights leading to the theorem

- The following are equivalent for any cycle $z \in Z_*(K_i)$:
 - ▶ z is lexicographically minimal in its homology class $[z] \in H_*(K_i)$
 - ▶ z is invariant under the flow induced by the persistence gradient
 - ▶ z is exhaustively reduced
 - ▶ z is supported only on death and essential simplices

Key insights leading to the theorem

- The following are equivalent for any cycle $z \in Z_*(K_i)$:
 - ▶ z is lexicographically minimal in its homology class $[z] \in H_*(K_i)$
 - ▶ z is invariant under the flow induced by the persistence gradient
 - ▶ z is exhaustively reduced
 - ▶ z is supported only on death and essential simplices
- Flow-invariant cycles are supported on the descending complex.

Key insights leading to the theorem

- The following are equivalent for any cycle $z \in Z_*(K_i)$:
 - ▶ z is lexicographically minimal in its homology class $[z] \in H_*(K_i)$
 - ▶ z is invariant under the flow induced by the persistence gradient
 - ▶ z is exhaustively reduced
 - ▶ z is supported only on death and essential simplices
- Flow-invariant cycles are supported on the descending complex.
- Smaller gradients have more flow-invariant cycles.

Key insights leading to the theorem

- The following are equivalent for any cycle $z \in Z_*(K_i)$:
 - ▶ z is lexicographically minimal in its homology class $[z] \in H_*(K_i)$
 - ▶ z is invariant under the flow induced by the persistence gradient
 - ▶ z is exhaustively reduced
 - ▶ z is supported only on death and essential simplices
- Flow-invariant cycles are supported on the descending complex.
- Smaller gradients have more flow-invariant cycles.
 - ▶ Thus, an invariant cycle for the algebraic flow (by the persistence gradient) is also invariant for the geometric flow (by apparent pairs of persistence zero).

Key insights leading to the theorem


- The following are equivalent for any cycle $z \in Z_*(K_i)$:
 - ▶ z is lexicographically minimal in its homology class $[z] \in H_*(K_i)$
 - ▶ z is invariant under the flow induced by the persistence gradient
 - ▶ z is exhaustively reduced
 - ▶ z is supported only on death and essential simplices
- Flow-invariant cycles are supported on the descending complex.
- Smaller gradients have more flow-invariant cycles.
 - ▶ Thus, an invariant cycle for the algebraic flow (by the persistence gradient) is also invariant for the geometric flow (by apparent pairs of persistence zero).
- Coarser gradients have larger descending complexes.

Key insights leading to the theorem

- The following are equivalent for any cycle $z \in Z_*(K_i)$:
 - ▶ z is lexicographically minimal in its homology class $[z] \in H_*(K_i)$
 - ▶ z is invariant under the flow induced by the persistence gradient
 - ▶ z is exhaustively reduced
 - ▶ z is supported only on death and essential simplices
- Flow-invariant cycles are supported on the descending complex.
- Smaller gradients have more flow-invariant cycles.
 - ▶ Thus, an invariant cycle for the algebraic flow (by the persistence gradient) is also invariant for the geometric flow (by apparent pairs of persistence zero).
- Coarser gradients have larger descending complexes.
 - ▶ Thus, the minimal cycles are also supported on the Wrap complex.

Point cloud reconstruction with most persistent features

The lexicographically minimal cycle corresponding to the interval in the persistence barcode of the Delaunay function with the largest death/birth ratio:



Summary

- Got reminded about discrete Morse theory and the Wrap complex
- Saw how persistent homology relates to algebraic gradient flows
- Learned that lexicographically minimal cycles of $\text{Del}_r(X)$ are supported on $\text{Wrap}_r(X)$

Funding: DFG – TRR109 *Discretization in Geometry and Dynamics*