

UNIVERSITÀ
DEGLI STUDI
DELL'AQUILA

Lesson 3: Regression

Technologies for Artificial Intelligence

Fabio Antonini, Università degli Studi dell'Aquila

16 October 2026

Before we start

- Exercise 2, which we discuss now
- The pipeline you built is the one we fit models into from now on

UNIVERSITÀ
DEGLI STUDI
DELL'AQUILA

Today: the first real model

- The model and what it claims
- The exact solution, and why it sometimes fails
- Gradient descent
- Curves, and the price of flexibility
- Ridge and Lasso

UNIVERSITÀ
DEGLI STUDI
DELL'AQUILA

One dataset, all lesson


600 houses. Six measurements. A price.

And we know the coefficients that generated it.

In code: `train_test_split(X, y, test_size=0.25)` keeps 150 houses for testing.

UNIVERSITÀ DEGLI STUDI DELL'AQUILA

What we are trying to predict



UNIVERSITÀ DEGLI STUDI DELL'AQUILA

What a linear model claims

Each feature contributes a fixed amount per unit, and the contributions add up: $\hat{y} = w_1x_1 + w_2x_2 + \ldots + w_nx_n + b$, one coefficient w per feature plus an intercept b.

- Every square metre: the same 2,400 €
- Every kilometre out: the same −6,500 €

UNIVERSITÀ DEGLI STUDI DELL'AQUILA

Fitting means minimising something

We need one number saying how badly a candidate model is doing.

Adding up the errors, $\Sigma_i (\hat{y}^{(i)} - y^{(i)})$, fails: +50,000 and −50,000 cancel to zero.

Every error must count as a positive amount:

- absolute error $\Sigma_i |\hat{y}^{(i)} - y^{(i)}|$
- squared error $\Sigma_i (\hat{y}^{(i)} - y^{(i)})^2$

UNIVERSITÀ DEGLI STUDI DELL'AQUILA

Why squared, not absolute

- Squared error $\Sigma_i (\hat{y}^{(i)} - y^{(i)})^2$: each error $e = \hat{y}^{(i)} - y^{(i)}$ counts as e^2, not as $|e|$
- **Differentiable everywhere**: e^2 is smooth, $|e|$ has a corner at $e = 0$
- **Punishes one large error more**: double an error and $|e|$ doubles, e^2 quadruples
- Under Gaussian noise it is **maximum likelihood**: it picks the coefficients that make the observed prices most probable

UNIVERSITÀ DEGLI STUDI DELL'AQUILA

The cost function: mean squared error

$$J(w,b) = \frac{1}{2m} \sum_{i=1}^{m} \left(\hat{y}^{(i)} - y^{(i)}\right)^2$$

UNIVERSITÀ DEGLI STUDI DELL'AQUILA

Trying $\hat{y} = 2{,}400 \cdot \text{area} + 45{,}000$

Area (m^2)	True (€)	Predicted (€)	Error $\hat{y} - y$	Error^2
80	240,000	237,000	−3,000	9.0×10^6
120	320,000	333,000	+13,000	1.69×10^8
200	540,000	525,000	−15,000	2.25×10^8

UNIVERSITÀ DEGLI STUDI DELL'AQUILA

Squared error spends its attention on the worst house

- The 3,000 € error contributes about **2%** of the total
- The two large errors contribute the other **98%**
- Halving the 15,000 € error cuts the cost by **42%**
- In code: `mean_squared_error(y, y_pred)`, the mean of the squared errors without the ½

UNIVERSITÀ DEGLI STUDI DELL'AQUILA

A cost needs something to be compared against

- Trivial answer: **always predict the mean**, `DummyRegressor()`
- Root mean squared error, RMSE = √MSE, on the 150 test houses: **96,440 €** for the mean, **20,341 €** for the model
- $R^2 = 1 - \Sigma(y - \hat{y})^2 \div \Sigma(y - \bar{y})^2 =$ **0.956** (`r2_score`): 0 is the mean, 1 is perfect
- Quote the euros too: R^2 depends on *this* test set's variance


UNIVERSITÀ DEGLI STUDI DELL'AQUILA

Fitting is finding where J is smallest

$$(w^*, b^*) = \arg\min_{w,\ b} J(w, b)$$


UNIVERSITÀ DEGLI STUDI DELL'AQUILA

The cost is a bowl



UNIVERSITÀ DEGLI STUDI DELL'AQUILA

The picture behind the exact solution



the closest point on the surface is the shadow, and the leftover error is perpendicular, or you could have done better

UNIVERSITÀ
DEGLI STUDI
DELL'AQUILA

The normal equation

$$\theta = (X^\top X)^{-1} X^\top y$$

UNIVERSITÀ
DEGLI STUDI
DELL'AQUILA

The same equation, by hand

Quantity	Value
the three houses	80, 120, 200 m² → 240k, 320k, 540k
X, and θ	a column of 1s (for b) beside the areas; θ = (b, w)
X^TX	rows (3, 400) and (400, 60,800)
X^Ty	(1,100, 165,600)
determinant	22,400
slope w	**2,536 €/m²** (truth 2,400)
intercept b	28,600 €

UNIVERSITÀ DEGLI STUDI DELL'AQUILA

And it really is the minimum

- The second derivative is X^TX, never negative in any direction
- So the cost is **convex**: one bottom, no local traps
- Any stationary point (zero gradient) is *the* answer, not *an* answer
- In code: `LinearRegression().fit(X, y)` solves the same problem

UNIVERSITÀ DEGLI STUDI DELL'AQUILA

So why learn anything else?

- $(X^TX)^{-1}$ costs about n^3 operations
- Two columns carrying one fact → **no inverse exists**
- Nearly-redundant columns → an inverse that is useless

UNIVERSITÀ DEGLI STUDI DELL'AQUILA

How stretched is the valley?

The **condition number** κ = steepest curvature ÷ shallowest curvature: how many times steeper the steepest direction of the bowl is than the shallowest.

- Housing features as recorded: **285**
- The same features, standardised: **3.4**
- Add `area_sqft` beside `area_sqm`: **2,286**

UNIVERSITÀ DEGLI STUDI DELL'AQUILA

The picture behind gradient descent

You are on a hillside in fog.

You cannot see the valley, but you can feel which way the ground slopes.

So: step against the slope, by an amount set by the **learning rate** α, and repeat until the cost stops falling.

UNIVERSITÀ DEGLI STUDI DELL'AQUILA

The update rule

$$w \leftarrow w - \alpha \frac{\partial J}{\partial w}, \qquad \frac{\partial J}{\partial w_j} = \frac{1}{m} \sum_i \left(\hat{y}^{(i)} - y^{(i)} \right) x_j^{(i)}$$


UNIVERSITÀ DEGLI STUDI DELL'AQUILA


One step on two houses, from w = 0 and b = 0

80 m^2 at 240k, 200 m^2 at 540k; $\alpha = 10^{-5}$	Gradient	After one step
slope w	−63,600	0.636
intercept b	−390	0.0039

UNIVERSITÀ DEGLI STUDI DELL'AQUILA


The path it takes





UNIVERSITÀ DEGLI STUDI DELL'AQUILA

Three learning rates



UNIVERSITÀ DEGLI STUDI DELL'AQUILA

Notebook 1, live

section, and its slides	you call (→ the number to come back with)
one feature, 4-5	`train_test_split` → 450 houses to fit, 150 to test
the exact answer, 15-18	the normal equation in NumPy, then `LinearRegression` → 2,785 €/m², agreeing to 1.82e-12
the iterative answer, 21-25	NumPy only: gradient descent → the same answer, slowly
all six features, 50-51	`LinearRegression` → area 2,421, bedrooms −16.6%
the baseline, 12	`DummyRegressor(strategy="mean")`, `r2_score` → 96,440 € against 20,341 €, R^2 0.956

Break

What if the relationship bends?


Energy consumption against temperature: heating in the cold, cooling in the heat.

No straight line follows that.

In code: `PolynomialFeatures(degree=2)` adds the squared temperature, t^2, as a second column.

UNIVERSITÀ DEGLI STUDI DELL'AQUILA

A relationship a line cannot follow




UNIVERSITÀ DEGLI STUDI DELL'AQUILA

“Linear” means linear in the coefficients

$$\hat{y} = w_1 t + w_2 t^2 + b$$


UNIVERSITÀ DEGLI STUDI DELL'AQUILA

The straight line’s residuals keep the shape




UNIVERSITÀ DEGLI STUDI DELL'AQUILA

What “nothing left” looks like



UNIVERSITÀ DEGLI STUDI DELL'AQUILA

So use more flexibility?




UNIVERSITÀ DEGLI STUDI DELL'AQUILA

Root mean squared error (RMSE), by degree

Degree	Train RMSE (kWh)	Test RMSE (kWh)
1	113.3	212.1
3	17.5	**16.9**
9	16.2	590.1
12	13.6	24,655.7

The gap is the overfitting



UNIVERSITÀ DEGLI STUDI DELL'AQUILA

Notebook 2, live

section, and its slides	you call (→ the number to come back with)
a relationship that bends, 28-29	`train_test_split` → 21 days to fit, 9 to test
linear in the coefficients, 30-32	`PolynomialFeatures`, `StandardScaler` and `LinearRegression`, chained by `make_pipeline`
too much flexibility, 33-35	`mean_squared_error`, square-rooted → test RMSE 16.9 at degree 3, 24,655.7 at 12
why the wiggles appear, 37	the fitted coefficients → largest 3,097,038,010 at degree 12

Why flexibility goes wrong

Largest coefficient, degree 2: **411**

Largest coefficient, degree 12: **3,097,038,010**

UNIVERSITÀ
DEGLI STUDI
DELL'AQUILA

Charge for size

$$J_{\text{ridge}} = \text{MSE} + \lambda \sum_j w_j^2 \qquad J_{\text{lasso}} = \text{MSE} + \lambda \sum_j |w_j|$$

UNIVERSITÀ
DEGLI STUDI
DELL'AQUILA

Two rules that are not optional

- The **intercept is never penalised**: it is not a claim about any feature
- Features **must be scaled first**, or the penalty is arbitrary
- In code: `make_pipeline(StandardScaler(), Ridge(alpha=0.01))`; scikit-learn calls λ `alpha`


UNIVERSITÀ DEGLI STUDI DELL'AQUILA

What a penalty buys

Model	Train RMSE	Test RMSE	max |w|
none	13.6	24,655.7	3,097,038,010
$\lambda = 0.01$	18.0	**22.8**	365
$\lambda = 1$	44.5	105.2	156
$\lambda = 100$	96.0	227.6	6


UNIVERSITÀ DEGLI STUDI DELL'AQUILA

The unpenalised fit leaves the picture



UNIVERSITÀ DEGLI STUDI DELL'AQUILA

Too flexible, too rigid



UNIVERSITÀ DEGLI STUDI DELL'AQUILA

Ridge has an exact solution too

$$\theta = (X^\top X + \lambda I)^{-1} X^\top y$$


UNIVERSITÀ
DEGLI STUDI
DELL'AQUILA

Why that one change fixes it

- Least squares fails when a direction costs nothing: the valley is flat
- The penalty makes **every** direction cost something
- Every eigenvalue of X^TX shifts up by λ, so none of them can be zero


UNIVERSITÀ DEGLI STUDI DELL'AQUILA

The picture behind Ridge vs Lasso



UNIVERSITÀ
DEGLI STUDI
DELL'AQUILA

Ridge shrinks, Lasso selects



UNIVERSITÀ DEGLI STUDI DELL'AQUILA

The order Lasso drops features

λ	Features kept
1,000	all six
10,000	five: `garage` dropped
20,000	three: `area_sqm`, `bedrooms`, `bathrooms`
40,000	one: `area_sqm`

The case Ridge was invented for

`area_sqm` and `area_sqft`: the same measurement, two units.

Correlation: **0.999999**

UNIVERSITÀ DEGLI STUDI DELL'AQUILA


The coefficients are noise

Split	area_sqm	area_sqft
0	9,261	−640
1	14,535	−1,127
2	45,307	−3,989
3	39,061	−3,402

What a coefficient actually says

The change in the prediction for a one-unit change in that feature, **with every other feature held fixed**.

When can you trust a coefficient?



UNIVERSITÀ DEGLI STUDI DELL'AQUILA

Notebook 3, live

section, and its slides	you call (→ the number to come back with)
Ridge on the degree-12 fit, 38-42	`Ridge(alpha=0.01)` after `PolynomialFeatures` and `StandardScaler` → largest coefficient 365
regularisation paths, 45-47	`make_pipeline(StandardScaler(), Ridge(alpha=...))`, the same with `Lasso` → only area left at 40,000
the collinear case, 48-49	`LinearRegression` against `Ridge(alpha=10)` on four splits → 9,261 to 45,307, then about 40,000

What we did today

- A model that claims a fixed amount per unit
- An exact solution (a projection), and when it fails
- An iterative one: walk downhill, mind your stride
- Flexibility helps until it does not
- Charging for size buys generalisation, and readable coefficients
- **365** against billions: what a penalty of 0.01 costs, and buys

UNIVERSITÀ DEGLI STUDI DELL'AQUILA

Homework: we discuss it on Friday 23 October

`Exercises/03_regression.md`

Fit, regularise, and **explain which coefficients you believe**.

UNIVERSITÀ DEGLI STUDI DELL'AQUILA