

SAPIENZA
UNIVERSITÀ DI ROMA

Combinatorial Optimization

Exercises 5 — Approximation algorithms

Worked exercises on the chapters of Volume 5 (approximation algorithms), each followed by its complete solution. The instances are those of the selected problems, cited by their code.

Fabio Furini

fabiofurini.github.io

Department of Computer, Control and Management Engineering — Sapienza University of Rome

Online version and downloadable material:
fabiofurini.github.io/combinatorial-optimization/
github.com/fabiofurini/combinatorial-optimization

Academic year 2026–2027

Contents

25 Approximation algorithms: definitions **2**

26 Approximation algorithms for bin packing **8**

26.1 Next Fit 8

26.2 First Fit 13

26.3 Best Fit 18

26.4 Worst Fit 23

27 Approximation algorithms for the knapsack problem **28**

27.1 Greedy algorithm 28

27.2 Extended greedy algorithm 33

27.3 The H^2 heuristic 38

27.4 The H^ℓ heuristic 42

27.5 The FPTAS 47

28 Approximation for $P\|C_{\max}$ **52**

29 Approximation for the metric TSP **57**

30 Limits of approximation: vertex coloring **62**

Approximation algorithms: definitions

Exercise 25.1 — Performance ratios and errors (1)

A heuristic algorithm $\mathscr{A}$ for a **minimization** problem and a heuristic algorithm $\mathscr{B}$ for a **maximization** problem give the following values on some instances:

$\mathscr{A}$ (min)	I_1	I_2	I_3	I_4	I_5
$\mathscr{A}(I)$	12	9	15	20	7
$z(I)$	10	9	10	16	6

$\mathscr{B}$ (max)	I_1	I_2	I_3
$\mathscr{B}(I)$	30	25	16
$z(I)$	36	25	24

Questions

1. Compute the performance ratio, the additive error and the relative error of $\mathscr{A}$ on its five instances.
2. Do the same for $\mathscr{B}$ on its three instances.
3. Using only these data, which of the constants $\alpha = \frac{7}{5}$ and $\alpha = \frac{3}{2}$ can be a relative performance guarantee of $\mathscr{A}$? What would a proof that $\mathscr{A}$ is a $\frac{3}{2}$-approximation algorithm need?
4. Suppose that $\mathscr{A}$ is proved to be a $\frac{3}{2}$-approximation algorithm. Is the guarantee tight? Give a lower bound on $r(\mathscr{A})$ and on $r(\mathscr{B})$.

Solution

Performance ratio: $\frac{\mathscr{A}(I)}{z(I)}$ (minimization), $\frac{z(I)}{\mathscr{A}(I)}$ (maximization). Additive error: $\mathscr{A}(I) - z(I)$ (minimization), $z(I) - \mathscr{A}(I)$ (maximization). Relative error: $\frac{\mathscr{A}(I)-z(I)}{z(I)}$ (minimization), $\frac{z(I)-\mathscr{A}(I)}{z(I)}$ (maximization). The worst-case performance ratio $r(\mathscr{A})$ is the supremum of the performance ratio over all the instances.

1. For a minimization problem the ratio is $\frac{\mathscr{A}(I)}{z(I)}$:

instance	$\mathscr{A}(I)$	$z(I)$	ratio	additive error	relative error
I_1	12	10	$\frac{6}{5}$	2	$\frac{1}{5}$
I_2	9	9	1	0	0
I_3	15	10	$\frac{3}{2}$	5	$\frac{1}{2}$
I_4	20	16	$\frac{5}{4}$	4	$\frac{1}{4}$
I_5	7	6	$\frac{7}{6}$	1	$\frac{1}{6}$

2. For a maximization problem the ratio is $\frac{z(I)}{\mathscr{B}(I)}$:

instance	$\mathscr{A}(I)$	$z(I)$	ratio	additive error	relative error
I_1	30	36	$\frac{6}{5}$	6	$\frac{1}{6}$
I_2	25	25	1	0	0
I_3	16	24	$\frac{3}{2}$	8	$\frac{1}{3}$

   All the ratios are at least 1, as they must be for feasible solutions.

3. On I_3 the ratio is $\frac{3}{2} > \frac{7}{5}$: $\alpha = \frac{7}{5}$ is not a relative performance guarantee. The data do not exclude $\alpha = \frac{3}{2}$, since every ratio is at most $\frac{3}{2}$. A proof must show $\mathscr{A}(I) \leq \frac{3}{2}\, z(I)$ for *every* instance I: finitely many examples can only disprove a guarantee.

4. Yes: the instance I_3 attains the ratio $\frac{3}{2}$, so no smaller constant is a guarantee, and $r(\mathscr{A}) = \frac{3}{2}$. Without a proof, the data give only lower bounds: $r(\mathscr{A}) \geq \frac{3}{2}$ and $r(\mathscr{B}) \geq \frac{3}{2}$ (instance I_3 of $\mathscr{B}$).

Exercise 25.2 — A tight guarantee for a simple algorithm (1)

For $P\|C_{\max}$ consider the algorithm ONE, which assigns all the jobs to machine 1, so that $\mathrm{ONE}(I) = \sum_{j \in \mathscr{J}} t_j$. Every instance satisfies $z(I) \geq \frac{\sum_j t_j}{m}$, since the m loads sum to $\sum_j t_j$. Consider the instances

I_1: $m = 2$ and $t = (7, 5, 4, 3, 3)$;

I_2: $m = 3$ and $t = (9, 2, 2, 2, 1)$;

I_3: $m = 3$ and $t = (5, 5, 4, 4, 3, 3, 3)$;

Questions

1. Compute $\mathrm{ONE}(I)$, $z(I)$ (by enumeration) and the performance ratio on the three instances.

2. Prove that ONE is an m-approximation algorithm for every fixed number m of machines.

3. Build, for every $m \geq 2$, an instance on which the ratio is exactly m, and check it with $m = 4$. What can you conclude?

4. Does ONE have a finite additive performance guarantee? Use the instance I_1 with all the processing times multiplied by 10.

Solution

Performance ratio: $\frac{\mathscr{A}(I)}{z(I)}$ (minimization), $\frac{z(I)}{\mathscr{A}(I)}$ (maximization). Additive error: $\mathscr{A}(I) - z(I)$ (minimization), $z(I) - \mathscr{A}(I)$ (maximization). Relative error: $\frac{\mathscr{A}(I)-z(I)}{z(I)}$ (minimization), $\frac{z(I)-\mathscr{A}(I)}{z(I)}$ (maximization). The worst-case performance ratio $r(\mathscr{A})$ is the supremum of the performance ratio over all the instances.

1.

instance	m	ONE(I)	$z(I)$	ratio	m
I_1	2	22	11	2	2
I_2	3	16	9	$\frac{16}{9}$	3
I_3	3	27	9	3	3

The ratio never exceeds m, and it is equal to m on I_1 and I_3.

2. For every instance, $z(I) \geq \frac{\sum_j t_j}{m}$ gives

$$\frac{\text{ONE}(I)}{z(I)} = \frac{\sum_j t_j}{z(I)} \leq \frac{\sum_j t_j}{\frac{\sum_j t_j}{m}} = m.$$

3. Take $n = 2m$ jobs of time 1. Two jobs per machine give $z(I) = 2$, while ONE$(I) = 2m$: the ratio is m. With $m = 4$: ONE$(I) = 8$, $z(I) = 2$ (enumeration) and the ratio is 4. The guarantee m is tight and $r(\text{ONE}) = m$.

4. Multiplying the times of I_1 by 10 gives $t = (70, 50, 40, 30, 30)$, with ONE$(I) = 220$ and $z(I) = 110$: the additive error goes from 11 to 110. Multiplying by any K multiplies the error by K, so no constant β bounds it: $e(\text{ONE}) = \infty$, while the ratio stays 2.

Exercise 25.3 — An algorithm without a guarantee (1)

For the BPP consider the algorithm SEP, which packs every item in a separate bin, so that SEP$(I) = n$. Consider the instance I_0 with $c = 10$ and weights $(6, 5, 4, 3, 2)$, and, for $k \in \mathbb{Z}_{\geq 2}$, the instances I_k with $c = k$ and $2k$ items of weight 1.

Questions

1. Compute SEP(I_0), $z(I_0)$, the performance ratio and the additive error on I_0.
2. Compute the performance ratio on I_k for $k = 3, 5, 10$ (use enumeration for $k = 3$), and its expression for every k.
3. Is SEP an α-approximation algorithm for some constant α? Give $r(\text{SEP})$ and $e(\text{SEP})$.
4. Using $z(I) \geq 2$, show that $\frac{\text{SEP}(I)}{z(I)} \leq \frac{n}{2}$. Why is $\frac{n}{2}$ not a relative performance guarantee?

Solution

Performance ratio: $\frac{\mathscr{A}(I)}{z(I)}$ (minimization), $\frac{z(I)}{\mathscr{A}(I)}$ (maximization). Additive error: $\mathscr{A}(I) - z(I)$ (minimization), $z(I) - \mathscr{A}(I)$ (maximization). Relative error: $\frac{\mathscr{A}(I)-z(I)}{z(I)}$ (minimization), $\frac{z(I)-\mathscr{A}(I)}{z(I)}$ (maximization). The worst-case performance ratio $r(\mathscr{A})$ is the supremum of the performance ratio over all the instances.

1. $\text{SEP}(I_0) = 5$ and $z(I_0) = 2$ (patterns $\{1,3\}$ and $\{2,4,5\}$). The ratio is $\frac{5}{2}$ and the additive error is $5 - 2 = 3$.

2. Two bins of capacity k hold the $2k$ unit items, and one bin is not enough, so $z(I_k) = 2$:

k	c	$\text{SEP}(I_k) = 2k$	$z(I_k)$	ratio
3	3	6	2	3
5	5	10	2	5
10	10	20	2	10

   In general the ratio is $\frac{2k}{2} = k$.

3. No: for every constant α, the instance I_k with $k > \alpha$ has ratio $k > \alpha$. Hence $r(\text{SEP}) = \infty$. The additive error on I_k is $2k - 2$, also unbounded: $e(\text{SEP}) = \infty$.

4. The assumption $\sum_j w_j > c$ gives $z(I) \geq 2$, so $\frac{\text{SEP}(I)}{z(I)} = \frac{n}{z(I)} \leq \frac{n}{2}$. The bound depends on the instance through n: a relative performance guarantee must be one constant α valid for all the instances.

Exercise 25.4 — Unbounded ratio of a greedy algorithm (1)

For the KP consider the algorithm GR: sort the items by non-increasing efficiency $\frac{p_j}{w_j}$ (ties: smallest index) and insert each item, in this order, if it fits in the residual capacity. Consider the instance I_0 with $c = 9$, $p = (10, 7, 6, 4)$, $w = (5, 4, 3, 3)$, and, for $M \in \mathbb{Z}_{\geq 3}$, the instances I_M with $c = M$ and two items: $p_1 = 2$, $w_1 = 1$ and $p_2 = M$, $w_2 = M$.

Questions

1. Apply GR to I_0, compute $z(I_0)$ by enumeration, the performance ratio and the relative error.

2. Apply GR to I_M and compute $z(I_M)$, the performance ratio and the additive error for $M = 3, 10, 100$.

3. Is GR an α-approximation algorithm for some constant α? Is it a β-additive approximation algorithm?

4. Which property of the instances I_M makes GR fail?

Solution

Performance ratio: $\frac{\mathscr{A}(I)}{z(I)}$ (minimization), $\frac{z(I)}{\mathscr{A}(I)}$ (maximization). Additive error: $\mathscr{A}(I) - z(I)$ (minimization), $z(I) - \mathscr{A}(I)$ (maximization). Relative error: $\frac{\mathscr{A}(I)-z(I)}{z(I)}$ (minimization), $\frac{z(I)-\mathscr{A}(I)}{z(I)}$ (maximization). The worst-case performance ratio $r(\mathscr{A})$ is the supremum of the performance ratio over all the instances.

1. The efficiencies are 2, $\frac{7}{4}$, 2, $\frac{4}{3}$: the order is $(1, 3, 2, 4)$. GR inserts items 1, 3 and gets $\mathrm{GR}(I_0) = 16$. Enumeration gives $z(I_0) = 17$ (items $\{1, 2\}$). The ratio is $\frac{17}{16} = \frac{17}{16}$ and the relative error is $\frac{17-16}{17} = \frac{1}{17}$.

2. Item 1 has efficiency 2 and item 2 has efficiency 1: GR inserts item 1, and then item 2 does not fit in the residual capacity $M - 1$. Hence $\mathrm{GR}(I_M) = 2$, while item 2 alone gives $z(I_M) = M$:

M	$\mathrm{GR}(I_M)$	$z(I_M)$	ratio	additive error
3	2	3	$\frac{3}{2}$	1
10	2	10	5	8
100	2	100	50	98

3. The ratio $\frac{M}{2}$ grows without limit, so no constant α works and $r(\mathrm{GR}) = \infty$. The additive error $M - 2$ is also unbounded: $e(\mathrm{GR}) = \infty$.

4. The efficient item is small and blocks the large item, whose profit is almost all the optimal value: the order by efficiency ignores how much profit an item brings in absolute terms.

Exercise 25.5 — Approximation schemes (1)

An approximation scheme $\{\mathscr{A}_\varepsilon\}_{\varepsilon\in(0,1)}$ is run on some instances. The table gives the type of problem, the accuracy parameter ε, the optimal value and the value returned:

case	problem	ε	$z(I)$	$\mathscr{A}_\varepsilon(I)$
(1)	minimization	$\frac{1}{10}$	100	108
(2)	minimization	$\frac{1}{10}$	60	67
(3)	maximization	$\frac{1}{5}$	50	41
(4)	maximization	$\frac{1}{5}$	50	38

Questions

1. For each case, compute the relative error and decide whether the value is compatible with the definition of an approximation scheme.

2. Translate $\varepsilon = \frac{1}{4}$ into a relative performance guarantee, for a minimization and for a maximization problem. What is the accuracy?

3. A scheme for a maximization problem must be a $\frac{5}{4}$-approximation algorithm. Which value of ε is needed?

4. Classify the following running times of an approximation scheme on instances of size n: $O(n^{1/\varepsilon})$, $O(n^3/\varepsilon)$, $O(n\,2^{1/\varepsilon})$, $O(n^2/\varepsilon^2)$, $O(2^n/\varepsilon)$.

Solution

1. (1) $\frac{108-100}{100} = \frac{2}{25} \leq \frac{1}{10}$: compatible.
   (2) $\frac{67-60}{60} = \frac{7}{60} > \frac{1}{10}$: not compatible, the scheme is wrong on this instance.
   (3) $\frac{50-41}{50} = \frac{9}{50} \leq \frac{1}{5}$: compatible.
   (4) $\frac{50-38}{50} = \frac{6}{25} > \frac{1}{5}$: not compatible, the scheme is wrong on this instance.

2. For a minimization problem $\mathscr{A}_\varepsilon(I) \leq (1+\varepsilon)\,z(I)$, so the guarantee is $1 + \frac{1}{4} = \frac{5}{4}$. For a maximization problem $\mathscr{A}_\varepsilon(I) \geq (1-\varepsilon)\,z(I)$, so the guarantee is $\frac{1}{1-\varepsilon} = \frac{4}{3}$. The accuracy is $\frac{1}{\varepsilon} = 4$.

3. We need $\frac{1}{1-\varepsilon} \leq \frac{5}{4}$, that is $1 - \varepsilon \geq \frac{4}{5}$ and $\varepsilon \leq \frac{1}{5}$: any $\varepsilon \leq \frac{1}{5}$ works, and $\varepsilon = \frac{1}{5}$ is the largest one.

4. - $O(n^{1/\varepsilon})$: PTAS, not FPTAS.
   - $O(n^3/\varepsilon)$: FPTAS.
   - $O(n\,2^{1/\varepsilon})$: PTAS, not FPTAS.
   - $O(n^2/\varepsilon^2)$: FPTAS.
   - $O(2^n/\varepsilon)$: not a PTAS.

   A PTAS needs a polynomial in n for every fixed ε; an FPTAS needs a polynomial in both n and $\frac{1}{\varepsilon}$. The terms $n^{1/\varepsilon}$ and $2^{1/\varepsilon}$ are not polynomial in $\frac{1}{\varepsilon}$, and 2^n is not polynomial in n.

Chapter 26

Approximation algorithms for bin packing

26.1 Next Fit

Exercise 26.1 — Next Fit 1

We consider a bin of capacity $c = 10$ and a set $\mathscr{I} = \{1, 2, 3, 4\}$ of 4 items ($n = 4$), with the following item weights:

	$i = 1$	$i = 2$	$i = 3$	$i = 4$
w_i	6	6	4	4

Questions

1. Apply the Next Fit (NF) algorithm to this instance. Show the execution step by step, indicating at each iteration whether the item is packed into the currently open bin or a new bin is opened, and the resulting residual capacity. State the final bin assignment and $NF(I)$.
2. Construct the packing patterns $P_1, \ldots, P_{NF(I)}$ from the bin vector obtained in Question 1.
3. Compute a lower bound on $z(I)$ and determine whether NF finds an optimal solution on this instance.
4. Verify whether the packing patterns found by NF satisfy the half-full property (i.e., at most one packing pattern has total weight at most $c/2$), and determine whether the solution found is optimal for this BPP instance, either by comparing $NF(I)$ with a lower bound on $z(I)$, or by exhibiting a feasible solution of strictly smaller value.

Solution

1. The NF algorithm produces the following feasible solution:

   (a) Item 1 (weight 6) is packed into bin 1. The residual capacity of bin 1 becomes $10 - 6 = 4$.

   (b) Item 2 (weight 6) does not fit into bin 1 (residual capacity 4). Thus bin 1 is closed and item 2 is packed into a new bin 2. The residual capacity of bin 2 becomes $10 - 6 = 4$.

   (c) Item 3 (weight 4) is packed into bin 2. The residual capacity of bin 2 becomes $4 - 4 = 0$.

   (d) Item 4 (weight 4) does not fit into bin 2 (residual capacity 0). Thus bin 2 is closed and item 4 is packed into a new bin 3. The residual capacity of bin 3 becomes $10 - 4 = 6$.

   Hence NF returns:

   $$bin = \begin{pmatrix} 1 & 2 & 2 & 3 \end{pmatrix}, \qquad nb = 3.$$

   and therefore

   $$NF(I) = nb = 3.$$

2. From the bin vector $bin = (1, 2, 2, 3)$, the packing patterns are

   $$P_1 = \{i \in \{1,2,3,4\} : bin_i = 1\} = \{1\},$$
   $$P_2 = \{i \in \{1,2,3,4\} : bin_i = 2\} = \{2,3\},$$
   $$P_3 = \{i \in \{1,2,3,4\} : bin_i = 3\} = \{4\}.$$

3. Accordingly, 3 is an upper bound on the optimal number of bins for this instance:

   $$z(I) \leq 3.$$

   Moreover,

   $$z(I) \geq \left\lceil \frac{\sum_{i \in \mathscr{I}} w_i}{c} \right\rceil = \left\lceil \frac{6+6+4+4}{10} \right\rceil = \left\lceil \frac{20}{10} \right\rceil = 2.$$

   In addition, the following is a feasible BPP solution:

   $$\tilde{P}_1 = \{1,3\}, \qquad \tilde{P}_2 = \{2,4\}$$

   which uses 2 bins; hence $z(I) \leq 2$. Therefore $z(I) = 2$, and NF does **not** compute an optimal solution on this instance.

4. From Question 2, the packing patterns found by NF have total weights

   $$\sum_{i \in P_1} w_i = 6, \qquad \sum_{i \in P_2} w_i = 6 + 4 = 10, \qquad \sum_{i \in P_3} w_i = 4.$$

   Since $c/2 = 5$, only P_3 has total weight at most $c/2$ (the other two exceed it); hence the solution **satisfies** the half-full property.

   As shown in Question 3, $NF(I) = 3$ while the feasible solution $\tilde{P}_1 = \{1,3\}$, $\tilde{P}_2 = \{2,4\}$ uses only 2 bins; since this is strictly smaller than $NF(I)$, the solution found by NF is **not optimal** (indeed $z(I) = 2 < NF(I) = 3$).

Exercise 26.2 — Next Fit 2

We consider a bin of capacity $c = 10$ and a set $\mathscr{I} = \{1, 2, 3, 4\}$ of 4 items ($n = 4$), with the following item weights:

	$i = 1$	$i = 2$	$i = 3$	$i = 4$
w_i	7	7	3	3

Questions

1. Apply the Next Fit (NF) algorithm to this instance. Show the execution step by step, indicating at each iteration whether the item is packed into the currently open bin or a new bin is opened, and the resulting residual capacity. State the final bin assignment and $NF(I)$.
2. Construct the packing patterns $P_1, \ldots, P_{NF(I)}$ from the bin vector obtained in Question 1.
3. Compute a lower bound on $z(I)$ and determine whether NF finds an optimal solution on this instance.
4. Verify whether the packing patterns found by NF satisfy the half-full property (i.e., at most one packing pattern has total weight at most $c/2$), and determine whether the solution found is optimal for this BPP instance, either by comparing $NF(I)$ with a lower bound on $z(I)$, or by exhibiting a feasible solution of strictly smaller value.

Solution

1. The NF algorithm produces the following feasible solution:
   (a) Item 1 (weight 7) is packed into bin 1. The residual capacity of bin 1 becomes $10 - 7 = 3$.
   (b) Item 2 (weight 7) does not fit into bin 1 (residual capacity 3). Thus bin 1 is closed and item 2 is packed into a new bin 2. The residual capacity of bin 2 becomes $10 - 7 = 3$.
   (c) Item 3 (weight 3) is packed into bin 2 (residual capacity 3). The residual capacity of bin 2 becomes $3 - 3 = 0$.
   (d) Item 4 (weight 3) does not fit into bin 2 (residual capacity 0). Thus bin 2 is closed and item 4 is packed into a new bin 3. The residual capacity of bin 3 becomes $10 - 3 = 7$.

   Hence NF returns:
   $$bin = \begin{pmatrix} 1 & 2 & 2 & 3 \end{pmatrix}, \qquad nb = 3.$$
   and therefore
   $$NF(I) = nb = 3.$$

2. From the bin vector $bin = (1, 2, 2, 3)$, the packing patterns are

$$P_1 = \{i \in \{1,2,3,4\} : bin_i = 1\} = \{1\},$$
$$P_2 = \{i \in \{1,2,3,4\} : bin_i = 2\} = \{2,3\},$$
$$P_3 = \{i \in \{1,2,3,4\} : bin_i = 3\} = \{4\}.$$

3. Accordingly, 3 is an upper bound on the optimal number of bins for this instance:

$$z(I) \leq 3.$$

Moreover,

$$z(I) \geq \left\lceil \frac{\sum_{i \in \mathscr{I}} w_i}{c} \right\rceil = \left\lceil \frac{7+7+3+3}{10} \right\rceil = \left\lceil \frac{20}{10} \right\rceil = 2.$$

In addition, the following is a feasible BPP solution:

$$\tilde{P}_1 = \{1,3\}, \qquad \tilde{P}_2 = \{2,4\}$$

which uses 2 bins (since $w_1 + w_3 = 7 + 3 = 10$ and $w_2 + w_4 = 7 + 3 = 10$); hence $z(I) \leq 2$. Therefore $z(I) = 2$, and NF does **not** compute an optimal solution on this instance.

4. From Question 2, the packing patterns found by NF have total weights

$$\sum_{i \in P_1} w_i = 7, \qquad \sum_{i \in P_2} w_i = 7 + 3 = 10, \qquad \sum_{i \in P_3} w_i = 3.$$

Since $c/2 = 5$, only P_3 has total weight at most $c/2$ (the other two exceed it); hence the solution **satisfies** the half-full property.

As shown in Question 3, $NF(I) = 3$ while the feasible solution $\tilde{P}_1 = \{1,3\}$, $\tilde{P}_2 = \{2,4\}$ uses only 2 bins; since this is strictly smaller than $NF(I)$, the solution found by NF is **not optimal** (indeed $z(I) = 2 < NF(I) = 3$).

Exercise 26.3 — Next Fit 3

We consider a bin of capacity $c = 10$ and a set $\mathscr{I} = \{1,2,3,4,5\}$ of 5 items ($n = 5$), with the following item weights:

	$i=1$	$i=2$	$i=3$	$i=4$	$i=5$
w_i	4	7	6	3	5

Questions

1. Apply the Next Fit (NF) algorithm to this instance. Show the execution step by step, indicating at each iteration whether the item is packed into the currently open bin or a new bin is opened, and the resulting residual capacity. State the final bin assignment and $NF(I)$.

2. Construct the packing patterns $P_1, \ldots, P_{NF(I)}$ from the bin vector obtained in Question 1.

3. Compute a lower bound on $z(I)$ and determine whether NF finds an optimal solution on this instance.

4. Verify whether the packing patterns found by NF satisfy the half-full property (i.e., at most one packing pattern has total weight at most $c/2$), and determine whether the solution found is optimal for this BPP instance, either by comparing $NF(I)$ with a lower bound on $z(I)$, or by exhibiting a feasible solution of strictly smaller value.

Solution

1. The NF algorithm produces the following feasible solution:

   (a) Item 1 (weight 4) is packed into bin 1. The residual capacity of bin 1 becomes $10 - 4 = 6$.

   (b) Item 2 (weight 7) does not fit into bin 1 (residual capacity 6). Thus bin 1 is closed and item 2 is packed into a new bin 2. The residual capacity of bin 2 becomes $10 - 7 = 3$.

   (c) Item 3 (weight 6) does not fit into bin 2 (residual capacity 3). Thus bin 2 is closed and item 3 is packed into a new bin 3. The residual capacity of bin 3 becomes $10 - 6 = 4$.

   (d) Item 4 (weight 3) is packed into bin 3 (residual capacity 4). The residual capacity of bin 3 becomes $4 - 3 = 1$.

   (e) Item 5 (weight 5) does not fit into bin 3 (residual capacity 1). Thus bin 3 is closed and item 5 is packed into a new bin 4. The residual capacity of bin 4 becomes $10 - 5 = 5$.

   Hence NF returns:

   $$bin = \begin{pmatrix} 1 & 2 & 3 & 3 & 4 \end{pmatrix}, \qquad nb = 4.$$

   and therefore

   $$NF(I) = nb = 4.$$

2. From the bin vector $bin = (1, 2, 3, 3, 4)$, the packing patterns are

   $$\begin{aligned} P_1 &= \{i : bin_i = 1\} = \{1\}, \\ P_2 &= \{i : bin_i = 2\} = \{2\}, \\ P_3 &= \{i : bin_i = 3\} = \{3, 4\}, \\ P_4 &= \{i : bin_i = 4\} = \{5\}. \end{aligned}$$

3. Accordingly, 4 is an upper bound on the optimal number of bins for this instance:

   $$z(I) \leq 4.$$

Moreover,

$$z(I) \geq \left\lceil \frac{\sum_{i \in \mathscr{I}} w_i}{c} \right\rceil = \left\lceil \frac{4+7+6+3+5}{10} \right\rceil = \left\lceil \frac{25}{10} \right\rceil = 3.$$

In addition, the following is a feasible BPP solution:

$$\tilde{P}_1 = \{2,4\}, \qquad \tilde{P}_2 = \{1,3\}, \qquad \tilde{P}_3 = \{5\}$$

which uses 3 bins (since $w_2 + w_4 = 7+3 = 10$ and $w_1 + w_3 = 4+6 = 10$); hence $z(I) \leq 3$. Therefore $z(I) = 3$, and NF does **not** compute an optimal solution on this instance.

4. From Question 2, the packing patterns found by NF have total weights

$$\sum_{i \in P_1} w_i = 4, \qquad \sum_{i \in P_2} w_i = 7, \qquad \sum_{i \in P_3} w_i = 6+3 = 9, \qquad \sum_{i \in P_4} w_i = 5.$$

Since $c/2 = 5$, *both* P_1 (weight 4) and P_4 (weight 5) have total weight at most $c/2$; hence the solution found by NF **does not satisfy** the half-full property (this is possible precisely because NF is not an Any Fit algorithm).

As shown in Question 3, $NF(I) = 4$ while the feasible solution $\tilde{P}_1 = \{2,4\}$, $\tilde{P}_2 = \{1,3\}$, $\tilde{P}_3 = \{5\}$ uses only 3 bins; since this is strictly smaller than $NF(I)$, the solution found by NF is **not optimal** (indeed $z(I) = 3 < NF(I) = 4$).

26.2 First Fit

Exercise 26.4 — First Fit 1

We consider a bin of capacity $c = 10$ and a set $\mathscr{I} = \{1,2,3,4\}$ of 4 items ($n = 4$), with the following item weights:

	$i=1$	$i=2$	$i=3$	$i=4$
w_i	6	6	4	4

Questions

1. Apply the First Fit (FF) algorithm to this instance. Show the execution step by step, indicating at each iteration which bin the item is packed into (scanning bins in the order they were opened) and the resulting residual capacity. State the final bin assignment and $FF(I)$.
2. Construct the packing patterns $P_1, \ldots, P_{FF(I)}$ from the bin vector obtained in Question 1.
3. Compute a lower bound on $z(I)$ and determine whether FF finds an optimal solution on this instance.

4. Verify whether the packing patterns found by FF satisfy the half-full property (i.e., at most one packing pattern has total weight at most $c/2$), and determine whether the solution found is optimal for this BPP instance, either by comparing $FF(I)$ with a lower bound on $z(I)$, or by exhibiting a feasible solution of strictly smaller value.

Solution

1. Recall the FF pseudocode: for each item i, $s := -1$ and, for $b := 1$ to nb, if $w[i] \leq rc[b]$ then $s \leftarrow b$ and stop the scan; if $s = -1$ after the scan, a new bin is opened.

   (a) Item 1 (weight 6): bin 1 is open at the start ($nb = 1$, $rc[1] = c = 10$) and $w[1] \leq rc[1]$, so $s = 1$ and FF packs item 1 into bin 1, and $rc[1] \leftarrow 10 - 6 = 4$.

   (b) Item 2 (weight 6): $b = 1$: $w[2] = 6 \leq rc[1] = 4$? No. No bin fits, so FF opens bin 2 ($rc[2] \leftarrow 10$). Item 2 is packed into bin 2, and $rc[2] \leftarrow 10 - 6 = 4$.

   (c) Item 3 (weight 4): $b = 1$: $w[3] = 4 \leq rc[1] = 4$? Yes, so $s \leftarrow 1$ and the scan stops. Item 3 is packed into bin 1, and $rc[1] \leftarrow 4 - 4 = 0$.

   (d) Item 4 (weight 4): $b = 1$: $w[4] = 4 \leq rc[1] = 0$? No. $b = 2$: $w[4] = 4 \leq rc[2] = 4$? Yes, so $s \leftarrow 2$. Item 4 is packed into bin 2, and $rc[2] \leftarrow 4 - 4 = 0$.

   Hence FF returns:

   $$bin = \begin{pmatrix} 1 & 2 & 1 & 2 \end{pmatrix}, \qquad nb = 2.$$

   and therefore

   $$FF(I) = nb = 2.$$

2. From the bin vector $bin = (1, 2, 1, 2)$, the packing patterns are

   $$P_1 = \{i \in \{1, 2, 3, 4\} : bin_i = 1\} = \{1, 3\},$$
   $$P_2 = \{i \in \{1, 2, 3, 4\} : bin_i = 2\} = \{2, 4\}.$$

3. Accordingly,

   $$z(I) \leq 2.$$

   Moreover,

   $$z(I) \geq \left\lceil \frac{\sum_{i \in \mathscr{I}} w_i}{c} \right\rceil = \left\lceil \frac{6 + 6 + 4 + 4}{10} \right\rceil = \left\lceil \frac{20}{10} \right\rceil = 2.$$

   Hence $z(I) = 2$ and FF computes an **optimal** solution on this instance.

4. From Question 2, the packing patterns found by FF have total weights

   $$\sum_{i \in P_1} w_i = 6 + 4 = 10, \qquad \sum_{i \in P_2} w_i = 6 + 4 = 10.$$

   Since $c/2 = 5$, no packing pattern has total weight at most $c/2$; in particular *at most one* does, so the solution **satisfies** the half-full property.

   Since $FF(I) = 2$ matches the lower bound $z(I) \geq 2$ found in Question 3, the solution found by FF is **optimal**.

Exercise 26.5 — First Fit 2

We consider a bin of capacity $c = 10$ and a set $\mathscr{I} = \{1, 2, 3, 4\}$ of 4 items ($n = 4$), with the following item weights:

	$i = 1$	$i = 2$	$i = 3$	$i = 4$
w_i	7	7	3	3

Questions

1. Apply the First Fit (FF) algorithm to this instance. Show the execution step by step, indicating at each iteration which bin the item is packed into (scanning bins in the order they were opened) and the resulting residual capacity. State the final bin assignment and $FF(I)$.
2. Construct the packing patterns $P_1, \ldots, P_{FF(I)}$ from the bin vector obtained in Question 1.
3. Compute a lower bound on $z(I)$ and determine whether FF finds an optimal solution on this instance.
4. Verify whether the packing patterns found by FF satisfy the half-full property (i.e., at most one packing pattern has total weight at most $c/2$), and determine whether the solution found is optimal for this BPP instance, either by comparing $FF(I)$ with a lower bound on $z(I)$, or by exhibiting a feasible solution of strictly smaller value.

Solution

1. Recall the FF pseudocode: for each item i, $s := -1$ and, for $b := 1$ to nb, if $w[i] \le rc[b]$ then $s \leftarrow b$ and stop the scan; if $s = -1$ after the scan, a new bin is opened.

   (a) Item 1 (weight 7): bin 1 is open at the start ($nb = 1$, $rc[1] = c = 10$) and $w[1] \le rc[1]$, so $s = 1$ and FF packs item 1 into bin 1, and $rc[1] \leftarrow 10 - 7 = 3$.

   (b) Item 2 (weight 7): $b = 1$: $w[2] = 7 \le rc[1] = 3$? No. No bin fits, so FF opens bin 2 ($rc[2] \leftarrow 10$). Item 2 is packed into bin 2, and $rc[2] \leftarrow 10 - 7 = 3$.

   (c) Item 3 (weight 3): $b = 1$: $w[3] = 3 \le rc[1] = 3$? Yes, so $s \leftarrow 1$ and the scan stops. Item 3 is packed into bin 1, and $rc[1] \leftarrow 3 - 3 = 0$.

   (d) Item 4 (weight 3): $b = 1$: $w[4] = 3 \le rc[1] = 0$? No. $b = 2$: $w[4] = 3 \le rc[2] = 3$? Yes, so $s \leftarrow 2$. Item 4 is packed into bin 2, and $rc[2] \leftarrow 3 - 3 = 0$.

   Hence FF returns:

   $$bin = \begin{pmatrix} 1 & 2 & 1 & 2 \end{pmatrix}, \qquad nb = 2.$$

   and therefore

   $$FF(I) = nb = 2.$$

2. From the bin vector $bin = (1, 2, 1, 2)$, the packing patterns are

$$P_1 = \{i \in \{1, 2, 3, 4\} : bin_i = 1\} = \{1, 3\},$$
$$P_2 = \{i \in \{1, 2, 3, 4\} : bin_i = 2\} = \{2, 4\}.$$

3. Accordingly,

$$z(I) \leq 2.$$

Moreover,

$$z(I) \geq \left\lceil \frac{\sum_{i \in \mathscr{I}} w_i}{c} \right\rceil = \left\lceil \frac{7 + 7 + 3 + 3}{10} \right\rceil = \left\lceil \frac{20}{10} \right\rceil = 2.$$

Hence $z(I) = 2$ and FF computes an **optimal** solution on this instance.

4. From Question 2, the packing patterns found by FF have total weights

$$\sum_{i \in P_1} w_i = 7 + 3 = 10, \qquad \sum_{i \in P_2} w_i = 7 + 3 = 10.$$

Since $c/2 = 5$, no packing pattern has total weight at most $c/2$; in particular *at most one* does, so the solution **satisfies** the half-full property.

Since $FF(I) = 2$ matches the lower bound $z(I) \geq 2$ found in Question 3, the solution found by FF is **optimal**.

Exercise 26.6 — First Fit 3

We consider a bin of capacity $c = 10$ and a set $\mathscr{I} = \{1, 2, 3, 4, 5\}$ of 5 items ($n = 5$), with the following item weights:

	$i = 1$	$i = 2$	$i = 3$	$i = 4$	$i = 5$
w_i	4	7	6	3	5

Questions

1. Apply the First Fit (FF) algorithm to this instance. Show the execution step by step, indicating at each iteration which bin the item is packed into (scanning bins in the order they were opened) and the resulting residual capacity. State the final bin assignment and $FF(I)$.

2. Construct the packing patterns $P_1, \ldots, P_{FF(I)}$ from the bin vector obtained in Question 1.

3. Compute a lower bound on $z(I)$ and determine whether FF finds an optimal solution on this instance.

4. Verify whether the packing patterns found by FF satisfy the half-full property (i.e., at most one packing pattern has total weight at most $c/2$), and determine whether

the solution found is optimal for this BPP instance, either by comparing $FF(I)$ with a lower bound on $z(I)$, or by exhibiting a feasible solution of strictly smaller value.

Solution

1. Recall the FF pseudocode: for each item i, $s := -1$ and, for $b := 1$ to nb, if $w[i] \le rc[b]$ then $s \leftarrow b$ and stop the scan; if $s = -1$ after the scan, a new bin is opened.

   (a) Item 1 (weight 4): bin 1 is open at the start ($nb = 1$, $rc[1] = c = 10$) and $w[1] \le rc[1]$, so $s = 1$ and FF packs item 1 into bin 1, and $rc[1] \leftarrow 10 - 4 = 6$.

   (b) Item 2 (weight 7): $b = 1$: $w[2] = 7 \le rc[1] = 6$? No. No bin fits, so FF opens bin 2 ($rc[2] \leftarrow 10$). Item 2 is packed into bin 2, and $rc[2] \leftarrow 10 - 7 = 3$.

   (c) Item 3 (weight 6): $b = 1$: $w[3] = 6 \le rc[1] = 6$? Yes, so $s \leftarrow 1$ and the scan stops. Item 3 is packed into bin 1, and $rc[1] \leftarrow 6 - 6 = 0$.

   (d) Item 4 (weight 3): $b = 1$: $w[4] = 3 \le rc[1] = 0$? No. $b = 2$: $w[4] = 3 \le rc[2] = 3$? Yes, so $s \leftarrow 2$. Item 4 is packed into bin 2, and $rc[2] \leftarrow 3 - 3 = 0$.

   (e) Item 5 (weight 5): $b = 1$: $w[5] = 5 \le rc[1] = 0$? No. $b = 2$: $w[5] = 5 \le rc[2] = 0$? No. No bin fits, so FF opens bin 3 ($rc[3] \leftarrow 10$). Item 5 is packed into bin 3, and $rc[3] \leftarrow 10 - 5 = 5$.

   Hence FF returns:

   $$bin = \begin{pmatrix} 1 & 2 & 1 & 2 & 3 \end{pmatrix}, \qquad nb = 3.$$

   and therefore

   $$FF(I) = nb = 3.$$

2. From the bin vector $bin = (1, 2, 1, 2, 3)$, the packing patterns are

   $$P_1 = \{i : bin_i = 1\} = \{1, 3\},$$
   $$P_2 = \{i : bin_i = 2\} = \{2, 4\},$$
   $$P_3 = \{i : bin_i = 3\} = \{5\}.$$

3. Accordingly,

   $$z(I) \le 3.$$

   Moreover,

   $$z(I) \ge \left\lceil \frac{\sum_{i \in \mathcal{I}} w_i}{c} \right\rceil = \left\lceil \frac{4 + 7 + 6 + 3 + 5}{10} \right\rceil = \left\lceil \frac{25}{10} \right\rceil = 3.$$

   Hence $z(I) = 3$ and FF computes an **optimal** solution on this instance.

4. From Question 2, the packing patterns found by FF have total weights

   $$\sum_{i \in P_1} w_i = 4 + 6 = 10, \qquad \sum_{i \in P_2} w_i = 7 + 3 = 10, \qquad \sum_{i \in P_3} w_i = 5.$$

   Since $c/2 = 5$, only P_3 has total weight at most $c/2$ (the other two exceed it); hence the solution **satisfies** the half-full property.

   Since $FF(I) = 3$ matches the lower bound $z(I) \ge 3$ found in Question 3, the solution found by FF is **optimal**.

26.3 Best Fit

Exercise 26.7 — Best Fit 1

We consider a bin of capacity $c = 10$ and a set $\mathscr{I} = \{1, 2, 3, 4\}$ of 4 items ($n = 4$), with the following item weights:

	$i = 1$	$i = 2$	$i = 3$	$i = 4$
w_i	6	6	4	4

Questions

1. Apply the Best Fit (BF) algorithm to this instance. Show the execution step by step: for each item, trace the pseudocode variables $minrc$ and s over the scan of the open bins, and state into which bin the item is packed and the resulting residual capacity.
2. Construct the packing patterns $P_1, \ldots, P_{BF(I)}$ from the bin vector obtained in Question 1.
3. Compute a lower bound on $z(I)$ and determine whether BF finds an optimal solution on this instance.
4. Verify whether the packing patterns found by BF satisfy the half-full property (i.e., at most one packing pattern has total weight at most $c/2$), and determine whether the solution found is optimal for this BPP instance, either by comparing $BF(I)$ with a lower bound on $z(I)$, or by exhibiting a feasible solution of strictly smaller value.

Solution

1. Recall the BF pseudocode: for each item, $minrc := c + 1$, $s := -1$, and for $b := 1$ to nb, if $w[i] \le rc[b]$ and $rc[b] < minrc$ then $s \leftarrow b$, $minrc \leftarrow rc[b]$.

   (a) Item 1 (weight 6): bin 1 is open at the start ($nb = 1$, $rc[1] = c$) and item 1 fits into it, so $s = 1$ and BF packs item 1 into bin 1. Residual capacity of bin 1: $10 - 6 = 4$.

   (b) Item 2 (weight 6): $minrc := 11$, $s := -1$. For $b = 1$: $w[2] = 6 \le rc[1] = 4$? No. No bin fits, so BF opens bin 2 and packs item 2 into it. Residual capacity of bin 2: $10 - 6 = 4$.

   (c) Item 3 (weight 4): $minrc := 11$, $s := -1$. For $b = 1$: $w[3] = 4 \le rc[1] = 4$ and $rc[1] = 4 < 11$, so $s \leftarrow 1$, $minrc \leftarrow 4$. For $b = 2$: $w[3] = 4 \le rc[2] = 4$, but $rc[2] = 4$ is *not* strictly smaller than $minrc = 4$, so s is not updated. Hence $s = 1$: item 3 is packed into bin 1 (the first bin opened, since the tie between bin 1 and bin 2 is broken by the strict inequality in the pseudocode). Residual capacity of bin 1: $4 - 4 = 0$.

(d) Item 4 (weight 4): $minrc := 11$, $s := -1$. For $b = 1$: $w[4] = 4 \leq rc[1] = 0$? No. For $b = 2$: $w[4] = 4 \leq rc[2] = 4$ and $rc[2] = 4 < 11$, so $s \leftarrow 2$, $minrc \leftarrow 4$. Hence $s = 2$: item 4 is packed into bin 2. Residual capacity of bin 2: $4 - 4 = 0$.

Hence BF returns:

$$bin = \begin{pmatrix} 1 & 2 & 1 & 2 \end{pmatrix}, \qquad nb = 2.$$

and therefore

$$BF(I) = nb = 2.$$

2. From the bin vector $bin = (1, 2, 1, 2)$, the packing patterns are

$$P_1 = \{i \in \{1, 2, 3, 4\} : bin_i = 1\} = \{1, 3\},$$
$$P_2 = \{i \in \{1, 2, 3, 4\} : bin_i = 2\} = \{2, 4\}.$$

3. Accordingly,

$$z(I) \leq 2.$$

Moreover,

$$z(I) \geq \left\lceil \frac{\sum_{i \in \mathscr{I}} w_i}{c} \right\rceil = \left\lceil \frac{6 + 6 + 4 + 4}{10} \right\rceil = \left\lceil \frac{20}{10} \right\rceil = 2.$$

Hence $z(I) = 2$ and BF computes an **optimal** solution on this instance.

4. From Question 2, the packing patterns found by BF have total weights

$$\sum_{i \in P_1} w_i = 6 + 4 = 10, \qquad \sum_{i \in P_2} w_i = 6 + 4 = 10.$$

Since $c/2 = 5$, no packing pattern has total weight at most $c/2$; in particular *at most one* does, so the solution **satisfies** the half-full property.

Since $BF(I) = 2$ matches the lower bound $z(I) \geq 2$ found in Question 3, the solution found by BF is **optimal**.

Exercise 26.8 — Best Fit 2

We consider a bin of capacity $c = 10$ and a set $\mathscr{I} = \{1, 2, 3, 4\}$ of 4 items ($n = 4$), with the following item weights:

	$i = 1$	$i = 2$	$i = 3$	$i = 4$
w_i	7	7	3	3

Questions

1. Apply the Best Fit (BF) algorithm to this instance. Show the execution step by step: for each item, trace the pseudocode variables $minrc$ and s over the scan of the open bins, and state into which bin the item is packed and the resulting residual capacity.

2. Construct the packing patterns $P_1, \ldots, P_{BF(I)}$ from the bin vector obtained in Question 1.

3. Compute a lower bound on $z(I)$ and determine whether BF finds an optimal solution on this instance.

4. Verify whether the packing patterns found by BF satisfy the half-full property (i.e., at most one packing pattern has total weight at most $c/2$), and determine whether the solution found is optimal for this BPP instance, either by comparing $BF(I)$ with a lower bound on $z(I)$, or by exhibiting a feasible solution of strictly smaller value.

Solution

1. Recall the BF pseudocode: for each item i, $minrc := c + 1$, $s := -1$, and for $b := 1$ to nb, if $w[i] \leq rc[b]$ and $rc[b] < minrc$ then $s \leftarrow b$, $minrc \leftarrow rc[b]$.

   (a) Item 1 (weight 7): bin 1 is open at the start ($nb = 1$, $rc[1] = c$) and item 1 fits into it, so $s = 1$ and BF packs item 1 into bin 1. Residual capacity of bin 1: $10 - 7 = 3$.

   (b) Item 2 (weight 7): $minrc := 11$, $s := -1$. For $b = 1$: $w[2] = 7 \leq rc[1] = 3$? No. No bin fits, so BF opens bin 2 and packs item 2 into it. Residual capacity of bin 2: $10 - 7 = 3$.

   (c) Item 3 (weight 3): $minrc := 11$, $s := -1$. For $b = 1$: $w[3] = 3 \leq rc[1] = 3$ and $rc[1] = 3 < 11$, so $s \leftarrow 1$, $minrc \leftarrow 3$. For $b = 2$: $w[3] = 3 \leq rc[2] = 3$, but $rc[2] = 3$ is *not* strictly smaller than $minrc = 3$, so s is not updated. Hence $s = 1$: item 3 is packed into bin 1 (the first bin opened, since the tie between bin 1 and bin 2 is broken by the strict inequality in the pseudocode). Residual capacity of bin 1: $3 - 3 = 0$.

   (d) Item 4 (weight 3): $minrc := 11$, $s := -1$. For $b = 1$: $w[4] = 3 \leq rc[1] = 0$? No. For $b = 2$: $w[4] = 3 \leq rc[2] = 3$ and $rc[2] = 3 < 11$, so $s \leftarrow 2$, $minrc \leftarrow 3$. Hence $s = 2$: item 4 is packed into bin 2. Residual capacity of bin 2: $3 - 3 = 0$.

   Hence BF returns:

$$bin = \begin{pmatrix} 1 & 2 & 1 & 2 \end{pmatrix}, \qquad nb = 2.$$

   and therefore

$$BF(I) = nb = 2.$$

2. From the bin vector $bin = (1, 2, 1, 2)$, the packing patterns are

$$P_1 = \{i \in \{1, 2, 3, 4\} : bin_i = 1\} = \{1, 3\},$$
$$P_2 = \{i \in \{1, 2, 3, 4\} : bin_i = 2\} = \{2, 4\}.$$

3. Accordingly,

$$z(I) \leq 2.$$

   Moreover,

$$z(I) \geq \left\lceil \frac{\sum_{i \in \mathscr{I}} w_i}{c} \right\rceil = \left\lceil \frac{7 + 7 + 3 + 3}{10} \right\rceil = \left\lceil \frac{20}{10} \right\rceil = 2.$$

Hence $z(I) = 2$ and BF computes an **optimal** solution on this instance.

4. From Question 2, the packing patterns found by BF have total weights

$$\sum_{i \in P_1} w_i = 7 + 3 = 10, \qquad \sum_{i \in P_2} w_i = 7 + 3 = 10.$$

Since $c/2 = 5$, no packing pattern has total weight at most $c/2$; in particular *at most one* does, so the solution **satisfies** the half-full property.

Since $BF(I) = 2$ matches the lower bound $z(I) \geq 2$ found in Question 3, the solution found by BF is **optimal**.

Exercise 26.9 — Best Fit 3

We consider a bin of capacity $c = 10$ and a set $\mathscr{I} = \{1, 2, 3, 4, 5\}$ of 5 items ($n = 5$), with the following item weights:

	$i = 1$	$i = 2$	$i = 3$	$i = 4$	$i = 5$
w_i	4	7	6	3	5

Questions

1. Apply the Best Fit (BF) algorithm to this instance. Show the execution step by step: for each item, trace the pseudocode variables $minrc$ and s over the scan of the open bins, and state into which bin the item is packed and the resulting residual capacity.
2. Construct the packing patterns $P_1, \ldots, P_{BF(I)}$ from the bin vector obtained in Question 1.
3. Compute a lower bound on $z(I)$ and determine whether BF finds an optimal solution on this instance.
4. Verify whether the packing patterns found by BF satisfy the half-full property (i.e., at most one packing pattern has total weight at most $c/2$), and determine whether the solution found is optimal for this BPP instance, either by comparing $BF(I)$ with a lower bound on $z(I)$, or by exhibiting a feasible solution of strictly smaller value.

Solution

1. Recall the BF pseudocode: for each item i, $minrc := c + 1$, $s := -1$, and for $b := 1$ to nb, if $w[i] \leq rc[b]$ and $rc[b] < minrc$ then $s \leftarrow b$, $minrc \leftarrow rc[b]$.

   (a) Item 1 (weight 4): bin 1 is open at the start ($nb = 1$, $rc[1] = c$) and item 1 fits into it, so $s = 1$ and BF packs item 1 into bin 1. Residual capacity of bin 1: $10 - 4 = 6$.

(b) Item 2 (weight 7): $minrc := 11$, $s := -1$. For $b = 1$: $w[2] = 7 \leq rc[1] = 6$? No. No bin fits, so BF opens bin 2 and packs item 2 into it. Residual capacity of bin 2: $10 - 7 = 3$.

(c) Item 3 (weight 6): $minrc := 11$, $s := -1$. For $b = 1$: $w[3] = 6 \leq rc[1] = 6$ and $rc[1] = 6 < 11$, so $s \leftarrow 1$, $minrc \leftarrow 6$. For $b = 2$: $w[3] = 6 \leq rc[2] = 3$? No. Hence $s = 1$: item 3 is packed into bin 1. Residual capacity of bin 1: $6 - 6 = 0$.

(d) Item 4 (weight 3): $minrc := 11$, $s := -1$. For $b = 1$: $w[4] = 3 \leq rc[1] = 0$? No. For $b = 2$: $w[4] = 3 \leq rc[2] = 3$ and $rc[2] = 3 < 11$, so $s \leftarrow 2$, $minrc \leftarrow 3$. Hence $s = 2$: item 4 is packed into bin 2. Residual capacity of bin 2: $3 - 3 = 0$.

(e) Item 5 (weight 5): $minrc := 11$, $s := -1$. For $b = 1$: $w[5] = 5 \leq rc[1] = 0$? No. For $b = 2$: $w[5] = 5 \leq rc[2] = 0$? No. No bin fits, so BF opens bin 3 and packs item 5 into it. Residual capacity of bin 3: $10 - 5 = 5$.

Hence BF returns:

$$bin = \begin{pmatrix} 1 & 2 & 1 & 2 & 3 \end{pmatrix}, \qquad nb = 3.$$

and therefore

$$BF(I) = nb = 3.$$

2. From the bin vector $bin = (1, 2, 1, 2, 3)$, the packing patterns are

$$P_1 = \{i : bin_i = 1\} = \{1, 3\},$$
$$P_2 = \{i : bin_i = 2\} = \{2, 4\},$$
$$P_3 = \{i : bin_i = 3\} = \{5\}.$$

3. Accordingly,

$$z(I) \leq 3.$$

Moreover,

$$z(I) \geq \left\lceil \frac{\sum_{i \in \mathscr{I}} w_i}{c} \right\rceil = \left\lceil \frac{4 + 7 + 6 + 3 + 5}{10} \right\rceil = \left\lceil \frac{25}{10} \right\rceil = 3.$$

Hence $z(I) = 3$ and BF computes an **optimal** solution on this instance.

4. From Question 2, the packing patterns found by BF have total weights

$$\sum_{i \in P_1} w_i = 4 + 6 = 10, \qquad \sum_{i \in P_2} w_i = 7 + 3 = 10, \qquad \sum_{i \in P_3} w_i = 5.$$

Since $c/2 = 5$, only P_3 has total weight at most $c/2$ (the other two exceed it); hence the solution **satisfies** the half-full property.

Since $BF(I) = 3$ matches the lower bound $z(I) \geq 3$ found in Question 3, the solution found by BF is **optimal**.

26.4 Worst Fit

Exercise 26.10 — Worst Fit 1

We consider a bin of capacity $c = 10$ and a set $\mathscr{I} = \{1, 2, 3, 4\}$ of 4 items ($n = 4$), with the following item weights:

	$i = 1$	$i = 2$	$i = 3$	$i = 4$
w_i	6	6	4	4

Questions

1. Apply the Worst Fit (WF) algorithm to this instance. Show the execution step by step: for each item, trace the pseudocode variables $maxrc$ and s over the scan of the open bins, and state into which bin the item is packed and the resulting residual capacity.
2. Construct the packing patterns $P_1, \ldots, P_{WF(I)}$ from the bin vector obtained in Question 1.
3. Compute a lower bound on $z(I)$ and determine whether WF finds an optimal solution on this instance.
4. Verify whether the packing patterns found by WF satisfy the half-full property (i.e., at most one packing pattern has total weight at most $c/2$), and determine whether the solution found is optimal for this BPP instance, either by comparing $WF(I)$ with a lower bound on $z(I)$, or by exhibiting a feasible solution of strictly smaller value.

Solution

1. Recall the WF pseudocode: for each item i, $maxrc := -1$, $s := -1$, and for $b := 1$ to nb, if $w[i] \leq rc[b]$ and $rc[b] > maxrc$ then $s \leftarrow b$, $maxrc \leftarrow rc[b]$.
   (a) Item 1 (weight 6): bin 1 is open at the start ($nb = 1$, $rc[1] = c$) and item 1 fits into it, so $s = 1$ and WF packs item 1 into bin 1. Residual capacity of bin 1: $10 - 6 = 4$.
   (b) Item 2 (weight 6): $maxrc := -1$, $s := -1$. For $b = 1$: $w[2] = 6 \leq rc[1] = 4$? No. No bin fits, so WF opens bin 2 and packs item 2 into it. Residual capacity of bin 2: $10 - 6 = 4$.
   (c) Item 3 (weight 4): $maxrc := -1$, $s := -1$. For $b = 1$: $w[3] = 4 \leq rc[1] = 4$ and $rc[1] = 4 > -1$, so $s \leftarrow 1$, $maxrc \leftarrow 4$. For $b = 2$: $w[3] = 4 \leq rc[2] = 4$, but $rc[2] = 4$ is *not* strictly greater than $maxrc = 4$, so s is not updated. Hence $s = 1$: item 3 is packed into bin 1 (the first bin opened, since the tie between bin 1 and bin 2 is broken by the strict inequality in the pseudocode). Residual capacity of bin 1: $4 - 4 = 0$.

(d) Item 4 (weight 4): $maxrc := -1$, $s := -1$. For $b = 1$: $w[4] = 4 \leq rc[1] = 0$? No. For $b = 2$: $w[4] = 4 \leq rc[2] = 4$ and $rc[2] = 4 > -1$, so $s \leftarrow 2$, $maxrc \leftarrow 4$. Hence $s = 2$: item 4 is packed into bin 2. Residual capacity of bin 2: $4 - 4 = 0$.

Hence WF returns:

$$bin = \begin{pmatrix} 1 & 2 & 1 & 2 \end{pmatrix}, \qquad nb = 2.$$

and therefore

$$WF(I) = nb = 2.$$

2. From the bin vector $bin = (1, 2, 1, 2)$, the packing patterns are

$$P_1 = \{i \in \{1, 2, 3, 4\} : bin_i = 1\} = \{1, 3\},$$
$$P_2 = \{i \in \{1, 2, 3, 4\} : bin_i = 2\} = \{2, 4\}.$$

3. Accordingly,

$$z(I) \leq 2.$$

Moreover,

$$z(I) \geq \left\lceil \frac{\sum_{i \in \mathscr{I}} w_i}{c} \right\rceil = \left\lceil \frac{6 + 6 + 4 + 4}{10} \right\rceil = \left\lceil \frac{20}{10} \right\rceil = 2.$$

Hence $z(I) = 2$ and WF computes an **optimal** solution on this instance.

4. From Question 2, the packing patterns found by WF have total weights

$$\sum_{i \in P_1} w_i = 6 + 4 = 10, \qquad \sum_{i \in P_2} w_i = 6 + 4 = 10.$$

Since $c/2 = 5$, no packing pattern has total weight at most $c/2$; in particular *at most one* does, so the solution **satisfies** the half-full property.

Since $WF(I) = 2$ matches the lower bound $z(I) \geq 2$ found in Question 3, the solution found by WF is **optimal**.

Exercise 26.11 — Worst Fit 2

We consider a bin of capacity $c = 10$ and a set $\mathscr{I} = \{1, 2, 3, 4\}$ of 4 items ($n = 4$), with the following item weights:

	$i = 1$	$i = 2$	$i = 3$	$i = 4$
w_i	7	7	3	3

Questions

1. Apply the Worst Fit (WF) algorithm to this instance. Show the execution step by step: for each item, trace the pseudocode variables $maxrc$ and s over the scan of

the open bins, and state into which bin the item is packed and the resulting residual capacity.

2. Construct the packing patterns $P_1, \ldots, P_{WF(I)}$ from the bin vector obtained in Question 1.

3. Compute a lower bound on $z(I)$ and determine whether WF finds an optimal solution on this instance.

4. Verify whether the packing patterns found by WF satisfy the half-full property (i.e., at most one packing pattern has total weight at most $c/2$), and determine whether the solution found is optimal for this BPP instance, either by comparing $WF(I)$ with a lower bound on $z(I)$, or by exhibiting a feasible solution of strictly smaller value.

Solution

1. Recall the WF pseudocode: for each item i, $maxrc := -1$, $s := -1$, and for $b := 1$ to nb, if $w[i] \leq rc[b]$ and $rc[b] > maxrc$ then $s \leftarrow b$, $maxrc \leftarrow rc[b]$.

   (a) Item 1 (weight 7): bin 1 is open at the start ($nb = 1$, $rc[1] = c$) and item 1 fits into it, so $s = 1$ and WF packs item 1 into bin 1. Residual capacity of bin 1: $10 - 7 = 3$.

   (b) Item 2 (weight 7): $maxrc := -1$, $s := -1$. For $b = 1$: $w[2] = 7 \leq rc[1] = 3$? No. No bin fits, so WF opens bin 2 and packs item 2 into it. Residual capacity of bin 2: $10 - 7 = 3$.

   (c) Item 3 (weight 3): $maxrc := -1$, $s := -1$. For $b = 1$: $w[3] = 3 \leq rc[1] = 3$ and $rc[1] = 3 > -1$, so $s \leftarrow 1$, $maxrc \leftarrow 3$. For $b = 2$: $w[3] = 3 \leq rc[2] = 3$, but $rc[2] = 3$ is *not* strictly greater than $maxrc = 3$, so s is not updated. Hence $s = 1$: item 3 is packed into bin 1 (the first bin opened, since the tie between bin 1 and bin 2 is broken by the strict inequality in the pseudocode). Residual capacity of bin 1: $3 - 3 = 0$.

   (d) Item 4 (weight 3): $maxrc := -1$, $s := -1$. For $b = 1$: $w[4] = 3 \leq rc[1] = 0$? No. For $b = 2$: $w[4] = 3 \leq rc[2] = 3$ and $rc[2] = 3 > -1$, so $s \leftarrow 2$, $maxrc \leftarrow 3$. Hence $s = 2$: item 4 is packed into bin 2. Residual capacity of bin 2: $3 - 3 = 0$.

   Hence WF returns:

   $$bin = \begin{pmatrix} 1 & 2 & 1 & 2 \end{pmatrix}, \qquad nb = 2.$$

   and therefore

   $$WF(I) = nb = 2.$$

2. From the bin vector $bin = (1, 2, 1, 2)$, the packing patterns are

   $$P_1 = \{i \in \{1, 2, 3, 4\} : bin_i = 1\} = \{1, 3\},$$
   $$P_2 = \{i \in \{1, 2, 3, 4\} : bin_i = 2\} = \{2, 4\}.$$

3. Accordingly,

   $$z(I) \leq 2.$$

Moreover,

$$z(I) \geq \left\lceil \frac{\sum_{i \in \mathscr{I}} w_i}{c} \right\rceil = \left\lceil \frac{7+7+3+3}{10} \right\rceil = \left\lceil \frac{20}{10} \right\rceil = 2.$$

Hence $z(I) = 2$ and WF computes an **optimal** solution on this instance.

4. From Question 2, the packing patterns found by WF have total weights

$$\sum_{i \in P_1} w_i = 7 + 3 = 10, \qquad \sum_{i \in P_2} w_i = 7 + 3 = 10.$$

Since $c/2 = 5$, no packing pattern has total weight at most $c/2$; in particular *at most one* does, so the solution **satisfies** the half-full property.

Since $WF(I) = 2$ matches the lower bound $z(I) \geq 2$ found in Question 3, the solution found by WF is **optimal**.

Exercise 26.12 — Worst Fit 3

We consider a bin of capacity $c = 10$ and a set $\mathscr{I} = \{1, 2, 3, 4, 5\}$ of 5 items ($n = 5$), with the following item weights:

	$i=1$	$i=2$	$i=3$	$i=4$	$i=5$
w_i	4	7	6	3	5

Questions

1. Apply the Worst Fit (WF) algorithm to this instance. Show the execution step by step: for each item, trace the pseudocode variables *maxrc* and s over the scan of the open bins, and state into which bin the item is packed and the resulting residual capacity.

2. Construct the packing patterns $P_1, \ldots, P_{WF(I)}$ from the bin vector obtained in Question 1.

3. Compute a lower bound on $z(I)$ and determine whether WF finds an optimal solution on this instance.

4. Verify whether the packing patterns found by WF satisfy the half-full property (i.e., at most one packing pattern has total weight at most $c/2$), and determine whether the solution found is optimal for this BPP instance, either by comparing $WF(I)$ with a lower bound on $z(I)$, or by exhibiting a feasible solution of strictly smaller value.

Solution

1. Recall the WF pseudocode: for each item i, $maxrc := -1$, $s := -1$, and for $b := 1$ to nb, if $w[i] \leq rc[b]$ and $rc[b] > maxrc$ then $s \leftarrow b$, $maxrc \leftarrow rc[b]$.

(a) Item 1 (weight 4): bin 1 is open at the start ($nb = 1$, $rc[1] = c$) and item 1 fits into it, so $s = 1$ and WF packs item 1 into bin 1. Residual capacity of bin 1: $10 - 4 = 6$.

(b) Item 2 (weight 7): $maxrc := -1$, $s := -1$. For $b = 1$: $w[2] = 7 \leq rc[1] = 6$? No. No bin fits, so WF opens bin 2 and packs item 2 into it. Residual capacity of bin 2: $10 - 7 = 3$.

(c) Item 3 (weight 6): $maxrc := -1$, $s := -1$. For $b = 1$: $w[3] = 6 \leq rc[1] = 6$ and $rc[1] = 6 > -1$, so $s \leftarrow 1$, $maxrc \leftarrow 6$. For $b = 2$: $w[3] = 6 \leq rc[2] = 3$? No. Hence $s = 1$: item 3 is packed into bin 1. Residual capacity of bin 1: $6 - 6 = 0$.

(d) Item 4 (weight 3): $maxrc := -1$, $s := -1$. For $b = 1$: $w[4] = 3 \leq rc[1] = 0$? No. For $b = 2$: $w[4] = 3 \leq rc[2] = 3$ and $rc[2] = 3 > -1$, so $s \leftarrow 2$, $maxrc \leftarrow 3$. Hence $s = 2$: item 4 is packed into bin 2. Residual capacity of bin 2: $3 - 3 = 0$.

(e) Item 5 (weight 5): $maxrc := -1$, $s := -1$. For $b = 1$: $w[5] = 5 \leq rc[1] = 0$? No. For $b = 2$: $w[5] = 5 \leq rc[2] = 0$? No. No bin fits, so WF opens bin 3 and packs item 5 into it. Residual capacity of bin 3: $10 - 5 = 5$.

Hence WF returns:

$$bin = \begin{pmatrix} 1 & 2 & 1 & 2 & 3 \end{pmatrix}, \qquad nb = 3.$$

and therefore

$$WF(I) = nb = 3.$$

2. From the bin vector $bin = (1, 2, 1, 2, 3)$, the packing patterns are

$$P_1 = \{i : bin_i = 1\} = \{1, 3\},$$
$$P_2 = \{i : bin_i = 2\} = \{2, 4\},$$
$$P_3 = \{i : bin_i = 3\} = \{5\}.$$

3. Accordingly,

$$z(I) \leq 3.$$

Moreover,

$$z(I) \geq \left\lceil \frac{\sum_{i \in \mathscr{I}} w_i}{c} \right\rceil = \left\lceil \frac{4 + 7 + 6 + 3 + 5}{10} \right\rceil = \left\lceil \frac{25}{10} \right\rceil = 3.$$

Hence $z(I) = 3$ and WF computes an **optimal** solution on this instance.

4. From Question 2, the packing patterns found by WF have total weights

$$\sum_{i \in P_1} w_i = 4 + 6 = 10, \qquad \sum_{i \in P_2} w_i = 7 + 3 = 10, \qquad \sum_{i \in P_3} w_i = 5.$$

Since $c/2 = 5$, only P_3 has total weight at most $c/2$ (the other two exceed it); hence the solution **satisfies** the half-full property.

Since $WF(I) = 3$ matches the lower bound $z(I) \geq 3$ found in Question 3, the solution found by WF is **optimal**.

Chapter 27

Approximation algorithms for the knapsack problem

27.1 Greedy algorithm

Exercise 27.1 — Greedy 1

We consider a knapsack of capacity $c = 15$ and a set $\mathscr{I} = \{1, 2, 3, 4, 5\}$ of 5 items ($n = 5$), with the following profits and weights:

	$i = 1$	$i = 2$	$i = 3$	$i = 4$	$i = 5$
p_i	12	6	6	8	6
w_i	3	2	3	9	7

Questions

1. Verify that the items are sorted by non-increasing efficiency, and determine the split item s and the residual capacity $\bar{c}$.
2. Apply the greedy algorithm (GR) to this instance. Show the execution step by step, indicating at each iteration whether the item is selected and the resulting residual capacity. State the solution S returned by GR and $GR(I)$.
3. Compute the upper bound $U = \sum_{i=1}^{s-1} p_i + p_s \frac{\bar{c}}{w_s}$ on $z(I)$ given by the LP relaxation, and deduce an upper bound on the performance ratio $\frac{z(I)}{GR(I)}$.
4. Determine whether GR finds an optimal solution on this instance, either by showing that $GR(I)$ attains the upper bound $\lfloor U \rfloor$, or by exhibiting a feasible solution of larger profit.

Solution

1. The efficiencies of the items are

$$\frac{p_1}{w_1} = \frac{12}{3} = 4, \quad \frac{p_2}{w_2} = \frac{6}{2} = 3, \quad \frac{p_3}{w_3} = \frac{6}{3} = 2, \quad \frac{p_4}{w_4} = \frac{8}{9} \approx 0.89, \quad \frac{p_5}{w_5} = \frac{6}{7} \approx 0.86$$

   hence the items are sorted by non-increasing efficiency. Since

$$\sum_{i=1}^{3} w_i = 3 + 2 + 3 = 8 \leq 15 \qquad \text{and} \qquad \sum_{i=1}^{4} w_i = 3 + 2 + 3 + 9 = 17 > 15,$$

   the split item is $s = 4$ and the residual capacity is $\bar{c} = 15 - 8 = 7$.

2. Initially $rc := c = 15$. The execution of GR proceeds as follows:

   (a) Item 1 (weight $w_1 = 3$): $w_1 = 3 \leq rc = 15$, so the item is selected and $rc \leftarrow 15 - 3 = 12$.

   (b) Item 2 (weight $w_2 = 2$): $w_2 = 2 \leq rc = 12$, so the item is selected and $rc \leftarrow 12 - 2 = 10$.

   (c) Item 3 (weight $w_3 = 3$): $w_3 = 3 \leq rc = 10$, so the item is selected and $rc \leftarrow 10 - 3 = 7$.

   (d) Item 4 (weight $w_4 = 9$): $w_4 = 9 > rc = 7$, so the item is not selected.

   (e) Item 5 (weight $w_5 = 7$): $w_5 = 7 \leq rc = 7$, so the item is selected and $rc \leftarrow 7 - 7 = 0$.

   Hence GR returns the solution $S = \{1, 2, 3, 5\}$ with

$$GR(I) = 12 + 6 + 6 + 6 = 30.$$

   Note that GR selects all the items preceding the split item, does not select the split item 4, and also selects item 5, which follows it.

3. The LP relaxation gives

$$z(I) \leq U = 12 + 6 + 6 + 8 \cdot \frac{7}{9} = \frac{272}{9} \approx 30.22,$$

   and since $z(I)$ is integer, $z(I) \leq \lfloor U \rfloor = 30$. Hence

$$\frac{z(I)}{GR(I)} \leq \frac{30}{30} \approx 1.$$

4. Since $GR(I) = 30 = \lfloor U \rfloor$, no feasible solution has a larger profit: GR finds an **optimal** solution on this instance, and $z(I) = 30$.

Exercise 27.2 — Greedy 2

We consider a knapsack of capacity $c = 13$ and a set $\mathscr{I} = \{1, 2, 3, 4, 5, 6\}$ of 6 items ($n = 6$), with the following profits and weights:

	$i=1$	$i=2$	$i=3$	$i=4$	$i=5$	$i=6$
p_i	10	6	3	7	11	2
w_i	1	1	1	4	8	3

Questions

1. Verify that the items are sorted by non-increasing efficiency, and determine the split item s and the residual capacity $\bar{c}$.
2. Apply the greedy algorithm (GR) to this instance. Show the execution step by step, indicating at each iteration whether the item is selected and the resulting residual capacity. State the solution S returned by GR and $GR(I)$.
3. Compute the upper bound $U = \sum_{i=1}^{s-1} p_i + p_s \frac{\bar{c}}{w_s}$ on $z(I)$ given by the LP relaxation, and deduce an upper bound on the performance ratio $\frac{z(I)}{GR(I)}$.
4. Determine whether GR finds an optimal solution on this instance, either by showing that $GR(I)$ attains the upper bound $\lfloor U \rfloor$, or by exhibiting a feasible solution of larger profit.

Solution

1. The efficiencies of the items are

$$\frac{p_1}{w_1} = \frac{10}{1} = 10, \quad \frac{p_2}{w_2} = \frac{6}{1} = 6, \quad \frac{p_3}{w_3} = \frac{3}{1} = 3, \quad \frac{p_4}{w_4} = \frac{7}{4} = 1.75, \quad \frac{p_5}{w_5} = \frac{11}{8} \approx 1.38, \quad \frac{p_6}{w_6} = \frac{2}{3} \approx 0.67,$$

   hence the items are sorted by non-increasing efficiency. Since

$$\sum_{i=1}^{4} w_i = 1 + 1 + 1 + 4 = 7 \leq 13 \qquad \text{and} \qquad \sum_{i=1}^{5} w_i = 1 + 1 + 1 + 4 + 8 = 15 > 13,$$

   the split item is $s = 5$ and the residual capacity is $\bar{c} = 13 - 7 = 6$.

2. Initially $rc := c = 13$. The execution of GR proceeds as follows:

   (a) Item 1 (weight $w_1 = 1$): $w_1 = 1 \leq rc = 13$, so the item is selected and $rc \leftarrow 13 - 1 = 12$.

   (b) Item 2 (weight $w_2 = 1$): $w_2 = 1 \leq rc = 12$, so the item is selected and $rc \leftarrow 12 - 1 = 11$.

(c) Item 3 (weight $w_3 = 1$): $w_3 = 1 \leq rc = 11$, so the item is selected and $rc \leftarrow 11 - 1 = 10$.

(d) Item 4 (weight $w_4 = 4$): $w_4 = 4 \leq rc = 10$, so the item is selected and $rc \leftarrow 10 - 4 = 6$.

(e) Item 5 (weight $w_5 = 8$): $w_5 = 8 > rc = 6$, so the item is not selected.

(f) Item 6 (weight $w_6 = 3$): $w_6 = 3 \leq rc = 6$, so the item is selected and $rc \leftarrow 6 - 3 = 3$.

Hence GR returns the solution $S = \{1, 2, 3, 4, 6\}$ with

$$GR(I) = 10 + 6 + 3 + 7 + 2 = 28.$$

Note that GR selects all the items preceding the split item, does not select the split item 5, and also selects item 6, which follows it.

3. The LP relaxation gives

$$z(I) \leq U = 10 + 6 + 3 + 7 + 11 \cdot \frac{6}{8} = \frac{137}{4} \approx 34.25,$$

and since $z(I)$ is integer, $z(I) \leq \lfloor U \rfloor = 34$. Hence

$$\frac{z(I)}{GR(I)} \leq \frac{34}{28} \approx 1.21.$$

4. The solution $S^* = \{1, 2, 3, 5\}$, of weight $1 + 1 + 1 + 8 = 11 \leq 13$ and profit $z(I) = 10 + 6 + 3 + 11 = 30$ is feasible and has a larger profit than $GR(I) = 28$. Hence GR does **not** find an optimal solution on this instance; the solution S^* is optimal (this can be checked by enumerating the feasible solutions), and the performance ratio is

$$\frac{z(I)}{GR(I)} = \frac{30}{28} \approx 1.07.$$

Note that the item of maximum profit, item 5 with $p_5 = 11$, is not selected by GR: its low efficiency puts it late in the order, when the residual capacity is no longer sufficient.

Exercise 27.3 — Greedy 3

We consider a knapsack of capacity $c = 12$ and a set $\mathscr{I} = \{1, 2, 3, 4, 5\}$ of 5 items ($n = 5$), with the following profits and weights:

	$i = 1$	$i = 2$	$i = 3$	$i = 4$	$i = 5$
p_i	8	7	18	14	10
w_i	3	3	9	8	7

Questions

1. Verify that the items are sorted by non-increasing efficiency, and determine the split item s and the residual capacity $\bar{c}$.
2. Apply the greedy algorithm (GR) to this instance. Show the execution step by step, indicating at each iteration whether the item is selected and the resulting residual capacity. State the solution S returned by GR and $GR(I)$.
3. Compute the upper bound $U = \sum_{i=1}^{s-1} p_i + p_s \frac{\bar{c}}{w_s}$ on $z(I)$ given by the LP relaxation, and deduce an upper bound on the performance ratio $\frac{z(I)}{GR(I)}$.
4. Determine whether GR finds an optimal solution on this instance, either by showing that $GR(I)$ attains the upper bound $\lfloor U \rfloor$, or by exhibiting a feasible solution of larger profit.

Solution

1. The efficiencies of the items are

$$\frac{p_1}{w_1} = \frac{8}{3} \approx 2.67, \quad \frac{p_2}{w_2} = \frac{7}{3} \approx 2.33, \quad \frac{p_3}{w_3} = \frac{18}{9} = 2, \quad \frac{p_4}{w_4} = \frac{14}{8} = 1.75, \quad \frac{p_5}{w_5} = \frac{10}{7} \approx 1.43,$$

   hence the items are sorted by non-increasing efficiency. Since

$$\sum_{i=1}^{2} w_i = 3 + 3 = 6 \leq 12 \qquad \text{and} \qquad \sum_{i=1}^{3} w_i = 3 + 3 + 9 = 15 > 12,$$

   the split item is $s = 3$ and the residual capacity is $\bar{c} = 12 - 6 = 6$.

2. Initially $rc := c = 12$. The execution of GR proceeds as follows:

   (a) Item 1 (weight $w_1 = 3$): $w_1 = 3 \leq rc = 12$, so the item is selected and $rc \leftarrow 12 - 3 = 9$.

   (b) Item 2 (weight $w_2 = 3$): $w_2 = 3 \leq rc = 9$, so the item is selected and $rc \leftarrow 9 - 3 = 6$.

   (c) Item 3 (weight $w_3 = 9$): $w_3 = 9 > rc = 6$, so the item is not selected.

   (d) Item 4 (weight $w_4 = 8$): $w_4 = 8 > rc = 6$, so the item is not selected.

   (e) Item 5 (weight $w_5 = 7$): $w_5 = 7 > rc = 6$, so the item is not selected.

   Hence GR returns the solution $S = \{1, 2\}$ with

$$GR(I) = 8 + 7 = 15.$$

   Here GR selects exactly the items preceding the split item.

3. The LP relaxation gives

$$z(I) \leq U = 8 + 7 + 18 \cdot \frac{6}{9} = 27 \approx 27,$$

and since $z(I)$ is integer, $z(I) \leq \lfloor U \rfloor = 27$. Hence

$$\frac{z(I)}{GR(I)} \leq \frac{27}{15} \approx 1.8.$$

4. The solution $S^* = \{1, 3\}$, of weight $3 + 9 = 12 \leq 12$ and profit $z(I) = 8 + 18 = 26$ is feasible and has a larger profit than $GR(I) = 15$. Hence GR does **not** find an optimal solution on this instance; the solution S^* is optimal (this can be checked by enumerating the feasible solutions), and the performance ratio is

$$\frac{z(I)}{GR(I)} = \frac{26}{15} \approx 1.73.$$

Note that the item of maximum profit, item 3 with $p_3 = 18$, is not selected by GR: its low efficiency puts it late in the order, when the residual capacity is no longer sufficient.

27.2 Extended greedy algorithm

Exercise 27.4 — Ext-Greedy 1

We consider a knapsack of capacity $c = 22$ and a set $\mathscr{I} = \{1, 2, 3, 4, 5\}$ of 5 items ($n = 5$), with the following profits and weights:

	$i=1$	$i=2$	$i=3$	$i=4$	$i=5$
p_i	11	5	3	4	3
w_i	8	4	3	9	8

Questions

1. Apply the greedy algorithm (GR) to this instance step by step, and state $GR(I)$.
2. Determine the maximum profit $\hat{p}$ and the best single-item feasible solution $\hat{S}$. State the solution returned by the Ext-Greedy algorithm (GR^e) and $GR^e(I)$.
3. Determine the split item s and verify the two inequalities $\sum_{i=1}^{s-1} p_i \leq GR^e(I)$ and $p_s \leq GR^e(I)$ used in the proof of the performance guarantee of GR^e. Deduce an upper bound on $z(I)$.
4. Determine whether GR^e finds an optimal solution on this instance, and compute the performance ratio $\frac{z(I)}{GR^e(I)}$, comparing it with the relative performance guarantee 2 of GR^e.

Solution

1. The efficiencies of the items are

$$\frac{p_1}{w_1} = \frac{11}{8} \approx 1.38, \quad \frac{p_2}{w_2} = \frac{5}{4} = 1.25, \quad \frac{p_3}{w_3} = \frac{3}{3} = 1, \quad \frac{p_4}{w_4} = \frac{4}{9} \approx 0.44, \quad \frac{p_5}{w_5} = \frac{3}{8} \approx 0.38$$

   so the items are sorted by non-increasing efficiency. Initially $rc := c = 22$, and GR proceeds as follows:

   (a) Item 1 (weight $w_1 = 8$): $w_1 = 8 \leq rc = 22$, so the item is selected and $rc \leftarrow 22 - 8 = 14$.

   (b) Item 2 (weight $w_2 = 4$): $w_2 = 4 \leq rc = 14$, so the item is selected and $rc \leftarrow 14 - 4 = 10$.

   (c) Item 3 (weight $w_3 = 3$): $w_3 = 3 \leq rc = 10$, so the item is selected and $rc \leftarrow 10 - 3 = 7$.

   (d) Item 4 (weight $w_4 = 9$): $w_4 = 9 > rc = 7$, so the item is not selected.

   (e) Item 5 (weight $w_5 = 8$): $w_5 = 8 > rc = 7$, so the item is not selected.

   Hence GR returns $S = \{1, 2, 3\}$ and $GR(I) = 11 + 5 + 3 = 19$.

2. The maximum profit is $\hat{p} = p_1 = 11$, and item 1 fits into the knapsack ($w_1 = 8 \leq 22$), so $\hat{S} = \{1\}$. Since $GR(I) = 19 \geq \hat{p} = 11$, GR^e keeps the greedy solution:

$$GR^e(I) = GR(I) = 19.$$

3. Since $\sum_{i=1}^{3} w_i = 15 \leq 22$ and $\sum_{i=1}^{4} w_i = 24 > 22$, the split item is $s = 4$. Then

$$\sum_{i=1}^{s-1} p_i = 11 + 5 + 3 = 19 \leq GR^e(I) = 19 \qquad \text{and} \qquad p_s = p_4 = 4 \leq GR^e(I) = 19,$$

and, as in the proof of the performance guarantee,

$$z(I) \;\leq\; \sum_{i=1}^{s-1} p_i + p_s \;=\; 23 \;\leq\; 2\,GR^e(I) = 38.$$

4. The solution $S^* = \{1, 2, 4\}$, of weight $8+4+9 = 21 \leq 22$ and profit $z(I) = 11+5+4 = 20$ is feasible and better than the one returned by GR^e, so GR^e does **not** find an optimal solution; S^* is optimal (this can be checked by enumerating the feasible solutions). The performance ratio is

$$\frac{z(I)}{GR^e(I)} = \frac{20}{19} \approx 1.05 \;\leq\; 2,$$

as guaranteed.

Exercise 27.5 — Ext-Greedy 2

We consider a knapsack of capacity $c = 8$ and a set $\mathscr{I} = \{1, 2, 3, 4, 5\}$ of 5 items ($n = 5$), with the following profits and weights:

	$i=1$	$i=2$	$i=3$	$i=4$	$i=5$
p_i	16	19	18	14	8
w_i	5	8	8	7	6

Questions

1. Apply the greedy algorithm (GR) to this instance step by step, and state $GR(I)$.
2. Determine the maximum profit $\hat{p}$ and the best single-item feasible solution $\hat{S}$. State the solution returned by the Ext-Greedy algorithm (GR^e) and $GR^e(I)$.
3. Determine the split item s and verify the two inequalities $\sum_{i=1}^{s-1} p_i \leq GR^e(I)$ and $p_s \leq GR^e(I)$ used in the proof of the performance guarantee of GR^e. Deduce an upper bound on $z(I)$.

4. Determine whether GR^e finds an optimal solution on this instance, and compute the performance ratio $\frac{z(I)}{GR^e(I)}$, comparing it with the relative performance guarantee 2 of GR^e.

Solution

1. The efficiencies of the items are

$$\frac{p_1}{w_1} = \frac{16}{5} = 3.2, \quad \frac{p_2}{w_2} = \frac{19}{8} \approx 2.38, \quad \frac{p_3}{w_3} = \frac{18}{8} = 2.25, \quad \frac{p_4}{w_4} = \frac{14}{7} = 2, \quad \frac{p_5}{w_5} = \frac{8}{6} \approx 1.33$$

   so the items are sorted by non-increasing efficiency. Initially $rc := c = 8$, and GR proceeds as follows:

   (a) Item 1 (weight $w_1 = 5$): $w_1 = 5 \leq rc = 8$, so the item is selected and $rc \leftarrow 8 - 5 = 3$.

   (b) Item 2 (weight $w_2 = 8$): $w_2 = 8 > rc = 3$, so the item is not selected.

   (c) Item 3 (weight $w_3 = 8$): $w_3 = 8 > rc = 3$, so the item is not selected.

   (d) Item 4 (weight $w_4 = 7$): $w_4 = 7 > rc = 3$, so the item is not selected.

   (e) Item 5 (weight $w_5 = 6$): $w_5 = 6 > rc = 3$, so the item is not selected.

   Hence GR returns $S = \{1\}$ and $GR(I) = 16 = 16$.

2. The maximum profit is $\hat{p} = p_2 = 19$, and item 2 fits into the knapsack ($w_2 = 8 \leq 8$), so $\hat{S} = \{2\}$. Since $GR(I) = 16 < \hat{p} = 19$, GR^e replaces the greedy solution with $\hat{S}$:

$$GR^e(I) = \hat{p} = 19.$$

3. Since $\sum_{i=1}^{1} w_i = 5 \leq 8$ and $\sum_{i=1}^{2} w_i = 13 > 8$, the split item is $s = 2$. Then

$$\sum_{i=1}^{s-1} p_i = 16 = 16 \leq GR^e(I) = 19 \qquad \text{and} \qquad p_s = p_2 = 19 \leq GR^e(I) = 19,$$

   and, as in the proof of the performance guarantee,

$$z(I) \;\leq\; \sum_{i=1}^{s-1} p_i + p_s \;=\; 35 \;\leq\; 2\,GR^e(I) = 38.$$

4. The solution returned by GR^e is **optimal**: $S^* = \{2\}$, of weight $8 = 8 \leq 8$ and profit $z(I) = 19 = 19$. The performance ratio is $\frac{z(I)}{GR^e(I)} = 1$.

Exercise 27.6 — Ext-Greedy 3

We consider a knapsack of capacity $c = 16$ and a set $\mathscr{I} = \{1, 2, 3\}$ of 3 items ($n = 3$), with

the following profits and weights:

	$i=1$	$i=2$	$i=3$
p_i	2	8	8
w_i	1	8	8

Questions

1. Apply the greedy algorithm (GR) to this instance step by step, and state $GR(I)$.
2. Determine the maximum profit $\hat{p}$ and the best single-item feasible solution $\hat{S}$. State the solution returned by the Ext-Greedy algorithm (GR^e) and $GR^e(I)$.
3. Determine the split item s and verify the two inequalities $\sum_{i=1}^{s-1} p_i \leq GR^e(I)$ and $p_s \leq GR^e(I)$ used in the proof of the performance guarantee of GR^e. Deduce an upper bound on $z(I)$.
4. Determine whether GR^e finds an optimal solution on this instance, and compute the performance ratio $\frac{z(I)}{GR^e(I)}$, comparing it with the relative performance guarantee 2 of GR^e.

Solution

1. The efficiencies of the items are

$$\frac{p_1}{w_1} = \frac{2}{1} = 2, \quad \frac{p_2}{w_2} = \frac{8}{8} = 1, \quad \frac{p_3}{w_3} = \frac{8}{8} = 1,$$

   so the items are sorted by non-increasing efficiency. Initially $rc := c = 16$, and GR proceeds as follows:

   (a) Item 1 (weight $w_1 = 1$): $w_1 = 1 \leq rc = 16$, so the item is selected and $rc \leftarrow 16 - 1 = 15$.

   (b) Item 2 (weight $w_2 = 8$): $w_2 = 8 \leq rc = 15$, so the item is selected and $rc \leftarrow 15 - 8 = 7$.

   (c) Item 3 (weight $w_3 = 8$): $w_3 = 8 > rc = 7$, so the item is not selected.

   Hence GR returns $S = \{1, 2\}$ and $GR(I) = 2 + 8 = 10$.

2. The maximum profit is $\hat{p} = p_2 = 8$, and item 2 fits into the knapsack ($w_2 = 8 \leq 16$), so $\hat{S} = \{2\}$. Since $GR(I) = 10 \geq \hat{p} = 8$, GR^e keeps the greedy solution:

$$GR^e(I) = GR(I) = 10.$$

3. Since $\sum_{i=1}^{2} w_i = 9 \leq 16$ and $\sum_{i=1}^{3} w_i = 17 > 16$, the split item is $s = 3$. Then

$$\sum_{i=1}^{s-1} p_i = 2 + 8 = 10 \leq GR^e(I) = 10 \qquad \text{and} \qquad p_s = p_3 = 8 \leq GR^e(I) = 10,$$

and, as in the proof of the performance guarantee,

$$z(I) \;\leq\; \sum_{i=1}^{s-1} p_i + p_s \;=\; 18 \;\leq\; 2\,GR^e(I) = 20.$$

4. The solution $S^* = \{2, 3\}$, of weight $8 + 8 = 16 \leq 16$ and profit $z(I) = 8 + 8 = 16$ is feasible and better than the one returned by GR^e, so GR^e does **not** find an optimal solution; S^* is optimal (this can be checked by enumerating the feasible solutions). The performance ratio is
$$\frac{z(I)}{GR^e(I)} = \frac{16}{10} \approx 1.6 \;\leq\; 2,$$
as guaranteed.

Remark. This instance belongs to the family of instances used to show that the relative performance guarantee 2 of GR^e is tight, with $\vartheta = 8$: there the ratio is $\frac{2\vartheta}{\vartheta+2}$, and here $\frac{16}{10} = 1.6$; it tends to 2 as ϑ grows.

27.3 The H^2 heuristic

Exercise 27.7 — H^2 1

We consider a knapsack of capacity $c = 11$ and a set $\mathscr{I} = \{1, 2, 3, 4, 5\}$ of 5 items ($n = 5$), with the following profits and weights:

	$i=1$	$i=2$	$i=3$	$i=4$	$i=5$
p_i	8	9	2	3	2
w_i	5	8	3	7	9

Questions

1. Verify that the items are sorted by non-increasing efficiency, and determine the set $\mathscr{P}$ of the feasible pairs of items.
2. Apply the H^2 algorithm to this instance: for each feasible pair (i, k) compute the residual capacity $\bar{c}$, the residual set $\mathscr{I}(i, k)$, the value of GR^e on the residual instance and the value $lb(i, k)$.
3. State the solution returned by H^2 and $H^2(I)$.
4. Compare $H^2(I)$ with $GR^e(I)$ and with $z(I)$, and compute the performance ratio $\frac{z(I)}{H^2(I)}$, comparing it with the relative performance guarantee $\frac{4}{3}$ of H^2.

Solution

1. The efficiencies of the items are

$$\frac{p_1}{w_1} = \frac{8}{5} = 1.6, \quad \frac{p_2}{w_2} = \frac{9}{8} \approx 1.12, \quad \frac{p_3}{w_3} = \frac{2}{3} \approx 0.67, \quad \frac{p_4}{w_4} = \frac{3}{7} \approx 0.43, \quad \frac{p_5}{w_5} = \frac{2}{9} \approx 0.22,$$

   so the items are sorted by non-increasing efficiency. A pair (i, k) with $i < k$ is feasible if $w_i + w_k \leq c = 11$, hence

$$\mathscr{P} = \{(1,3), (2,3), (3,4)\}.$$

2. The algorithm starts from the best single item, $lb = p_{\max} = p_2 = 9$. For each feasible pair, the residual set keeps the items $j \notin \{i, k\}$ with $p_j \leq \min\{p_i, p_k\}$ and $w_j \leq \bar{c}$:

pair	weight	$\bar{c}$	residual set	GR^e on the residual	value
$(1,2)$	$13 > 11$			not feasible	
$(1,3)$	8	3	$\varnothing$	0	10
$(1,4)$	$12 > 11$			not feasible	
$(1,5)$	$14 > 11$			not feasible	
$(2,3)$	11	0	$\varnothing$	0	11
$(2,4)$	$15 > 11$			not feasible	
$(2,5)$	$17 > 11$			not feasible	
$(3,4)$	10	1	$\varnothing$	0	5
$(3,5)$	$12 > 11$			not feasible	
$(4,5)$	$16 > 11$			not feasible	

3. The largest value is 11, obtained with the solution $\{2, 3\}$. Hence

$$H^2(I) = 11.$$

4. On this instance $GR^e(I) = 10$, so H^2 improves on GR^e. The solution returned by H^2 is **optimal**: $S^* = \{2, 3\}$, of weight $8 + 3 = 11 \leq 11$ and profit $z(I) = 9 + 2 = 11$, and the performance ratio is 1.

Exercise 27.8 — H^2 2

We consider a knapsack of capacity $c = 15$ and a set $\mathscr{I} = \{1, 2, 3, 4, 5\}$ of 5 items ($n = 5$), with the following profits and weights:

	$i = 1$	$i = 2$	$i = 3$	$i = 4$	$i = 5$
p_i	14	9	5	10	15
w_i	1	1	2	5	8

Questions

1. Verify that the items are sorted by non-increasing efficiency, and determine the set $\mathscr{P}$ of the feasible pairs of items.
2. Apply the H^2 algorithm to this instance: for each feasible pair (i,k) compute the residual capacity $\bar{c}$, the residual set $\mathscr{I}(i,k)$, the value of GR^e on the residual instance and the value $lb(i,k)$.
3. State the solution returned by H^2 and $H^2(I)$.
4. Compare $H^2(I)$ with $GR^e(I)$ and with $z(I)$, and compute the performance ratio $\frac{z(I)}{H^2(I)}$, comparing it with the relative performance guarantee $\frac{4}{3}$ of H^2.

Solution

1. The efficiencies of the items are

$$\frac{p_1}{w_1} = \frac{14}{1} = 14, \quad \frac{p_2}{w_2} = \frac{9}{1} = 9, \quad \frac{p_3}{w_3} = \frac{5}{2} = 2.5, \quad \frac{p_4}{w_4} = \frac{10}{5} = 2, \quad \frac{p_5}{w_5} = \frac{15}{8} \approx 1.88$$

   so the items are sorted by non-increasing efficiency. A pair (i,k) with $i<k$ is feasible if $w_i + w_k \le c = 15$, hence

$$\mathscr{P} = \{(1,2),(1,3),(1,4),(1,5),(2,3),(2,4),(2,5),(3,4),(3,5),(4,5)\}.$$

2. The algorithm starts from the best single item, $lb = p_{\max} = p_5 = 15$. For each feasible pair, the residual set keeps the items $j \notin \{i,k\}$ with $p_j \le \min\{p_i, p_k\}$ and $w_j \le \bar{c}$:

pair	weight	$\bar{c}$	residual set	GR^e on the residual	value
$(1,2)$	2	13	$\{3\}$	5	28
$(1,3)$	3	12	$\varnothing$	0	19
$(1,4)$	6	9	$\{2,3\}$	14	38
$(1,5)$	9	6	$\{2,3,4\}$	14	43
$(2,3)$	3	12	$\varnothing$	0	14
$(2,4)$	6	9	$\{3\}$	5	24
$(2,5)$	9	6	$\{3\}$	5	29
$(3,4)$	7	8	$\varnothing$	0	15
$(3,5)$	10	5	$\varnothing$	0	20
$(4,5)$	13	2	$\{2,3\}$	9	34

3. The largest value is 43, obtained with the solution $\{1,2,3,5\}$. Hence

$$H^2(I) = 43.$$

4. On this instance $GR^e(I) = 38$, so H^2 improves on GR^e. The solution $S^* = \{1,2,4,5\}$, of weight $1+1+5+8 = 15 \le 15$ and profit $z(I) = 14+9+10+15 = 48$ is better, so H^2

does **not** find an optimal solution; S^* is optimal (this can be checked by enumerating the feasible solutions). The performance ratio is

$$\frac{z(I)}{H^2(I)} = \frac{48}{43} \approx 1.12 \;\leq\; \frac{4}{3} \approx 1.33,$$

as guaranteed.

Exercise 27.9 — H^2 3

We consider a knapsack of capacity $c = 10$ and a set $\mathscr{I} = \{1, 2, 3, 4, 5\}$ of 5 items ($n = 5$), with the following profits and weights:

	$i=1$	$i=2$	$i=3$	$i=4$	$i=5$
p_i	13	5	8	5	7
w_i	1	1	3	2	5

Questions

1. Verify that the items are sorted by non-increasing efficiency, and determine the set $\mathscr{P}$ of the feasible pairs of items.
2. Apply the H^2 algorithm to this instance: for each feasible pair (i, k) compute the residual capacity $\bar{c}$, the residual set $\mathscr{I}(i, k)$, the value of GR^e on the residual instance and the value $lb(i, k)$.
3. State the solution returned by H^2 and $H^2(I)$.
4. Compare $H^2(I)$ with $GR^e(I)$ and with $z(I)$, and compute the performance ratio $\frac{z(I)}{H^2(I)}$, comparing it with the relative performance guarantee $\frac{4}{3}$ of H^2.

Solution

1. The efficiencies of the items are

$$\frac{p_1}{w_1} = \frac{13}{1} = 13, \quad \frac{p_2}{w_2} = \frac{5}{1} = 5, \quad \frac{p_3}{w_3} = \frac{8}{3} \approx 2.67, \quad \frac{p_4}{w_4} = \frac{5}{2} = 2.5, \quad \frac{p_5}{w_5} = \frac{7}{5} = 1.4,$$

   so the items are sorted by non-increasing efficiency. A pair (i, k) with $i < k$ is feasible if $w_i + w_k \leq c = 10$, hence

$$\mathscr{P} = \big\{(1,2), (1,3), (1,4), (1,5), (2,3), (2,4), (2,5), (3,4), (3,5), (4,5)\big\}.$$

2. The algorithm starts from the best single item, $lb = p_{\max} = p_1 = 13$. For each feasible pair, the residual set keeps the items $j \notin \{i, k\}$ with $p_j \leq \min\{p_i, p_k\}$ and $w_j \leq \bar{c}$:

pair	weight	$\bar{c}$	residual set	GR^e on the residual	value
$(1,2)$	2	8	$\{4\}$	5	23
$(1,3)$	4	6	$\{2,4,5\}$	10	31
$(1,4)$	3	7	$\{2\}$	5	23
$(1,5)$	6	4	$\{2,4\}$	10	30
$(2,3)$	4	6	$\{4\}$	5	18
$(2,4)$	3	7	$\varnothing$	0	10
$(2,5)$	6	4	$\{4\}$	5	17
$(3,4)$	5	5	$\{2\}$	5	18
$(3,5)$	8	2	$\{2,4\}$	5	20
$(4,5)$	7	3	$\{2\}$	5	17

3. The largest value is 31, obtained with the solution $\{1,2,3,4\}$. Hence

$$H^2(I) = 31.$$

4. On this instance $GR^e(I) = 31$, the same value as H^2. The solution $S^* = \{1,2,3,5\}$, of weight $1+1+3+5 = 10 \leq 10$ and profit $z(I) = 13+5+8+7 = 33$ is better, so H^2 does **not** find an optimal solution; S^* is optimal (this can be checked by enumerating the feasible solutions). The performance ratio is

$$\frac{z(I)}{H^2(I)} = \frac{33}{31} \approx 1.06 \;\leq\; \frac{4}{3} \approx 1.33,$$

as guaranteed.

27.4 The H^ℓ heuristic

Exercise 27.10 — H^ℓ 1

We consider a knapsack of capacity $c = 16$ and a set $\mathscr{I} = \{1,2,3,4,5\}$ of 5 items ($n = 5$), with the following profits and weights:

	$i=1$	$i=2$	$i=3$	$i=4$	$i=5$
p_i	14	13	13	10	11
w_i	1	2	3	3	9

Questions

1. We apply the H^ℓ algorithm with $\ell = 3$. Compute the best feasible solution made of at most $\ell - 1 = 2$ items.

2. For each feasible set L of exactly $\ell = 3$ items, compute the residual capacity $\bar{c}$, the residual set $\mathscr{I}(L)$, the value of GR^e on the residual instance and the value $lb(L)$.

3. State the solution returned by H^3 and $H^3(I)$.

4. Determine whether H^3 finds an optimal solution on this instance, and compute the performance ratio $\frac{z(I)}{H^3(I)}$, comparing it with the relative performance guarantee $\frac{\ell+2}{\ell+1} = \frac{5}{4}$.

Solution

1. The best feasible solution with at most 2 items is $\{1, 2\}$, of weight $1+2 = 3 \leq 16$ and profit $14 + 13 = 27$.

2. For each set L of 3 items, the residual set keeps the items $j \notin L$ with $p_j \leq \min\{p_i : i \in L\}$ and $w_j \leq \bar{c}$:

set L	weight	$\bar{c}$	residual set	GR^e on the residual	value
$\{1,2,3\}$	6	10	$\{4,5\}$	11	51
$\{1,2,4\}$	6	10	$\varnothing$	0	37
$\{1,2,5\}$	12	4	$\{4\}$	10	48
$\{1,3,4\}$	7	9	$\varnothing$	0	37
$\{1,3,5\}$	13	3	$\{4\}$	10	48
$\{1,4,5\}$	13	3	$\varnothing$	0	35
$\{2,3,4\}$	8	8	$\varnothing$	0	36
$\{2,3,5\}$	14	2	$\varnothing$	0	37
$\{2,4,5\}$	14	2	$\varnothing$	0	34
$\{3,4,5\}$	15	1	$\varnothing$	0	34

3. The largest value is 51, obtained with the solution $\{1, 2, 3, 5\}$. Hence

$$H^3(I) = 51.$$

4. The solution returned by H^3 is **optimal**: $S^* = \{1, 2, 3, 5\}$, of weight $1 + 2 + 3 + 9 = 15 \leq 16$ and profit $z(I) = 14 + 13 + 13 + 11 = 51$, and the performance ratio is 1. For comparison, on this instance $H^2(I) = 50$: guessing three items instead of two is enough to find the optimal solution.

Exercise 27.11 — H^ℓ 2

We consider a knapsack of capacity $c = 20$ and a set $\mathscr{I} = \{1, 2, 3, 4, 5, 6\}$ of 6 items ($n = 6$), with the following profits and weights:

	$i=1$	$i=2$	$i=3$	$i=4$	$i=5$	$i=6$
p_i	2	4	4	4	4	4
w_i	1	4	4	4	4	4

Questions

1. We apply the H^ℓ algorithm with $\ell = 3$. Compute the best feasible solution made of at most $\ell - 1 = 2$ items.
2. For each feasible set L of exactly $\ell = 3$ items, compute the residual capacity $\bar{c}$, the residual set $\mathscr{I}(L)$, the value of GR^e on the residual instance and the value $lb(L)$.
3. State the solution returned by H^3 and $H^3(I)$.
4. Determine whether H^3 finds an optimal solution on this instance, and compute the performance ratio $\frac{z(I)}{H^3(I)}$, comparing it with the relative performance guarantee $\frac{\ell+2}{\ell+1} = \frac{5}{4}$.

Solution

1. The best feasible solution with at most 2 items is $\{2, 3\}$, of weight $4+4 = 8 \leq 20$ and profit $4 + 4 = 8$.
2. For each set L of 3 items, the residual set keeps the items $j \notin L$ with $p_j \leq \min\{p_i : i \in L\}$ and $w_j \leq \bar{c}$:

set L	weight	$\bar{c}$	residual set	GR^e on the residual	value
$\{1,2,3\}$	9	11	$\varnothing$	0	10
$\{1,2,4\}$	9	11	$\varnothing$	0	10
$\{1,2,5\}$	9	11	$\varnothing$	0	10
$\{1,2,6\}$	9	11	$\varnothing$	0	10
$\{1,3,4\}$	9	11	$\varnothing$	0	10
$\{1,3,5\}$	9	11	$\varnothing$	0	10
$\{1,3,6\}$	9	11	$\varnothing$	0	10
$\{1,4,5\}$	9	11	$\varnothing$	0	10
$\{1,4,6\}$	9	11	$\varnothing$	0	10
$\{1,5,6\}$	9	11	$\varnothing$	0	10
$\{2,3,4\}$	12	8	$\{1,5,6\}$	6	18
$\{2,3,5\}$	12	8	$\{1,4,6\}$	6	18
$\{2,3,6\}$	12	8	$\{1,4,5\}$	6	18
$\{2,4,5\}$	12	8	$\{1,3,6\}$	6	18
$\{2,4,6\}$	12	8	$\{1,3,5\}$	6	18
$\{2,5,6\}$	12	8	$\{1,3,4\}$	6	18
$\{3,4,5\}$	12	8	$\{1,2,6\}$	6	18
$\{3,4,6\}$	12	8	$\{1,2,5\}$	6	18
$\{3,5,6\}$	12	8	$\{1,2,4\}$	6	18
$\{4,5,6\}$	12	8	$\{1,2,3\}$	6	18

3. The largest value is 18, obtained with the solution $\{1, 2, 3, 4, 5\}$. Hence

$$H^3(I) = 18.$$

4. The solution $S^* = \{2, 3, 4, 5, 6\}$, of weight $4 + 4 + 4 + 4 + 4 = 20 \leq 20$ and profit $z(I) = 4+4+4+4+4 = 20$ is better, so H^3 does **not** find an optimal solution; S^* is optimal (this can be checked by enumerating the feasible solutions). The performance ratio is
$$\frac{z(I)}{H^3(I)} = \frac{20}{18} \approx 1.11 \leq \frac{5}{4} = 1.25,$$
as guaranteed. For comparison, on this instance $H^2(I) = 18$.

Remark. This instance belongs to the family of instances used to show that the relative performance guarantee $\frac{\ell+2}{\ell+1}$ of H^ℓ is tight, with $\ell = 3$ and $M = 4$: there $H^\ell(I_M) = (\ell + 1)\, M + 2$ and $z(I_M) = (\ell + 2)\, M$, here 18 and 20.

Exercise 27.12 — H^ℓ 3

We consider a knapsack of capacity $c = 18$ and a set $\mathscr{I} = \{1, 2, 3, 4, 5, 6\}$ of 6 items ($n = 6$), with the following profits and weights:

	$i=1$	$i=2$	$i=3$	$i=4$	$i=5$	$i=6$
p_i	15	15	14	8	12	13
w_i	1	2	3	2	6	9

Questions

1. We apply the H^ℓ algorithm with $\ell = 3$. Compute the best feasible solution made of at most $\ell - 1 = 2$ items.
2. For each feasible set L of exactly $\ell = 3$ items, compute the residual capacity $\bar{c}$, the residual set $\mathscr{I}(L)$, the value of GR^e on the residual instance and the value $lb(L)$.
3. State the solution returned by H^3 and $H^3(I)$.
4. Determine whether H^3 finds an optimal solution on this instance, and compute the performance ratio $\frac{z(I)}{H^3(I)}$, comparing it with the relative performance guarantee $\frac{\ell+2}{\ell+1} = \frac{5}{4}$.

Solution

1. The best feasible solution with at most 2 items is $\{1, 2\}$, of weight $1 + 2 = 3 \leq 18$ and profit $15 + 15 = 30$.
2. For each set L of 3 items, the residual set keeps the items $j \notin L$ with $p_j \leq \min\{p_i : i \in L\}$ and $w_j \leq \bar{c}$:

set L	weight	$\bar{c}$	residual set	GR^e on the residual	value
$\{1,2,3\}$	6	12	$\{4,5,6\}$	20	64
$\{1,2,4\}$	5	13	$\varnothing$	0	38
$\{1,2,5\}$	9	9	$\{4\}$	8	50
$\{1,2,6\}$	12	6	$\{4,5\}$	12	55
$\{1,3,4\}$	6	12	$\varnothing$	0	37
$\{1,3,5\}$	10	8	$\{4\}$	8	49
$\{1,3,6\}$	13	5	$\{4\}$	8	50
$\{1,4,5\}$	9	9	$\varnothing$	0	35
$\{1,4,6\}$	12	6	$\varnothing$	0	36
$\{1,5,6\}$	16	2	$\{4\}$	8	48
$\{2,3,4\}$	7	11	$\varnothing$	0	37
$\{2,3,5\}$	11	7	$\{4\}$	8	49
$\{2,3,6\}$	14	4	$\{4\}$	8	50
$\{2,4,5\}$	10	8	$\varnothing$	0	35
$\{2,4,6\}$	13	5	$\varnothing$	0	36
$\{2,5,6\}$	17	1	$\varnothing$	0	40
$\{3,4,5\}$	11	7	$\varnothing$	0	34
$\{3,4,6\}$	14	4	$\varnothing$	0	35
$\{3,5,6\}$	18	0	$\varnothing$	0	39
$\{4,5,6\}$	17	1	$\varnothing$	0	33

3. The largest value is 64, obtained with the solution $\{1,2,3,4,5\}$. Hence

$$H^3(I) = 64.$$

4. The solution $S^* = \{1,2,3,4,6\}$, of weight $1+2+3+2+9 = 17 \leq 18$ and profit $z(I) = 15{+}15{+}14{+}8{+}13 = 65$ is better, so H^3 does **not** find an optimal solution; S^* is optimal (this can be checked by enumerating the feasible solutions). The performance ratio is

$$\frac{z(I)}{H^3(I)} = \frac{65}{64} \approx 1.02 \leq \frac{5}{4} = 1.25,$$

as guaranteed. For comparison, on this instance $H^2(I) = 64$.

27.5 The FPTAS

Exercise 27.13 — Basic-FPTAS 1

We consider a knapsack of capacity $c = 13$ and a set $\mathscr{I} = \{1, 2, 3, 4\}$ of 4 items ($n = 4$), with the following profits and weights:

	$i = 1$	$i = 2$	$i = 3$	$i = 4$
p_i	24	11	23	5
w_i	2	3	9	3

We apply the Basic-FPTAS algorithm with accuracy $\varepsilon = \frac{1}{2}$.

Questions

1. Compute the lower bound $GR^e(I)$ used by the Basic-FPTAS algorithm.
2. Compute the scaling factor and the scaled profits
$$K = \frac{\varepsilon\, GR^e(I)}{n} \qquad \text{and} \qquad p'_j = \left\lfloor \frac{p_j}{K} \right\rfloor \quad \text{for all } j \in \mathscr{I}.$$
3. Solve the scaled instance by dynamic programming by profits, i.e., compute the minimum weights $u(i, q)$ and find the largest profit q such that $u(4, q) \leq c$.
4. State the solution returned by FP^ε, its profit $FP^\varepsilon(I)$ in the original instance, and compare it with $GR^e(I)$ and $z(I)$. Compute the performance ratio and compare it with the guarantee $\frac{1}{1-\varepsilon}$.

Solution

1. The items are sorted by non-increasing efficiency. GR selects the items $\{1, 2, 4\}$, with $GR(I) = 24 + 11 + 5 = 40$; the maximum profit is $\hat{p} = p_1 = 24$. Hence
$$GR^e(I) = 40.$$
2. The scaling factor is
$$K = \frac{\varepsilon\, GR^e(I)}{n} = \frac{\frac{1}{2} \cdot 40}{4} = 5,$$
and the scaled profits are
$$p'_1 = \left\lfloor \frac{24}{5} \right\rfloor = 4, \quad p'_2 = \left\lfloor \frac{11}{5} \right\rfloor = 2, \quad p'_3 = \left\lfloor \frac{23}{5} \right\rfloor = 4, \quad p'_4 = \left\lfloor \frac{5}{5} \right\rfloor = 1.$$
The total scaled profit is $P' = 4 + 2 + 4 + 1 = 11$.
3. The minimum weights $u(i, q)$ of a subset of $\{1, \ldots, i\}$ with scaled profit exactly q are (only the values of q that can be attained are reported; ∞ means that q cannot be attained with the first i items):

q	0	1	2	3	4	5	6	7	8	9	10
$u(0,q)$	0	∞	∞	∞	∞	∞	∞	∞	∞	∞	∞
$u(1,q)$	0	∞	∞	∞	2	∞	∞	∞	∞	∞	∞
$u(2,q)$	0	∞	3	∞	2	∞	5	∞	∞	∞	∞
$u(3,q)$	0	∞	3	∞	2	∞	5	∞	11	∞	14
$u(4,q)$	0	3	3	6	2	5	5	8	11	14	14

q	11
$u(0,q)$	∞
$u(1,q)$	∞
$u(2,q)$	∞
$u(3,q)$	∞
$u(4,q)$	17

The largest q with $u(4,q) \leq c = 13$ is $q = 8$, of weight $u(4,8) = 11$, attained by the items $\{1,3\}$.

4. FP^{ε} returns $S = \{1,3\}$, of profit

$$FP^{\varepsilon}(I) = 24 + 23 = 47$$

in the original instance, better than $GR^{e}(I) = 40$. The solution is **optimal**: $S^* = \{1,3\}$, of weight $2+9 = 11 \leq 13$ and profit $z(I) = 24+23 = 47$, and the performance ratio is 1.

Exercise 27.14 — Basic-FPTAS 2

We consider a knapsack of capacity $c = 16$ and a set $\mathscr{I} = \{1,2,3,4\}$ of 4 items ($n = 4$), with the following profits and weights:

	$i=1$	$i=2$	$i=3$	$i=4$
p_i	14	12	14	18
w_i	3	4	5	7

We apply the Basic-FPTAS algorithm with accuracy $\varepsilon = \frac{1}{2}$.

Questions

1. Compute the lower bound $GR^{e}(I)$ used by the Basic-FPTAS algorithm.
2. Compute the scaling factor and the scaled profits

$$K = \frac{\varepsilon\, GR^{e}(I)}{n} \qquad \text{and} \qquad p'_j = \left\lfloor \frac{p_j}{K} \right\rfloor \quad \text{for all } j \in \mathscr{I}.$$

3. Solve the scaled instance by dynamic programming by profits, i.e., compute the minimum weights $u(i, q)$ and find the largest profit q such that $u(4, q) \leq c$.

4. State the solution returned by FP^{ε}, its profit $FP^{\varepsilon}(I)$ in the original instance, and compare it with $GR^{e}(I)$ and $z(I)$. Compute the performance ratio and compare it with the guarantee $\frac{1}{1-\varepsilon}$.

Solution

1. The items are sorted by non-increasing efficiency. GR selects the items $\{1, 2, 3\}$, with $GR(I) = 14 + 12 + 14 = 40$; the maximum profit is $\hat{p} = p_4 = 18$. Hence
$$GR^{e}(I) = 40.$$

2. The scaling factor is
$$K = \frac{\varepsilon\, GR^{e}(I)}{n} = \frac{\frac{1}{2} \cdot 40}{4} = 5,$$
and the scaled profits are
$$p'_1 = \left\lfloor \frac{14}{5} \right\rfloor = 2, \quad p'_2 = \left\lfloor \frac{12}{5} \right\rfloor = 2, \quad p'_3 = \left\lfloor \frac{14}{5} \right\rfloor = 2, \quad p'_4 = \left\lfloor \frac{18}{5} \right\rfloor = 3.$$
The total scaled profit is $P' = 2 + 2 + 2 + 3 = 9$.

3. The minimum weights $u(i, q)$ of a subset of $\{1, \ldots, i\}$ with scaled profit exactly q are (only the values of q that can be attained are reported; ∞ means that q cannot be attained with the first i items):

q	0	2	3	4	5	6	7	9
$u(0, q)$	0	∞	∞	∞	∞	∞	∞	∞
$u(1, q)$	0	3	∞	∞	∞	∞	∞	∞
$u(2, q)$	0	3	∞	7	∞	∞	∞	∞
$u(3, q)$	0	3	∞	7	∞	12	∞	∞
$u(4, q)$	0	3	7	7	10	12	14	19

The largest q with $u(4, q) \leq c = 16$ is $q = 7$, of weight $u(4, 7) = 14$, attained by the items $\{1, 2, 4\}$.

4. FP^{ε} returns $S = \{1, 2, 4\}$, of profit
$$FP^{\varepsilon}(I) = 14 + 12 + 18 = 44$$
in the original instance, better than $GR^{e}(I) = 40$. The solution $S^* = \{1, 3, 4\}$, of weight $3 + 5 + 7 = 15 \leq 16$ and profit $z(I) = 14 + 14 + 18 = 46$ is better, so FP^{ε} does **not** find an optimal solution; S^* is optimal (this can be checked by enumerating the feasible solutions). The performance ratio is
$$\frac{z(I)}{FP^{\varepsilon}(I)} = \frac{46}{44} \approx 1.05 \leq \frac{1}{1-\varepsilon} = 2,$$
as guaranteed.

Exercise 27.15 — Basic-FPTAS 3

We consider a knapsack of capacity $c = 16$ and a set $\mathscr{I} = \{1, 2, 3, 4\}$ of 4 items ($n = 4$), with the following profits and weights:

	$i = 1$	$i = 2$	$i = 3$	$i = 4$
p_i	14	12	14	18
w_i	3	4	5	7

We apply the Basic-FPTAS algorithm with accuracy $\varepsilon = \frac{1}{4}$.

Questions

1. Compute the lower bound $GR^e(I)$ used by the Basic-FPTAS algorithm.
2. Compute the scaling factor and the scaled profits
$$K = \frac{\varepsilon\, GR^e(I)}{n} \qquad \text{and} \qquad p'_j = \left\lfloor \frac{p_j}{K} \right\rfloor \quad \text{for all } j \in \mathscr{I}.$$
3. Solve the scaled instance by dynamic programming by profits, i.e., compute the minimum weights $u(i, q)$ and find the largest profit q such that $u(4, q) \leq c$.
4. State the solution returned by FP^ε, its profit $FP^\varepsilon(I)$ in the original instance, and compare it with $GR^e(I)$ and $z(I)$. Compute the performance ratio and compare it with the guarantee $\frac{1}{1-\varepsilon}$.

Solution

1. The items are sorted by non-increasing efficiency. GR selects the items $\{1, 2, 3\}$, with $GR(I) = 14 + 12 + 14 = 40$; the maximum profit is $\hat{p} = p_4 = 18$. Hence
$$GR^e(I) = 40.$$
2. The scaling factor is
$$K = \frac{\varepsilon\, GR^e(I)}{n} = \frac{\frac{1}{4} \cdot 40}{4} = \frac{5}{2} = 2.5,$$
and the scaled profits are
$$p'_1 = \left\lfloor \frac{14}{\frac{5}{2}} \right\rfloor = 5, \quad p'_2 = \left\lfloor \frac{12}{\frac{5}{2}} \right\rfloor = 4, \quad p'_3 = \left\lfloor \frac{14}{\frac{5}{2}} \right\rfloor = 5, \quad p'_4 = \left\lfloor \frac{18}{\frac{5}{2}} \right\rfloor = 7.$$
The total scaled profit is $P' = 5 + 4 + 5 + 7 = 21$.
3. The minimum weights $u(i, q)$ of a subset of $\{1, \ldots, i\}$ with scaled profit exactly q are (only the values of q that can be attained are reported; ∞ means that q cannot be attained with the first i items):

q	0	4	5	7	9	10	11	12	14	16	17
$u(0,q)$	0	∞	∞	∞	∞	∞	∞	∞	∞	∞	∞
$u(1,q)$	0	∞	3	∞	∞	∞	∞	∞	∞	∞	∞
$u(2,q)$	0	4	3	∞	7	∞	∞	∞	∞	∞	∞
$u(3,q)$	0	4	3	∞	7	8	∞	∞	12	∞	∞
$u(4,q)$	0	4	3	7	7	8	11	10	12	14	15

q	21
$u(0,q)$	∞
$u(1,q)$	∞
$u(2,q)$	∞
$u(3,q)$	∞
$u(4,q)$	19

The largest q with $u(4,q) \leq c = 16$ is $q = 17$, of weight $u(4,17) = 15$, attained by the items $\{1,3,4\}$.

4. FP^{ε} returns $S = \{1,3,4\}$, of profit

$$FP^{\varepsilon}(I) = 14 + 14 + 18 = 46$$

in the original instance, better than $GR^{e}(I) = 40$. The solution is **optimal**: $S^{*} = \{1,3,4\}$, of weight $3 + 5 + 7 = 15 \leq 16$ and profit $z(I) = 14 + 14 + 18 = 46$, and the performance ratio is 1.

Remark. This is the instance of the exercise Basic-FPTAS 2, solved there with $\varepsilon = \frac{1}{2}$. Halving ε halves the scaling factor K: the scaled profits are finer, the dynamic programming table is longer (P' grows), and the optimal solution is found.

Approximation for $P||C_{\max}$

Exercise 28.1 — $P\,||\,C_{\max}$ approximation 1

We consider $m = 3$ identical machines and a set $\mathscr{J} = \{1, \ldots, 7\}$ of 7 jobs ($n = 7$), with the following processing times:

	$j=1$	$j=2$	$j=3$	$j=4$	$j=5$	$j=6$	$j=7$
t_j	2	2	4	8	8	6	6

Questions

1. Apply $LIST$ with the list $\lambda = (1, 2, \ldots, 7)$. Show, for each job, the loads before the assignment and the chosen machine, and give $LIST(I)$.
2. Identify a job ℓ that ends last in the schedule of Question 1 and check the inequality $S_\ell \leq \frac{1}{m}\sum_{j\neq\ell} t_j$ used in the proof of the guarantee of $LIST$.
3. Apply LPT and give $LPT(I)$.
4. Compute the lower bound $\max\{\max_j t_j, \frac{1}{m}\sum_j t_j\}$ on $z(I)$ and determine $z(I)$. Compute the two ratios $\frac{LIST(I)}{z(I)}$ and $\frac{LPT(I)}{z(I)}$ and compare them with the guarantees $2 - \frac{1}{m}$ and $\frac{4}{3} - \frac{1}{3m}$.

Solution

1. Each job goes to a machine with the smallest load (smallest index in case of ties):

job j	t_j	loads (L_1, L_2, L_3) before	machine
1	2	$(0, 0, 0)$	1
2	2	$(2, 0, 0)$	2
3	4	$(2, 2, 0)$	3
4	8	$(2, 2, 4)$	1
5	8	$(10, 2, 4)$	2
6	6	$(10, 10, 4)$	3
7	6	$(10, 10, 10)$	1

Hence $mach = (1, 2, 3, 1, 2, 3, 1)$, $J_1 = \{1, 4, 7\}$, $J_2 = \{2, 5\}$, $J_3 = \{3, 6\}$, the loads are $(16, 10, 10)$ and

$$LIST(I) = 16.$$

2. Job $\ell = 7$ ends last: it starts at $S_7 = 10$ on machine 1 and ends at 16. The bound gives

$$S_7 = 10 \;\leq\; \frac{1}{3}\sum_{j \neq 7} t_j = \frac{2+2+4+8+8+6}{3} = \frac{30}{3} = 10,$$

which holds with equality: when job 7 arrives the three loads are equal, so the smallest one coincides with the average.

3. The jobs sorted by non-increasing processing time (ties by index) are $\lambda = (4, 5, 6, 7, 3, 1, 2)$:

job j	t_j	loads (L_1, L_2, L_3) before	machine
4	8	$(0, 0, 0)$	1
5	8	$(8, 0, 0)$	2
6	6	$(8, 8, 0)$	3
7	6	$(8, 8, 6)$	3
3	4	$(8, 8, 12)$	1
1	2	$(12, 8, 12)$	2
2	2	$(12, 10, 12)$	2

Hence $J_1 = \{3, 4\}$, $J_2 = \{1, 2, 5\}$, $J_3 = \{6, 7\}$, the loads are $(12, 12, 12)$ and

$$LPT(I) = 12.$$

4. The lower bound is

$$z(I) \;\geq\; \max\left\{8,\; \frac{2+2+4+8+8+6+6}{3}\right\} = \max\left\{8,\; \frac{36}{3}\right\} = 12.$$

LPT attains it, so $z(I) = 12$ and the schedule of LPT is optimal. The ratios are

$$\frac{LIST(I)}{z(I)} = \frac{16}{12} = \frac{4}{3} \;\leq\; 2 - \frac{1}{3} = \frac{5}{3}, \qquad \frac{LPT(I)}{z(I)} = \frac{12}{12} = 1 \;\leq\; \frac{4}{3} - \frac{1}{9} = \frac{11}{9}.$$

Both guarantees hold. With the list in the input order, the two longest jobs arrive when the machines are already loaded, and $LIST$ is 4 units above the optimum; sorting the list puts them first and closes the gap.

Exercise 28.2 — $P\,||\,C_{\max}$ approximation 2

We consider $m = 3$ identical machines and a set $\mathscr{J} = \{1, \dots, 7\}$ of 7 jobs ($n = 7$), with the following processing times:

	$j=1$	$j=2$	$j=3$	$j=4$	$j=5$	$j=6$	$j=7$
t_j	1	1	1	1	1	1	3

Questions

1. Apply $LIST$ with the list $\lambda = (1, 2, \dots, 7)$ and give $LIST(I)$.
2. Determine $z(I)$, by exhibiting an optimal schedule and a matching lower bound.
3. Compute $\frac{LIST(I)}{z(I)}$ and compare it with the guarantee of $LIST$. Which instance of the notes is this?
4. Apply LPT and give $LPT(I)$. Explain why sorting the list removes the bad case.

Solution

1.

job j	t_j	loads (L_1, L_2, L_3) before	machine
1	1	$(0,0,0)$	1
2	1	$(1,0,0)$	2
3	1	$(1,1,0)$	3
4	1	$(1,1,1)$	1
5	1	$(2,1,1)$	2
6	1	$(2,2,1)$	3
7	3	$(2,2,2)$	1

The loads are $(5, 2, 2)$, with $J_1 = \{1, 4, 7\}$, $J_2 = \{2, 5\}$, $J_3 = \{3, 6\}$, and

$$LIST(I) = 5.$$

2. The schedule $J_1 = \{7\}$, $J_2 = \{1, 2, 3\}$, $J_3 = \{4, 5, 6\}$ has loads $(3, 3, 3)$ and makespan 3. The lower bound gives

$$z(I) \;\geq\; \max\left\{3,\; \frac{6 \cdot 1 + 3}{3}\right\} = \max\{3, 3\} = 3.$$

Hence $z(I) = 3$.

3.
$$\frac{LIST(I)}{z(I)} = \frac{5}{3} = 2 - \frac{1}{3},$$

exactly the guarantee $2 - \frac{1}{m}$ for $m = 3$: the bound of $LIST$ is attained. This is the tight instance I_m of the notes for $m = 3$: $m(m-1) = 6$ unit jobs followed by one job

of length $m = 3$. The unit jobs fill all the machines up to $m - 1 = 2$, and the long job starts on top of one of them.

4. The sorted list is $\lambda = (7, 1, 2, 3, 4, 5, 6)$. Job 7 goes to machine 1, with loads $(3, 0, 0)$; the unit jobs then go to machines $2, 3, 2, 3, 2, 3$ (machine 1 is never the least loaded again, since its load 3 is never smaller than the others'), giving the loads $(3, 3, 3)$. Hence
$$LPT(I) = 3 = z(I).$$
The long job is the one that hurts when it comes last; LPT schedules it first, and the short jobs level the loads around it.

Exercise 28.3 — $P\,||\,C_{\max}$ approximation 3

We consider $m = 4$ identical machines and a set $\mathscr{J} = \{1, \ldots, 9\}$ of 9 jobs ($n = 9$), with the following processing times:

	$j=1$	$j=2$	$j=3$	$j=4$	$j=5$	$j=6$	$j=7$	$j=8$	$j=9$
t_j	7	7	6	6	5	5	4	4	4

Questions

1. Apply LPT and give $LPT(I)$.
2. Determine $z(I)$.
3. Compute $\frac{LPT(I)}{z(I)}$ and compare it with the guarantee $\frac{4}{3} - \frac{1}{3m}$.
4. In the proof of the guarantee of LPT, which of the two cases applies to the job ℓ that ends last? Check that every inequality of that case holds with equality on this instance.

Solution

1. The jobs are already sorted, $\lambda = (1, 2, \ldots, 9)$:

job j	t_j	loads (L_1, L_2, L_3, L_4) before	machine
1	7	$(0,0,0,0)$	1
2	7	$(7,0,0,0)$	2
3	6	$(7,7,0,0)$	3
4	6	$(7,7,6,0)$	4
5	5	$(7,7,6,6)$	3
6	5	$(7,7,11,6)$	4
7	4	$(7,7,11,11)$	1
8	4	$(11,7,11,11)$	2
9	4	$(11,11,11,11)$	1

Hence $J_1 = \{1,7,9\}$, $J_2 = \{2,8\}$, $J_3 = \{3,5\}$, $J_4 = \{4,6\}$, the loads are $(15,11,11,11)$ and

$$LPT(I) = 15.$$

2. The total processing time is 48, so $z(I) \geq \max\{7, \frac{48}{4}\} = 12$. The schedule $J_1 = \{1,5\}$, $J_2 = \{2,6\}$, $J_3 = \{3,4\}$, $J_4 = \{7,8,9\}$ has loads $(12,12,12,12)$. Hence $z(I) = 12$.

3. $$\frac{LPT(I)}{z(I)} = \frac{15}{12} = \frac{5}{4} = \frac{4}{3} - \frac{1}{12} = \frac{4}{3} - \frac{1}{3 \cdot 4}.$$

   The guarantee of LPT is attained: this is the tight instance of the notes for $m = 4$, with two jobs of each length $2m-1 = 7$, 6, $m+1 = 5$ and three jobs of length $m = 4$; after $2m = 8$ jobs all the loads are $3m - 1 = 11$, and the last job gives $4m - 1 = 15$, while $z(I) = 3m = 12$.

4. The last job is $\ell = 9$, with $t_9 = 4 = \frac{z(I)}{3}$, so Case 1 ($t_\ell \leq \frac{z}{3}$) applies, at its border. Its inequalities are:

$$\begin{aligned} LPT(I) = S_9 + t_9 &= 11 + 4 = 15, \\ \frac{1}{m}\sum_j t_j + \Big(1 - \frac{1}{m}\Big)t_9 &= \frac{48}{4} + \frac{3}{4} \cdot 4 = 15, \\ z(I) + \Big(1 - \frac{1}{m}\Big)\frac{z(I)}{3} &= 12 + \frac{3}{4} \cdot 4 = 15, \\ \Big(\frac{4}{3} - \frac{1}{3m}\Big)z(I) &= \frac{5}{4} \cdot 12 = 15. \end{aligned}$$

   All the quantities are equal: the starting time $S_9 = 11$ equals the average $\frac{48-4}{4} = 11$ of the other loads (the four machines are perfectly balanced), $\frac{1}{m}\sum_j t_j = z(I)$, and $t_9 = \frac{z(I)}{3}$.

Approximation for the metric TSP

Exercise 29.1 — Metric TSP approximation 1

We consider a set $\mathscr{V} = \{1, \ldots, 6\}$ of 6 cities with coordinates 1: $(3, 4)$, 2: $(6, 1)$, 3: $(3, 8)$, 4: $(8, 8)$, 5: $(0, 4)$, 6: $(8, 0)$ and distances equal to the Euclidean distances rounded to the nearest integer (the instance is metric):

d_{ij}	1	2	3	4	5	6
1	—	4	4	6	3	6
2	4	—	8	7	7	2
3	4	8	—	5	5	9
4	6	7	5	—	9	8
5	3	7	5	9	—	9
6	6	2	9	8	9	—

Questions

1. Compute a minimum spanning tree T with Kruskal's algorithm (edges by non-decreasing distance, ties by the pair of indices) and its weight $d(T)$.
2. Apply the double tree algorithm DT: give the Eulerian circuit of the doubled tree from city 1 (Fleury's rule), the tour obtained by shortcutting, and $DT(I)$.
3. Apply the Christofides algorithm CH: give the set O of odd-degree cities, the minimum weight perfect matching M (compare all the perfect matchings of O), the Eulerian circuit of $T \cup M$ from city 1, the tour and $CH(I)$.
4. The optimal tour has length $z(I) = 27$. Compute the two ratios and check the inequalities $DT(I) \leq 2\, d(T) \leq 2\, z(I)$ and $CH(I) \leq d(T) + d(M) \leq \frac{3}{2}\, z(I)$.

Solution

1. Kruskal's algorithm examines $\{2,6\}$ (2, taken), $\{1,5\}$ (3, taken), $\{1,2\}$ (4, taken), $\{1,3\}$ (4, taken), $\{3,4\}$ (5, taken). After five edges the tree spans the six cities:

$$T = \big\{\{2,6\},\{1,5\},\{1,2\},\{1,3\},\{3,4\}\big\}, \qquad d(T) = 2+3+4+4+5 = 18.$$

   It is the unique MST of the instance.

2. The neighbours in T are $1\colon\{2,3,5\}$, $2\colon\{1,6\}$, $3\colon\{1,4\}$, $4\colon\{3\}$, $5\colon\{1\}$, $6\colon\{2\}$. The Eulerian circuit of the doubled tree from city 1 is the depth-first visit with the children in increasing order:

$$W = (1,2,6,2,1,3,4,3,1,5,1), \qquad 4+2+2+4+4+5+5+4+3+3 = 36 = 2\,d(T).$$

   Keeping each city at its first occurrence gives $\sigma = (1,2,6,3,4,5)$, of length

$$DT(I) = d_{12} + d_{26} + d_{63} + d_{34} + d_{45} + d_{51} = 4+2+9+5+9+3 = 32.$$

3. The degrees in T are $\deg(1) = 3$, $\deg(2) = 2$, $\deg(3) = 2$, $\deg(4) = 1$, $\deg(5) = 1$, $\deg(6) = 1$, so $O = \{1,4,5,6\}$. Its three perfect matchings weigh

$$\{\{1,4\},\{5,6\}\}\colon 6+9 = 15, \quad \{\{1,5\},\{4,6\}\}\colon 3+8 = 11, \quad \{\{1,6\},\{4,5\}\}\colon 6+9 = 15,$$

   so $M = \{\{1,5\},\{4,6\}\}$, with $d(M) = 11$. The multigraph $T \cup M$ has two copies of $\{1,5\}$. Fleury's rule from city 1 (neighbours $2,3,5,5$) goes to 2, then 6, 4 (along M), 3, back to 1, and finally 5 and 1 along the two copies of $\{1,5\}$:

$$W = (1,2,6,4,3,1,5,1), \qquad 4+2+8+5+4+3+3 = 29 = d(T) + d(M).$$

   City 1 is repeated in the middle; shortcutting it gives $\sigma = (1,2,6,4,3,5)$, of length

$$CH(I) = d_{12} + d_{26} + d_{64} + d_{43} + d_{35} + d_{51} = 4+2+8+5+5+3 = 27.$$

   The shortcut replaces $3 \to 1 \to 5$ $(4+3)$ by $3 \to 5$ (5) and saves 2.

4. The ratios are

$$\frac{DT(I)}{z(I)} = \frac{32}{27}, \qquad \frac{CH(I)}{z(I)} = \frac{27}{27} = 1,$$

   so CH finds the optimal tour, which is unique. The inequalities of the proofs are

$$DT(I) = 32 \le 2\,d(T) = 36 \le 2\,z(I) = 54,$$
$$CH(I) = 27 \le d(T) + d(M) = 29 \le \frac{3}{2}\,z(I) = \frac{81}{2}.$$

Exercise 29.2 — Metric TSP approximation 2

We consider a set $\mathcal{V} = \{1,\dots,6\}$ of 6 cities with coordinates $1\colon(2,4)$, $2\colon(6,0)$, $3\colon(7,3)$, $4\colon(0,7)$, $5\colon(0,1)$, $6\colon(2,1)$ and distances equal to the Euclidean distances rounded to the

nearest integer (the instance is metric):

d_{ij}	1	2	3	4	5	6
1	—	6	5	4	4	3
2	6	—	3	9	6	4
3	5	3	—	8	7	5
4	4	9	8	—	6	6
5	4	6	7	6	—	2
6	3	4	5	6	2	—

Questions

1. Compute a minimum spanning tree T with Kruskal's algorithm and its weight $d(T)$.
2. Apply CH: give O, the weights of all the perfect matchings of O, the matching M, the Eulerian circuit of $T \cup M$ from city 1 (Fleury's rule), the tour and $CH(I)$.
3. Apply DT and give $DT(I)$.
4. The unique optimal tour is $(1, 3, 2, 6, 5, 4)$. Compute $z(I)$ and the two ratios. Is CH optimal on this instance? Check the bound $d(M) \leq \frac{z(I)}{2}$ by shortcutting the optimal tour to the cities of O, as in the proof of the notes.

Solution

1. Kruskal's algorithm examines $\{5,6\}$ (2, taken), $\{1,6\}$ (3, taken), $\{2,3\}$ (3, taken), $\{1,4\}$ (4, taken), $\{1,5\}$ (4, it closes the cycle $1,5,6$: rejected), $\{2,6\}$ (4, taken). Hence

$$T = \big\{\{5,6\},\{1,6\},\{2,3\},\{1,4\},\{2,6\}\big\}, \qquad d(T) = 2+3+3+4+4 = 16,$$

the unique MST of the instance.

2. The degrees in T are $\deg(1) = 2$, $\deg(2) = 2$, $\deg(3) = 1$, $\deg(4) = 1$, $\deg(5) = 1$, $\deg(6) = 3$, so $O = \{3,4,5,6\}$. Its perfect matchings weigh

$$\{\{3,4\},\{5,6\}\}\text{: } 8+2=10, \quad \{\{3,5\},\{4,6\}\}\text{: } 7+6=13, \quad \{\{3,6\},\{4,5\}\}\text{: } 5+6=11,$$

so $M = \{\{3,4\},\{5,6\}\}$, with $d(M) = 10$; $T \cup M$ has two copies of $\{5,6\}$. Fleury's rule from city 1 (neighbours 4, 6) goes to 4, then to 3 along M, to 2 and to 6. At city 6 the unused edges are $\{6,1\}$ and the two copies of $\{6,5\}$: city 1 has the smallest index, but $\{6,1\}$ is a bridge of the unused edges (removing it leaves city 1 isolated from the two copies of $\{5,6\}$), so the walk goes to 5, back to 6 and finally to 1:

$$W = (1,4,3,2,6,5,6,1), \qquad 4+8+3+4+2+2+3 = 26 = d(T) + d(M).$$

Shortcutting the second visit of city 6 gives $\sigma = (1,4,3,2,6,5)$, of length

$$CH(I) = d_{14} + d_{43} + d_{32} + d_{26} + d_{65} + d_{51} = 4+8+3+4+2+4 = 25.$$

3. The neighbours in T are $1{:}\{4,6\}$, $2{:}\{3,6\}$, $3{:}\{2\}$, $4{:}\{1\}$, $5{:}\{6\}$, $6{:}\{1,2,5\}$. The depth-first visit from city 1 is

$$W = (1,4,1,6,2,3,2,6,5,6,1), \qquad \text{of length } 32 = 2\,d(T),$$

and shortcutting gives $\sigma = (1,4,6,2,3,5)$, of length

$$DT(I) = 4+6+4+3+7+4 = 28.$$

4. $z(I) = d_{13} + d_{32} + d_{26} + d_{65} + d_{54} + d_{41} = 5+3+4+2+6+4 = 24$, and

$$\frac{CH(I)}{z(I)} = \frac{25}{24}, \qquad \frac{DT(I)}{z(I)} = \frac{28}{24} = \frac{7}{6}.$$

CH is not optimal here, but it is only one unit above the optimum. Shortcutting the optimal tour to $O = \{3,4,5,6\}$ gives the cycle $(3,6,5,4)$, of length $d_{36} + d_{65} + d_{54} + d_{43} = 5+2+6+8 = 21 \le 24$. Its alternate edges form the matchings $\{\{3,6\},\{5,4\}\}$, of weight 11, and $\{\{6,5\},\{4,3\}\}$, of weight 10, and indeed

$$d(M) = 10 \le \min\{11,10\} \le \frac{21}{2} \le \frac{z(I)}{2} = 12.$$

Exercise 29.3 — Metric TSP approximation 3

We consider two graphs with 5 vertices, $V = \{1,2,3,4,5\}$:

$$G_1: \; E_1 = \big\{\{1,2\},\{2,3\},\{3,4\},\{4,5\},\{5,1\},\{1,3\}\big\},$$

$$G_2: \; E_2 = \big\{\{1,3\},\{1,4\},\{1,5\},\{2,3\},\{2,4\},\{2,5\}\big\}.$$

Suppose that a polynomial algorithm $\mathscr{A}$ for the TSP had the guarantee $\frac{\mathscr{A}(I)}{z(I)} \le \alpha$ with $\alpha = 2$ on every TSP instance.

Questions

1. Build the TSP instances I_{G_1} and I_{G_2} of the proof of the notes: $d_{ij} = 1$ if $\{i,j\}$ is an edge and $d_{ij} = \lceil \alpha n \rceil + 1$ otherwise.
2. Determine whether G_1 and G_2 are Hamiltonian, and compute $z(I_{G_1})$ and $z(I_{G_2})$.
3. What can $\mathscr{A}$ return on the two instances? Explain how $\mathscr{A}$ would decide whether a graph is Hamiltonian.
4. Is I_{G_2} metric? Could $\mathscr{A}$ be the Christofides algorithm?

Solution

1. With $n = 5$ and $\alpha = 2$, the non-edges get distance $\lceil 2 \cdot 5 \rceil + 1 = 11$:

I_{G_1}	1	2	3	4	5
1	—	1	1	11	1
2	1	—	1	11	11
3	1	1	—	1	11
4	11	11	1	—	1
5	1	11	11	1	—

I_{G_2}	1	2	3	4	5
1	—	11	1	1	1
2	11	—	1	1	1
3	1	1	—	11	11
4	1	1	11	—	11
5	1	1	11	11	—

2. G_1 contains the cycle $(1, 2, 3, 4, 5)$, which is Hamiltonian, so the tour $(1, 2, 3, 4, 5)$ has length 5 and $z(I_{G_1}) = 5$.

   G_2 is the complete bipartite graph with sides $\{1, 2\}$ and $\{3, 4, 5\}$. A cycle in a bipartite graph alternates between the two sides, so it has as many vertices on each side: no cycle can contain the three vertices $3, 4, 5$ and only two vertices of the other side, and G_2 is not Hamiltonian. Every tour of I_{G_2} uses at least one distance 11. The tour $(1, 3, 2, 4, 5)$ uses the edges $\{1, 3\}, \{3, 2\}, \{2, 4\}, \{1, 5\}$ and the non-edge $\{4, 5\}$, so its length is $4 \cdot 1 + 11 = 15$, and

   $$z(I_{G_2}) = 15.$$

   (Enumerating the 12 tours, their lengths are only 15 and 35.)

3. On I_{G_1}: $\mathscr{A}(I_{G_1}) \leq 2 \cdot 5 = 10$. A tour that uses a distance 11 has length at least $4 + 11 = 15 > 10$, so $\mathscr{A}$ must return a tour of length 5, i.e., a Hamiltonian cycle of G_1.

   On I_{G_2}: $\mathscr{A}(I_{G_2}) \geq z(I_{G_2}) = 15 > 10$.

   So the test "$\mathscr{A}(I_G) \leq \alpha n = 10$" answers "yes" on G_1 and "no" on G_2. In general, building I_G and running $\mathscr{A}$ takes polynomial time and gives the right answer on every graph, so the NP-complete Hamiltonian cycle problem would be solved in polynomial time: such an $\mathscr{A}$ cannot exist unless $P = NP$.

4. I_{G_2} is not metric: for example $d_{12} = 11 > d_{13} + d_{32} = 1 + 1 = 2$. The guarantee $\frac{3}{2}$ of the Christofides algorithm holds only for metric instances, so it says nothing on I_{G_2}, and there is no contradiction with the proposition of the notes: it is the triangle inequality that makes the TSP approximable.


Chapter 30

Limits of approximation: vertex coloring

Exercise 30.1 — Vertex coloring approximation 1

We consider the graph $G = (V, E)$ with $V = \{1, \ldots, 7\}$ ($n = 7$) and $|E| = 11$ edges:

$$E = \big\{\{1,2\},\{1,3\},\{1,4\},\{2,4\},\{2,6\},\{3,7\},\{4,5\},\{4,6\},\{5,6\},\{5,7\},\{6,7\}\big\}.$$



Questions

1. Compute the degrees and $\Delta(G)$, and the bound $\Delta(G)+1$ on the number of colors of GC.
2. Apply GC with the list $\lambda = (1, 2, \ldots, 7)$: for each vertex give the set U of colors of its colored neighbours and the color assigned.
3. Apply $DSATUR$: for each step give the vertex chosen, its saturation degree sat, its degree among the uncolored vertices du, and its color.
4. Find a clique of maximum size and determine $\chi(G)$. Which of the two colorings is optimal?

Solution

1. The neighbourhoods are $N(1) = \{2,3,4\}$, $N(2) = \{1,4,6\}$, $N(3) = \{1,7\}$, $N(4) = \{1,2,5,6\}$, $N(5) = \{4,6,7\}$, $N(6) = \{2,4,5,7\}$, $N(7) = \{3,5,6\}$. The degrees are $(3,3,2,4,3,4,3)$, so $\Delta(G) = 4$ and GC uses at most $\Delta(G)+1 = 5$ colors, with any

list.

2.

v	1	2	3	4	5	6	7
U	$\emptyset$	$\{1\}$	$\{1\}$	$\{1,2\}$	$\{3\}$	$\{1,2,3\}$	$\{1,2,4\}$
$col[v]$	1	2	2	3	1	4	3

GC uses 4 colors: $Q_1 = \{1,5\}$, $Q_2 = \{2,3\}$, $Q_3 = \{4,7\}$, $Q_4 = \{6\}$. The trouble starts at vertex 5: it takes color 1, and vertex 6, adjacent to 2, 4 and 5, then sees the colors $1, 2, 3$.

3. At step 1 all the saturation degrees are 0, and the vertices of largest degree are 4 and 6: vertex 4 is chosen.

step	1	2	3	4	5	6	7
v	4	6	2	1	5	7	3
$sat[v]$	0	1	2	2	2	2	2
$du[v]$	4	3	1	1	1	1	0
$col[v]$	1	2	3	2	3	1	3

At step 2 the neighbours of 4 have saturation degree 1, and vertex 6 has the most uncolored neighbours (5, 7 and 2). At step 3 the vertices 2 and 5 (both adjacent to 4 and 6) have saturation degree 2 and one uncolored neighbour each, and 2 is chosen. $DSATUR$ uses 3 colors: $Q_1 = \{4,7\}$, $Q_2 = \{1,6\}$, $Q_3 = \{2,3,5\}$.

4. $\{1,2,4\}$ is a clique of size 3 ($\{1,2\}, \{1,4\}, \{2,4\} \in E$), so $\chi(G) \geq \omega(G) \geq 3$; there are no cliques of size 4, but this is not needed: the $DSATUR$ coloring uses 3 colors, hence $\chi(G) = 3$. The $DSATUR$ coloring is optimal, the one of GC with the list $(1, \ldots, 7)$ uses one color more:

$$\frac{GC(I)}{\chi(G)} = \frac{4}{3}, \qquad \frac{DSATUR(I)}{\chi(G)} = 1.$$

Exercise 30.2 — Vertex coloring approximation 2

We consider the crown graph G_5 with $n = 10$ vertices: the vertices $1, \ldots, 5$ on one side, the vertices $6, \ldots, 10$ on the other, and vertex $i \in \{1, \ldots, 5\}$ adjacent to vertex $j + 5$, $j \in \{1, \ldots, 5\}$, if and only if $i \neq j$:

$$E = \big\{\{i, j+5\} : i, j \in \{1, \ldots, 5\},\ i \neq j\big\}, \qquad |E| = 20.$$

Questions

1. Compute $\Delta(G_5)$ and $\chi(G_5)$.
2. Apply GC with the list $\lambda = (1, 6, 2, 7, 3, 8, 4, 9, 5, 10)$ and give $GC(I)$. Compare it with $\Delta(G_5) + 1$ and with $\chi(G_5)$.
3. Give a list for which GC is optimal.

4. Apply $DSATUR$ and give $DSATUR(I)$.

Solution

1. Each vertex $i \leq 5$ is adjacent to the 4 vertices $j+5$ with $j \neq i$, and symmetrically: every degree is 4 and $\Delta(G_5) = 4$. The graph is bipartite (all the edges join $\{1,\dots,5\}$ and $\{6,\dots,10\}$) and has edges, so $\chi(G_5) = 2$.

2. With $u_i = i$ and $v_i = i+5$ the list is $(u_1, v_1, u_2, v_2, \dots, u_5, v_5)$, the bad list of the notes:

v	1	6	2	7	3	8	4	9	5	10
U	$\emptyset$	$\emptyset$	$\{1\}$	$\{1\}$	$\{1,2\}$	$\{1,2\}$	$\{1,2,3\}$	$\{1,2,3\}$	$\{1,2,3,4\}$	$\{1,2,3,4\}$
$col[v]$	1	1	2	2	3	3	4	4	5	5

   Vertices i and $i+5$ are not adjacent and both get color i, so $GC(I) = 5 = \frac{n}{2}$. This is exactly the bound $\Delta(G_5)+1 = 5$, and

$$\frac{GC(I)}{\chi(G_5)} = \frac{5}{2}.$$

3. The list $(1,2,3,4,5,6,7,8,9,10)$, one side first: the vertices $1,\dots,5$ are pairwise non-adjacent and all get color 1; each vertex $6,\dots,10$ has only neighbours of color 1 and gets color 2. Hence $GC(I) = 2 = \chi(G_5)$.

4. At step 1 all the vertices have degree 4 and vertex 1 is chosen (color 1). Then:


step	1	2	3	4	5	6	7	8	9	10
v	1	7	3	4	5	6	2	8	9	10
$sat[v]$	0	1	1	1	1	1	1	1	1	1
$du[v]$	4	3	3	3	3	1	3	0	0	0
$col[v]$	1	2	1	1	1	2	1	2	2	2

   At step 2 the neighbours $7,8,9,10$ of vertex 1 have saturation degree 1 and 3 uncolored neighbours each: vertex 7 is chosen and gets color 2. From then on, every vertex adjacent to a colored vertex has saturation degree 1 (its colored neighbours are all on the other side, with the same color); the vertices of the side $\{1,\dots,5\}$ get color 1 and those of the side $\{6,\dots,10\}$ color 2. Hence $DSATUR(I) = 2 = \chi(G_5)$, as guaranteed on bipartite graphs.

Exercise 30.3 — Vertex coloring approximation 3

We consider the graph $G = (V, E)$ with $V = \{1,\dots,7\}$ ($n = 7$) and $|E| = 10$ edges:

$$E = \{\{1,3\},\{1,5\},\{1,6\},\{2,4\},\{2,5\},\{2,7\},\{3,5\},\{4,6\},\{4,7\},\{6,7\}\}.$$



Questions

1. Apply $DSATUR$ (ties: largest degree among the uncolored vertices, then smallest index) and give $DSATUR(I)$.
2. Show that $\chi(G) = 3$.
3. Suppose that a polynomial algorithm $\mathscr{A}$ had $\frac{\mathscr{A}(I)}{\chi(G)} < \frac{4}{3}$ on every graph. What would $\mathscr{A}$ return on this graph? Why does this show that such an algorithm cannot exist unless $P = NP$?
4. Is the result of Question 1 in contradiction with the bound $\Delta(G) + 1$?

Solution

1. The neighbourhoods are $N(1) = \{3, 5, 6\}$, $N(2) = \{4, 5, 7\}$, $N(3) = \{1, 5\}$, $N(4) = \{2, 6, 7\}$, $N(5) = \{1, 2, 3\}$, $N(6) = \{1, 4, 7\}$, $N(7) = \{2, 4, 6\}$.

step	1	2	3	4	5	6	7
v	1	5	3	2	4	6	7
$sat[v]$	0	1	2	1	1	2	3
$du[v]$	3	2	0	2	2	1	0
$col[v]$	1	2	3	1	2	3	4

   - Step 1: vertex 1 (degree 3, smallest index among the vertices of degree 3), color 1.
   - Step 2: the neighbours 3, 5, 6 of vertex 1 have saturation degree 1; their uncolored neighbours are 1, 2, 2 respectively, and vertex 5 is chosen: color 2.
   - Step 3: vertex 3 sees colors 1 and 2 (saturation degree 2): color 3.
   - Step 4: the vertices 2 (neighbour 5) and 6 (neighbour 1) have saturation degree 1 and two uncolored neighbours each; vertex 2 is chosen and gets color 1.
   - Step 5: the vertices 4, 6, 7 have saturation degree 1 and two uncolored neighbours each; vertex 4 gets color 2.
   - Step 6: vertex 6 sees colors 1 (vertex 1) and 2 (vertex 4), vertex 7 sees 1 (vertex 2) and 2 (vertex 4); both have saturation degree 2 and one uncolored neighbour, and vertex 6 gets color 3.
   - Step 7: vertex 7 sees colors 1, 2, 3 and gets color 4.

   Hence $DSATUR(I) = 4$.

2. $\{1,3,5\}$ is a triangle, so $\chi(G) \geq 3$. The coloring $Q_1 = \{1,4\}$, $Q_2 = \{2,3,6\}$, $Q_3 = \{5,7\}$ is feasible (no edge inside a class: $\{1,4\}$, $\{2,3\}$, $\{2,6\}$, $\{3,6\}$, $\{5,7\} \notin E$), so $\chi(G) = 3$ and
$$\frac{DSATUR(I)}{\chi(G)} = \frac{4}{3}.$$
The damage is done by step 5: vertices 6 and 7 already see color 1 (from vertices 1 and 2), and vertex 4, their common neighbour, takes color 2. The two adjacent vertices 6 and 7 are left with color 3 only, and one of them needs a fourth color. In the optimal coloring vertex 4 has the same color as vertex 1.

3. On this graph $\mathscr{A}(I) < \frac{4}{3} \cdot 3 = 4$, so $\mathscr{A}(I) \leq 3$: $\mathscr{A}$ would return a 3-coloring. On a graph with $\chi(G) \geq 4$, instead, $\mathscr{A}(I) \geq 4$. Comparing $\mathscr{A}(I)$ with 3 would decide 3-colorability, an NP-complete problem, in polynomial time. In particular, $DSATUR$ is not such an algorithm: on this graph its ratio is exactly $\frac{4}{3}$.

4. No: the degrees are $(3,3,2,3,3,3,3)$, so $\Delta(G)+1 = 4$ and $DSATUR(I) = 4$ attains the bound. The bound $\Delta(G)+1$ limits the number of colors, it does not compare it with $\chi(G)$.