

SAPIENZA
UNIVERSITÀ DI ROMA

Combinatorial Optimization

Volume 2 — Formulations and their properties

Chapters 8–13 of the module «Combinatorial Optimization»: formulations and their properties. Definitions, results with proofs and worked examples; exercises with complete solutions are in Exercises 2.

Fabio Furini

fabiofurini.github.io

Department of Computer, Control and Management Engineering — Sapienza University of Rome

Online version and downloadable material:
fabiofurini.github.io/combinatorial-optimization/
github.com/fabiofurini/combinatorial-optimization

Academic year 2026–2027

Contents

8 Formulations of the knapsack problem **3**

8.1 The problem in brief . . . 4
8.2 An ILP model for the Knapsack Problem . . . 4
8.3 A first Linear Programming relaxation . . . 6
8.4 A second Linear Programming relaxation . . . 8
8.5 Knapsack Problem upper bounds . . . 17
8.6 Typical mistakes . . . 19
8.7 In summary . . . 19

9 Formulations of the bin packing problem **21**

9.1 The problem in brief . . . 22
9.2 An ILP model for the Bin Packing Problem . . . 22
9.3 A first Linear Programming relaxation . . . 25
9.4 A second Linear Programming relaxation . . . 29
9.5 Typical mistakes . . . 36
9.6 In summary . . . 37

10 Formulations of the vertex coloring problem **38**

10.1 The problem in brief . . . 39
10.2 An ILP model for the Vertex Coloring Problem . . . 40
10.3 A first Linear Programming relaxation . . . 42
10.4 A second Linear Programming relaxation . . . 47
10.5 Typical mistakes . . . 50
10.6 In summary . . . 51

11 Formulations of the travelling salesman problem **52**

11.1 The problem in brief . . . 53
11.2 A model with position variables (MTZ) . . . 54
11.3 A model with subtour elimination constraints (DFJ) . . . 56
11.4 Linear Programming relaxations . . . 57
11.5 Typical mistakes . . . 58
11.6 In summary . . . 59

12 Formulations of parallel machine scheduling **60**
12.1 The problem in brief . . . 61
12.2 An ILP model for $P\|C_{\max}$ . . . 61
12.3 A first Linear Programming relaxation . . . 63
12.4 Additional considerations . . . 66
12.5 Typical mistakes . . . 66
12.6 In summary . . . 67

13 Formulations of capacitated lot sizing **68**
13.1 The problem in brief . . . 69
13.2 A formulation with a big-M link . . . 70
13.3 A shortest-path formulation . . . 72
13.4 Comparing the formulations . . . 74
13.5 Several products, setup times and initial inventory . . . 74
13.6 Typical mistakes . . . 76
13.7 In summary . . . 76

Chapter 8

Formulations of the knapsack problem

Prerequisites

- From the module *Linear Programming*, Chapter 9 «Duality theory»: the dual of an LP problem, weak and strong duality.
- From the module *Integer linear programming*, the chapters «Integer linear programming» and «LP relaxation and dual bounds»: ILP problems, LP relaxations and upper bounds.
- From this module, the chapter «Knapsack problems»: the problem, its assumptions and the instance of Example 1.

Objectives

At the end of this chapter you will be able to:

- write the ILP model of the Knapsack Problem and explain each of its constraints;
- write the two LP relaxations and their duals, and compute their optimal solutions with closed formulas;
- compute the split item, the residual capacity and the value $z(\mathrm{LP_{KP}})$;
- compute the Dantzig and the Martello-Toth upper bounds, rounding them correctly;
- explain why $z(\mathrm{LP_{KP}})$ is never larger than twice $z(\mathrm{ILP_{KP}})$.

8.1 The problem in brief

Where we were

In the chapter «Knapsack problems» we met the Knapsack Problem as a *problem*: definition, variants and the instance of Example 1 (capacity 9, six items). Here we look at it as a *model* and we compute upper bounds on its optimal value. The same instance returns with dynamic programming, heuristic algorithms, approximation algorithms and the branch-and-bound algorithm for the KP.

Definition 8.1: Knapsack Problem

Given:

1. a knapsack of *capacity* $c \in \mathbb{Z}_{\geq 1}$,
2. a set $\mathscr{I} = \{1, 2, \ldots, n\}$ of n items,
3. for each item $j \in \mathscr{I}$, a *profit* $p_j \in \mathbb{Z}_{\geq 1}$ and a *weight* $w_j \in \mathbb{Z}_{\geq 1}$,

the **Knapsack Problem** (KP) asks to decide which items to insert into the knapsack, so that the total weight of the selected items does not exceed the knapsack capacity, and the total profit of the selected items is maximized.

- We assume that any KP instance satisfies the problem assumptions, i.e.,

$$w_j \leq c \quad \text{for all} \quad j \in \mathscr{I} \qquad \text{and} \qquad \sum_{j \in \mathscr{I}} w_j > c$$

The first assumption guarantees that every item can be selected, the second one that not all the items fit together into the knapsack.

8.2 An ILP model for the Knapsack Problem

- We introduce the following n **binary variables**:

$$x_j = \begin{cases} 1 & \text{if item } j \text{ is selected in the knapsack} \\ 0 & \text{otherwise} \end{cases} \qquad \forall j \in \underbrace{\{1, 2, \ldots, n\}}_{=\mathscr{I}}. \tag{8.1}$$

Using these variables, an Integer Linear Programming (ILP) model for the Knapsack Problem (KP) reads as follows:

$$\max \sum_{j \in \mathscr{I}} p_j \, x_j \tag{8.2a}$$

$$\text{subject to} \sum_{j \in \mathscr{I}} w_j \, x_j \leq c, \tag{8.2b}$$

$$x_j \in \{0,1\}, \qquad \forall j \in \mathscr{I}. \tag{8.2c}$$

This ILP model is referred to as the ILP_{KP} model.

- Description of the objective function and constraints of the ILP_{KP} model:
  - the linear objective function (8.2a) maximizes the total profit of the items selected in the knapsack;
  - constraint (8.2b) ensures that the total weight of the selected items does not exceed the knapsack capacity (one linear constraint);
  - finally, constraints (8.2c) define the variables of the ILP model.

Example 8.1: ILP_{KP} model

We consider a knapsack of capacity $c = 9$ and a set $\mathscr{I} = \{1,2,3,4,5,6\}$ of 6 items ($n = 6$), with the following profits and weights:

	$j=1$	$j=2$	$j=3$	$j=4$	$j=5$	$j=6$
p_j	6	5	8	9	6	7
w_j	2	3	6	7	5	9

We refer to this KP instance as Example 1. The ILP_{KP} model for this KP instance is:

$$\begin{aligned}
\max \quad & 6\,x_1 + 5\,x_2 + 8\,x_3 + 9\,x_4 + 6\,x_5 + 7\,x_6 \\
\text{subject to} \quad & 2\,x_1 + 3\,x_2 + 6\,x_3 + 7\,x_4 + 5\,x_5 + 9\,x_6 \leq 9 \\
& x_1, \; x_2, \; x_3, \; x_4, \; x_5, \; x_6 \in \{0,1\}
\end{aligned}$$

An optimal solution $\tilde{\boldsymbol{x}}(\text{ILP}_{\text{KP}})$ is:

$$\tilde{x}_1 = 1, \quad \tilde{x}_2 = 0, \quad \tilde{x}_3 = 0, \quad \tilde{x}_4 = 1, \quad \tilde{x}_5 = 0, \quad \tilde{x}_6 = 0$$

The optimal value $z(\text{ILP}_{\text{KP}})$ is 15. The associated optimal solution of the KP is $\tilde{S} = \{1,4\}$, and it is the unique optimal solution.

8.3 A first Linear Programming relaxation

To obtain a Linear Programming (LP) relaxation of the ILP_{KP} model, we relax constraints (8.2c) with the following ones:

$$x_j \geq 0, \qquad \forall j \in \mathscr{J}, \tag{8.3}$$

This LP model is referred to as the LP-0_{KP} model.

Example 8.2: LP-0_{KP} model

The LP-0_{KP} model for the KP instance of Example 1 is:

$$\begin{array}{rrrrrrrrrrrrrrr} \max & 6 & x_1 & +5 & x_2 & +8 & x_3 & +9 & x_4 & +6 & x_5 & +7 & x_6 & & \\ \text{subject to} & 2 & x_1 & +3 & x_2 & +6 & x_3 & +7 & x_4 & +5 & x_5 & +9 & x_6 & \leq & 9 \\ & & x_1, & & x_2, & & x_3, & & x_4, & & x_5, & & x_6 & \geq & 0 \end{array}$$

An optimal solution $\tilde{\boldsymbol{x}}(\text{LP-}0_{\text{KP}})$ is:

$$\tilde{x}_1 = \frac{9}{2}, \quad \tilde{x}_2 = \tilde{x}_3 = \tilde{x}_4 = \tilde{x}_5 = \tilde{x}_6 = 0$$

The optimal value $z(\text{LP-}0_{\text{KP}})$ is 27.

Accordingly we have the upper bound:

$$z(\text{ILP}_{\text{KP}}) \leq 27$$

Using a non-negative dual variable π for the constraint (8.2b), the dual model of the LP-0_{KP} model is:

$$\min \quad c\,\pi \tag{8.4a}$$

$$\text{subject to} \quad w_j\,\pi \geq p_j, \qquad \forall j \in \mathscr{J}, \tag{8.4b}$$

$$\pi \geq 0. \tag{8.4c}$$

The obtained LP model is referred to as the $D(\text{LP-}0_{\text{KP}})$ model.

Example 8.3: D(LP-0_{KP}) model

The dual $D(\text{LP-}0_{\text{KP}})$ model for the KP instance of Example 1 is:

$$
\begin{array}{rrcl}
\min & 9\ \pi & & \\
\text{subject to} & 2\ \pi & \geq & 6 \\
& 3\ \pi & \geq & 5 \\
& 6\ \pi & \geq & 8 \\
& 7\ \pi & \geq & 9 \\
& 5\ \pi & \geq & 6 \\
& 9\ \pi & \geq & 7 \\
& \pi & \geq & 0
\end{array}
$$

An optimal solution $\tilde{\pi}(D(\text{LP-0}_{\text{KP}}))$ is:

$$\tilde{\pi} = 3$$

The optimal value $z(D(\text{LP-0}_{\text{KP}}))$ is 27.

- The following proposition provides formulas to compute an optimal solution of the primal LP-0_{KP} model and of the dual D(LP-0_{KP}) model.

Proposition 8.2

If the first item $j = 1$ has the largest efficiency, an optimal solution of the primal LP-0_{KP} model is:

$$\tilde{x}_1 = \frac{c}{w_1} \quad \text{and} \quad \tilde{x}_j = 0, \quad \forall j \in \{2, 3, \dots, n\}, \tag{8.5}$$

and an optimal solution of the dual D(LP-0_{KP}) model is:

$$\tilde{\pi} = \frac{p_1}{w_1}. \tag{8.6}$$

Corollary 8.3

The optimal value of the primal LP-0_{KP} and dual D(LP-0_{KP}) models is:

$$z(\text{LP-0}_{\text{KP}}) = p_1 \frac{c}{w_1} = z(D(\text{LP-0}_{\text{KP}})) \tag{8.7}$$

- The proof proceeds in three steps:
  1. We show that (8.5) is a feasible solution for the primal LP-0_{KP} model;
  2. We show that (8.6) is a feasible solution for the dual D(LP-0_{KP}) model;

3. We show that both feasible solutions have the same objective function value.

Therefore, by LP Strong Duality, (8.5) and (8.6) are a pair of optimal primal and dual solutions.

Proof.

1. We have:
$$\sum_{j \in \mathscr{I}} w_j \, \tilde{x}_j = w_1 \, \frac{c}{w_1} = c$$
thus constraint (8.2b) is satisfied.
Since $\tilde{x}_j \geq 0$, for every $j \in \mathscr{I}$, constraints (8.3) are satisfied.
Therefore, (8.5) is a feasible solution for the primal LP-0_{KP} model.

2. For every $j \in \mathscr{I}$, we have:
$$w_j \, \tilde{\pi} = w_j \, \frac{p_1}{w_1} \geq p_j \qquad \text{since} \qquad \frac{p_1}{w_1} \geq \frac{p_j}{w_j}$$
thus constraints (8.4b) are satisfied.
We have $\tilde{\pi} \geq 0$, thus constraint (8.4c) is satisfied.
Therefore, (8.6) is a feasible solution for the dual $D(\text{LP-}0_{\text{KP}})$ model.

3. For the objective functions, we have:
$$\sum_{j \in \mathscr{I}} p_j \, \tilde{x}_j = p_1 \, \frac{c}{w_1} = c \, \tilde{\pi}$$
thus the primal and dual objective function values are equal.

Therefore, by LP Strong Duality, (8.5) and (8.6) are a pair of optimal primal and dual solutions.

□

8.4 A second Linear Programming relaxation

To obtain a LP relaxation of the ILP model, we relax constraints (8.2c) with the following ones:

$$x_j \leq 1, \qquad \forall j \in \mathscr{I}, \tag{8.8a}$$

$$x_j \geq 0, \qquad \forall j \in \mathscr{I}. \tag{8.8b}$$

The obtained LP model is referred to as the LP_{KP} model.

Example 8.4: $\mathrm{LP_{KP}}$ model

The $\mathrm{LP_{KP}}$ model for the KP instance of Example 1 is:

$$
\begin{array}{rrrrrrrrrrrrrrr}
\max & 6 & x_1 & +5 & x_2 & +8 & x_3 & +9 & x_4 & +6 & x_5 & +7 & x_6 & & \\
\text{subject to} & 2 & x_1 & +3 & x_2 & +6 & x_3 & +7 & x_4 & +5 & x_5 & +9 & x_6 & \leq & 9 \\
 & & x_1 & & & & & & & & & & & \leq & 1 \\
 & & & & x_2 & & & & & & & & & \leq & 1 \\
 & & & & & & x_3 & & & & & & & \leq & 1 \\
 & & & & & & & & x_4 & & & & & \leq & 1 \\
 & & & & & & & & & & x_5 & & & \leq & 1 \\
 & & & & & & & & & & & & x_6 & \leq & 1 \\
 & & x_1, & & x_2, & & x_3, & & x_4, & & x_5, & & x_6 & \geq & 0
\end{array}
$$

An optimal solution $\tilde{\boldsymbol{x}}(\mathrm{LP_{KP}})$ is:

$$\tilde{x}_1 = \tilde{x}_2 = 1, \quad \tilde{x}_3 = \frac{2}{3}, \quad \tilde{x}_4 = \tilde{x}_5 = \tilde{x}_6 = 0$$

The optimal value $z(\mathrm{LP_{KP}})$ is $\frac{49}{3}$.

Accordingly we have the upper bound:

$$z(\mathrm{ILP_{KP}}) \leq \frac{49}{3}$$

Using a non-negative dual variable π for the constraint (8.2b) and a non-negative dual variable μ_j for each constraint (8.8a), the dual model of the $\mathrm{LP_{KP}}$ model is:

$$\min \quad c\,\pi + \sum_{j \in \mathscr{I}} \mu_j \tag{8.9a}$$

$$\text{subject to} \quad w_j\,\pi + \mu_j \geq p_j, \qquad \forall j \in \mathscr{I}, \tag{8.9b}$$

$$\pi \geq 0, \tag{8.9c}$$

$$\mu_j \geq 0, \qquad \forall j \in \mathscr{I}. \tag{8.9d}$$

This LP model is referred to as the dual $D(\mathrm{LP_{KP}})$ model.

Example 8.5: $D(\text{LP}_{\text{KP}})$ model

The dual $D(\text{LP}_{\text{KP}})$ model for the KP instance of Example 1 is:

$$
\begin{array}{lrrrrrrrrrr}
\min & 9 & \pi & +\mu_1 & +\mu_2 & +\mu_3 & +\mu_4 & +\mu_5 & +\mu_6 & & \\
& 2 & \pi & +\mu_1 & & & & & & \geq & 6 \\
& 3 & \pi & & +\mu_2 & & & & & \geq & 5 \\
& 6 & \pi & & & +\mu_3 & & & & \geq & 8 \\
& 7 & \pi & & & & +\mu_4 & & & \geq & 9 \\
& 5 & \pi & & & & & +\mu_5 & & \geq & 6 \\
& 9 & \pi & & & & & & +\mu_6 & \geq & 7 \\
& & \pi, & \mu_1, & \mu_2, & \mu_3, & \mu_4, & \mu_5, & \mu_6 & \geq & 0
\end{array}
$$

An optimal solution $(\tilde{\pi}, \tilde{\boldsymbol{\mu}})(D(\text{LP}_{\text{KP}}))$ is:

$$\tilde{\pi} = \frac{4}{3}, \quad \tilde{\mu}_1 = \frac{10}{3}, \quad \tilde{\mu}_2 = 1, \quad \tilde{\mu}_3 = \tilde{\mu}_4 = \tilde{\mu}_5 = \tilde{\mu}_6 = 0$$

The optimal value $z(D(\text{LP}_{\text{KP}}))$ is $\frac{49}{3}$.

8.4.1 Item efficiency

Definition 8.4: efficiency

The **efficiency** of an item $j \in \mathscr{I}$ is defined as the ratio between its profit and its weight:

$$\frac{p_j}{w_j} \tag{8.10}$$

- The efficiency of an item is a measure of its profitability per unit of weight: it indicates how much profit is gained for each unit of weight.

Example 8.6: item efficiencies

Let us consider the KP instance of Example 1. We have:

Items		$j=1$	$j=2$	$j=3$	$j=4$	$j=5$	$j=6$
Profits	p_j	6	5	8	9	6	7
Weights	w_j	2	3	6	7	5	9
Efficiencies	$\frac{p_j}{w_j}$	3	$\frac{5}{3}$	$\frac{4}{3}$	$\frac{9}{7}$	$\frac{6}{5}$	$\frac{7}{9}$

The items are sorted by non-increasing efficiency, since:

$$3 \geq \frac{5}{3} \geq \frac{4}{3} \geq \frac{9}{7} \geq \frac{6}{5} \geq \frac{7}{9}$$

Sorting the items by non-increasing efficiency requires a sorting algorithm with running time $O(n \log n)$, which is a polynomial function of the number n of items.

8.4.2 Split item and residual capacity

Definition 8.5: split item and residual capacity

Given a set $\mathscr{I} = \{1, 2, \ldots, n\}$ of n items ordered by non-increasing efficiency, the **split item** is the item

$$s \in \mathscr{I} \quad \text{such that} \quad \sum_{j=1}^{s-1} w_j \leq c \quad \text{and} \quad \sum_{j=1}^{s} w_j > c \tag{8.11}$$

and the **residual capacity** is

$$rc = c - \sum_{j=1}^{s-1} w_j \tag{8.12}$$

- The split item is the first one that does not fit entirely in the knapsack if all the previous ones are selected. It is also called **break item** or **critical item**, and it can be written as

$$s = \min \left\{ j \in \mathscr{I} : \sum_{k=1}^{j} w_k > c \right\}$$

- The residual capacity is the knapsack capacity remaining when all the items that precede the split item are selected. It satisfies $0 \leq rc < w_s$.

Example 8.7: split item and residual capacity

For the KP instance of Example 1, the split item is $s = 3$ since:

$$\sum_{j=1}^{2} w_j = 2 + 3 = 5 \leq 9 \quad \text{and} \quad \sum_{j=1}^{3} w_j = 2 + 3 + 6 = 11 > 9$$

and the residual capacity is:

$$rc = c - \sum_{j=1}^{2} w_j = 9 - 2 - 3 = 4$$

- The following proposition provides formulas to compute an optimal solution of the primal LP_{KP} model and of the dual $D(\text{LP}_{\text{KP}})$ model.

Proposition 8.6

If the items are sorted by non-increasing efficiency, an optimal solution of the primal $\mathrm{LP_{KP}}$ model is:

$$\tilde{x}_j = 1, \quad \forall j \in \{1, 2, \ldots, s-1\}, \qquad \tilde{x}_s = \frac{rc}{w_s} \quad \text{and} \quad \tilde{x}_j = 0, \quad \forall j \in \{s+1, s+2, \ldots, n\}, \tag{8.13}$$

and an optimal solution of the dual $D(\mathrm{LP_{KP}})$ model is:

$$\tilde{\pi} = \frac{p_s}{w_s}, \qquad \tilde{\mu}_j = p_j - w_j \frac{p_s}{w_s}, \ \forall j \in \{1, 2, \ldots, s-1\} \quad \text{and} \quad \tilde{\mu}_j = 0, \ \forall j \in \{s, s+1, \ldots, n\}. \tag{8.14}$$

Corollary 8.7

The optimal value of the primal $\mathrm{LP_{KP}}$ and dual $D(\mathrm{LP_{KP}})$ models is:

$$z(\mathrm{LP_{KP}}) = \sum_{j=1}^{s-1} p_j + p_s \frac{rc}{w_s} = z(D(\mathrm{LP_{KP}})) \tag{8.15}$$

- The proof proceeds in three steps:
  1. We show that (8.13) is a feasible solution for the primal $\mathrm{LP_{KP}}$ model;
  2. We show that (8.14) is a feasible solution for the dual $D(\mathrm{LP_{KP}})$ model;
  3. We show that both feasible solutions have the same objective function value.

  Therefore, by LP Strong Duality, (8.13) and (8.14) are a pair of optimal primal and dual solutions.

Proof.

1. We have:
$$\sum_{j \in \mathscr{I}} w_j \tilde{x}_j = \sum_{j=1}^{s-1} w_j + w_s \frac{rc}{w_s} = \sum_{j=1}^{s-1} w_j + c - \sum_{j=1}^{s-1} w_j = c$$
thus, constraint (8.2b) is satisfied.

Since $rc < w_s$, we have $0 \leq \tilde{x}_j \leq 1$, for every $j \in \mathscr{I}$. Accordingly, constraints (8.8a) and (8.8b) are satisfied.

Therefore, (8.13) is a feasible solution for the primal $\mathrm{LP_{KP}}$ model.

□

Proof.

2. For every $j \in \{1, 2, \ldots, s-1\}$, we have:

$$w_j\,\tilde{\pi} + \tilde{\mu}_j = w_j\,\frac{p_s}{w_s} + \left(p_j - w_j\,\frac{p_s}{w_s}\right) = p_j$$

and, for every $j \in \{s, s+1 \ldots, n\}$, we have:

$$w_j\,\tilde{\pi} + \tilde{\mu}_j = w_j\,\frac{p_s}{w_s} \geq p_j \qquad \text{since} \qquad \frac{p_s}{w_s} \geq \frac{p_j}{w_j}$$

thus constraints (8.9b) are satisfied.

We have $\tilde{\pi} \geq 0$. For every $j \in \{1, 2, \ldots, s-1\}$, we have:

$$p_j - w_j\,\frac{p_s}{w_s} \geq 0 \qquad \text{since} \qquad \frac{p_j}{w_j} \geq \frac{p_s}{w_s}$$

accordingly, $\tilde{\mu}_j \geq 0$ for every $j \in \mathscr{I}$. Thus constraint (8.9c) and constraints (8.9d) are satisfied.

Therefore, (8.14) is a feasible solution for the dual $D(\mathrm{LP_{KP}})$ model.

3. For the objective functions we have:

$$\begin{aligned}
\sum_{j \in \mathscr{I}} p_j\,\tilde{x}_j &= \sum_{j=1}^{s-1} p_j + p_s\,\frac{rc}{w_s} \\
&= \sum_{j=1}^{s-1} p_j + \frac{p_s}{w_s}\left(c - \sum_{j=1}^{s-1} w_j\right) \\
&= c\,\frac{p_s}{w_s} + \sum_{j=1}^{s-1}\left(p_j - w_j\,\frac{p_s}{w_s}\right) \\
&= c\,\tilde{\pi} + \sum_{j \in \mathscr{I}} \tilde{\mu}_j
\end{aligned}$$

thus the primal and dual objective function values are equal.

Therefore, by LP Strong Duality, (8.13) and (8.14) are a pair of optimal primal and dual solutions.

□

In the majority of KP instances the ratio between $z(\mathrm{LP_{KP}})$ and $z(\mathrm{ILP_{KP}})$ slightly exceeds one: $z(\mathrm{LP_{KP}})$ typically is a tight upper bound for $z(\mathrm{ILP_{KP}})$.

- We report an alternative proof for the formulas (8.13). We proceed by contradiction, proving that a different solution cannot be optimal.
- In this proof we assume that items with the same efficiency are merged into a single item. As we are allowed to choose any fraction of an item, this does not change the problem,

and we avoid discussing symmetric solutions.

Proof. Assume $\bar{\boldsymbol{x}} = (\bar{x}_1, \bar{x}_2, \ldots, \bar{x}_n)$ is an optimal solution of LP_{KP} different from $\tilde{\boldsymbol{x}} = (\tilde{x}_1, \tilde{x}_2, \ldots, \tilde{x}_n)$ determined with formulas (8.13). Every optimal solution of LP_{KP} has a total weight equal to c, i.e., constraint (8.2b) is satisfied at equality (otherwise it can be improved).

Assume that no pair of items $h < k$ exists with $\bar{x}_h < 1$ and $\bar{x}_k > 0$. Then $\bar{\boldsymbol{x}}$ selects the items in order of efficiency, filling the knapsack item by item up to the weight c: this is exactly $\tilde{\boldsymbol{x}}$, a contradiction. Hence such a pair of items $h < k$ exists (the case $k = s$ is included).

Let:

$$d = \min \left\{ \; w_k \, \bar{x}_k \; , \; w_h \, (1 - \bar{x}_h) \; \right\} > 0$$

Construct a new feasible solution $\hat{\boldsymbol{x}} = (\hat{x}_1, \hat{x}_2, \ldots, \hat{x}_n)$ which is identical to $\bar{\boldsymbol{x}}$ except that:

$$\hat{x}_k = \bar{x}_k - \frac{d}{w_k} \geq 0 \qquad \text{and} \qquad \hat{x}_h = \bar{x}_h + \frac{d}{w_h} \leq 1$$

Notice that $\hat{x}_k \geq 0$ and $\hat{x}_h \leq 1$ due to the definition of d, and $\hat{\boldsymbol{x}}$ is a feasible solution since:

$$\sum_{j \in \mathscr{I}} w_j \, \hat{x}_j = \sum_{j \in \mathscr{I}} w_j \, \bar{x}_j + w_h \, \frac{d}{w_h} - w_k \, \frac{d}{w_k} = \sum_{j \in \mathscr{I}} w_j \, \bar{x}_j = c$$

The objective function value of $\hat{\boldsymbol{x}}$ is:

$$\sum_{j \in \mathscr{I}} p_j \, \hat{x}_j = \sum_{j \in \mathscr{I}} p_j \, \bar{x}_j + d \underbrace{\left(\frac{p_h}{w_h} - \frac{p_k}{w_k} \right)}_{>0 \;\; \text{since} \;\; \frac{p_h}{w_h} > \frac{p_k}{w_k}} > \sum_{j \in \mathscr{I}} p_j \, \bar{x}_j$$

Then the objective function value of $\hat{\boldsymbol{x}}$ is larger than the objective function value of $\bar{\boldsymbol{x}}$, contradicting the fact that $\bar{\boldsymbol{x}}$ is an optimal solution.
The strict inequality $p_h/w_h > p_k/w_k$ holds for every $h < k$, since items with the same efficiency have been merged. □

Example 8.8: Primal and dual LP solutions

For the KP instance of Example 1, we have $s = 3$ and $rc = 4$.

Using formulas (8.13), we can compute the following optimal solution of the LP_{KP} model:

$$\tilde{x}_1 = \tilde{x}_2 = 1, \quad \tilde{x}_3 = \frac{2}{3}, \quad \tilde{x}_4 = \tilde{x}_5 = \tilde{x}_6 = 0$$

Using formulas (8.14), we can compute the following optimal solution of the $D(\text{LP}_{\text{KP}})$ model:

$$\tilde{\pi} = \frac{4}{3}, \qquad \tilde{\mu}_1 = 6 - 2\,\frac{4}{3} = \frac{10}{3}, \quad \tilde{\mu}_2 = 5 - 3\,\frac{4}{3} = 1, \quad \tilde{\mu}_3 = \tilde{\mu}_4 = \tilde{\mu}_5 = \tilde{\mu}_6 = 0$$

Using formula (8.15), the optimal value of both LP models, primal and dual, is:

$$\sum_{j=1}^{2} p_j + p_s \frac{rc}{w_s} = 6 + 5 + 8\,\frac{2}{3} = \frac{49}{3}$$

8.4.3 Some important theoretical properties

- The following two propositions show that the ratio between $z(\text{LP}_{\text{KP}})$ and $z(\text{ILP}_{\text{KP}})$ is at most two for all KP instances. Furthermore, there exist instances for which the value two can be asymptotically achieved as the capacity $c \to +\infty$.

Proposition 8.8

For every KP instance, we have:

$$\frac{z(\text{LP}_{\text{KP}})}{z(\text{ILP}_{\text{KP}})} \leq 2 \tag{8.16}$$

Proof. We consider two feasible solutions of the ILP_{KP} model:

1. The solution selecting all items preceding the split item:

   $$\bar{x}_j = 1, \quad \forall j \in \{1, 2, \ldots, s-1\} \quad \text{and} \quad \bar{x}_j = 0, \quad \forall j \in \{s, s+1, \ldots, n\}$$

   which is feasible by definition of the split item, and provides the valid lower bound:

   $$\sum_{j=1}^{s-1} p_j \leq z(\text{ILP}_{\text{KP}})$$

2. The solution selecting only the split item:

   $$\bar{x}_j = 0, \quad \forall j \in \mathscr{I} \setminus \{s\} \quad \text{and} \quad \bar{x}_s = 1$$

   which is feasible by assumption that $w_s \leq c$, and provides the valid lower bound:

   $$p_s \leq z(\text{ILP}_{\text{KP}})$$

Adding the two inequalities yields:

$$2\, z(\text{ILP}_{\text{KP}}) \geq \sum_{j=1}^{s-1} p_j + p_s \geq z(\text{LP}_{\text{KP}})$$

and thus:

$$\frac{z(\mathrm{LP_{KP}})}{z(\mathrm{ILP_{KP}})} \leq 2$$

□

Proposition 8.9

There are KP instances such that:

$$\frac{z(\mathrm{LP_{KP}})}{z(\mathrm{ILP_{KP}})} \to 2 \quad \text{as} \quad c \to +\infty \tag{8.17}$$

Proof. Given $\vartheta \in \mathbb{Z}_{\geq 1}$, consider the KP instance with knapsack capacity $c = 2\vartheta$ and $n = 2$ items having the following profits and weights:

$$p_1 = 1, \quad w_1 = \vartheta + 1 \qquad \text{and} \qquad p_2 = 1, \quad w_2 = \vartheta + 1$$

Since only one item can be selected, an optimal solution $\tilde{\boldsymbol{x}}(\mathrm{ILP_{KP}})$ of the $\mathrm{ILP_{KP}}$ model is:

$$\tilde{x}_1 = 1, \qquad \tilde{x}_2 = 0 \qquad \text{and} \qquad z(\mathrm{ILP_{KP}}) = 1$$

The split item is $s = 2$, and the residual capacity is:

$$rc = 2\vartheta - (\vartheta + 1) = \vartheta - 1$$

Accordingly, an optimal solution $\tilde{\boldsymbol{x}}(\mathrm{LP_{KP}})$ of the $\mathrm{LP_{KP}}$ model is:

$$\tilde{x}_1 = 1, \qquad \tilde{x}_2 = \frac{\vartheta - 1}{\vartheta + 1} \qquad \text{and} \qquad z(\mathrm{LP_{KP}}) = 1 + \frac{\vartheta - 1}{\vartheta + 1} = \frac{2\vartheta}{\vartheta + 1}$$

It follows that:

$$\lim_{\vartheta \to +\infty} \frac{z(\mathrm{LP_{KP}})}{z(\mathrm{ILP_{KP}})} = \lim_{\vartheta \to +\infty} \frac{\frac{2\vartheta}{\vartheta+1}}{1} = \lim_{\vartheta \to +\infty} \frac{2(\vartheta + 1) - 2}{\vartheta + 1} = \lim_{\vartheta \to +\infty} 2 - \underbrace{\frac{2}{\vartheta + 1}}_{\to 0} = 2$$

Accordingly, we conclude that:

$$\frac{z(\mathrm{LP_{KP}})}{z(\mathrm{ILP_{KP}})} \to 2 \qquad \text{as} \qquad c \to +\infty$$

□

8.5 Knapsack Problem upper bounds

- **The following two propositions require that the items are sorted by non-increasing efficiency**.

Proposition 8.10: Dantzig upper bound

For every KP instance, we have:

$$z(\mathrm{ILP_{KP}}) \leq \underbrace{\sum_{j=1}^{s-1} p_j + \left\lfloor p_s \frac{rc}{w_s} \right\rfloor}_{\mathrm{UB_D}} \tag{8.18}$$

and $\mathrm{UB_D}$ is called the **Dantzig upper bound** for the KP.

Proof. We have:

$$\lfloor z(\mathrm{LP_{KP}}) \rfloor = \left\lfloor \sum_{j=1}^{s-1} p_j + p_s \frac{rc}{w_s} \right\rfloor = \sum_{j=1}^{s-1} p_j + \left\lfloor p_s \frac{rc}{w_s} \right\rfloor$$

which is a valid upper bound for $z(\mathrm{ILP_{KP}})$ since $p_j \in \mathbb{Z}_{\geq 1}$, for every $j \in \mathscr{I}$. □

- The Martello-Toth bound uses the items $s-1$ and $s+1$. Since $w_1 \leq c$ and $\sum_j w_j > c$ we have $s \geq 2$, so the item $s-1$ always exists. If $s = n$ the item $s+1$ does not exist, and we set $p_{n+1}/w_{n+1} := 0$: in this case $\mathrm{UB_0} = \sum_{j=1}^{s-1} p_j$.

Proposition 8.11: Martello-Toth upper bound

For every KP instance, we have:

$$z(\mathrm{ILP_{KP}}) \leq \mathrm{UB_{MT}} = \max \left\{ \mathrm{UB_0}, \ \mathrm{UB_1} \right\} \tag{8.19}$$

where

$$\mathrm{UB_0} = \sum_{j=1}^{s-1} p_j + \left\lfloor rc \frac{p_{s+1}}{w_{s+1}} \right\rfloor$$

$$\mathrm{UB_1} = \sum_{j=1}^{s-1} p_j + \left\lfloor p_s - (w_s - rc) \frac{p_{s-1}}{w_{s-1}} \right\rfloor$$

and $\mathrm{UB_{MT}}$ is called the **Martello-Toth upper bound** for the KP.

Proof. Since x_s cannot take a fractional value, an optimal solution $\tilde{\boldsymbol{x}}(\text{ILP}_{\text{KP}})$ can be obtained from an optimal solution $\tilde{\boldsymbol{x}}(\text{LP}_{\text{KP}})$ either without the split item s or with the split item s and hence removing at least one of the first $s-1$ items.

1. If the split item is out, we have $z(\text{ILP}_{\text{KP}}) \leq \text{UB}_0$ since the residual capacity rc is filled with items having the largest possible efficiency $\frac{p_{s+1}}{w_{s+1}}$
2. If the split item is in, we have $z(\text{ILP}_{\text{KP}}) \leq \text{UB}_1$ since it is supposed that the item to be removed has exactly the minimum necessary weight $w_s - rc$ and the smallest possible efficiency $\frac{p_{s-1}}{w_{s-1}}$

□

Example 8.9: Dantzig and Martello-Toth upper bound

For the KP instance of Example 1, using formula (8.18), we can compute the Dantzig upper bound:

$$\text{UB}_{\text{D}} = \sum_{j=1}^{2} p_j + \left\lfloor p_3 \frac{rc}{w_3} \right\rfloor = 6 + 5 + \left\lfloor 8 \frac{2}{3} \right\rfloor = 11 + \left\lfloor \frac{16}{3} \right\rfloor = 16$$

Using formula (8.19), we can compute the Martello-Toth upper bound:

$$\text{UB}_0 = \sum_{j=1}^{2} p_j + \left\lfloor rc \frac{p_4}{w_4} \right\rfloor = 6 + 5 + \left\lfloor 4 \frac{9}{7} \right\rfloor = 11 + \left\lfloor \frac{36}{7} \right\rfloor = 16$$

$$\text{UB}_1 = \sum_{j=1}^{2} p_j + \left\lfloor p_3 - (w_3 - rc) \frac{p_2}{w_2} \right\rfloor = 6 + 5 + \left\lfloor 8 - (6-4) \frac{5}{3} \right\rfloor = 11 + \left\lfloor \frac{14}{3} \right\rfloor = 15$$

$$\text{UB}_{\text{MT}} = \max \left\{ \text{UB}_0, \ \text{UB}_1 \right\} = \max \left\{ 16, \ 15 \right\} = 16$$

For this KP instance, the Martello-Toth upper bound is as strong as the Dantzig upper bound.

8.6 Typical mistakes

Warning

- **Forgetting to sort the items.** The formulas (8.13), (8.14) and the bounds hold only if the items are sorted by non-increasing efficiency: with another order the vector (8.13) is feasible but not optimal.
- **Using z(LP-0$_{\text{KP}}$) as the bound.** The first relaxation has no upper bounds on the variables, and it is much weaker: in Example 1 it gives 27, against $\frac{49}{3}$ for the second relaxation.
- **Rounding when the profits are not integer.** The bounds UB_{D} and UB_{MT} round down the fractional part: this is valid only because all the profits are integer.
- **Confusing the split item with the last selected item.** The split item s is the first item that does not fit: in Example 1 it is $s = 3$, while the items 1 and 2 are selected entirely.
- **Assuming that the LP optimum is unique.** With items of equal efficiency the LP_{KP} problem has other optimal solutions; the formulas (8.13) give one of them.

8.7 In summary

Summary

- The ILP_{KP} model has one binary variable per item, one capacity constraint and the objective of maximizing the total profit.
- The relaxation LP_{KP} is solved by sorting the items by non-increasing efficiency: the items before the split item are selected, the split item is taken fractionally, the others are discarded.
- The dual of LP_{KP} has the optimal solution $\tilde{\pi} = p_s/w_s$, and $z(\text{LP}_{\text{KP}})$ is never larger than twice $z(\text{ILP}_{\text{KP}})$.
- Rounding down $z(\text{LP}_{\text{KP}})$ gives the Dantzig bound; the Martello-Toth bound considers separately the split item out and in.

Dashboard of Example 1. Upper bounds on $z(\text{ILP}_{\text{KP}})$ computed in this chapter:

Bound	Value	From
First relaxation	27	$z(\text{LP-0}_{\text{KP}})$, item 1 with $\frac{9}{2}$ copies
Second relaxation	$\frac{49}{3}$	$z(\text{LP}_{\text{KP}})$, split item $s = 3$
Dantzig bound	16	$\lfloor \frac{49}{3} \rfloor$
Martello-Toth bound	16	$\max\{16, 15\}$
Optimal value	15	solution $\{1, 4\}$ of the ILP_{KP} model

The gap between the best upper bound (16) and the optimal value (15) is closed by the exact algorithms of the following chapters.

Chapter 9

Formulations of the bin packing problem

Prerequisites

- From the module *Linear Programming*, Chapter 9 «Duality theory»: the dual of an LP problem, weak and strong duality.
- From the module *Integer linear programming*, the chapters «Integer linear programming» and «LP relaxation and dual bounds»: ILP problems, LP relaxations and bounds.
- From this module, the chapter «Bin packing and cutting stock»: the problem and its assumptions.

Objectives

At the end of this chapter you will be able to:

- write the ILP model of the Bin Packing Problem and explain the link between the assignment variables and the activation variables;
- write the two LP relaxations and their duals, and prove that their optimal value is $\sum_j w_j/c$;
- round the LP bound up to obtain a lower bound on the number of bins;
- prove that the LP bound is at least one half of the optimal value, and that the factor is tight.

9.1 The problem in brief

Where we were

In the chapter «Bin packing and cutting stock» we met the Bin Packing Problem as a *problem*: definition, applications and a first bound on the number of bins. Here we look at it as a *model* and we compute lower bounds on its optimal value. The same problem returns with heuristic algorithms and approximation algorithms for the BPP.

Definition 9.1: Bin Packing Problem

Given:

1. an infinite number of bins, each with *capacity* $c \in \mathbb{Z}_{\geq 1}$,
2. a set $\mathscr{I} = \{1, 2, \ldots, n\}$ of n items,
3. for each item $j \in \mathscr{I}$, a *weight* $w_j \in \mathbb{Z}_{\geq 1}$,

the **Bin Packing Problem** (BPP) asks to decide in which bin to place each item, so that the total weight of the items in any bin does not exceed the bin capacity, and the total number of bins used is minimized.

- We assume that any BPP instance satisfies the problem assumptions, i.e.,

$$w_j \leq c \quad \text{for all} \quad j \in \mathscr{I} \qquad \text{and} \qquad \sum_{j \in \mathscr{I}} w_j > c$$

The first assumption guarantees that every item can be packed into a bin, the second one that more than one bin is needed.

9.2 An ILP model for the Bin Packing Problem

- Given an upper bound $m \in \mathbb{Z}_{\geq 2}$ on the number of bins used in any optimal BPP solution, we introduce the following mn **binary variables**:

$$x_{jb} = \begin{cases} 1 & \text{if item } j \text{ is packed in bin } b \\ 0 & \text{otherwise} \end{cases} \qquad \forall j \in \underbrace{\{1, 2, \ldots, n\}}_{=\mathscr{I}}, \quad \forall b \in \underbrace{\{1, 2, \ldots, m\}}_{=\mathscr{B}},$$

and the following m **binary variables**:

$$y_b = \begin{cases} 1 & \text{if bin } b \text{ is used} \\ 0 & \text{otherwise} \end{cases} \qquad \forall b \in \underbrace{\{1, 2, \ldots, m\}}_{=\mathscr{B}}.$$

The number n of items is a valid upper bound on the minimum number of bins needed to pack all the items since, in the worst possible case, each item is packed in a different bin. Also the number of bins used in any feasible BPP solution is a valid upper bound.

Using these variables, an Integer Linear Programming (ILP) model for the Bin Packing Problem (BPP) reads as follows:

$$\min \quad \sum_{b \in \mathscr{B}} y_b \tag{9.1a}$$

$$\text{subject to} \quad \sum_{b \in \mathscr{B}} x_{jb} = 1, \qquad \forall j \in \mathscr{I}, \tag{9.1b}$$

$$-\sum_{j \in \mathscr{I}} w_j \, x_{jb} + c \, y_b \geq 0, \qquad \forall b \in \mathscr{B}, \tag{9.1c}$$

$$x_{jb} \in \{0,1\}, \qquad \forall j \in \mathscr{I}, \ \forall b \in \mathscr{B}, \tag{9.1d}$$

$$y_b \in \{0,1\}, \qquad \forall b \in \mathscr{B}. \tag{9.1e}$$

This ILP model is referred to as the ILP_{BPP} model.

- Description of the objective function and constraints of the ILP_{BPP} model:
  - the linear objective function (9.1a) minimizes the total number of used bins;
  - constraints (9.1b) ensure that each item is packed in exactly one bin, so that all items are packed (n linear constraints);
  - constraints (9.1c) link the assignment variables with the activation variables and impose the capacity restrictions: if bin b is used, the weight of its items is at most c; if it is not used, it contains no item (m linear constraints);
  - finally, constraints (9.1d) and (9.1e) define the variables of the ILP model.

Explanation of the link between the variables

- **If at least one item is packed in bin b, then the bin is used.** The logical implication is

  $$(x_{1b} \ \texttt{OR} \ x_{2b} \ \texttt{OR} \ \ldots \ \texttt{OR} \ x_{nb}) \quad \Longrightarrow \quad y_b.$$

  Since $w_j > 0$, packing an item in bin b gives $\sum_{j \in \mathscr{I}} w_j x_{jb} > 0$, and the constraint $\sum_{j \in \mathscr{I}} w_j x_{jb} \leq c \, y_b$ forces $y_b = 1$.

- **If bin b is not used, then it is empty.** This is the contrapositive of the previous implication. For $y_b = 0$ the constraint reads $\sum_{j \in \mathscr{I}} w_j x_{jb} \leq 0$, and since $w_j > 0$ we obtain $x_{jb} = 0$ for every item j.

- **If bin b is used, then it contains at least one item, but only in an optimal solution.** The constraints do not impose it. However, if $x_{jb} = 0$ for all j and $y_b = 1$, setting $y_b = 0$ keeps all the constraints satisfied and decreases the objective value by one. Hence, in every optimal solution, empty bins are not counted as used.

Example 9.1: ILP_{BPP} model

We consider a bin of capacity $c = 10$ and a set $\mathscr{I} = \{1, 2, 3\}$ of 3 items ($n = 3$), with the following item weights:

	$j = 1$	$j = 2$	$j = 3$
w_j	7	6	3

We refer to this BPP instance as Example 1.

The ILP_{BPP} model with $m = 3$ for this BPP instance is:

$$
\begin{aligned}
\min \quad & y_1 + y_2 + y_3 \\
\text{subject to} \quad & x_{11} + x_{12} + x_{13} = 1 \\
& x_{21} + x_{22} + x_{23} = 1 \\
& x_{31} + x_{32} + x_{33} = 1 \\
& -7x_{11} - 6x_{21} - 3x_{31} + 10y_1 \geq 0 \\
& -7x_{12} - 6x_{22} - 3x_{32} + 10y_2 \geq 0 \\
& -7x_{13} - 6x_{23} - 3x_{33} + 10y_3 \geq 0 \\
& x_{11}, x_{12}, x_{13}, x_{21}, x_{22}, x_{23}, x_{31}, x_{32}, x_{33}, y_1, y_2, y_3 \in \{0, 1\}
\end{aligned}
$$

An optimal solution $\tilde{\boldsymbol{x}}(\text{ILP}_{\text{BPP}})$ is:

$\tilde{x}_{11} = \tilde{x}_{31} = 1,\ \tilde{x}_{12} = \tilde{x}_{13} = \tilde{x}_{21} = 0,\ \tilde{x}_{22} = 1,\ \tilde{x}_{23} = \tilde{x}_{32} = \tilde{x}_{33} = 0,\ \tilde{y}_1 = \tilde{y}_2 = 1,\ \tilde{y}_3 = 0$

Another optimal solution, obtained by permuting the indices of bins 1 and 2, is:

$\tilde{x}_{12} = \tilde{x}_{32} = 1,\ \tilde{x}_{11} = \tilde{x}_{13} = \tilde{x}_{22} = 0,\ \tilde{x}_{21} = 1,\ \tilde{x}_{23} = \tilde{x}_{31} = \tilde{x}_{33} = 0,\ \tilde{y}_1 = \tilde{y}_2 = 1,\ \tilde{y}_3 = 0.$

The optimal value is:

$$z(\text{ILP}_{\text{BPP}}) = 2$$

The associated optimal solution of the BPP is given by the packing patterns

$$\tilde{P}_1 = \{1, 3\} \quad \text{and} \quad \tilde{P}_2 = \{2\}$$

The optimal solution is not unique: also the packing patterns $\{1\}$ and $\{2, 3\}$ use two bins.

9.3 A first Linear Programming relaxation

To obtain a Linear Programming (LP) relaxation of the ILP_{BPP} model, we relax constraints (9.1d) and (9.1e) with the following ones:

$$x_{jb} \geq 0, \qquad \forall j \in \mathscr{I},\ \forall b \in \mathscr{B}, \tag{9.2a}$$

$$y_b \geq 0, \qquad \forall b \in \mathscr{B}. \tag{9.2b}$$

The obtained LP model is referred to as the LP-0_{BPP} model.

Example 9.2: LP-0_{BPP} model

The LP-0_{BPP} model for the BPP instance of Example 1 is:

$$\begin{aligned}
\min \quad & y_1 + y_2 + y_3 \\
\text{subject to} \quad & x_{11} + x_{12} + x_{13} = 1 \\
& x_{21} + x_{22} + x_{23} = 1 \\
& x_{31} + x_{32} + x_{33} = 1 \\
& -7x_{11} - 6x_{21} - 3x_{31} + 10y_1 \geq 0 \\
& -7x_{12} - 6x_{22} - 3x_{32} + 10y_2 \geq 0 \\
& -7x_{13} - 6x_{23} - 3x_{33} + 10y_3 \geq 0 \\
& x_{11}, x_{12}, x_{13}, x_{21}, x_{22}, x_{23}, x_{31}, x_{32}, x_{33}, y_1, y_2, y_3 \geq 0
\end{aligned}$$

An optimal solution $\tilde{\boldsymbol{x}}(\text{LP-}0_{\text{BPP}})$ is:

$$\tilde{x}_{11} = \tilde{x}_{12} = \tilde{x}_{13} = \tilde{x}_{21} = \tilde{x}_{22} = \tilde{x}_{23} = \tilde{x}_{31} = \tilde{x}_{32} = \tilde{x}_{33} = \frac{1}{3}, \quad \tilde{y}_1 = \tilde{y}_2 = \tilde{y}_3 = \frac{8}{15}$$

The optimal value $z(\text{LP-}0_{\text{BPP}})$ is $\frac{8}{5}$.

Accordingly, we have the lower bound:

$$\frac{8}{5} \leq z(\text{ILP}_{\text{BPP}})$$

and, since $z(\text{ILP}_{\text{BPP}})$ is an integer, $z(\text{ILP}_{\text{BPP}}) \geq \lceil \frac{8}{5} \rceil = 2$.

Using

- a free dual variable μ_j for each constraint (9.1b) ($j \in \mathscr{I}$)
- a non-negative dual variable π_b for each constraint (9.1c) ($b \in \mathscr{B}$)

the dual model of the LP-0_{BPP} model reads as follows:

$$\max \sum_{j \in \mathscr{I}} \mu_j \tag{9.3a}$$

$$\text{subject to} \quad \mu_j - w_j \, \pi_b \leq 0, \qquad \forall j \in \mathscr{I}, \ \forall b \in \mathscr{B}, \tag{9.3b}$$

$$c \, \pi_b \leq 1, \qquad \forall b \in \mathscr{B}, \tag{9.3c}$$

$$\mu_j \gtreqless 0, \qquad \forall j \in \mathscr{I}, \tag{9.3d}$$

$$\pi_b \geq 0, \qquad \forall b \in \mathscr{B}. \tag{9.3e}$$

The obtained model is referred to as the dual D(LP-0_{BPP}) model.

Example 9.3: D(LP-0_{BPP}) model

The dual D(LP-0_{BPP}) model for the BPP instance of Example 1 is:

$$\begin{array}{rccccccccc}
\max & \mu_1 & + \mu_2 & + \mu_3 & & & & & & \\
\text{subject to} & \mu_1 & & & - 7\pi_1 & & & & & \leq 0 \\
& \mu_1 & & & & - 7\pi_2 & & & & \leq 0 \\
& \mu_1 & & & & & - 7\pi_3 & & & \leq 0 \\
& & \mu_2 & & - 6\pi_1 & & & & & \leq 0 \\
& & \mu_2 & & & - 6\pi_2 & & & & \leq 0 \\
& & \mu_2 & & & & - 6\pi_3 & & & \leq 0 \\
& & & \mu_3 & - 3\pi_1 & & & & & \leq 0 \\
& & & \mu_3 & & - 3\pi_2 & & & & \leq 0 \\
& & & \mu_3 & & & - 3\pi_3 & & & \leq 0 \\
& & & & 10\pi_1 & & & & & \leq 1 \\
& & & & & 10\pi_2 & & & & \leq 1 \\
& & & & & & 10\pi_3 & & & \leq 1 \\
& \mu_1, & \mu_2, & \mu_3 & & & & & & \gtreqless 0 \\
& & & & \pi_1, & \pi_2, & \pi_3 & & & \geq 0
\end{array}$$

An optimal solution $(\tilde{\boldsymbol{\mu}}, \tilde{\boldsymbol{\pi}})$ of D(LP-0_{BPP}) is:

$$\tilde{\mu}_1 = \frac{7}{10}, \quad \tilde{\mu}_2 = \frac{3}{5}, \quad \tilde{\mu}_3 = \frac{3}{10}, \quad \tilde{\pi}_1 = \tilde{\pi}_2 = \tilde{\pi}_3 = \frac{1}{10}$$

The optimal value $z(D(\text{LP-}0_{\text{BPP}}))$ is $\frac{8}{5}$.

Proposition 9.2

An optimal solution of the primal LP-0_{BPP} model is:

$$\begin{aligned} \tilde{x}_{jb} &= \frac{1}{m}, && \forall j \in \mathscr{I},\ \forall b \in \mathscr{B}, \\ \tilde{y}_b &= \frac{\sum_{j \in \mathscr{I}} w_j}{m\, c}, && \forall b \in \mathscr{B}, \end{aligned} \tag{9.4}$$

and an optimal solution of the dual $D(\text{LP-}0_{\text{BPP}})$ model is:

$$\begin{aligned} \tilde{\mu}_j &= \frac{w_j}{c}, && \forall j \in \mathscr{I}, \\ \tilde{\pi}_b &= \frac{1}{c}, && \forall b \in \mathscr{B}. \end{aligned} \tag{9.5}$$

Corollary 9.3

The optimal value of the primal LP-0_{BPP} and dual $D(\text{LP-}0_{\text{BPP}})$ models is:

$$z(\text{LP-}0_{\text{BPP}}) = \frac{\sum_{j \in \mathscr{I}} w_j}{c} = z(D(\text{LP-}0_{\text{BPP}})) \tag{9.6}$$

- The proof proceeds in three steps:
  1. We show that (9.4) is a feasible solution for the primal LP-0_{BPP} model;
  2. We show that (9.5) is a feasible solution for the dual $D(\text{LP-}0_{\text{BPP}})$ model;
  3. We show that both feasible solutions have the same objective function value.

  Therefore, by LP Weak Duality, (9.4) and (9.5) are a pair of optimal primal and dual solutions.

Proof.

1. For every $j \in \mathscr{I}$, we have:

$$\sum_{b \in \mathscr{B}} \tilde{x}_{jb} = \sum_{b \in \mathscr{B}} \frac{1}{m} = \frac{m}{m} = 1$$

   thus constraints (9.1b) are satisfied.

   For every $b \in \mathscr{B}$, we have:

$$-\sum_{j \in \mathscr{I}} w_j\, \tilde{x}_{jb} + c\, \tilde{y}_b = -\sum_{j \in \mathscr{I}} w_j\, \frac{1}{m} + c\, \frac{\sum_{j \in \mathscr{I}} w_j}{m\, c} = 0$$

   thus constraints (9.1c) are satisfied.

   We have $\tilde{x}_{jb} \geq 0$ and $\tilde{y}_b \geq 0$, for every $j \in \mathscr{I}$ and every $b \in \mathscr{B}$, thus constraints (9.2a) and (9.2b) are satisfied.

   Therefore, (9.4) provides a feasible solution for the primal LP-0_{BPP} model.

2. For every $b \in \mathscr{B}$ and every $j \in \mathscr{I}$, we have:

$$\tilde{\mu}_j - w_j \, \tilde{\pi}_b = \frac{w_j}{c} - \frac{w_j}{c} = 0$$

thus constraints (9.3b) are satisfied.

For every $b \in \mathscr{B}$, we have:

$$c \, \tilde{\pi}_b = \frac{c}{c} = 1$$

thus constraints (9.3c) are satisfied.

We have $\tilde{\mu}_j \geq 0$ and $\tilde{\pi}_b \geq 0$, for every $b \in \mathscr{B}$ and every $j \in \mathscr{I}$, thus constraints (9.3d) and (9.3e) are satisfied.

Therefore, (9.5) provides a feasible solution for the dual $D(\text{LP-0}_{\text{BPP}})$ model.

3. For the objective functions, we have:

$$\sum_{b \in \mathscr{B}} \tilde{y}_b = \sum_{b \in \mathscr{B}} \frac{\sum_{j \in \mathscr{I}} w_j}{m \, c} = \frac{\sum_{j \in \mathscr{I}} w_j}{c} = \sum_{j \in \mathscr{I}} \tilde{\mu}_j$$

thus the primal and dual objective function values are equal.

Therefore, by LP Weak Duality, (9.4) and (9.5) are a pair of optimal primal and dual solutions.

□

- This is an alternative proof that does not rely on the dual formulation but instead reasons directly on the minimum objective function value imposed by the constraints.

Proposition 9.4

An optimal solution of the primal LP-0_{BPP} model is:

$$\begin{aligned} \tilde{x}_{jb} &= \frac{1}{m}, && \forall j \in \mathscr{I}, \, \forall b \in \mathscr{B}, \\ \tilde{y}_b &= \frac{\sum_{j \in \mathscr{I}} w_j}{m \, c}, && \forall b \in \mathscr{B}, \end{aligned} \tag{9.7}$$

and the optimal value is

$$z(\text{LP-0}_{\text{BPP}}) = \frac{\sum_{j \in \mathscr{I}} w_j}{c}. \tag{9.8}$$

Proof. Summing the capacity constraints (9.1c) over all $b \in \mathscr{B}$ gives

$$c \sum_{b \in \mathscr{B}} y_b \geq \sum_{b \in \mathscr{B}} \sum_{j \in \mathscr{I}} w_j x_{jb} = \sum_{j \in \mathscr{I}} w_j \underbrace{\sum_{b \in \mathscr{B}} x_{jb}}_{\text{by (9.1b)}=1} = \sum_{j \in \mathscr{I}} w_j,$$

hence

$$\sum_{b \in \mathscr{B}} y_b \geq \frac{1}{c} \sum_{j \in \mathscr{I}} w_j. \tag{9.9}$$

Therefore every feasible solution of LP-0_{BPP} satisfies (9.9).

Now consider the point defined in (9.7). For every $j \in \mathscr{I}$, we have

$$\sum_{b \in \mathscr{B}} \tilde{x}_{jb} = m \, \tfrac{1}{m} = 1,$$

so constraints (9.1b) hold. Moreover, for every $b \in \mathscr{B}$, we have

$$\sum_{j \in \mathscr{I}} w_j \tilde{x}_{jb} = \sum_{j \in \mathscr{I}} w_j \, \tfrac{1}{m} = \frac{1}{m} \sum_{j \in \mathscr{I}} w_j = c \, \tilde{y}_b,$$

so constraints (9.1c) hold with equality. Finally, constraints (9.2a) and (9.2b) are satisfied by construction. Hence the point defined in (9.7) is a feasible solution for LP-0_{BPP}.

The objective value of (9.7) is

$$\sum_{b \in \mathscr{B}} \tilde{y}_b = m \, \frac{\sum_{j \in \mathscr{I}} w_j}{m \, c} = \frac{1}{c} \sum_{j \in \mathscr{I}} w_j,$$

which matches the lower bound (9.9). Therefore the point (9.7) is an optimal solution with objective value (9.8). □

9.4 A second Linear Programming relaxation

To obtain a Linear Programming (LP) relaxation of the ILP_{BPP} model, we relax constraints (9.1d) and (9.1e) with the following ones:

$$x_{jb} \geq 0, \qquad \forall j \in \mathscr{I}, \, \forall b \in \mathscr{B}, \tag{9.10a}$$

$$y_b \leq 1, \qquad \forall b \in \mathscr{B}, \tag{9.10b}$$

$$y_b \geq 0, \qquad \forall b \in \mathscr{B}. \tag{9.10c}$$

The obtained LP model is referred to as the LP_{BPP} model.

Example 9.4: LP_{BPP} model

The LP_{BPP} model for the BPP instance of Example 1 is:

$$
\begin{array}{lrrr

$$\pi_b \geq 0, \qquad \forall b \in \mathscr{B}, \tag{9.11e}$$

$$\lambda_b \leq 0, \qquad \forall b \in \mathscr{B}. \tag{9.11f}$$

This LP model is referred to as the dual $D(\text{LP}_{\text{BPP}})$ model.

Example 9.5: $D(\text{LP}_{\text{BPP}})$ model

The $D(\text{LP}_{\text{BPP}})$ model of Example 1 is:

$$\begin{array}{rrrrrrrrrrrr}
\max & \mu_1 & +\mu_2 & +\mu_3 & & & & \lambda_1 & +\lambda_2 & +\lambda_3 & \\
\text{subject to} & \mu_1 & & & -7\pi_1 & & & & & & \leq 0 \\
& \mu_1 & & & & -7\pi_2 & & & & & \leq 0 \\
& \mu_1 & & & & & -7\pi_3 & & & & \leq 0 \\
& & \mu_2 & & -6\pi_1 & & & & & & \leq 0 \\
& & \mu_2 & & & -6\pi_2 & & & & & \leq 0 \\
& & \mu_2 & & & & -6\pi_3 & & & & \leq 0 \\
& & & \mu_3 & -3\pi_1 & & & & & & \leq 0 \\
& & & \mu_3 & & -3\pi_2 & & & & & \leq 0 \\
& & & \mu_3 & & & -3\pi_3 & & & & \leq 0 \\
& & & & 10\pi_1 & & & +\lambda_1 & & & \leq 1 \\
& & & & & 10\pi_2 & & & +\lambda_2 & & \leq 1 \\
& & & & & & 10\pi_3 & & & +\lambda_3 & \leq 1 \\
& \mu_1, & \mu_2, & \mu_3 & & & & & & & \gtreqless 0 \\
& & & & \pi_1, & \pi_2, & \pi_3 & & & & \geq 0 \\
& & & & & & & \lambda_1, & \lambda_2, & \lambda_3 & \leq 0
\end{array}$$

The optimal value $z(D(\text{LP}_{\text{BPP}}))$ is $\frac{8}{5}$. An optimal solution $(\tilde{\boldsymbol{\mu}}, \tilde{\boldsymbol{\pi}}, \tilde{\boldsymbol{\lambda}})$ of $D(\text{LP}_{\text{BPP}})$ is:

$$\tilde{\mu}_1 = \frac{7}{10}, \quad \tilde{\mu}_2 = \frac{3}{5}, \quad \tilde{\mu}_3 = \frac{3}{10}, \quad \tilde{\pi}_1 = \tilde{\pi}_2 = \tilde{\pi}_3 = \frac{1}{10}, \quad \tilde{\lambda}_1 = \tilde{\lambda}_2 = \tilde{\lambda}_3 = 0$$

- The constraints:

  $$x_{jb} \leq 1, \qquad \forall j \in \mathscr{I},\ \forall b \in \mathscr{B},$$

  are also valid since, for every $j \in \mathscr{I}$ and every $b \in \mathscr{B}$, we have:

  $$x_{jb} \in \{0, 1\} \qquad \Rightarrow \qquad x_{jb} \leq 1$$

  but they are implied by constraints (9.1b).

Proposition 9.5

An optimal solution of the primal LP_{BPP} model is:

$$\begin{aligned} \tilde{x}_{jb} &= \frac{1}{m}, && \forall j \in \mathscr{I},\ \forall b \in \mathscr{B}, \\ \tilde{y}_b &= \frac{\sum_{j \in \mathscr{I}} w_j}{m\, c}, && \forall b \in \mathscr{B}, \end{aligned} \tag{9.12}$$

and an optimal solution of the dual $D(\text{LP}_{\text{BPP}})$ model is:

$$\begin{aligned} \tilde{\mu}_j &= \frac{w_j}{c}, && \forall j \in \mathscr{I}, \\ \tilde{\pi}_b &= \frac{1}{c}, && \forall b \in \mathscr{B}, \\ \tilde{\lambda}_b &= 0, && \forall b \in \mathscr{B}. \end{aligned} \tag{9.13}$$

Corollary 9.6

The optimal value of the primal LP_{BPP} and dual $D(\text{LP}_{\text{BPP}})$ models is:

$$z(\text{LP}_{\text{BPP}}) = \frac{\sum_{j \in \mathscr{I}} w_j}{c} = z(D(\text{LP}_{\text{BPP}})) \tag{9.14}$$

Proof. The proof is similar to the previous one. In addition, it is necessary to observe that

$$\tilde{y}_b = \frac{\sum_{j \in \mathscr{I}} w_j}{m\, c} \leq 1, \quad \forall b \in \mathscr{B},$$

since $\sum_j w_j / c \leq z(\text{ILP}_{\text{BPP}}) \leq m$, where the last inequality holds because m is an upper bound on the number of bins in any optimal BPP solution. Accordingly, constraints (9.10b) are also satisfied.

□

9.4.1 Some important theoretical properties

- The following two propositions show that the ratio between $z(\mathrm{LP}_{\mathrm{BPP}})$ and $z(\mathrm{ILP}_{\mathrm{BPP}})$ is at least one half for every BPP instance (**by hypothesis** $z(\mathrm{ILP}_{\mathrm{BPP}}) > 1$ since all items do not fit in a single bin). Furthermore, there exist instances for which the value one half can be asymptotically achieved as the capacity $c \to +\infty$.

Proposition 9.7

For every BPP instance, we have:

$$\frac{z(\mathrm{LP}_{\mathrm{BPP}})}{z(\mathrm{ILP}_{\mathrm{BPP}})} \geq \frac{1}{2} \tag{9.15}$$

Proof. Consider an optimal solution $\tilde{\boldsymbol{x}}(\mathrm{ILP}_{\mathrm{BPP}})$ of value $z = z(\mathrm{ILP}_{\mathrm{BPP}})$ (number of used bins). We can assume, without loss of generality, that bins $1, 2, \ldots, z$ are the non-empty ones.

In an optimal solution, no two bins can have total item weight at most $c/2$, since otherwise their contents could be merged into a single bin without exceeding the bin capacity c (contradicting optimality).

If all non-empty bins have total item weight greater than $c/2$, then

$$\sum_{j\in\mathscr{I}} w_j = \sum_{b=1}^{z}\sum_{j\in\mathscr{I}} w_j\,\tilde{x}_{jb} > z\,\frac{c}{2} \quad\Longrightarrow\quad \frac{\frac{\sum_{j\in\mathscr{I}} w_j}{c}}{z} > \frac{1}{2} \quad\Longrightarrow\quad \frac{z(\mathrm{LP}_{\mathrm{BPP}})}{z(\mathrm{ILP}_{\mathrm{BPP}})} \geq \frac{1}{2}.$$

Otherwise, suppose that there exists a bin with total item weight at most $c/2$. Renumbering bins if necessary, let it be bin z, so that

$$\sum_{j\in\mathscr{I}} w_j\,\tilde{x}_{jz} \leq \frac{c}{2}.$$

Since $z > 1$, there exists another bin say bin 1; by optimality, we must have

$$\sum_{j\in\mathscr{I}} w_j\,\tilde{x}_{j1} + \sum_{j\in\mathscr{I}} w_j\,\tilde{x}_{jz} \;>\; c,$$

and, by optimality, every bin in $\{2, 3, \ldots, z-1\}$ has total item weight $> c/2$. Hence,

$$\sum_{j\in\mathscr{I}} w_j = \sum_{b=1}^{z}\sum_{j\in\mathscr{I}} w_j\,\tilde{x}_{jb} = \underbrace{\sum_{b=2}^{z-1}\sum_{j\in\mathscr{I}} w_j\,\tilde{x}_{jb}}_{>\frac{c}{2}(z-2)} + \underbrace{\sum_{j\in\mathscr{I}} w_j\,\tilde{x}_{j1} + \sum_{j\in\mathscr{I}} w_j\,\tilde{x}_{jz}}_{>c}$$

$$> \frac{c}{2}(z-2) + c = z\,\frac{c}{2} \quad\Longrightarrow\quad \frac{\frac{\sum_{j\in\mathscr{I}} w_j}{c}}{z} > \frac{1}{2} \quad\Longrightarrow\quad \frac{z(\mathrm{LP}_{\mathrm{BPP}})}{z(\mathrm{ILP}_{\mathrm{BPP}})} \geq \frac{1}{2}.$$

□

Proposition 9.8

There are BPP instances such that:

$$\frac{z(\mathrm{LP}_{\mathrm{BPP}})}{z(\mathrm{ILP}_{\mathrm{BPP}})} \to \frac{1}{2} \quad \text{as} \quad c \to +\infty \tag{9.16}$$

Proof. Given $\vartheta \in \mathbb{Z}_{\geq 1}$, consider the BPP instance with bin capacity $c = 2\vartheta$ and n items having the following weights:

$$w_j = \vartheta + 1, \quad \forall j \in \mathscr{J}.$$

No bin can contain more than one item, thus each item must be packed alone in its own bin. Therefore, the BPP optimal value is

$$z(\mathrm{ILP}_{\mathrm{BPP}}) = n.$$

For the LP relaxation we have:

$$z(\mathrm{LP}_{\mathrm{BPP}}) = \frac{n(\vartheta+1)}{2\,\vartheta}.$$

It follows that:

$$\lim_{\vartheta \to +\infty} \frac{z(\mathrm{LP}_{\mathrm{BPP}})}{z(\mathrm{ILP}_{\mathrm{BPP}})} = \lim_{\vartheta \to +\infty} \frac{\frac{n(\vartheta+1)}{2\,\vartheta}}{n} = \lim_{\vartheta \to +\infty} \left(\frac{1}{2} + \underbrace{\frac{1}{2\,\vartheta}}_{\to 0} \right) = \frac{1}{2}.$$

Accordingly, we conclude that:

$$\frac{z(\mathrm{LP}_{\mathrm{BPP}})}{z(\mathrm{ILP}_{\mathrm{BPP}})} \to \frac{1}{2} \quad \text{as} \quad c \to +\infty.$$

□

- The following two propositions show that the ratio between $\lceil z(\mathrm{LP}_{\mathrm{BPP}}) \rceil$ and $z(\mathrm{ILP}_{\mathrm{BPP}})$ is at least one half for every BPP instance. Furthermore, there exist instances for which the value one half can be asymptotically achieved as the number of items $n \to +\infty$.

Proposition 9.9

For every BPP instance, we have:

$$\frac{\lceil z(\mathrm{LP}_{\mathrm{BPP}}) \rceil}{z(\mathrm{ILP}_{\mathrm{BPP}})} \geq \frac{1}{2} \tag{9.17}$$

Proof. Consider an optimal solution $\tilde{\boldsymbol{x}}(\mathrm{ILP}_{\mathrm{BPP}})$ of value $z = z(\mathrm{ILP}_{\mathrm{BPP}})$ (number of used bins). We can assume, without loss of generality, that bins $1, 2, \ldots, z$ are the non-empty

ones.

In an optimal solution, no two bins can have total item weight at most $c/2$, since otherwise their contents could be merged into a single bin without exceeding the bin capacity c (contradicting optimality).

If all non-empty bins have total item weight greater than $c/2$, then

$$\sum_{j\in\mathscr{I}} w_j = \sum_{b=1}^{z}\sum_{j\in\mathscr{I}} w_j\,\tilde{x}_{jb} > \frac{z\,c}{2} \quad\Longrightarrow\quad \frac{\sum_{j\in\mathscr{I}} w_j}{c} > \frac{z}{2} \quad\Longrightarrow\quad \left\lceil\frac{\sum_{j\in\mathscr{I}} w_j}{c}\right\rceil \geq \left\lceil\frac{z}{2}\right\rceil \geq \frac{z}{2}$$

$$\Longrightarrow\quad \frac{\left\lceil\frac{\sum_{j\in\mathscr{I}} w_j}{c}\right\rceil}{z} \geq \frac{1}{2} \quad\Longrightarrow\quad \frac{\lceil z(\mathrm{LP_{BPP}})\rceil}{z(\mathrm{ILP_{BPP}})} \geq \frac{1}{2}.$$

Otherwise, suppose that there exists a bin with total item weight at most $c/2$. Renumbering bins if necessary, let it be bin z. Then we have:

$$\sum_{j\in\mathscr{I}} w_j = \sum_{b=1}^{z}\sum_{j\in\mathscr{I}} w_j\tilde{x}_{jb} > \sum_{b=1}^{z-1}\sum_{j\in\mathscr{I}} w_j\tilde{x}_{jb} > (z-1)\frac{c}{2} \quad\Longrightarrow\quad z < 2\sum_{j\in\mathscr{I}}\frac{w_j}{c} + 1.$$

Since z must be an integer, this implies:

$$z \leq \left\lceil 2\sum_{j\in\mathscr{I}}\frac{w_j}{c}\right\rceil \leq 2\left\lceil\sum_{j\in\mathscr{I}}\frac{w_j}{c}\right\rceil.$$

Therefore:

$$\frac{\left\lceil\sum_{j\in\mathscr{I}}\frac{w_j}{c}\right\rceil}{z} \geq \frac{1}{2} \quad\Longrightarrow\quad \frac{\lceil z(\mathrm{LP_{BPP}})\rceil}{z(\mathrm{ILP_{BPP}})} \geq \frac{1}{2}.$$

□

Proposition 9.10

There are BPP instances such that:

$$\frac{\lceil z(\mathrm{LP_{BPP}})\rceil}{z(\mathrm{ILP_{BPP}})} \to \frac{1}{2} \quad \text{as} \quad n\to+\infty \tag{9.18}$$

Proof. Given $n\in\mathbb{Z}_{\geq 2}$ (take n even), consider the BPP instance with bin capacity $c=n$ and n items having the following weights:

$$w_j = \frac{c}{2}+1 = \frac{n}{2}+1, \qquad \forall j\in\mathscr{I}.$$

No bin can contain more than one item, since $w_j > c/2$ for every j. Thus each item must be packed alone in its own bin. Therefore, the BPP optimal value is

$$z(\mathrm{ILP_{BPP}}) = n.$$

For the LP relaxation we have:

$$z(\mathrm{LP}_{\mathrm{BPP}}) = \frac{\sum_{j \in \mathscr{I}} w_j}{c} = \frac{n\left(\frac{n}{2}+1\right)}{n} = \frac{n}{2}+1,$$

hence $\lceil z(\mathrm{LP}_{\mathrm{BPP}}) \rceil = \frac{n}{2}+1$ (since n is even). It follows that:

$$\lim_{n \to +\infty} \frac{\lceil z(\mathrm{LP}_{\mathrm{BPP}}) \rceil}{z(\mathrm{ILP}_{\mathrm{BPP}})} = \lim_{n \to +\infty} \frac{\frac{n}{2}+1}{n} = \lim_{n \to +\infty} \left(\frac{1}{2} + \underbrace{\frac{1}{n}}_{\to 0} \right) = \frac{1}{2}.$$

Accordingly, we conclude that:

$$\frac{\lceil z(\mathrm{LP}_{\mathrm{BPP}}) \rceil}{z(\mathrm{ILP}_{\mathrm{BPP}})} \to \frac{1}{2} \qquad \text{as} \qquad n \to +\infty.$$

□

9.5 Typical mistakes

Warning

- **Choosing m too small.** The number m of bins must be an upper bound on the number of bins of an optimal solution: with $m < z(\mathrm{ILP}_{\mathrm{BPP}})$ the model has no feasible solution. The value $m = n$ is always valid.
- **Forgetting to round up.** The optimal value of the $\mathrm{ILP}_{\mathrm{BPP}}$ model is an integer, so the lower bound from the relaxations is $\lceil \sum_j w_j / c \rceil$; in Example 1 it is $\lceil \frac{8}{5} \rceil = 2$, not $\frac{8}{5}$.
- **Believing that the model counts only non-empty bins.** The constraints allow $y_b = 1$ for an empty bin: it is the objective function that switches such bins off, in every optimal solution.
- **Ignoring the symmetry.** Permuting the indices of the bins gives another optimal solution with the same packing: the $\mathrm{ILP}_{\mathrm{BPP}}$ model has many equivalent optimal solutions.
- **Expecting a better relaxation from $y_b \leq 1$.** The second relaxation has the same optimal value as the first one: the upper bounds $y_b \leq 1$ do not cut off the optimal solution of the first relaxation.

9.6 In summary

Summary

- The $\mathrm{ILP}_{\mathrm{BPP}}$ model has $m \cdot n$ assignment variables x_{jb} and m activation variables y_b; the capacity constraints link them.
- Both LP relaxations have optimal value $z(\mathrm{LP}_{\mathrm{BPP}}) = \sum_j w_j/c$, with dual solution $\tilde{\mu}_j = w_j/c$ and $\tilde{\pi}_b = 1/c$.
- Since the optimal value is an integer, $\lceil z(\mathrm{LP}_{\mathrm{BPP}}) \rceil$ is a lower bound on the number of bins.
- The ratio between $z(\mathrm{LP}_{\mathrm{BPP}})$ and $z(\mathrm{ILP}_{\mathrm{BPP}})$ is at least $\frac{1}{2}$, and the value $\frac{1}{2}$ is asymptotically attained; the same holds for $\lceil z(\mathrm{LP}_{\mathrm{BPP}}) \rceil$.

Dashboard of Example 1. Bounds on $z(\mathrm{ILP}_{\mathrm{BPP}})$ computed in this chapter:

Bound	Value	From
First relaxation	$\frac{8}{5}$	$z(\text{LP-0}_{\mathrm{BPP}})$
Second relaxation	$\frac{8}{5}$	$z(\mathrm{LP}_{\mathrm{BPP}})$
Lower bound	2	$\lceil \frac{8}{5} \rceil$
Optimal value	2	patterns $\{1, 3\}$ and $\{2\}$

Here the rounded bound is already equal to the optimal value.

Chapter 10

Formulations of the vertex coloring problem

Prerequisites

- From the module *Linear Programming*, Chapter 9 «Duality theory»: the dual of an LP problem, weak and strong duality.
- From the module *Integer linear programming*, the chapters «Integer linear programming» and «LP relaxation and dual bounds»: ILP problems, LP relaxations and bounds.
- From this module, the chapter «Vertex coloring»: the problem and the notation on graphs.

Objectives

At the end of this chapter you will be able to:

- write the ILP model of the Vertex Coloring Problem and explain the role of the activation variables y_c;
- write the two LP relaxations and their duals, and compute their optimal solutions;
- prove that the optimal value of the LP relaxations is 2 for every graph with an edge;
- explain why this bound is very weak, and why the model is not enough to color a graph.

10.1 The problem in brief

Where we were

In the chapter «Vertex coloring» we met the Vertex Coloring Problem as a *problem*: definition, notation on graphs and first bounds. Here we look at it as a *model* and we study its LP relaxations. The problem returns with heuristic algorithms and approximation algorithms for the VCP.

Definition 10.1: Vertex Coloring Problem

Given:

1. a simple *graph* $G = (V, E)$, where

$$V = \{1, 2, \ldots, n\}$$

is the set of n *vertices*, and

$$E \subseteq \big\{\, \{u, v\} : u, v \in V,\ u \neq v \,\big\}$$

is the set of m *edges*,

the **Vertex Coloring Problem** (VCP) asks to decide how to assign a color to each vertex, so that no two adjacent vertices share the same color, and the total number of colors used is minimized.

- For a vertex $v \in V$ we denote by $\delta(v)$ the set of the edges incident to v, by $\deg(v) = |\delta(v)|$ its degree and by $N(v)$ the set of the vertices adjacent to v.

- We assume that any VCP instance satisfies the problem assumptions, i.e.,

$$\deg(v) \geq 1 \quad \text{for all} \quad v \in V \qquad \text{and} \qquad N(u) \not\subseteq N(v) \quad \text{for all} \quad u, v \in V,\ u \neq v$$

- The first assumption guarantees that the graph has no isolated vertices: they can always take any color used by the rest of the graph.

- The second assumption guarantees that no vertex can be removed without affecting the optimal value: if $N(u) \subseteq N(v)$, any coloring of $G \setminus \{u\}$ can be extended to u by assigning it the color of v.

10.2 An ILP model for the Vertex Coloring Problem

- Given an upper bound $k \in \mathbb{Z}_{\geq 2}$ on the number of colors used in any optimal VCP solution, we introduce the following $k \underbrace{|V|}_{=n}$ binary variables:

$$x_{vc} = \begin{cases} 1 & \text{if vertex } v \text{ takes color } c, \\ 0 & \text{otherwise} \end{cases} \qquad \forall v \in \underbrace{\{1, 2, \ldots, n\}}_{=V},\ \forall c \in \underbrace{\{1, 2, \ldots, k\}}_{=\mathscr{C}}, \tag{10.1}$$

and the following k binary variables:

$$y_c = \begin{cases} 1 & \text{if color } c \text{ is used}, \\ 0 & \text{otherwise} \end{cases} \qquad \forall c \in \underbrace{\{1, 2, \ldots, k\}}_{=\mathscr{C}}. \tag{10.2}$$

The number n of vertices is a valid upper bound on the minimum number of colors needed to color all the vertices since, in the worst possible case, each vertex takes a different color. Also the number of colors used in any feasible VCP solution is a valid upper bound.

Using these variables, an Integer Linear Programming (ILP) model for the Vertex Coloring Problem (VCP) reads as follows:

$$\min \quad \sum_{c \in \mathscr{C}} y_c \tag{10.3a}$$

$$\text{subject to} \quad \sum_{c \in \mathscr{C}} x_{vc} = 1, \qquad \forall v \in V, \tag{10.3b}$$

$$-x_{uc} - x_{vc} + y_c \geq 0, \qquad \forall c \in \mathscr{C},\ \forall \{u, v\} \in E, \tag{10.3c}$$

$$x_{vc} \in \{0, 1\}, \qquad \forall v \in V,\ \forall c \in \mathscr{C}, \tag{10.3d}$$

$$y_c \in \{0, 1\}, \qquad \forall c \in \mathscr{C}. \tag{10.3e}$$

This ILP model is referred to as the ILP_{VCP} model.

- Description of the objective function and constraints of the ILP_{VCP} model:
  - the linear objective function (10.3a) minimizes the total number of used colors;
  - constraints (10.3b) ensure that each vertex takes exactly one color (n linear constraints);
  - constraints (10.3c) ensure that, for each color and each edge, the two adjacent vertices cannot both take that color, and that an unused color is taken by no vertex ($k \underbrace{|E|}_{=m}$ linear constraints);
  - finally, constraints (10.3d) and (10.3e) define the variables of the ILP model.

Explanation of the link between the variables

- **If color c is assigned to a vertex, then it is used.** The logical implication is

$$(x_{1c} \texttt{ OR } x_{2c} \texttt{ OR } \ldots \texttt{ OR } x_{nc}) \quad \Longrightarrow \quad y_c.$$


  Every vertex has an incident edge $\{u, v\}$, and the constraint $x_{uc} + x_{vc} \leq y_c$ forces $y_c = 1$ as soon as $x_{vc} = 1$.

- **If color c is not used, then no vertex takes it.** This is the contrapositive of the previous implication. For $y_c = 0$ the edge constraint gives $x_{uc} = x_{vc} = 0$ for every edge $\{u, v\} \in E$.

- **If color c is used, then some vertex takes it, but only in an optimal solution.** The constraints do not impose it. However, if $x_{vc} = 0$ for all v and $y_c = 1$, setting $y_c = 0$ keeps all the constraints satisfied and decreases the objective value by one. Hence, in every optimal solution, unused colors are not counted.

Example 10.1: ILP_{VCP} model

We consider the following graph $G = (V, E)$ with 3 vertices ($n = 3$) and 3 edges ($m = 3$):



The vertex set V and the edge set E are:

$$V = \{\ 1,\ 2,\ 3\ \}, \qquad E = \{\ \{1,2\},\ \{1,3\},\ \{2,3\}\ \}.$$

We refer to this VCP instance as Example 1.

The ILP_{VCP} model with $k = 3$ for this instance is:

$$
\begin{array}{lccccccccccccc}
\min & & & & & & & & & & y_1 & + y_2 & + y_3 & \\
\text{subject to} & x_{11} & + x_{12} & + x_{13} & & & & & & & & & & = 1 \\
& & & & x_{21} & + x_{22} & + x_{23} & & & & & & & = 1 \\
& & & & & & & x_{31} & + x_{32} & + x_{33} & & & & = 1 \\
& -x_{11} & & & -x_{21} & & & & & & + y_1 & & & \geq 0 \\
& -x_{11} & & & & & & -x_{31} & & & + y_1 & & & \geq 0 \\
& & & & -x_{21} & & & -x_{31} & & & + y_1 & & & \geq 0 \\
& & -x_{12} & & & -x_{22} & & & & & & + y_2 & & \geq 0 \\
& & -x_{12} & & & & & & -x_{32} & & & + y_2 & & \geq 0 \\
& & & & & -x_{22} & & & -x_{32} & & & + y_2 & & \geq 0 \\
& & & -x_{13} & & & -x_{23} & & & & & & + y_3 & \geq 0 \\
& & & -x_{13} & & & & & & -x_{33} & & & + y_3 & \geq 0 \\
& & & & & & -x_{23} & & & -x_{33} & & & + y_3 & \geq 0 \\
& x_{11}, & x_{12}, & x_{13}, & x_{21}, & x_{22}, & x_{23}, & x_{31}, & x_{32}, & x_{33}, & y_1, & y_2, & y_3 & \in \{0,1\}
\end{array}
$$

An optimal solution $\tilde{\boldsymbol{x}}(\text{ILP}_{\text{VCP}})$ is:

$\tilde{x}_{11} = 1,\ \tilde{x}_{12} = \tilde{x}_{13} = \tilde{x}_{21} = 0,\ \tilde{x}_{22} = 1,\ \tilde{x}_{23} = \tilde{x}_{31} = \tilde{x}_{32} = 0,\ \tilde{x}_{33} = 1,\ \tilde{y}_1 = \tilde{y}_2 = \tilde{y}_3 = 1$

The optimal value $z(\text{ILP}_{\text{VCP}})$ is 3.

The associated optimal solution of the VCP is given by the color classes $\tilde{Q}_1 = \{1\}$, $\tilde{Q}_2 = \{2\}$ and $\tilde{Q}_3 = \{3\}$. The optimal partition is unique, but the 3! permutations of the colors give 6 different optimal solutions.

10.3 A first Linear Programming relaxation

To obtain a Linear Programming (LP) relaxation of the ILP_{VCP} model, we relax constraints (10.3d) and (10.3e) with the following ones:

$$x_{vc} \geq 0, \qquad \forall v \in V,\ \forall c \in \mathscr{C}, \tag{10.4a}$$

$$y_c \geq 0, \qquad \forall c \in \mathscr{C}. \tag{10.4b}$$

The obtained LP model is referred to as the LP-0_{VCP} model.

Example 10.2: LP-0_{VCP} **model**

The LP-0_{VCP} model for the VCP instance of Example 1 is:

$$
\begin{array}{lcccccccccccccc}
\min & & & & & & & & & & y_1 & + y_2 & + y_3 & & \\
\text{subject to} & x_{11} & + x_{12} & + x_{13} & & & & & & & & & & & = 1 \\
& & & & x_{21} & + x_{22} & + x_{23} & & & & & & & & = 1 \\
& & & & & & & x_{31} & + x_{32} & + x_{33} & & & & & = 1 \\
& -x_{11} & & & -x_{21} & & & & & & + y_1 & & & & \geq 0 \\
& -x_{11} & & & & & & -x_{31} & & & + y_1 & & & & \geq 0 \\
& & & & -x_{21} & & & -x_{31} & & & + y_1 & & & & \geq 0 \\
& & -x_{12} & & & -x_{22} & & & & & & + y_2 & & & \geq 0 \\
& & -x_{12} & & & & & & -x_{32} & & & + y_2 & & & \geq 0 \\
& & & & & -x_{22} & & & -x_{32} & & & + y_2 & & & \geq 0 \\
& & & -x_{13} & & & -x_{23} & & & & & & + y_3 & & \geq 0 \\
& & & -x_{13} & & & & & & -x_{33} & & & + y_3 & & \geq 0 \\
& & & & & & -x_{23} & & & -x_{33} & & & + y_3 & & \geq 0 \\
& x_{11}, & x_{12}, & x_{13}, & x_{21}, & x_{22}, & x_{23}, & x_{31}, & x_{32}, & x_{33}, & y_1, & y_2, & y_3 & & \geq 0
\end{array}
$$

An optimal solution $\tilde{\boldsymbol{x}}(\text{LP-0}_{\text{VCP}})$ is:

$$\tilde{x}_{11} = \tilde{x}_{12} = \tilde{x}_{21} = \tilde{x}_{22} = \tilde{x}_{31} = \tilde{x}_{32} = \frac{1}{2}, \quad \tilde{x}_{13} = \tilde{x}_{23} = \tilde{x}_{33} = 0, \quad \tilde{y}_1 = \tilde{y}_2 = 1, \quad \tilde{y}_3 = 0$$

The optimal value $z(\text{LP-0}_{\text{VCP}})$ is 2.

Accordingly we have the lower bound:

$$2 \leq z(\text{ILP}_{\text{VCP}})$$

Using

- a free dual variable μ_v for each constraint (10.3b) ($v \in V$)
- a non-negative dual variable π_{ce} for each constraint (10.3c) ($c \in \mathscr{C},\ e = \{u, v\} \in E$)

the dual model of the LP-0_{VCP} model reads as follows:

$$\max \quad \sum_{v \in V} \mu_v \tag{10.5a}$$

$$\text{subject to} \quad \mu_v - \sum_{e \in \delta(v)} \pi_{ce} \leq 0, \qquad \forall v \in V,\ \forall c \in \mathscr{C}, \tag{10.5b}$$

$$\sum_{e \in E} \pi_{ce} \leq 1, \qquad \forall c \in \mathscr{C}, \tag{10.5c}$$

$$\mu_v \gtreqless 0, \qquad \forall v \in V, \tag{10.5d}$$

$$\pi_{ce} \geq 0, \qquad \forall c \in \mathscr{C},\ \forall e \in E. \tag{10.5e}$$

The obtained LP model is referred to as the dual $D(\text{LP-0}_{\text{VCP}})$ model.

Example 10.3: $D(\text{LP-0}_{\text{VCP}})$ model

The dual $D(\text{LP-0}_{\text{VCP}})$ model for the VCP instance of Example 1 is:

$$\begin{array}{rrrrrrrrrrrrrrrrrrrrrr}
\max & \mu_1 & +\mu_2 & +\mu_3 & & & & & & & & & & & & & & & & & & \\
\text{subject to} & \mu_1 & & & -\pi_{112} & -\pi_{113} & & & & & & & & & & & & & & & \leq 0 \\
& \mu_1 & & & & & & -\pi_{212} & -\pi_{213} & & & & & & & & & & & & \leq 0 \\
& \mu_1 & & & & & & & & & & & & -\pi_{312} & -\pi_{313} & & & & & & \leq 0 \\
& & \mu_2 & & -\pi_{112} & & -\pi_{123} & & & & & & & & & & & & & & \leq 0 \\
& & \mu_2 & & & & & -\pi_{212} & & -\pi_{223} & & & & & & & & & & & \leq 0 \\
& & \mu_2 & & & & & & & & & & & -\pi_{312} & & -\pi_{323} & & & & & \leq 0 \\
& & & \mu_3 & & -\pi_{113} & -\pi_{123} & & & & & & & & & & & & & & \leq 0 \\
& & & \mu_3 & & & & & -\pi_{213} & -\pi_{223} & & & & & & & & & & & \leq 0 \\
& & & \mu_3 & & & & & & & & & & & -\pi_{313} & -\pi_{323} & & & & & \leq 0 \\
& & & & \pi_{112} & +\pi_{113} & +\pi_{123} & & & & & & & & & & & & & & \leq 1 \\
& & & & & & & \pi_{212} & +\pi_{213} & +\pi_{223} & & & & & & & & & & & \leq 1 \\
& & & & & & & & & & & & & \pi_{312} & +\pi_{313} & +\pi_{323} & & & & & \leq 1 \\
& \mu_1, & \mu_2, & \mu_3 & & & & & & & & & & & & & & & & & \gtreqless 0 \\
& & & & \pi_{112}, & \pi_{113}, & \pi_{123}, & \pi_{212}, & \pi_{213}, & \pi_{223}, & \pi_{312}, & \pi_{313}, & \pi_{323} & & & & & & & & \geq 0
\end{array}$$

The optimal value $z(D(\text{LP-0}_{\text{VCP}}))$ is 2. An optimal solution $(\tilde{\boldsymbol{\mu}}, \tilde{\boldsymbol{\pi}})$ of $D(\text{LP-0}_{\text{VCP}})$ is:

$$\tilde{\mu}_1 = \tilde{\mu}_2 = 1, \quad \tilde{\mu}_3 = 0, \qquad \tilde{\pi}_{112} = \tilde{\pi}_{212} = \tilde{\pi}_{312} = 1, \qquad \text{all the other } \tilde{\pi}_{ce} = 0$$

Proposition 10.2

An optimal solution of the primal LP-0_{VCP} model is:

$$\tilde{x}_{v1} = \tilde{x}_{v2} = \frac{1}{2}, \quad \forall v \in V \quad \text{and} \quad \tilde{x}_{vc} = 0, \quad \forall v \in V, \forall c \in \{3, 4, \ldots, k\}, \tag{10.6a}$$

$$\tilde{y}_1 = \tilde{y}_2 = 1 \quad \text{and} \quad \tilde{y}_c = 0, \quad \forall c \in \{3, 4, \ldots, k\}, \tag{10.6b}$$

and an optimal solution of the dual $D(\text{LP-0}_{\text{VCP}})$ model is:

$$\tilde{\mu}_u = \tilde{\mu}_v = 1 \quad \text{and} \quad \tilde{\mu}_w = 0, \quad \forall w \in V \setminus \{u, v\}, \tag{10.7a}$$

$$\tilde{\pi}_{c\{u,v\}} = 1, \quad \forall c \in \mathscr{C} \quad \text{and} \quad \tilde{\pi}_{ce} = 0, \quad \forall e \in E \setminus \{\{u, v\}\}, \tag{10.7b}$$

where $\{u, v\}$ is any edge in E.

Corollary 10.3

The optimal value of the primal LP-0_{VCP} and dual D(LP-0_{VCP}) models is:

$$z(\text{LP-}0_{\text{VCP}}) = 2 = z(D(\text{LP-}0_{\text{VCP}})) \tag{10.8}$$

- The proof proceeds in three steps:
  1. We show that (10.6) is a feasible solution for the primal LP-0_{VCP} model;
  2. We show that (10.7) is a feasible solution for the dual D(LP-0_{VCP}) model;
  3. We show that both feasible solutions have the same objective function value.

  Therefore, by LP Weak Duality, (10.6) and (10.7) are a pair of optimal primal and dual solutions.

Proof.

1. For every $v \in V$, we have:
$$\sum_{c \in \mathscr{C}} \tilde{x}_{vc} = \frac{1}{2} + \frac{1}{2} = 1$$
Thus constraints (10.3b) are satisfied.

For $c \in \{1, 2\}$ and every $\{u, v\} \in E$, we have:
$$\tilde{y}_c - \tilde{x}_{uc} - \tilde{x}_{vc} = 1 - \frac{1}{2} - \frac{1}{2} = 0$$
and for $c \in \{3, 4, \dots, k\}$ we have $\tilde{y}_c - \tilde{x}_{uc} - \tilde{x}_{vc} = 0 - 0 - 0 = 0$. Thus constraints (10.3c) are satisfied.

We have $\tilde{x}_{vc} \geq 0$ and $\tilde{y}_c \geq 0$, for every $v \in V$ and every $c \in \mathscr{C}$, thus constraints (10.4a) and (10.4b) are satisfied.

Therefore, (10.6) provides a feasible solution for the primal LP-0_{VCP} model.

2. For every $c \in \mathscr{C}$, we have:
$$\sum_{e \in E} \tilde{\pi}_{ce} = 1$$
Thus constraints (10.5c) are satisfied.

For every $c \in \mathscr{C}$ and $w \in \{u, v\}$, we have:
$$\tilde{\mu}_w - \sum_{e \in \delta(w)} \tilde{\pi}_{ce} = 1 - 1 = 0$$
For every $c \in \mathscr{C}$ and $w \in V \setminus \{u, v\}$, we have $\tilde{\mu}_w = 0$ and $\tilde{\pi}_{ce} = 0$ for every $e \in \delta(w)$ (since $\tilde{\pi}_{ce}$ is nonzero only for $e = \{u, v\}$, which is not incident to w), so
$$\tilde{\mu}_w - \sum_{e \in \delta(w)} \tilde{\pi}_{ce} = 0 - 0 = 0.$$
Thus the constraints (10.5b) are satisfied.

We have $\tilde{\mu}_v \geq 0$, $\forall v \in V$ and $\tilde{\pi}_{ce} \geq 0$ for every $c \in \mathscr{C}$ and every $e \in E$. Thus constraints (10.5d) and (10.5e) are satisfied.

Therefore, (10.7) provides a feasible solution for the dual D(LP-0_{VCP}) model.

3. For the objective functions we have:

$$\sum_{c\in\mathscr{C}} \tilde{y}_c = 1 + 1 = \sum_{v\in V} \tilde{\mu}_v$$

thus the primal and dual objective function values are equal.

Therefore, by LP Weak Duality, (10.6) and (10.7) are a pair of optimal primal and dual solutions.

□

- This is an alternative proof that does not rely on the dual formulation but instead reasons directly on the minimum objective function value imposed by the constraints.

Proposition 10.4

An optimal solution of the primal LP-0_{VCP} model is:

$$\tilde{x}_{v1} = \tilde{x}_{v2} = \frac{1}{2}, \quad \forall v \in V \quad \text{and} \quad \tilde{x}_{vc} = 0, \quad \forall v \in V, \forall c \in \{3, 4, \dots, k\}, \tag{10.9a}$$

$$\tilde{y}_1 = \tilde{y}_2 = 1 \quad \text{and} \quad \tilde{y}_c = 0, \quad \forall c \in \{3, 4, \dots, k\}, \tag{10.9b}$$

and the optimal value is

$$z(\text{LP-}0_{\text{VCP}}) = 2. \tag{10.10}$$

Proof. Summing the edge constraints (10.3c) over all $c \in \mathscr{C}$, for any edge $\{u, v\} \in E$, gives

$$\sum_{c\in\mathscr{C}} y_c - \sum_{c\in\mathscr{C}} x_{uc} - \sum_{c\in\mathscr{C}} x_{vc} \geq 0 \implies \sum_{c\in\mathscr{C}} y_c \geq \underbrace{\sum_{c\in\mathscr{C}} x_{uc}}_{\text{by (10.3b)=1}} + \underbrace{\sum_{c\in\mathscr{C}} x_{vc}}_{\text{by (10.3b)=1}} = 2,$$

hence

$$\sum_{c\in\mathscr{C}} y_c \geq 2. \tag{10.11}$$

Therefore every feasible solution of LP-0_{VCP} satisfies (10.11).

Now consider the point defined in (10.9). For every $v \in V$, we have

$$\sum_{c\in\mathscr{C}} \tilde{x}_{vc} = \tfrac{1}{2} + \tfrac{1}{2} = 1,$$

so constraints (10.3b) hold. Moreover, for every $\{u, v\} \in E$ and $c \in \{1, 2\}$,

$$\tilde{y}_c - \tilde{x}_{uc} - \tilde{x}_{vc} = 1 - \tfrac{1}{2} - \tfrac{1}{2} = 0,$$

so constraints (10.3c) hold with equality, while for all $c \geq 3$ both sides are zero. Finally, constraints (10.4a) and (10.4b) are satisfied by construction. Hence the point defined

in (10.9) is a feasible solution for LP-0$_{\text{VCP}}$.

The objective value of (10.9) is

$$\sum_{c \in \mathscr{C}} \tilde{y}_c = 1 + 1 = 2,$$

which matches the lower bound (10.11). Therefore the point (10.9) is an optimal solution with objective value (10.10).

□

10.4 A second Linear Programming relaxation

To obtain a Linear Programming (LP) relaxation of the ILP$_{\text{VCP}}$ model, we relax constraints (10.3d) and (10.3e) with the following ones:

$$x_{vc} \geq 0, \qquad \forall v \in V, \forall c \in \mathscr{C}, \tag{10.12a}$$

$$y_c \leq 1, \qquad \forall c \in \mathscr{C}, \tag{10.12b}$$

$$y_c \geq 0, \qquad \forall c \in \mathscr{C}. \tag{10.12c}$$

The obtained LP model is referred to as the LP$_{\text{VCP}}$ model.

Example 10.4: LP$_{\text{VCP}}$ model

The LP$_{\text{VCP}}$ model for the VCP instance of Example 1 is:

$$
\begin{array}{rcccccccccccccc}
\min & & & & & & & & & & y_1 & + y_2 & + y_3 & \\
\text{subject to} & x_{11} & + x_{12} & + x_{13} & & & & & & & & & & = 1 \\
& & & & x_{21} & + x_{22} & + x_{23} & & & & & & & = 1 \\
& & & & & & & x_{31} & + x_{32} & + x_{33} & & & & = 1 \\
& -x_{11} & & & -x_{21} & & & & & & + y_1 & & & \geq 0 \\
& -x_{11} & & & & & & -x_{31} & & & + y_1 & & & \geq 0 \\
& & & & -x_{21} & & & -x_{31} & & & + y_1 & & & \geq 0 \\
& & -x_{12} & & & -x_{22} & & & & & & + y_2 & & \geq 0 \\
& & -x_{12} & & & & & & -x_{32} & & & + y_2 & & \geq 0 \\
& & & & & -x_{22} & & & -x_{32} & & & + y_2 & & \geq 0 \\
& & & -x_{13} & & & -x_{23} & & & & & & + y_3 & \geq 0 \\
& & & -x_{13} & & & & & & -x_{33} & & & + y_3 & \geq 0 \\
& & & & & & -x_{23} & & & -x_{33} & & & + y_3 & \geq 0 \\
& & & & & & & & & & y_1 & & & \leq 1 \\
& & & & & & & & & & & y_2 & & \leq 1 \\
& & & & & & & & & & & & y_3 & \leq 1 \\
& x_{11}, & x_{12}, & x_{13}, & x_{21}, & x_{22}, & x_{23}, & x_{31}, & x_{32}, & x_{33}, & y_1, & y_2, & y_3 & \geq 0
\end{array}
$$

An optimal solution $\tilde{\boldsymbol{x}}(\mathrm{LP}_{\mathrm{VCP}})$ is:

$$\tilde{x}_{11} = \tilde{x}_{12} = \tilde{x}_{21} = \tilde{x}_{22} = \tilde{x}_{31} = \tilde{x}_{32} = \frac{1}{2}, \quad \tilde{x}_{13} = \tilde{x}_{23} = \tilde{x}_{33} = 0, \quad \tilde{y}_1 = \tilde{y}_2 = 1, \quad \tilde{y}_3 = 0$$

The optimal value $z(\mathrm{LP}_{\mathrm{VCP}})$ is 2. Accordingly we have the same lower bound.

Using

- a free dual variable μ_v for each constraint (10.3b) ($v \in V$)
- a non-negative dual variable π_{ce} for each constraint (10.3c) ($c \in \mathscr{C},\ e = \{u, v\} \in E$)
- a non-positive dual variable λ_c for each constraint (10.12b) ($\forall c \in \mathscr{C}$)

the dual model of the $\mathrm{LP}_{\mathrm{VCP}}$ model reads as follows:

$$\max \quad \sum_{v \in V} \mu_v + \sum_{c \in \mathscr{C}} \lambda_c \tag{10.13a}$$

$$\text{subject to} \quad \mu_v - \sum_{e \in \delta(v)} \pi_{ce} \leq 0, \qquad \forall v \in V,\ \forall c \in \mathscr{C}, \tag{10.13b}$$

$$\sum_{e \in E} \pi_{ce} + \lambda_c \leq 1, \qquad \forall c \in \mathscr{C}, \tag{10.13c}$$

$$\mu_v \gtreqless 0, \qquad \forall v \in V, \tag{10.13d}$$

$$\pi_{ce} \geq 0, \qquad \forall c \in \mathscr{C},\ \forall e \in E, \tag{10.13e}$$

$$\lambda_c \leq 0, \qquad \forall c \in \mathscr{C}. \tag{10.13f}$$

The obtained LP model is referred to as the dual $D(\text{LP}_{\text{VCP}})$ model.

Example 10.5: $D(\text{LP}_{\text{VCP}})$ model

The dual $D(\text{LP}_{\text{VCP}})$ model for the VCP instance of Example 1 is:

$$\begin{array}{rrrr

Proposition 10.5

An optimal solution of the primal LP_{VCP} model is:

$$\tilde{x}_{v1} = \tilde{x}_{v2} = \frac{1}{2}, \quad \forall v \in V \quad \text{and} \quad \tilde{x}_{vc} = 0, \quad \forall v \in V, \forall c \in \{3, 4, \dots, k\}, \tag{10.14a}$$

$$\tilde{y}_1 = \tilde{y}_2 = 1 \quad \text{and} \quad \tilde{y}_c = 0, \quad \forall c \in \{3, 4, \dots, k\}, \tag{10.14b}$$

and an optimal solution of the dual $D(\text{LP}_{\text{VCP}})$ model is:

$$\tilde{\mu}_u = \tilde{\mu}_v = 1 \quad \text{and} \quad \tilde{\mu}_w = 0, \quad \forall w \in V \setminus \{u, v\}, \tag{10.15a}$$

$$\tilde{\pi}_{c\{u,v\}} = 1, \ \forall c \in \mathscr{C}, \quad \tilde{\pi}_{ce} = 0, \ \forall e \in E \setminus \{\{u, v\}\}, \quad \text{and} \quad \tilde{\lambda}_c = 0, \ \forall c \in \mathscr{C}, \tag{10.15b}$$

where $\{u, v\}$ is any edge in E.

Proof. The proof is similar to the previous one. In addition it is necessary to observe that:

$$\tilde{y}_c \leq 1, \quad \forall c \in \mathscr{C}.$$

Accordingly, also constraints (10.12b) are satisfied. □

Corollary 10.6

The optimal value of the primal LP_{VCP} and dual $D(\text{LP}_{\text{VCP}})$ models is:

$$z(\text{LP}_{\text{VCP}}) = 2 = z(D(\text{LP}_{\text{VCP}})) \tag{10.16}$$

10.5 Typical mistakes

Warning

- **Choosing k too small.** The number k of colors must be at least the chromatic number of the graph: with fewer colors the model has no feasible solution. The value $k = n$ is always valid.
- **Trusting the LP bound.** The optimal value of both relaxations is 2 for every graph with at least one edge, even for the complete graph on n vertices, whose optimal value is n.
- **Believing that the model counts only the colors that are really used.** The constraints allow $y_c = 1$ for a color that no vertex takes: it is the objective function that switches such colors off.
- **Ignoring the symmetry.** Permuting the colors of an optimal solution gives another optimal solution: the ILP_{VCP} model has $k!$ equivalent copies of every coloring.
- **Writing the dual solution with a wrong edge.** The dual values $\tilde{\pi}_{ce}$ are non-zero

only for one edge $e = \{u, v\}$, and for every color c: all the other edges have value zero.

10.6 In summary

Summary

- The ILP_{VCP} model has $k \cdot n$ variables x_{vc} (vertex v takes color c) and k variables y_c (color c is used); the edge constraints forbid equal colors on adjacent vertices.
- The two LP relaxations have the same optimal value 2 for every graph with an edge: put $\frac{1}{2}$ on two colors for all the vertices.
- The dual solution selects one edge $\{u, v\}$ and gives value 1 to μ_u, μ_v and $\pi_{c\{u,v\}}$ for every color.
- The bound 2 is very weak: a formulation that must color graphs needs additional valid inequalities.

Dashboard of Example 1 (triangle). Bounds on $z(\text{ILP}_{\text{VCP}})$ computed in this chapter:

Bound	Value	From
First relaxation	2	$z(\text{LP-0}_{\text{VCP}})$
Second relaxation	2	$z(\text{LP}_{\text{VCP}})$
Optimal value	3	color classes {1}, {2}, {3}

The gap between 2 and 3 is small here, but it grows with the size of the complete graph: 2 against n.

Chapter 11

Formulations of the travelling salesman problem

Prerequisites

- From the module *Integer linear programming*, the chapters «Integer linear programming» and «LP relaxation and dual bounds»: ILP problems, LP relaxations and bounds.
- From this module, the chapter «The Travelling Salesman Problem»: the problem, tours and the distance matrix.
- Later, the chapter «Branch-and-cut algorithms» of the module *Integer linear programming* shows how the constraints of Section 3 are added only when they are needed.

Objectives

At the end of this chapter you will be able to:

- write the ILP model of the Travelling Salesman Problem with the position variables of Miller-Tucker-Zemlin (MTZ) and explain how they eliminate subtours;
- write the ILP model with the subtour elimination constraints of Dantzig-Fulkerson-Johnson (DFJ), and prove that it is correct;
- count the variables and the constraints of the two models;
- compare the LP relaxations of the two models on the same instance.

11.1 The problem in brief

Where we were

In the chapter «The Travelling Salesman Problem» we met the TSP as a *problem*: definition, tours, the number of tours and the instance of this chapter. Here we look at it as a *model*, in two ways, and we compare the bounds that the two models give. The TSP returns with heuristic algorithms and approximation algorithms.

Definition 11.1: Travelling Salesman Problem

Given:

1. a set $\mathscr{V} = \{1, 2, \ldots, n\}$ of n *cities*,
2. for each pair of distinct cities $i, j \in \mathscr{V}$, a *distance* $d_{ij} \in \mathbb{Z}_{\geq 1}$, with $d_{ij} = d_{ji}$,

the **Travelling Salesman Problem** (TSP) asks to find a closed route that starts from a city, visits every other city exactly once and returns to the starting city, so that the total distance travelled is minimized.

- We assume that every TSP instance has at least three cities and symmetric distances, i.e.,

$$n \geq 3 \qquad \text{and} \qquad d_{ij} = d_{ji} \quad \forall i, j \in \mathscr{V},\ i \neq j.$$

- The formulations below use directed arcs. This does not change the symmetric problem: each edge $\{i, j\}$ is represented by the two opposite arcs (i, j) and (j, i), which have the same cost.

Example 11.1: the instance of the chapter

Consider the instance with $\mathscr{V} = \{1, 2, 3, 4, 5\}$ and distance matrix

d_{ij}	1	2	3	4	5
1	− − −	5	2	2	9
2	5	− − −	4	3	4
3	2	4	− − −	4	7
4	2	3	4	− − −	7
5	9	4	7	7	− − −

The tour $\sigma = (1, 3, 5, 2, 4)$ has length $L(\sigma) = d_{13}+d_{35}+d_{52}+d_{24}+d_{41} = 2+7+4+3+2 = 18$, and no shorter tour exists: the optimal value is 18 and σ is the unique optimal tour, up to the direction.

11.2 A model with position variables (MTZ)

- Let
$$A = \{(i,j) : i,j \in \mathscr{V},\ i \neq j\}$$
be the set of directed arcs. We introduce the following $n(n-1)$ **binary variables**:
$$x_{ij} = \begin{cases} 1 & \text{if arc } (i,j) \text{ is traversed by the tour,} \\ 0 & \text{otherwise,} \end{cases} \qquad \forall (i,j) \in A,$$
and the following $n-1$ **continuous variables**:
$$u_i = \text{position of city } i \text{ after leaving city 1,} \qquad \forall i \in \mathscr{V} \setminus \{1\}.$$
The city 1 is used only as a reference city: a tour has no preferred starting city.

Using these variables, the **MTZ model** of the Travelling Salesman Problem (TSP) reads as follows:

$$\min \quad \sum_{i \in \mathscr{V}} \sum_{\substack{j \in \mathscr{V} \\ j \neq i}} d_{ij} x_{ij} \tag{11.1a}$$

$$\text{subject to} \quad \sum_{\substack{j \in \mathscr{V} \\ j \neq i}} x_{ij} = 1, \qquad \forall i \in \mathscr{V}, \tag{11.1b}$$

$$\sum_{\substack{i \in \mathscr{V} \\ i \neq j}} x_{ij} = 1, \qquad \forall j \in \mathscr{V}, \tag{11.1c}$$

$$u_i - u_j + (n-1)x_{ij} \leq n-2, \qquad \forall i,j \in \mathscr{V} \setminus \{1\},\ i \neq j, \tag{11.1d}$$

$$x_{ij} \in \{0,1\}, \qquad \forall (i,j) \in A, \tag{11.1e}$$

$$1 \leq u_i \leq n-1, \qquad \forall i \in \mathscr{V} \setminus \{1\}. \tag{11.1f}$$

This model is referred to as the $\text{ILP}_{\text{TSP}}^{\text{MTZ}}$ model.

- Description of the objective function and constraints of the $\text{ILP}_{\text{TSP}}^{\text{MTZ}}$ model:
  - the linear objective function (11.1a) minimizes the total distance travelled by the tour;
  - constraints (11.1b) ensure that exactly one arc leaves each city (n linear constraints);
  - constraints (11.1c) ensure that exactly one arc enters each city (n linear constraints);
  - constraints (11.1d) link the arc variables with the position variables and eliminate the subtours that do not contain city 1 ($(n-1)(n-2)$ linear constraints);
  - constraints (11.1e) and (11.1f) define the domains of the variables of the model.

Explanation of the link between the variables

- For each pair of cities i, j different from city 1, if arc (i, j) is traversed, then the position of city j is at least one unit greater than the position of city i. This logical relation is the implication
$$x_{ij} = 1 \quad \Longrightarrow \quad u_j \geq u_i + 1.$$
- Its contrapositive is
$$u_j < u_i + 1 \quad \Longrightarrow \quad x_{ij} = 0.$$
In particular, an arc from a city to a city in the same or in an earlier position cannot be traversed.
- Both implications are imposed by the linear constraints (11.1d). If $x_{ij} = 1$, then the constraint gives $u_i - u_j + (n-1) \leq n-2$, that is $u_j \geq u_i + 1$.
- Conversely, if $u_j < u_i + 1$, then the constraint gives $(n-1)x_{ij} \leq n-2+u_j-u_i < n-1$. Since x_{ij} is binary, this forces $x_{ij} = 0$.
- When $x_{ij} = 0$, the constraint becomes $u_i - u_j \leq n-2$, which is always satisfied because both positions belong to $[1, n-1]$. Accordingly, $n-1$ is the smallest possible big-M coefficient for these bounds.

Proposition 11.2: MTZ constraints eliminate subtours

The constraints (11.1b)–(11.1f) are satisfied by an integer solution if and only if its selected arcs form one tour through all the cities.

Proof. Constraints (11.1b) and (11.1c) make the selected arcs a collection of vertex-disjoint directed cycles covering all cities. Suppose that one such cycle does not contain city 1 and visits, in order, $i_1, i_2, \ldots, i_k$. Since every arc (i_r, i_{r+1}) is selected, the implication above gives
$$u_{i_{r+1}} \geq u_{i_r} + 1 \qquad r = 1, \ldots, k,$$
where $i_{k+1} = i_1$. Summing the inequalities gives $u_{i_1} \geq u_{i_1} + k$, a contradiction. Thus no cycle can avoid city 1, and the collection is one tour.

Conversely, consider any tour and start it from city 1. For every city $i \neq 1$, set u_i equal to the position at which city i is visited after leaving city 1. The bounds (11.1f) hold. If a selected arc (i, j) has $i, j \neq 1$, then city j immediately follows city i, so that $u_j = u_i + 1$ and (11.1d) holds. If the arc is not selected, constraint (11.1d) holds by the bounds on the positions. Hence the tour defines a feasible integer solution. □

Example 11.2: $\mathrm{ILP}_{\mathrm{TSP}}^{\mathrm{MTZ}}$ model

For the instance of the chapter, the tour $\sigma = (1, 3, 5, 2, 4)$ is represented by
$$x_{13} = x_{35} = x_{52} = x_{24} = x_{41} = 1$$

and all the other arc variables equal to zero. Its position variables are

$$u_3 = 1, \qquad u_5 = 2, \qquad u_2 = 3, \qquad u_4 = 4.$$

The value of the objective function is $L(\sigma) = 18$, hence $z(\mathrm{ILP}_{\mathrm{TSP}}^{\mathrm{MTZ}}) = 18$.

11.3 A model with subtour elimination constraints (DFJ)

- The second model uses only the $n(n-1)$ arc variables x_{ij} of the MTZ model: there are no position variables. The subtours are forbidden directly, with one constraint for every subset of cities.

The **DFJ model** of the Travelling Salesman Problem reads as follows:

$$\min \quad \sum_{i \in \mathcal{V}} \sum_{\substack{j \in \mathcal{V} \\ j \neq i}} d_{ij} x_{ij} \tag{11.2a}$$

$$\text{subject to} \quad \sum_{\substack{j \in \mathcal{V} \\ j \neq i}} x_{ij} = 1, \qquad \forall i \in \mathcal{V}, \tag{11.2b}$$

$$\sum_{\substack{i \in \mathcal{V} \\ i \neq j}} x_{ij} = 1, \qquad \forall j \in \mathcal{V}, \tag{11.2c}$$

$$\sum_{i \in S} \sum_{\substack{j \in S \\ j \neq i}} x_{ij} \leq |S| - 1, \qquad \forall S \subseteq \mathcal{V} \setminus \{1\},\ |S| \geq 2, \tag{11.2d}$$

$$x_{ij} \in \{0, 1\}, \qquad \forall (i,j) \in A. \tag{11.2e}$$

This model is referred to as the $\mathrm{ILP}_{\mathrm{TSP}}^{\mathrm{DFJ}}$ model.

- Constraints (11.2d) are called **subtour elimination constraints**: a tour cannot contain all the $|S|$ arcs of a cycle on the cities of S, so it uses at most $|S| - 1$ arcs inside S.
- The number of constraints (11.2d) is $2^{n-1} - n$, exponential in n: for $n = 5$ they are 11, for $n = 20$ they are 524 268.

Proposition 11.3: Subtour elimination constraints

The constraints (11.2b)–(11.2e) are satisfied by an integer solution if and only if its selected arcs form one tour through all the cities.

Proof. Let a tour be given and let $S \subseteq \mathcal{V} \setminus \{1\}$ with $|S| \geq 2$. If the tour used $|S|$ arcs inside S, these arcs would form a cycle on the cities of S, which is not the whole tour since $1 \notin S$: a contradiction. Hence (11.2d) holds, and the degree constraints hold by definition of a tour.

Conversely, constraints (11.2b) and (11.2c) make the selected arcs a collection of vertex-disjoint directed cycles. If there were more than one cycle, one of them would not contain city 1. Let S be its set of cities: then $|S| \geq 2$ and the cycle uses $|S|$ arcs inside S, which violates (11.2d). Hence there is one cycle, a tour. □

Example 11.3: $\mathrm{ILP}_{\mathrm{TSP}}^{\mathrm{DFJ}}$ model

For the instance of the chapter, the constraints (11.2d) are 11, one for every subset $S \subseteq \{2,3,4,5\}$ with at least two cities. The subset $S = \{2,5\}$ gives $x_{25} + x_{52} \leq 1$, and $S = \{2,4,5\}$ gives

$$x_{24} + x_{25} + x_{42} + x_{45} + x_{52} + x_{54} \leq 2.$$

The subtour $2 \to 5 \to 2$ together with the subtour $1 \to 3 \to 4 \to 1$ satisfies the degree constraints but violates the first one. The tour σ satisfies all of them, and $z(\mathrm{ILP}_{\mathrm{TSP}}^{\mathrm{DFJ}}) = 18$.

11.4 Linear Programming relaxations

- We obtain the LP relaxation of each model by replacing the binary requirements with non-negativity constraints. In the MTZ model the position variables remain continuous and keep their bounds. The relaxations are called $\mathrm{LP}_{\mathrm{TSP}}^{\mathrm{MTZ}}$ and $\mathrm{LP}_{\mathrm{TSP}}^{\mathrm{DFJ}}$.
- Adding the upper bounds $x_{ij} \leq 1$ does not change the relaxations: for every arc (i,j), non-negativity and the degree equation (11.1b) give

$$x_{ij} \leq \sum_{\substack{k \in \mathscr{V} \\ k \neq i}} x_{ik} = 1.$$

- The dual of the relaxations has one variable for every ordering constraint, or for every subset S, and it does not provide a useful closed-form solution. The relevant theoretical point is the strength of the relaxation.

11.4.1 A weak region of the MTZ relaxation

Proposition 11.4: A weak region of the MTZ relaxation

Let (x,u) satisfy the degree equations (11.1b)–(11.1c), with $x_{ij} \geq 0$. If

$$x_{ij} \leq \frac{n-2}{n-1} \qquad \forall i,j \in \mathscr{V} \setminus \{1\},\ i \neq j,$$

then (x,u) is feasible for $\mathrm{LP}_{\mathrm{TSP}}^{\mathrm{MTZ}}$ after setting $u_i = 1$ for every $i \neq 1$.

Proof. With $u_i = u_j = 1$, each MTZ constraint (11.1d) becomes

$$(n-1)x_{ij} \leq n-2,$$

which is exactly the assumed upper bound. The bounds on the position variables also hold. □

- The proposition shows why the compact MTZ formulation can have a weak lower bound: many fractional solutions of the assignment relaxation already satisfy the ordering constraints.
- The MTZ model is valuable because it has only $O(n^2)$ variables and constraints. The DFJ model has a stronger relaxation, but exponentially many constraints: in practice, the subtour elimination constraints are added only when they are violated by the current solution of the relaxation.

Example 11.4: comparing the relaxations

For the instance of the chapter, with Gurobi we obtain three lower bounds on the optimal value 18:

Relaxation	Optimal value
degree constraints only (assignment relaxation)	16
$\mathrm{LP}_{\mathrm{TSP}}^{\mathrm{MTZ}}$	17
$\mathrm{LP}_{\mathrm{TSP}}^{\mathrm{DFJ}}$	18

The ordering constraints raise the bound from 16 to 17; the subtour elimination constraints close the gap on this instance.

11.5 Typical mistakes

Warning

- **Writing the MTZ constraints for the city 1.** The constraints (11.1d) are written only for the pairs of cities different from city 1: the city 1 has no position variable.
- **Using a smaller coefficient than $n-1$.** With a coefficient smaller than $n-1$ the constraint cuts off feasible tours; $n-1$ is the smallest valid value.
- **Taking $S = \mathscr{V}$ in the subtour elimination constraints.** A tour uses $|\mathscr{V}|$ arcs inside $\mathscr{V}$: the constraint would cut off every tour. The sets S are proper subsets of the cities, with at least two cities.
- **Trusting the MTZ bound.** By Proposition 11.4, many fractional solutions of the assignment relaxation are feasible for the MTZ relaxation: the compact formulation is not a strong one.
- **Listing all the subtour elimination constraints.** Their number is $2^{n-1}-n$: they are not written all at once, but added when violated.

11.6 In summary

Summary

- The two models have the same arc variables x_{ij} and the same degree constraints; they differ in how they forbid the subtours.
- The MTZ model adds $n-1$ position variables and $(n-1)(n-2)$ constraints: it is compact, but its relaxation is weak.
- The DFJ model adds $2^{n-1}-n$ subtour elimination constraints: the relaxation is stronger, and the constraints are generated when needed.
- Both models are correct: an integer solution is feasible if and only if its arcs form one tour.

Dashboard of the instance of the chapter. Lower bounds on the optimal value 18 computed in this chapter:

Bound	Value	From
Assignment relaxation	16	degree constraints only
MTZ relaxation	17	$z(\mathrm{LP}_{\mathrm{TSP}}^{\mathrm{MTZ}})$
DFJ relaxation	18	$z(\mathrm{LP}_{\mathrm{TSP}}^{\mathrm{DFJ}})$
Optimal value	18	tour $\sigma = (1,3,5,2,4)$

Chapter 12

Formulations of parallel machine scheduling

Prerequisites

- From the module *Linear Programming*, Chapter 9 «Duality theory»: the dual of an LP problem, weak and strong duality.
- From the module *Integer linear programming*, the chapters «Integer linear programming» and «LP relaxation and dual bounds»: ILP problems, LP relaxations and bounds.
- From this module, the chapter «Parallel machine scheduling»: the problem, the makespan and the instance of Example 1.

Objectives

At the end of this chapter you will be able to:

- write the ILP model of $P\|C_{\max}$ and explain how the makespan variable is linked to the machine loads;
- write the LP relaxation and its dual, and prove that its optimal value is $\sum_j t_j/m$;
- prove that the LP bound is at least $1/m$ of the optimal value, and that the factor is tight;
- use the valid inequalities and the symmetry-breaking constraints of the model.

12.1 The problem in brief

Where we were

In the chapter «Parallel machine scheduling» we met the problem $P\|C_{\max}$ as a *problem*: definition, Gantt charts and the lower bound on the makespan. Here we look at it as a *model* and we study its LP relaxations. The problem returns with heuristic algorithms and approximation algorithms.

Definition 12.1: Scheduling on identical parallel machines

Given:

1. a set of m identical *machines*,
2. a set $\mathscr{J} = \{1, 2, \dots, n\}$ of n *jobs*,
3. for each job $j \in \mathscr{J}$, a *processing time* $t_j \in \mathbb{Z}_{\geq 1}$,

the problem $P\|C_{\max}$ asks to assign each job to a machine, so that the largest total processing time assigned to a machine is minimized.

- We assume that an instance has more jobs than machines, i.e., $n > m$. Otherwise each job can be assigned to a different machine and the optimal makespan is simply $\max_{j \in \mathscr{J}} t_j$.

12.2 An ILP model for $P\|C_{\max}$

- Let $\mathscr{H} = \{1, 2, \dots, m\}$ be the set of machines. We introduce the following nm **binary variables**:

$$x_{jh} = \begin{cases} 1 & \text{if job } j \text{ is assigned to machine } h, \\ 0 & \text{otherwise,} \end{cases} \qquad \forall j \in \mathscr{J},\ \forall h \in \mathscr{H},$$

  and one **non-negative continuous variable**:

$$C_{\max} = \text{makespan of the schedule.}$$

Using these variables, an Integer Linear Programming (ILP) model for $P\|C_{\max}$ reads as follows:

$$\min \quad C_{\max} \tag{12.1a}$$

$$\text{subject to} \quad \sum_{h \in \mathscr{H}} x_{jh} = 1, \qquad \forall j \in \mathscr{J}, \tag{12.1b}$$

$$-\sum_{j \in \mathscr{J}} t_j x_{jh} + C_{\max} \geq 0, \qquad \forall h \in \mathscr{H}, \tag{12.1c}$$

$$x_{jh} \in \{0,1\}, \qquad \forall j \in \mathscr{J},\ \forall h \in \mathscr{H}, \tag{12.1d}$$

$$C_{\max} \geq 0. \tag{12.1e}$$

This ILP model is referred to as the $\text{ILP}_{P\|C_{\max}}$ model.

- Description of the objective function and constraints of the $\text{ILP}_{P\|C_{\max}}$ model:
  - the linear objective function (12.1a) minimizes the makespan;
  - constraints (12.1b) ensure that each job is assigned to exactly one machine (n linear constraints);
  - constraints (12.1c) link the assignment variables with the makespan variable. In particular, they ensure that the makespan is greater than or equal to the load of every machine (m linear constraints);
  - constraints (12.1d) and (12.1e) define the domains of the variables of the ILP model.

Explanation of the link between the variables

- For every machine h, the makespan and its load must satisfy the direct relation

$$C_{\max} \geq \sum_{j \in \mathscr{J}} t_j x_{jh},$$

  which is imposed by constraint (12.1c).

- Since the constraint holds for every machine, it imposes

$$C_{\max} \geq \max_{h \in \mathscr{H}} \left\{ \sum_{j \in \mathscr{J}} t_j x_{jh} \right\}.$$

- In every optimal solution equality holds. Indeed, if $C_{\max}$ were larger than the maximum load, replacing $C_{\max}$ with the maximum load would keep all the constraints (12.1c) satisfied and would decrease the objective value by a positive amount, which contradicts optimality.

- When no job is assigned to a machine, its load is zero.

Example 12.1: $\text{ILP}_{P\|C_{\max}}$ model

Consider $m = 3$ machines and the seven jobs with processing times

$$(t_1, t_2, t_3, t_4, t_5, t_6, t_7) = (5, 5, 4, 4, 3, 3, 3).$$

The assignment

$$\{1, 3\} \text{ to machine } 1, \qquad \{2, 4\} \text{ to machine } 2, \qquad \{5, 6, 7\} \text{ to machine } 3$$

has loads 9, 9, and 9. Hence its binary variables have value one precisely for

$$x_{11} = x_{31} = x_{22} = x_{42} = x_{53} = x_{63} = x_{73} = 1,$$

and the makespan is $C_{\max} = 9$. Since the lower bound

$$\max\left\{\max_{j\in\mathscr{J}} t_j,\ \left\lceil\frac{\sum_{j\in\mathscr{J}} t_j}{m}\right\rceil\right\} = \max\{5, \lceil 27/3\rceil\} = 9$$

is attained, $z(\text{ILP}_{P\|C_{\max}}) = 9$.

12.3 A first Linear Programming relaxation

- We obtain LP-$0_{P\|C_{\max}}$ by replacing the binary requirements (12.1d) with non-negativity constraints:

$$\min \quad C_{\max} \tag{12.2a}$$

$$\text{subject to} \quad \sum_{h\in\mathscr{H}} x_{jh} = 1, \qquad \forall j \in \mathscr{J}, \tag{12.2b}$$

$$-\sum_{j\in\mathscr{J}} t_j x_{jh} + C_{\max} \geq 0, \qquad \forall h \in \mathscr{H}, \tag{12.2c}$$

$$x_{jh} \geq 0, \qquad \forall j \in \mathscr{J},\ \forall h \in \mathscr{H}, \tag{12.2d}$$

$$C_{\max} \geq 0. \tag{12.2e}$$

This LP relaxation is referred to as the LP-$0_{P\|C_{\max}}$ model.

- The dual variables associated with constraints (12.2b) are free variables μ_j, and the dual variables associated with constraints (12.2c) are non-negative variables π_h. The dual model is:

$$\max \quad \sum_{j\in\mathscr{J}} \mu_j \tag{12.3a}$$

$$\text{subject to} \quad \mu_j - t_j\pi_h \leq 0, \qquad \forall j \in \mathscr{J},\ \forall h \in \mathscr{H}, \tag{12.3b}$$

$$\sum_{h\in\mathscr{H}} \pi_h \leq 1, \tag{12.3c}$$

$$\mu_j \gtreqless 0, \qquad \forall j \in \mathscr{J}, \tag{12.3d}$$

$$\pi_h \geq 0, \qquad \forall h \in \mathscr{H}. \tag{12.3e}$$

This dual LP model is referred to as $D(\text{LP-}0_{P\|C_{\max}})$.

Proposition 12.2: Optimal value of LP-0$_{P\|C_{\max}}$

For every $P\|C_{\max}$ instance,

$$z(\text{LP-0}_{P\|C_{\max}}) = \frac{\sum_{j\in\mathscr{J}} t_j}{m}.$$

Proof. Set $x_{jh} = 1/m$ for every job and every machine, and set

$$C_{\max} = \frac{\sum_{j\in\mathscr{J}} t_j}{m}.$$

Each job-assignment equation is satisfied and every machine has the displayed load, hence this is a feasible primal solution. In the dual, set

$$\pi_h = \frac{1}{m} \quad \forall h \in \mathscr{H}, \qquad \mu_j = \frac{t_j}{m} \quad \forall j \in \mathscr{J}.$$

The dual inequalities hold at equality, $\sum_h \pi_h = 1$, and the dual objective value is

$$\sum_{j\in\mathscr{J}} \mu_j = \frac{\sum_{j\in\mathscr{J}} t_j}{m}.$$

By LP Weak Duality, the two solutions are optimal, and the optimal value is the common value of the objective functions. □

12.3.1 A second Linear Programming relaxation

- Adding the upper bounds $x_{jh} \le 1$ gives LP$_{P\|C_{\max}}$. They do not change the relaxation: by (12.2b) and non-negativity, each x_{jh} is at most one. Thus

$$z(\text{LP}_{P\|C_{\max}}) = z(\text{LP-0}_{P\|C_{\max}}) = \frac{\sum_{j\in\mathscr{J}} t_j}{m}.$$

  The dual is consequently the same as $D(\text{LP-0}_{P\|C_{\max}})$, with zero dual multipliers for the redundant upper bounds.

12.3.2 Some important theoretical properties

Proposition 12.3: Worst-case ratio of the LP bound

For every $P\|C_{\max}$ instance,

$$\frac{z(\mathrm{LP}_{P\|C_{\max}})}{z(\mathrm{ILP}_{P\|C_{\max}})} \geq \frac{1}{m}.$$

Proof. Assigning every job to one machine gives a feasible schedule of makespan $\sum_j t_j$. Hence $z(\mathrm{ILP}_{P\|C_{\max}}) \leq \sum_j t_j$. By Proposition 12.2,

$$\frac{z(\mathrm{LP}_{P\|C_{\max}})}{z(\mathrm{ILP}_{P\|C_{\max}})} \geq \frac{\frac{\sum_j t_j}{m}}{\sum_j t_j} = \frac{1}{m}.$$

□

Proposition 12.4: The ratio $1/m$ is asymptotically attained

For a fixed number $m \geq 2$ of machines, there are instances for which

$$\frac{z(\mathrm{LP}_{P\|C_{\max}})}{z(\mathrm{ILP}_{P\|C_{\max}})} \longrightarrow \frac{1}{m}.$$

Proof. For $M \in \mathbb{Z}_{\geq 2}$, consider one job of processing time M and m jobs of processing time one. The long job alone forces $z(\mathrm{ILP}_{P\|C_{\max}}) \geq M$. The schedule that assigns the long job alone to machine 1 and the m unit jobs to the other $m-1$ machines has loads at most $2 \leq M$, hence the makespan M is feasible. Thus $z(\mathrm{ILP}_{P\|C_{\max}}) = M$. Proposition 12.2 gives

$$z(\mathrm{LP}_{P\|C_{\max}}) = \frac{M+m}{m}.$$

Therefore,

$$\lim_{M\to+\infty} \frac{z(\mathrm{LP}_{P\|C_{\max}})}{z(\mathrm{ILP}_{P\|C_{\max}})} = \lim_{M\to+\infty} \left(\frac{1}{m} + \frac{1}{M}\right) = \frac{1}{m}.$$

□

Example 12.2: an integrality gap

Consider two machines and three jobs with processing times $(3, 3, 2)$. Proposition 12.2 gives $z(\mathrm{LP}_{P\|C_{\max}}) = 8/2 = 4$. In every integer schedule, the job of length two must share a machine with one job of length three, so $z(\mathrm{ILP}_{P\|C_{\max}}) = 5$. The fractional assignment

$$x_{j1} = x_{j2} = \frac{1}{2} \qquad \forall j \in \{1, 2, 3\}$$

has load four on both machines, whereas the integer optimum has loads five and three.

12.4 Additional considerations

- For every job j, the inequality
$$C_{\max} \geq t_j \tag{12.4}$$
is a **valid inequality** for model (12.1). In every integer schedule, job j is assigned to one machine, whose load and therefore makespan are at least t_j (n linear constraints);

- The inequality
$$C_{\max} \geq \left\lceil \frac{\sum_{j \in \mathscr{J}} t_j}{m} \right\rceil \tag{12.5}$$
is also a **valid inequality** for model (12.1): the m machine loads sum to $\sum_j t_j$, so their maximum is at least their average and is an integer in every integer schedule (one linear constraint);

- The machines are identical. The constraints
$$\sum_{j \in \mathscr{J}} t_j x_{jh} \geq \sum_{j \in \mathscr{J}} t_j x_{j,h+1} \qquad \forall h \in \{1, \ldots, m-1\} \tag{12.6}$$
are **optimality-preserving constraints**: they are not valid for every feasible assignment, but every schedule can be relabelled so that its machine loads are non-increasing, without changing the makespan.

- The constraints (12.6) remove the equivalent relabellings of the machines only when the loads are distinct. With equal loads they remove none: in the instance of the chapter all the optimal schedules have loads 9, 9, 9, and all the 12 optimal solutions of the model satisfy them.

12.5 Typical mistakes

Warning

- **Using the LP bound alone.** The value $\sum_j t_j/m$ ignores the longest job: with one very long job it is much smaller than the optimal value. The valid inequality $C_{\max} \geq t_j$ must be added.
- **Forgetting to round up.** All the processing times are integer, so $C_{\max}$ is an integer in every integer schedule: the lower bound is $\lceil \sum_j t_j/m \rceil$.
- **Believing that the symmetry constraints remove all the equivalent solutions.** They remove the relabellings of the machines only when the loads are distinct; with equal loads they remove none.
- **Forgetting the direction of the load constraints.** The constraints (12.1c) bound $C_{\max}$ from below: it is the minimization that makes $C_{\max}$ equal to the largest load.
- **Mixing the indices.** The variable x_{jh} has the job j first and the machine h second.

12.6 In summary

Summary

- The $\text{ILP}_{P\|C_{\max}}$ model has $n \cdot m$ assignment variables x_{jh} and the makespan variable $C_{\max}$, linked by one load constraint for each machine.
- The LP relaxation has optimal value $z(\text{LP}_{P\|C_{\max}}) = \sum_j t_j/m$, attained by assigning every job with value $1/m$ to every machine; the dual solution is $\tilde{\pi}_h = 1/m$ and $\tilde{\mu}_j = t_j/m$.
- The ratio between $z(\text{LP}_{P\|C_{\max}})$ and $z(\text{ILP}_{P\|C_{\max}})$ is at least $1/m$, and $1/m$ is asymptotically attained.
- The valid inequalities $C_{\max} \geq t_j$ and $C_{\max} \geq \lceil \sum_j t_j/m \rceil$ strengthen the model.

Dashboard of the instance of the chapter ($m = 3$, $t = (5, 5, 4, 4, 3, 3, 3)$). Lower bounds on the optimal value 9:

Bound	Value	From
LP relaxation	9	$\sum_j t_j/m = 27/3$
Valid inequalities	9	$\max\{5, \lceil 27/3 \rceil\}$
Optimal value	9	loads 9, 9, 9

On the instance of the integrality gap ($m = 2$, $t = (3, 3, 2)$) the LP bound is 4 and the optimal value is 5.

Chapter 13

Formulations of capacitated lot sizing

Prerequisites

- From the module *Integer linear programming*, the chapters «Integer linear programming» and «LP relaxation and dual bounds»: ILP problems, LP relaxations and lower bounds.
- From this module, the chapter «Capacitated lot sizing»: the problem, the plans and the reference instance.
- From this module, the chapter «Dynamic programming for lot sizing»: the demand $D_{j,t}$ of a range of periods and the plans that produce only when the stock is empty.

Objectives

At the end of this chapter you will be able to:

- write the big-M formulation of the capacitated lot-sizing problem and compute a weak and a strong big-M value;
- write the shortest-path formulation and interpret its variables as the arcs of a network;
- represent a production plan as a combination of paths of the network;
- compare the LP bounds of the formulations on the reference instance;
- extend the formulations to several products, with setup times and initial inventory.

13.1 The problem in brief

Where we were

In the chapter «Capacitated lot sizing» we met the lot-sizing problem as a *problem*: plans, setups, counting and a first bound, on a reference instance of five periods. Here we look at it as a *model*: the natural formulation is weak, and we study how to make it stronger. The case without capacity returns with the DP algorithm of Wagner and Whitin.

- In the **capacitated lot-sizing problem** (CLS) a set $T = \{1, 2, \dots, m\}$ of m periods is given. In every period $t \in T$ we know the demand $d_t \in \mathbb{Z}_{\geq 1}$, the capacity $cap_t \in \mathbb{Z}_{\geq 1}$, the setup cost sc_t, the variable production cost vc_t per unit and the holding cost hc_t per unit in stock at the end of the period, all in $\mathbb{Z}_{\geq 0}$.

- We decide the integer quantity $x_t \in \mathbb{Z}_{\geq 0}$ produced in every period t, with $x_t \leq cap_t$, so that the demands are satisfied, the stock s_t at the end of period t is non-negative, and $s_0 = s_m = 0$. A **setup** is paid in the periods with $x_t > 0$.

- For $j \leq t$ we write $D_{j,t} = \sum_{k=j}^{t} d_k$ for the total demand of the periods $j, j+1, \dots, t$.

Example 13.1: the reference instance

We consider $m = 5$ periods, with the following data:

period t	1	2	3	4	5
demand d_t	2	1	3	4	1
capacity cap_t	4	4	4	4	4
setup cost sc_t	50	50	50	50	50
production cost vc_t	0	0	0	0	0
holding cost hc_t	10	10	10	10	10

This is the reference instance of the chapter «Capacitated lot sizing», with optimal plan $\boldsymbol{x} = (3, 0, 4, 4, 0)$ of cost 180.

13.2 A formulation with a big-M link

- We introduce the following **integer variables** and **binary variables**, for every period $t \in T$:

$$x_t \in \mathbb{Z}_{\geq 0} \text{ (production)}, \qquad s_t \in \mathbb{Z}_{\geq 0} \text{ (inventory at the end of the period)},$$

$$y_t = \begin{cases} 1 & \text{if there is a setup in period } t, \\ 0 & \text{otherwise.} \end{cases}$$

- The setup variables are linked to the quantities by the logical implication

$$x_t > 0 \quad \Longrightarrow \quad y_t = 1,$$

  modeled by the **big-M constraint** $x_t \leq M_t\, y_t$, where M_t is an upper bound on the production of period t.

- If the setups are ignored, the best plan produces in every period exactly its demand: it has no inventory and, in the reference instance, cost 0. This is a very weak lower bound: the setup costs must be modeled.

The **big-M formulation** of the CLS reads as follows:

$$\min \quad \sum_{t \in T} (sc_t\, y_t + vc_t\, x_t + hc_t\, s_t) \tag{13.1a}$$

$$s_{t-1} + x_t = d_t + s_t, \qquad \forall t \in T, \tag{13.1b}$$

$$x_t \leq M_t\, y_t, \qquad \forall t \in T, \tag{13.1c}$$

$$x_t \leq cap_t, \qquad \forall t \in T, \tag{13.1d}$$

$$s_0 = s_m = 0, \tag{13.1e}$$

$$x_t,\ s_t \in \mathbb{Z}_{\geq 0}, \qquad \forall t \in T, \tag{13.1f}$$

$$y_t \in \{0, 1\}, \qquad \forall t \in T. \tag{13.1g}$$

This model is referred to as the ILP_{CLSP} model.

- Description of the objective function and constraints of the ILP_{CLSP} model:
  - the objective function (13.1a) minimizes the cost of the setups, of the production and of the holding;
  - constraints (13.1b) are the **flow balancing constraints**: the demand of period t is satisfied by the production of the period and by the inventory of the previous period, and the surplus becomes the inventory at the end of period t;
  - constraints (13.1c) link the quantities with the setups, and constraints (13.1d) are the capacity constraints;
  - constraints (13.1e) impose an empty inventory at the beginning and at the end of the horizon.

13.2.1 Weak and strong big-M values

- **A weak value.** Since $s_m = 0$ and the inventories are non-negative, the production of period t cannot exceed the demand of the periods $t, \dots, m$: $M_t^{\text{dem}} = D_{t,m}$ is a valid value.

- **A second value.** By the capacity constraints (13.1d), $M_t^{\text{cap}} = cap_t$ is also valid.

- **A strong value.** The best value is the smallest of the two:

$$M_t = \min\{cap_t,\ D_{t,m}\}.$$

The smaller the big-M value, the stronger the LP relaxation: the constraint $x_t \leq M_t y_t$ cuts off the fractional values of y_t below x_t/M_t. A value that is too small, however, cuts off feasible plans.

Example 13.2: big-M values and LP bounds

For the reference instance we obtain the following values:

period t	1	2	3	4	5
$M_t^{\text{dem}} = D_{t,5}$	11	9	8	5	1
$M_t^{\text{cap}} = cap_t$	4	4	4	4	4
M_t	4	4	4	4	1

The three choices give the following optimal values of the LP relaxation, called $\text{LP}^{\text{dem}}_{\text{CLSP}}$, $\text{LP}^{\text{cap}}_{\text{CLSP}}$ and $\text{LP}^{\text{strong}}_{\text{CLSP}}$:

$$z(\text{LP}^{\text{dem}}_{\text{CLSP}}) = \frac{9865}{99} \approx 99.65,$$
$$z(\text{LP}^{\text{cap}}_{\text{CLSP}}) = \frac{275}{2} = 137.5,$$
$$z(\text{LP}^{\text{strong}}_{\text{CLSP}}) = \frac{315}{2} = 157.5.$$

The integer optimal solution is the same for all the choices: $\tilde{\boldsymbol{x}} = (3, 0, 4, 4, 0)$ with $\tilde{\boldsymbol{y}} = (1, 0, 1, 1, 0)$, and $z(\text{ILP}_{\text{CLSP}}) = 180$.

13.3 A shortest-path formulation

- The second formulation does not decide the quantities directly. For every $t, k \in T$, with $k \geq t$, we introduce the following **continuous variable**: $zv_{tk} \geq 0$, the fraction of the demand $D_{t,k}$ of the periods $t, \ldots, k$ that is produced in period t.
- Producing in period t the demand of the periods $t, \ldots, k$ costs the production of $D_{t,k}$ units and the holding of the units that wait for their period:

$$cv_{tk} = vc_t \, D_{t,k} + \sum_{s=t+1}^{k} \sum_{u=t}^{s-1} hc_u \, d_s.$$

The **shortest-path formulation** of the CLS reads as follows:

$$\min \quad \sum_{t \in T} sc_t \, y_t + \sum_{t \in T} \sum_{k=t}^{m} cv_{tk} \, zv_{tk} \tag{13.2a}$$

$$\sum_{k=1}^{m} zv_{1k} = 1, \tag{13.2b}$$

$$\sum_{u=1}^{t-1} zv_{u,t-1} = \sum_{k=t}^{m} zv_{tk}, \qquad \forall t \in T \setminus \{1\}, \tag{13.2c}$$

$$\sum_{k=t}^{m} zv_{tk} \leq y_t, \qquad \forall t \in T, \tag{13.2d}$$

$$\sum_{k=t}^{m} D_{t,k} \, zv_{tk} \leq cap_t, \qquad \forall t \in T, \tag{13.2e}$$

$$zv_{tk} \geq 0, \qquad \forall t, k \in T, \ k \geq t, \tag{13.2f}$$

$$y_t \in \{0, 1\}, \qquad \forall t \in T. \tag{13.2g}$$

This model is referred to as the shortest-path formulation; its LP relaxation is the $\text{LP}^{\text{SP}}_{\text{CLSP}}$ model.

- Constraints (13.2b) and (13.2c) are the flow constraints of a network, drawn in Figure 13.1: the nodes are $0, 1, \ldots, m$, and the variable zv_{tk} is the arc from node $t-1$ to node k, with cost cv_{tk}. A unit of flow leaves node 0 (constraint (13.2b)), and the flow is conserved at the other nodes (constraint (13.2c)).
- A path from node 0 to node m is a plan that produces only when the stock is empty: each arc $(t-1, k)$ is a production in period t for the periods $t, \ldots, k$. This is why the formulation is called a shortest-path formulation: with the capacities relaxed, the optimal plan is a shortest path.
- Constraints (13.2d) force the setup if the product is produced in period t, with the coefficient 1 instead of a big-M value. Constraints (13.2e) impose the capacity: the quantity produced in period t is $x_t = \sum_k D_{t,k} \, zv_{tk}$.

- The paths are the building blocks, and a plan that is not a path is a **combination of paths**: the quantities are integer, but the variables zv_{tk} may be fractional.




Figure 13.1: Network of the variables zv_{tk} for $m = 3$: the variable zv_{tk} is the arc from node $t-1$ to node k.

Example 13.3: costs and combinations of paths

For the reference instance the demands $D_{t,k}$ and the costs cv_{tk} are (the production cost is zero, so cv_{tk} is only holding cost):

	$D_{t,k}$					cv_{tk}				
$t\backslash k$	1	2	3	4	5	1	2	3	4	5
1	2	3	6	10	11	0	10	70	190	230
2		1	4	8	9		0	30	110	140
3			3	7	8			0	40	60
4				4	5				0	10
5					1					0

The optimal plan $\tilde{\boldsymbol{x}} = (3, 0, 4, 4, 0)$ is not a path: in period 3 it produces 4 units, more than the demand 3 of period 3, and less than the demand 7 of the periods 3 and 4. It is the following combination of paths:

$$zv_{12} = 1, \qquad zv_{33} = \tfrac{4}{5}, \quad zv_{35} = \tfrac{1}{5}, \qquad zv_{45} = \tfrac{4}{5},$$

and all the other variables zv_{tk} are zero: it is the combination, with weights $\frac{4}{5}$ and $\frac{1}{5}$, of the paths $0 \to 2 \to 3 \to 5$ (plan $(3, 0, 3, 5, 0)$) and $0 \to 2 \to 5$ (plan $(3, 0, 8, 0, 0)$). The quantities are $x_1 = D_{1,2}\, zv_{12} = 3$, $x_3 = 3 \cdot \frac{4}{5} + 8 \cdot \frac{1}{5} = 4$ and $x_4 = 5 \cdot \frac{4}{5} = 4$. The cost is

$$\underbrace{3 \cdot 50}_{\text{setups}} + \underbrace{10 \cdot 1 + 0 \cdot \tfrac{4}{5} + 60 \cdot \tfrac{1}{5} + 10 \cdot \tfrac{4}{5}}_{=30} = 180.$$

13.4 Comparing the formulations

- The formulations have the same optimal value $z(\text{ILP}_{\text{CLSP}}) = 180$, but their LP relaxations have different optimal values. On the reference instance the lower bounds, computed with the exact simplex algorithm and checked with Gurobi, are the following; the shortest-path formulation, solved with integer quantities, gives again the optimal value 180.

Model	Lower bound	Gap from 180
Without setups	0	100%
$\text{LP}^{\text{dem}}_{\text{CLSP}}$ $(M_t = D_{t,m})$	99.65	44.6%
$\text{LP}^{\text{cap}}_{\text{CLSP}}$ $(M_t = cap_t)$	137.50	23.6%
$\text{LP}^{\text{strong}}_{\text{CLSP}}$ $(M_t = \min\{cap_t, D_{t,m}\})$	157.50	12.5%
$\text{LP}^{\text{SP}}_{\text{CLSP}}$ (shortest path)	170	5.6%

- The bound 170 of the shortest-path formulation is the optimal value of the problem without capacity (plan $(3, 0, 8, 0, 0)$, of cost 170): the capacity constraints (13.2e) act only on the quantities, and a combination of paths can satisfy them.

- The price of the shortest-path formulation is its size: there are $O(m^2)$ variables zv_{tk}, against the $O(m)$ variables of the big-M formulation.

13.5 Several products, setup times and initial inventory

- In the general problem a set P of products shares the resource of each period. We write d_{pt}, sc_{pt}, vc_{pt} and hc_{pt} for the data of product $p \in P$ in period $t \in T$. Producing a unit of p takes the time $vt_{pt} \in \mathbb{Z}_{\geq 1}$, a setup takes the time $st_{pt} \in \mathbb{Z}_{\geq 0}$, and the capacity cap_t is a time. The initial inventory $si_p \in \mathbb{Z}_{\geq 0}$ of product p is a decision, bought at the cost fc_p per unit.

- The variables are x_{pt}, s_{pt}, si_p and y_{pt}. The balance constraints become $si_p + x_{p1} = d_{p1} + s_{p1}$ for the first period and $s_{p,t-1} + x_{pt} = d_{pt} + s_{pt}$ for the others. The capacity constraints and the big-M link become

$$\sum_{p \in P} (st_{pt}\, y_{pt} + vt_{pt}\, x_{pt}) \leq cap_t, \qquad \forall t \in T,$$

$$x_{pt} \leq M_{pt}\, y_{pt}, \qquad \forall p \in P,\ \forall t \in T,$$

  with the strong big-M value

$$M_{pt} = \min\left\{ \left\lfloor \frac{cap_t - st_{pt}}{vt_{pt}} \right\rfloor,\ sd_{ptm} \right\}, \qquad sd_{ptk} = \sum_{j=t}^{k} d_{pj}.$$

- In the shortest-path formulation each product has its own network, with an additional arc w_{pt} from node 0 to node t for the fraction of the initial inventory that covers the demands of the first t periods; its cost is $ci_{pt} = fc_p\, sd_{p1t} + \sum_{s=2}^{t} \sum_{u=1}^{s-1} hc_{pu}\, d_{ps}$. The constraint (13.2b) becomes $\sum_k (w_{pk} + zv_{p1k}) = 1$, the constraint (13.2c) has the arc $w_{p,t-1}$ on the left-hand side, and the capacity constraint is $\sum_p st_{pt}\, y_{pt} + \sum_p \sum_{k=t}^{m} vt_{pt}\, sd_{ptk}\, zv_{ptk} \leq cap_t$.

Example 13.4: two products

We consider $n = 2$ products and $m = 3$ periods, with capacities $cap_1 = 2500$, $cap_2 = 2100$, $cap_3 = 2500$, the initial inventory costs $fc_1 = fc_2 = 3000$ and the setup costs $sc_{pt} = 10000$. The other data are:

	product 1			product 2		
period t	1	2	3	1	2	3
d_{pt}	100	200	10	10	100	150
vt_{pt}	20	10	10	10	5	5
st_{pt}	100	100	100	110	110	110
hc_{pt}	110	100	100	105	105	105
vc_{pt}	1100	1000	1000	900	800	800

The big-M values are:

period t	1	2	3	1	2	3
sd_{ptm}	310	210	10	260	250	150
$\lfloor (cap_t - st_{pt})/vt_{pt} \rfloor$	120	200	240	239	398	478
M_{pt}	120	200	10	239	250	150

Ignoring the setups (in the costs and in the capacities), the optimal value is 575400, attained by the integer solution $x_{11} = 120$, $x_{12} = 160$, $x_{13} = 10$, $x_{21} = 10$, $x_{22} = 100$, $x_{23} = 150$, $s_{11} = 40$, $si_1 = 20$. The lower bounds and the optimal value of the formulations are:

Model	Value
Without setups	575 400.00
$M_{pt} = sd_{ptm}$	638 882.20
strong M_{pt}	651 267.37
shortest path	668 928.57
integer optimum	677 400

The optimal solution is unique: $x_{11} = 59$, $x_{12} = 200$, $x_{13} = 10$, $x_{21} = 110$, $x_{22} = 0$, $x_{23} = 150$, $si_1 = 41$, with a setup in every period except for product 2 in period 2.

13.6 Typical mistakes

Warning

- **Taking a big-M value that is too small.** A value smaller than the production of a feasible plan cuts off that plan: the values $D_{t,m}$ and cap_t are both valid.
- **Taking a big-M value that is too large.** Every valid value gives a correct model, but a large value gives a weak relaxation: the setup variables become almost zero in the LP solutions, as in the bound 99.65 of the reference instance.
- **Forgetting the setup times in the capacity constraints.** With several products the constraints contain $st_{pt}\, y_{pt}$ besides $vt_{pt}\, x_{pt}$.
- **Counting the holding cost of the demand of the production period.** In cv_{tk} the units for period t are not stored: the holding sum starts from the demand of period $t+1$.
- **Believing that the shortest-path formulation needs binary paths.** The variables zv_{tk} are continuous: a feasible plan that is not a path is a combination of paths.

13.7 In summary

Summary

- Without the setups the CLS is a simple flow problem; the setups need the binary variables y_t and a link with the quantities.
- In the big-M formulation the link $x_t \leq M_t\, y_t$ is correct for every valid M_t, but the LP bound improves when M_t decreases: the strong value is $\min\{cap_t, D_{t,m}\}$.
- The shortest-path formulation writes the plans as paths in a network, with the link $\sum_k zv_{tk} \leq y_t$ and no big-M value: its relaxation is the strongest.
- The three formulations have the same optimal value; they differ in the lower bound and in the size of the model.

Dashboard of the reference instance. Lower bounds on the optimal value computed in this chapter:

Bound	Value	From
Without setups	0	lot-for-lot plan
Big-M, $M_t = D_{t,m}$	99.65	$z(\mathrm{LP}^{\mathrm{dem}}_{\mathrm{CLSP}}) = \frac{9865}{99}$
Big-M, $M_t = cap_t$	137.5	$z(\mathrm{LP}^{\mathrm{cap}}_{\mathrm{CLSP}})$
Strong big-M	157.5	$z(\mathrm{LP}^{\mathrm{strong}}_{\mathrm{CLSP}})$
Shortest path	170	$z(\mathrm{LP}^{\mathrm{SP}}_{\mathrm{CLSP}})$
Optimal value	180	plan $(3, 0, 4, 4, 0)$