



SAPIENZA
UNIVERSITÀ DI ROMA

Mathematical Analysis 1

Volume 1 — Numbers and logic

The first volume: the language and the numbers on which analysis is built. Sets, logic and proof techniques, number sets and binary relations, ordered fields, supremum and the completeness axiom, roots, powers and logarithms, summations and geometric progressions, mathematical induction, factorials and binomial coefficients, the Fibonacci sequence, infinite sets, complex numbers. Definitions, theorems with proofs, worked examples; at the end, the exercises of the part with solutions.

Fabio Furini

sites.google.com/view/fabiofurini

Department of Computer, Control and Management Engineering — Sapienza University of Rome

Online version and downloadable material:
fabiofurini.github.io/mathematical-analysis-1/
github.com/fabiofurini/mathematical-analysis-1

Academic year 2026–2027

Contents

1	Sets	3
1.1	Informal introduction to set theory	3
1.2	Number sets	5
1.3	Relations between sets	7
1.4	Operations on sets	10
1.5	Further topics	17
2	Basics of logic and proof techniques	20
2.1	Logical symbols	20
2.2	Universal implications and proofs	20
2.3	Law of contraposition	24
2.4	Sufficient conditions and necessary conditions	26
2.5	Proofs by contradiction	30
2.6	Logic and sets	31
3	Number sets and intervals	34
3.1	The five main number sets	34
3.2	Intervals	38
4	Binary relations	40
4.1	Binary relations	40
4.2	Partial order relations	41
4.3	Total order relations	44
4.4	Functions	45
5	Ordered fields, supremum/infimum and the completeness axiom	47
5.1	Properties R_1 and R_2	47
5.2	Commensurable magnitudes	48
5.3	Geometric representation of $\mathbb{Q}$	50
5.4	Total order relations	50
5.5	Property R_3	51
5.6	Ordered fields	51
5.7	Bounded sets, maxima and minima	53
5.8	Upper/lower bounds and suprema/infima	56
5.9	Axiomatic definition of the real numbers	57
5.10	Geometric representation of $\mathbb{R}$	58

6	Roots, powers, logarithms and modular arithmetic	60
6.1	Roots, powers, logarithms	60
6.2	Modular arithmetic	67
7	Summations and geometric progressions	71
7.1	Summations	71
7.2	Geometric progressions	75
8	Principle of mathematical induction	79
8.1	The principle of mathematical induction	79
8.2	Proofs based on the principle of mathematical induction	80
9	Factorials, binomial coefficients and triangle inequality	85
9.1	Factorials	85
9.2	Binomial coefficients	86
9.3	Absolute value	90
10	The Fibonacci sequence	92
10.1	The Fibonacci sequence	92
11	Cardinality of infinite sets	97
11.1	Countable cardinality	97
11.2	Cardinality of the continuum	100
12	Complex numbers	103
12.1	Definition of $\mathbb{C}$ and field structure	103
12.2	Algebraic form of complex numbers	105
12.3	Trigonometric form of complex numbers	111
12.4	Quadratic equations	119
13	Exercises	123
13.1	Maxima, minima, suprema and infima	123
13.2	Equations and inequalities in one real variable	129
13.3	Summations	137
13.4	Principle of mathematical induction	139
13.5	Complex numbers	144

Sets

1.1 Informal introduction to set theory

- Set theory is based on the following three key concepts:

1. Sets

The notion of set is generally taken as **primitive** (that is, not reducible to more elementary concepts). The words *collection*, *class*, *aggregate*, *family* are used as synonyms of set.

Example 1.1: sets

For example, we have the set of points of a plane, the set of students of a university, or the set of stars of a galaxy.

2. Elements

A set is determined by its *elements*, in the sense that a set is defined when we have a **criterion** to establish whether a given element is or is not an element of this set.

There are sets with a *finite number of elements* (for example the set of students of a university) and sets with an *infinite number of elements* (for example the set of points of the plane).

Sets are usually denoted by capital letters. The elements of a set are usually denoted by lowercase letters.

3. Membership

The concept of *membership* links elements to sets. When an object is an element of a set, we say that the element *belongs* to the set. To indicate that an element x belongs to a set A we write:

$$x \in A$$

1.1.1 Informal definition of sets

1. A first way to define sets is **definition by listing** (roster notation)

A set can be defined **by listing**, that is, by listing the elements that belong to it between curly brackets. This technique assumes that the set has a finite number of elements.

Example 1.2: definition by listing

The notation:

$$A = \{a, b, c\}$$

means that the set A has as elements the three letters a , b and c . For example, a belongs to A , that is, $a \in A$.

2. A second way to define sets is **definition by a property**:

A set can be defined **by a property** as follows:

$$A = \{ x \in U : p(x) \text{ is true} \}$$

where $p(x)$ is the property that the element x of the set U must have in order to belong to the set A . This technique can be used to define sets with a finite or even an infinite number of elements.

Example 1.3: definition by a property

Consider for example the set of letters of the Latin alphabet:

$$U = \{a, b, c, d, e, f, g, h, i, j, k, l, m, n, o, p, q, r, s, t, u, v, w, x, y, z\}$$

Using for example the property $p(x)$ defined as “ x is a vowel” we can define the following set of vowels:

$$A = \underbrace{\{x \in U : x \text{ is a vowel}\}}_{\{a, e, i, o, u\}}$$

Note that to define a set A by a property we need a set U to which all the elements of the set A we want to define belong. The set U plays the role of the **universal set**.

It is important that the property $p(x)$ we use makes sense for every x of the set U (universal set), and therefore is true or false (without ambiguity of meaning) for every particular $x \in U$; the set A will then consist of exactly those x belonging to U for which the property $p(x)$ is true.

One must be careful when defining sets, since contradictions may arise. There is a formal definition of the concept of set, developed to avoid contradictions, but it is beyond the scope of this course.

- For example, the set of all sets that do not contain themselves is not a set in the formal definition of sets. Admitting this set would generate the contradiction: “the set of all sets that do not belong to themselves belongs to itself if and only if it does not belong to itself” (**Russell’s paradox** – section 1.5.1).

1.2 Number sets

Definition 1.1: numeral system

A **positional numeral system** is a way of encoding numbers using a **sequence of digits**, where each digit contributes differently to the number depending on its position. The number of distinct digits is the **base** of the system.

The **decimal expansion** of a number is a numerical representation in base 10; it expresses a number as a sum of powers of 10, and it can be finite or infinite depending on the type of number.

- Informal definition of the main number sets:
 1. We denote by $\mathbb{N}$ the set of **natural numbers**, that is, the set of numbers that can be written as **decimal expansions without decimal point and without sign**. We will use the informal notation:

$$\mathbb{N} = \{0, 1, 2, 3, 4, \dots\}$$

2. We denote by $\mathbb{Z}$ the set of **integers**, that is, the set of numbers that can be written as **decimal expansions without decimal point and with a sign**. We will use the informal notation:

$$\mathbb{Z} = \{0, \pm 1, \pm 2, \pm 3, \pm 4, \dots\}$$

3. We denote by $\mathbb{Q}$ the set of **rational numbers**, that is, the set of numbers that can be written as **finite or infinite periodic decimal expansions**. In other words, it is the set of numbers that can be written as a fraction $\frac{p}{q}$ where p is an integer and q is a natural number different from zero.

Example 1.4: rational numbers

- For example, with $p = 2$ and $q = 5$ we have the fraction $\frac{2}{5}$, whose decimal expansion is 0.4.
- For example, with $p = 4$ and $q = 10$ we have the fraction $\frac{4}{10}$, whose decimal expansion is again 0.4. Note that a rational number can be written with more than one fraction.
- For example, with $p = 13$ and $q = 30$ we have the fraction $\frac{13}{30}$, whose decimal expansion is $0.4\bar{3} = 0.4333\dots$

However, we can represent every rational number different from 0 by a single fraction $\frac{p}{q}$ by choosing $p \in \mathbb{Z}$ and $q \in \mathbb{N}$ coprime (that is, relatively prime: p and q are not both divisible by the same integer greater than 1).

Observation 1.2

$$0,\bar{9} = 1$$

Proof. There are different proofs of this remark, based on different mathematical techniques.

- (a) A simple proof follows directly from the definition of 1 divided by 3; indeed, we have:

$$\begin{aligned}\frac{1}{3} &= 0,\bar{3} \\ \frac{1}{3} \cdot 3 &= 0,\bar{3} \cdot 3 \\ 1 &= 0,\bar{9}\end{aligned}$$

- (b) Using algebraic arguments we can write:

$$\begin{array}{ll}x = 0.999\dots & \\10x = 9.999\dots & \text{multiplying by 10} \\10x = 9 + 0.999\dots & \text{separating the integer part} \\ & \text{from the fractional part} \\10x = 9 + x & \text{by definition of } x \\9x = 9 & \text{subtracting } x \\x = 1 & \text{dividing by 9}\end{array}$$

- (c) A proof by contradiction is the following:

$$\begin{aligned}0,\bar{9} &\neq 1 \\ 0,\bar{9} \cdot 9 &\neq 1 \cdot 9 \\ 0,\bar{9} \cdot 9 + 0,\bar{9} &\neq 1 \cdot 9 + 0,\bar{9} \\ 0,\bar{9} \cdot 9 + 0,\bar{9} &\neq 9,\bar{9} \\ 0,\bar{9} \cdot (9 + 1) &\neq 9,\bar{9} \\ 0,\bar{9} \cdot (10) &\neq 9,\bar{9} \\ 9,\bar{9} &\neq 9,\bar{9} \quad \text{contradiction!}\end{aligned}$$

□

4. We denote by $\mathbb{R}$ the set of **real numbers**, that is, the set of numbers identified with finite or infinite decimal expansions, periodic or non-periodic.

Example 1.5: real numbers

- Consider for example the number

$$0,10110111011110\dots$$

obtained by putting after the decimal point one digit equal to 1, then 0, then two digits equal to 1, then 0, then three digits equal to 1 ... and so on. The string of nonzero digits after the decimal point is neither finite nor periodic: therefore this number is real but not rational.

- Other examples of real but not rational numbers are $\sqrt{2}$ and $\sqrt{3}$, or π and Euler's number e (Napier's constant), which have infinite non-periodic decimal expansions and are therefore real but not rational numbers.

- There are formal definitions of the sets of natural, integer and rational numbers that are beyond the scope of this course. Later we will give a formal definition of the real numbers.

1.3 Relations between sets

1. Equality

Two sets A and B are equal when they have the same elements. We write

$$A = B$$

and this means that every element that belongs to A also belongs to B and every element that belongs to B also belongs to A .

2. Inclusion

It may happen that only one of the two requirements expressed by the equality relation holds. For example, if we only know that every element of A is also an element of B , we can say that A is contained in B . We write:

$$A \subseteq B \quad \text{or} \quad A \subset B$$

and we read “ A is contained in B ” or “ A is a **subset** of B ”. If we state that $A \subseteq B$, we do not exclude that $A = B$.

Saying that $A = B$ is equivalent to saying that $A \subseteq B$ and $B \subseteq A$.

Be careful not to confuse “belongs to” and “is contained in” (in symbols, $\in$ and $\subseteq$). In a sense, they are two ways of indicating that “something is inside something else”, but they have a fundamental logical difference:

- a set is contained in another set;
- an element belongs to a set.

Example 1.6: “belongs to” vs “is contained in”

For example:

$$3 \in \{1, 3, 4\};$$

$$\{3\} \subseteq \{1, 3, 4\};$$

$$\{1, 4\} \subseteq \{1, 3, 4\}.$$

In particular, do not confuse 3 (which is a number) with $\{3\}$, which is the set that contains the number 3 as its only element.

Sometimes we consider sets whose elements are other sets. In this case too, the symbols $\in$ and $\subseteq$ are not interchangeable, but must be used correctly.

Example 1.7: “belongs to” vs “is contained in”

For example, if we define the set

$$A = \left\{ \{1\}, \{2\}, \{1, 2\} \right\},$$

then it is correct to state that $\{2\} \in A$, because $\{2\}$ is a set that plays the role of an element of A , whereas it would be incorrect to say that $\{2\} \subseteq A$, because this would mean that $2 \in A$, while A has $\{2\}$ as an element but not 2. Instead, the subset of A containing only the element $\{2\}$ is denoted by $\{\{2\}\}$ and we have $\{\{2\}\} \subset A$.

1.3.1 Empty set, cardinality and power set**Definition 1.3: of empty set**

The **empty set** is the set that contains no elements. It is denoted by $\emptyset$.

Observation 1.4

Given a set A , we have:

$$\emptyset \subseteq A$$

Proof. To prove it, we should show that every element that belongs to $\emptyset$ also belongs to A ; but no element belongs to $\emptyset$, so the thesis holds. $\square$

Definition 1.5: of cardinality of a set

The number of elements of a set A is the **cardinality** of the set. It is denoted by $|A|$.

Definition 1.6: of power set

Given a set A , the set whose elements are all the subsets of A is called the **power set** of A and is denoted by the symbol $\mathcal{P}(A)$.

- Every set A has two trivial subsets, namely A itself and the empty set $\emptyset$ (they could coincide, if A is empty).

Example 1.8: power set

For example, given

$$A = \{1, 2, 3\},$$

then

$$\mathcal{P}(A) = \left\{ \emptyset, \{1\}, \{2\}, \{3\}, \{1, 2\}, \{2, 3\}, \{1, 3\}, \{1, 2, 3\} \right\}$$

Observation 1.7


Given a set A with n elements, the power set $\mathcal{P}(A)$ has 2^n elements:

$$|\mathcal{P}(A)| = 2^n$$

Proof. For each of the elements of A , the subsets of A may or may not contain that element. Hence we must make a choice between two options n times. The total number of possible subsets is therefore 2^n . $\square$

Example 1.9: construction of the power set with a binary tree

Given the set $A = \{1, 2, 3\}$ with $n = 3$ elements, the cardinality of its power set is $2^3 = 8$. The construction of the power set $\mathcal{P}(A)$ can be visualized through the following **binary tree** (an undirected, connected and acyclic graph) in which at each level we decide whether or not to include the object in the subset:



1.4 Operations on sets

Definition 1.8: of intersection of sets

The *intersection* of two sets $A, B \subseteq U$ is the set defined by:

$$A \cap B = \{x \in U : x \in A \text{ and } x \in B\}$$

It is the set of elements that belong both to the first and to the second set.

Definition 1.9: of union of sets

The **union** of two sets $A, B \subseteq U$ is the set defined by:

$$A \cup B = \{x \in U : x \in A \text{ or } x \in B\}$$

It is the set of elements that belong to the first or to the second set, where “or” is meant in the non-exclusive sense (the set of elements that belong to A or to B or to both).

Definition 1.10: of difference of sets

The **difference** of two sets $A, B \subseteq U$ is the set defined by:

$$A \setminus B = \{x \in A : x \notin B\}$$

It is the set of elements that belong to the first but not to the second set. The symbol “ $\setminus$ ” can also be written “ $-$ ” by analogy with arithmetic subtraction.

1.4.1 Complementary sets and disjoint sets

Example 1.10: universal sets

For example, in questions of arithmetic we could have $U = \mathbb{N}$, while in questions of analysis we could have $U = \mathbb{R}$.

Definition 1.11: of set complementation and complementary sets

The **complementation** of a set $A \subseteq U$ is the set defined by:

$$\overline{A} = \{x \in U : x \notin A\}$$

This set is called the **complement** (complementary set) of A with respect to U

- For every set $A \subseteq U$, where U is the universal set, we have the following relations:

$$\begin{aligned}\overline{\overline{A}} &= A \\ A \cap \overline{A} &= \emptyset \\ A \cup \overline{A} &= U\end{aligned}$$

$$\begin{aligned}A \cap \emptyset &= \emptyset \\ A \cap U &= A \\ A \cup \emptyset &= A \\ A \cup U &= U\end{aligned}$$

$$\begin{aligned}\overline{U} &= \emptyset \\ \overline{\emptyset} &= U\end{aligned}$$

Definition 1.12: of disjoint sets


Two sets A and B are **disjoint** if they have no elements in common:

$$A \cap B = \emptyset$$


1.4.2 Venn diagrams

Venn diagrams are graphical representations in which sets are represented as regions of the plane

- Venn diagrams of intersection, union and difference:



- Venn diagram of the complement of a set:



1.4.3 Cartesian product

There is another operation on sets, which can be performed on any two sets (i.e., two sets not necessarily contained in the same universal set):

Definition 1.13: of Cartesian product

Given two (not necessarily distinct) sets A and B , the set consisting of all ordered pairs (a, b) , with $a \in A$ and $b \in B$, is called the **Cartesian product** of A and B and is denoted by the symbol $A \times B$.

When A and B are two finite sets, the cardinality of their Cartesian product is:

$$|A \times B| = |A| \cdot |B|.$$

Example 1.11: Cartesian product

$$\{a, b\} \times \{a, b, c\} = \{(a, a), (a, b), (a, c), (b, a), (b, b), (b, c)\}$$

$$|\{a, b\}| = 2, \quad |\{a, b, c\}| = 3, \quad |\{a, b\} \times \{a, b, c\}| = 2 \cdot 3 = 6$$

- A typical use of the Cartesian product is $\mathbb{R} \times \mathbb{R}$, which is abbreviated by the symbol $\mathbb{R}^2$ and denotes the set of ordered pairs of real numbers.
- Similarly, $\mathbb{R}^n$ (abbreviation of the Cartesian product of n sets equal to $\mathbb{R}$) is the set of *ordered n -tuples of real numbers*

$$\mathbb{R}^n = \{ (x_1, x_2, \dots, x_n) : x_i \in \mathbb{R}, i \in \{1, 2, \dots, n\} \}$$

1.4.4 Properties of operations on sets

Observation 1.14: properties of intersection

Given three sets A, B and C , intersection has the following properties:

- **Commutative:**

$$A \cap B = B \cap A$$


- **Associative:**

$$A \cap (B \cap C) = (A \cap B) \cap C$$

- **Idempotence:**

$$A \cap A = A$$

Proof. The graphical proof of the associative property of intersection is the following:



□

Observation 1.15: properties of union

Given three sets A, B and C , union has the following properties:

- **Commutative:**

$$A \cup B = B \cup A$$


- **Associative:**

$$A \cup (B \cup C) = (A \cup B) \cup C$$

- **Idempotence:**

$$A \cup A = A$$

Proof. The graphical proof of the associative property of union is the following:



□

Observation 1.16: distributive properties (linking union and intersection)


Given three sets A, B and C we have:

$$A \cap (B \cup C) = (A \cap B) \cup (A \cap C)$$


$$A \cup (B \cap C) = (A \cup B) \cap (A \cup C).$$

The distributive properties link union and intersection to each other.

Proof. The graphical proof of the first property is the following:



The graphical proof of the second property is the following:



□


Proposition 1.17: De Morgan's laws (first version)

Given three sets A, B and C we have:


$$A \setminus (B \cap C) = (A \setminus B) \cup (A \setminus C)$$

$$A \setminus (B \cup C) = (A \setminus B) \cap (A \setminus C)$$

Proof. The graphical proof of the first property is the following:



The graphical proof of the second property is the following:


Proposition 1.18: De Morgan's laws (second version)

Given the sets $B, C \subseteq U$, we have

$$\overline{B \cap C} = \overline{B} \cup \overline{C}$$


$$\overline{B \cup C} = \overline{B} \cap \overline{C}$$

Proof. These laws can be derived by setting A equal to U in the previous De Morgan's laws, as follows:


$$\underbrace{U \setminus (B \cap C)}_{=\overline{B \cap C}} = \underbrace{(U \setminus B)}_{=\overline{B}} \cup \underbrace{(U \setminus C)}_{=\overline{C}}$$

$$\underbrace{U \setminus (B \cup C)}_{=\overline{B \cup C}} = \underbrace{(U \setminus B)}_{=\overline{B}} \cap \underbrace{(U \setminus C)}_{=\overline{C}}$$


Proof. The graphical proof of the first property is the following:



The graphical proof of the second property is the following:



- The set of elements that belong to a set B or to a set C but not to both (meaning “or” in the exclusive sense) is obtained by setting $U = B \cup C$ and taking the complement of the intersection:



Relation between inclusion and the operations of union and intersection:

$$A \subseteq B \text{ if and only if } A \cap B = A \text{ and } A \cup B = B.$$

1.4.5 Properties of the cardinality of sets

- For any two finite sets A and B , we have

$$|A \cup B| = |A| + |B| - |A \cap B|$$

from which we conclude that

$$|A \cup B| \leq |A| + |B|$$

- If A and B are disjoint then

$$|A \cap B| = 0 \quad \text{and hence} \quad |A \cup B| = |A| + |B|$$

- If $A \subsetneq B$, then $|A| < |B|$

1.5 Further topics

1.5.1 Russell's paradox

“can a set be an element of itself or not?”

- For example, the set of all books in a library is not an element of itself (a set of books is not a book). On the other hand, the set of all sets with more than 20 elements is an element of itself.
- Following this reasoning, two categories of sets can be defined:
 1. sets that are not elements of themselves
 2. sets that are elements of themselves

If we consider the set of all sets that are not elements of themselves, is it an element of itself or not?

Let us call this set S ; two hypotheses can be made:

- 1 If we suppose $S \in S$, then S contains itself as an element and therefore does not belong to S (since by definition a set belongs to S only if it does not contain itself as an element). Hence $S \notin S$, and we have a contradiction. We conclude that the hypothesis must be wrong.
- 2 If we suppose $S \notin S$, then S does not contain itself as an element and therefore belongs to S (since by definition a set belongs to S if it does not contain itself as an element). Hence $S \in S$ and we have another contradiction. We conclude that this hypothesis must be wrong too!

Russell's paradox: The set of all sets that do not belong to themselves belongs to itself if and only if it does not belong to itself.

- The formal definition of the concept of set is based on the **Zermelo-Fraenkel system of axioms**, abbreviated as **ZF**. This system of axioms includes the standard axioms of axiomatic set theory on which, together with the *axiom of choice*, all of ordinary mathematics is based.
- The **axiom of regularity** states that “Every non-empty set A contains an element disjoint from A ”.
- The **axiom of pairing** states that “Given two objects, there exists a set whose elements are the two objects”

Observation 1.19

No set is an element of itself

Proof. Given a set A , we apply the axiom of regularity to $\{A\}$, which is a set by the axiom of pairing. We thus obtain the set $\{A, A\}$, which we abbreviate to $\{A\}$ since sets cannot contain repeated objects (it is a special case of a pair). By the axiom of regularity there must exist an element of $\{A\}$ disjoint from $\{A\}$. Since the only element of $\{A\}$ is A , A is disjoint from $\{A\}$. Hence, since $A \cap \{A\} = \emptyset$, we cannot have $A \in A$ (by the definition of disjoint). $\square$

- The other ZF axioms are not part of the basic course in Mathematical Analysis.

- A further proof of

$$0, \bar{9} = 1$$

starts from the assumption that two numbers are equal if and only if their difference is equal to zero, and it is based on computing the value of $1 - 0, \bar{9}$.

- This proof is based on the fact that 0 is the only non-negative number less than all the reciprocals of the positive integers, or equivalently that there is no number greater than every integer. This is the **Archimedean property**, which holds for the rational and the real numbers.

Proof. We write the number $0,999\dots$ with n digits after the decimal point as $0, (9)_n$, hence $0, (9)_1 = 0.9$, $0, (9)_2 = 0.99$, $0, (9)_3 = 0.999$, and so on.

Given $\frac{1}{10^n} = 0,0\dots01$, with n digits after the decimal point, the addition rules for decimal numbers imply

$$0, (9)_n + \frac{1}{10^n} = 1 \quad \text{moreover} \quad 0, (9)_n < 1, \forall n \in \mathbb{N}.$$

We must prove that 1 is the smallest number that is not less than all the $0, (9)_n$. For this it is enough to prove that, if a number x is not greater than 1 and not less than all the $0, (9)_n$, then $x = 1$.

So let x be such that

$$0, (9)_n \leq x \leq 1$$

for every positive integer n . Hence

$$1 - 1 \leq 1 - x \leq 1 - 0, (9)_n$$

which, using basic arithmetic and the first equality established above, simplifies to

$$0 \leq 1 - x \leq \frac{1}{10^n}$$

This implies that the difference between 1 and x is less than the reciprocal of any positive integer. Hence this difference must be zero, and therefore $x = 1$; which in turn implies

$$0.999 \dots = 1$$

□

Basics of logic and proof techniques

2.1 Logical symbols

- a) the symbol “ $\forall$ ” is called the *universal quantifier* and is read “for every”, “for all”, “for each”
- b) the symbol “ $\exists$ ” is called the *existential quantifier* and is read “there exists”, “there exist”
- c) the symbol “ $\Rightarrow$ ” is called *logical implication* and is read “implies” or “if ... then”

- a) the symbol “ $:$ ” is read “such that”
- b) the symbol “ $\in$ ” is read “belongs to”
- c) the symbol “ $\notin$ ” is read “does not belong to”

- a) the symbol “ $\vee$ ” is called *logical disjunction* and is read “or”
- b) the symbol “ $\wedge$ ” is called *logical conjunction* and is read “and”
- c) the symbol “ $\neg$ ” is called *logical negation* and is read “not”.

2.2 Universal implications and proofs

Predicates (or properties) and **propositions** (or statements)

- Consider the following assertion:

$$\text{“the natural number } n \text{ is odd”} \quad (2.1)$$

and let us ask whether it is true. Obviously the answer is: “it depends on n ”. Indeed, in (2.1) the symbol n represents a variable that can take different values and make the assertion **true** or **false**.

A sentence of this kind is called a **predicate** (or *property*): its truth or falsity depends on the values of the **variable(s)** appearing in it.

- Now consider the following assertion:

$$\text{“for every natural number } n, \text{ if } n \text{ is odd then } n^2 \text{ is odd”} \quad (2.2)$$

which we can write more formally as follows:

$$\forall n \in \mathbb{N} \ (n \text{ odd} \Rightarrow n^2 \text{ odd}) \quad (2.3)$$

In this case we say that the variable n is not free, but bound by the *quantifier* $\forall$. As a consequence, (2.3) is true or false “once and for all” and is called a **proposition** (or *statement*).

In particular, the components of (2.3) are the set $\mathbb{N}$, two predicates defined on $\mathbb{N}$ given by

$$p(n) : \text{“} n \text{ odd”} \quad \text{and} \quad q(n) : \text{“} n^2 \text{ odd”}$$

and the implication

$$p(n) \Rightarrow q(n).$$

Definition 2.1: of universal implication

In general, a statement involving a **set** A , **two predicates** $p(x)$ and $q(x)$ whose argument x ranges over A , and the **logical structure**:

$$\forall x \in A \ (p(x) \Rightarrow q(x)) \quad (2.4)$$

is called a **universal implication**.

Most **theorems** consist of **universal implications**, in which the predicate $p(x)$ plays the role of the **hypothesis** and the predicate $q(x)$ plays the role of the **thesis** (conclusion).

- In particular, (2.3) is a proposition (or a statement):

Proposition 2.2

$$\forall n \in \mathbb{N} \quad (n \text{ odd} \Rightarrow n^2 \text{ odd}) \quad (2.5)$$

In this case one is easily convinced that the proposition is true, but how can we prove it rigorously?

- For example, is it enough to observe that 3 is odd and $3^2 = 9$ is odd to claim that the proposition is true? Certainly not, since the proposition requires the universal implication to hold for every natural number. However, there are infinitely many odd numbers: how can we prove a universal implication for infinitely many numbers?
- The key procedure is this: we consider a generic n satisfying the hypothesis (being odd) and we prove that n satisfies the thesis (its square is odd).

Let us see how to proceed to prove the previous proposition:

Proof. Let n be odd; we prove that then n^2 is odd.

Any odd number can be written in the form $2k + 1$, for a suitable $k \in \mathbb{N}$. We also observe that $2k$ is an even number for any $k \in \mathbb{N}$.

So let $n = 2k + 1$ be an odd number ($k \in \mathbb{N}$); we therefore need to write n^2 as an even integer plus one. We have:

$$n^2 = (2k + 1)^2 = 4k^2 + 4k + 1 = 2(2k^2 + 2k) + 1.$$

Since $2(2k^2 + 2k)$ is an even integer, n^2 is odd. □

To prove the correctness of a universal implication such as (2.4), we consider a generic x satisfying the hypothesis $p(x)$ and we try to prove that the thesis $q(x)$ is true.

- Let us now prove a similar relation for even numbers:

Proposition 2.3

$$\forall n \in \mathbb{N} \quad (n \text{ even} \Rightarrow n^2 \text{ even}) \quad (2.6)$$

Proof. Let n be even; we prove that then n^2 is even.

So let $n = 2k$ be an even number ($k \in \mathbb{N}$); we therefore need to write n^2 as an even integer. We have:

$$n^2 = (2k)^2 = 4k^2 = 2(2k^2)$$

Since $2(2k^2)$ is an even integer, n^2 is even. □

2.2.1 Counterexamples

Counterexamples are an important technique for proving the **falsity** of a universal implication.

- Let us ask, for example, whether the following universal implication is true or false:

$$\forall n \in \mathbb{N} \quad (n \text{ prime} \Rightarrow n \text{ odd}) \quad (2.7)$$

A moment's reflection shows that **this proposition is false**. Indeed, the number 2 is prime but it is even.

To claim that a universal implication is true, a proof is needed (an example is not enough), whereas to prove that a universal implication is false, one counterexample is enough.

- The universal implication requires that every x satisfying the hypothesis also satisfies the thesis: therefore, if we find even a single example of x that satisfies the hypothesis but not the thesis, this means that the universal implication is false. Not “false in one case”, but simply “false”, because the universal implication is true or false once and for all.

Definition 2.4: of counterexample

*In general, an example that satisfies the hypothesis but not the thesis of a universal implication, and that therefore proves its falsity, is called a **counterexample**.*

- The formal proof that the previous universal implication is false is the following:

Proof. The number 2 is a counterexample to the universal implication: "For every natural number n , if n is prime then n is odd".

□

The negation of the proposition

"for every $x \in A$, if $p(x)$ holds then $q(x)$ holds"

$$\forall x \in A \ (p(x) \Rightarrow q(x))$$

is the proposition

"there exists $x \in A$ for which $p(x)$ holds and $q(x)$ does not hold"

$$\exists x \in A \ (p(x) \wedge \neg q(x))$$

This particular x is a **counterexample**.

2.3 Law of contraposition

- It is an indirect proof technique

The universal implication

$$\forall x \in A \ (p(x) \Rightarrow q(x)) \tag{2.8}$$

is logically equivalent to

$$\forall x \in A \ (\text{not } q(x) \Rightarrow \text{not } p(x)) \tag{2.9}$$

The second implication is called the **contrapositive** of the first.

- For example: since we know that the universal implication (proposition 2.5) holds

$$\forall n \in \mathbb{N} \ (n \text{ odd} \Rightarrow n^2 \text{ odd})$$

the following proposition holds:

Proposition 2.5

$$\forall n \in \mathbb{N} \quad (n^2 \text{ even} \Rightarrow n \text{ even}) \quad (2.10)$$

Proof. Let n^2 be even; we prove that then n is even. We prove (2.10) relying on the truth of (2.5).

If n is not even then it is odd and therefore n^2 is odd by (2.5). This case contradicts the hypothesis that n^2 is even (hence it cannot occur).

Consequently n is even and (2.10) is proved. □

- The reasoning used in the previous proof has general validity, and shows precisely that if (2.8) is true then (2.9) is true; moreover, if the second is true then the first is true (because “not not $p(x)$ ” is logically equivalent to $p(x)$), so the two are logically equivalent.

The equivalence between (2.8) and (2.9) is called the **law of contraposition**. It is a method of *indirect proof* that consists in proving (2.9) to show that (2.8) is true (it requires proving that the negation of the thesis implies the negation of the hypothesis).

When using the law of contraposition, one must be able to construct the **correct negation** of a given proposition or property.

- Given any two predicates or properties $p(x)$ and $q(x)$, we list schematically some **rules** for constructing the **negation** of a proposition or property.

The negation of

“for every $x \in A$, $p(x)$ holds”

$$\forall x \in A \quad (p(x))$$

is

“there exists $x \in A$ for which $p(x)$ does not hold”

$$\exists x \in A \quad (\neg p(x))$$

The negation of

“there exists $x \in A$ for which $p(x)$ holds”

$$\exists x \in A \quad (p(x))$$

is

“for every $x \in A$, $p(x)$ does not hold”

$$\forall x \in A \quad (\neg p(x))$$

The negation of

$$\begin{aligned} & \text{"}p(x) \text{ holds and } q(x) \text{ holds"} \\ & (p(x) \wedge q(x)) \end{aligned}$$

is

$$\begin{aligned} & \text{"}p(x) \text{ does not hold or } q(x) \text{ does not hold"} \\ & (\neg p(x) \vee \neg q(x)) \end{aligned}$$

The negation of

$$\begin{aligned} & \text{"}p(x) \text{ holds or } q(x) \text{ holds"} \\ & (p(x) \vee q(x)) \end{aligned}$$

is

$$\begin{aligned} & \text{"}p(x) \text{ does not hold and } q(x) \text{ does not hold"} \\ & (\neg p(x) \wedge \neg q(x)) \end{aligned}$$

2.4 Sufficient conditions and necessary conditions

Definition 2.6: of sufficient condition

A **sufficient condition** is one that, if satisfied, guarantees the truth of the proposition.

Definition 2.7: of necessary condition

A **necessary condition** is one that must be satisfied for the proposition to be true.

Let $p(x)$ and $q(x)$ be any two predicates or properties. If $p(x)$ implies $q(x)$, formally:

$$p(x) \Rightarrow q(x)$$

then:

- $p(x)$ is a **sufficient condition** for $q(x)$
- $q(x)$ is a **necessary condition** for $p(x)$

- We have seen that the following universal implication is true (proposition (2.5)):

$$\forall n \in \mathbb{N} \ (n \text{ odd} \Rightarrow n^2 \text{ odd})$$

Hence “ n odd” is a sufficient condition for “ n^2 odd” and “ n^2 odd” is a necessary condition for “ n odd”.

- Let us now prove that the following proposition is also true:

Proposition 2.8

$$\forall n \in \mathbb{N} \ (n^2 \text{ odd} \Rightarrow n \text{ odd}) \quad (2.11)$$

Proof. Let n^2 be odd; we prove that then n is odd.

So let $n^2 = 2(2k^2 + 2k) + 1$ be an odd number ($k \in \mathbb{N}$), since $2(2k^2 + 2k)$ is an even number). We therefore need to write n as an even integer plus one. We have:

$$n = \sqrt{2(2k^2 + 2k) + 1} = \sqrt{4k^2 + 4k + 1} = \sqrt{(2k + 1)^2} = 2k + 1.$$

Since $2k + 1$ is an odd integer, n is odd.

□

In this way we have proved that “ n^2 odd” is a necessary and sufficient condition for “ n odd” and also that “ n odd” is a necessary and sufficient condition for “ n^2 odd”:

Proposition 2.9

$$\forall n \in \mathbb{N} \ (n \text{ odd} \iff n^2 \text{ odd}) \quad (2.12)$$

- We have seen that the following universal implication is true (proposition (2.6)):

$$\forall n \in \mathbb{N} \ (n \text{ even} \Rightarrow n^2 \text{ even})$$

Hence “ n even” is a sufficient condition for “ n^2 even” and “ n^2 even” is a necessary condition for “ n even”.

- Let us now prove that the following proposition is also true (direct proof, without using the law of contraposition as seen previously in (2.10)):

$$\forall n \in \mathbb{N} \ (n^2 \text{ even} \Rightarrow n \text{ even})$$

Proof. Let n^2 be even; we prove that then n is even.

So let $n^2 = 2 (2 k^2)$ be an even number ($k \in \mathbb{N}$); we therefore need to write n as an even integer. We have:

$$n = \sqrt{2 (2 k^2)} = \sqrt{4 k^2} = 2 k.$$

Since $2 k$ is an even integer, n is even. □

In this way we have proved that “ n^2 even” is a necessary and sufficient condition for “ n even” and also that “ n even” is a necessary and sufficient condition for “ n^2 even”:

Proposition 2.10

$$\forall n \in \mathbb{N} \quad (n \text{ even} \iff n^2 \text{ even}) \quad (2.13)$$

Example 2.1: necessary and sufficient conditions

For example, for a square matrix of real numbers, the fact that its determinant is different from zero is a necessary and sufficient condition for it to be invertible.

Proposition 2.11

$$\forall n \in \mathbb{N} \quad (n \text{ prime} > 2 \Rightarrow n \text{ odd}) \quad (2.14)$$

Proof. Let $n > 2$ be a prime number; we prove that then n is odd.

If n is not odd, it is even. But no even number greater than two is prime, a fact that contradicts the hypothesis that n is prime and greater than two (hence this case cannot occur).

Consequently n is odd. □

- Hence “ n odd” is a necessary condition for “ n prime > 2 ” and “ n prime > 2 ” is a sufficient condition for “ n odd”.

- However, “ n odd” does not imply “ n prime > 2 ”, since for example the number 9 is not prime (counterexample). That is:

$$\forall n \in \mathbb{N} \quad (n \text{ odd} \not\Rightarrow n \text{ prime number greater than } 2)$$

Hence “ n odd” is a necessary but not sufficient condition for “ n prime > 2 ” and “ n prime > 2 ” is a sufficient but not necessary condition for “ n odd”.

Proposition 2.12

$$\forall n \in \mathbb{N} \quad (n \text{ divisible by } 6 \Rightarrow n \text{ even}) \quad (2.15)$$

Proof. Let n be a number divisible by six; we prove that then n is even.

If n is not even, it is odd. But no odd number is divisible by six, a fact that contradicts the hypothesis that n is divisible by six (hence this case cannot occur).

Consequently n is even. □

- Hence “ n even” is a necessary condition for “ n divisible by 6” and “ n divisible by 6” is a sufficient condition for “ n even”.
- However, “ n even” does not imply “ n divisible by 6”, since for example the number 2 is even but not divisible by six (counterexample). That is:

$$\forall n \in \mathbb{N} \quad (n \text{ even} \not\Rightarrow n \text{ divisible by } 6)$$

Hence “ n even” is a necessary but not sufficient condition for “ n divisible by 6” and “ n divisible by 6” is a sufficient but not necessary condition for “ n even”.

Example 2.2: necessary/sufficient but not sufficient/necessary conditions

Being a square implies being a rectangle:

$$\text{being a square} \Rightarrow \text{being a rectangle}$$

since all squares are rectangles.

Hence “being a rectangle” is a necessary condition for “being a square” and “being a square” is a sufficient condition for “being a rectangle”.

But being a rectangle does not imply being a square

$$\text{being a rectangle} \not\Rightarrow \text{being a square}$$

because there are rectangles that are not squares.

Hence “being a rectangle” is not a sufficient condition (but it is a necessary one) for “being a square” and “being a square” is not a necessary condition (but it is a sufficient one) for “being a rectangle”.

2.5 Proofs by contradiction

- It is an indirect proof technique

Definition 2.13: of proof by contradiction

*In general, a **proof by contradiction** consists in assuming that the hypothesis of the theorem and the negation of the thesis are true, and deducing from these facts a contradiction of any kind.*

- We illustrate **proof by contradiction** with the following theorem.

Theorem 2.14

There is no rational number whose square is 2.

Proof. Suppose, by contradiction, that there exists a number $r \in \mathbb{Q}$ such that $r^2 = 2$.

We can write $r = \frac{n}{m}$ with $n, m \in \mathbb{Z}$, $m \neq 0$.

Moreover, we can assume that the fraction $\frac{n}{m}$ is already reduced to lowest terms, that is, “simplified” (in other words: n , m have no common factors).

We then have the chain of implications:

$$\left(\frac{n}{m}\right)^2 = 2$$

$$n^2 = 2m^2$$

so n^2 is even; but then by (2.13) n is also even and we can write $n = 2k$ for some $k \in \mathbb{Z}$.

Hence the relation $n^2 = 2 m^2$ can be rewritten as:

$$(2 k)^2 = 2 m^2$$

$$4 k^2 = 2 m^2$$

$$m^2 = 2 k^2.$$

So m^2 is even. But then by (2.13) m is also even.

Therefore both n and m are even, and this is a **contradiction**, because we had assumed that the fraction $\frac{n}{m}$ had already been simplified.

The proof can be found in Euclid's Elements (around 300 BC). □

2.6 Logic and sets

The language of logic and the language of sets are two sides of the same coin.

2.6.1 Logical implication and set inclusion

- There is a parallel between the relation of set inclusion and logical implication. To explain it, consider the universal implication:

$$\forall n \in \mathbb{N} \quad (n \text{ divisible by } 4 \Rightarrow n \text{ divisible by } 2) \quad (2.16)$$

If we denote by:

$$D_4 = \{ n \in \mathbb{N} : n \text{ is divisible by } 4 \} \text{ and } D_2 = \{ n \in \mathbb{N} : n \text{ is divisible by } 2 \}$$

we can observe that the universal implication written above is equivalent to the assertion:

$$D_4 \subseteq D_2$$

Indeed, this inclusion means that every element belonging to D_4 also belongs to D_2 , that is, that every natural number divisible by 4 is also divisible by 2.

The universal implication:

$$\text{“for every } x \in A, \text{ if } p(x) \text{ holds then } q(x) \text{ holds”}$$

is equivalent to the set inclusion:

$$\{ x \in A : p(x) \text{ is true} \} \subseteq \{ x \in A : q(x) \text{ is true} \}.$$

2.6.2 Equality of sets and universal implications

- Proving the equality of two sets, i.e., $A = B$, requires proving two universal implications. Formally:

$$\forall x (x \in A \Rightarrow x \in B) \quad \text{and} \quad \forall x (x \in B \Rightarrow x \in A). \quad (2.17)$$

- Stating that $A \subsetneq B$ means stating that “Every element that belongs to A also belongs to B , and there exists an element of B that does not belong to A ”. Formally:

$$\forall x (x \in A \Rightarrow x \in B) \quad \text{and} \quad \exists x \in B : x \notin A. \quad (2.18)$$

2.6.3 Set operations and logical operations

There is a relation between operations on sets and logical operations. Precisely:

1. **Set intersection** is defined by means of “and” (**logical conjunction**).
 2. **Set union** is defined by means of “or” (**logical disjunction**).
 3. **Set difference** and the **complement operation** are defined by means of “not” (**logical negation**).
- The distributive properties of union and intersection of sets:

Observation 2.15: distributive properties (sets)

Given three sets A, B and C we have:

$$A \cap (B \cup C) = (A \cap B) \cup (A \cap C)$$

$$A \cup (B \cap C) = (A \cup B) \cap (A \cup C).$$

- They can be rewritten in terms of predicates by observing that the intersection symbol $\cap$ corresponds to the conjunction $\wedge$ (“and”) and that the union symbol $\cup$ corresponds to the disjunction $\vee$ (“or”).

Observation 2.16: distributive properties (predicates)

Given three predicates $p(x)$, $q(x)$ and $r(x)$ we have:

$$p(x) \wedge (q(x) \vee r(x)) = (p(x) \wedge q(x)) \vee (p(x) \wedge r(x))$$

$$p(x) \vee (q(x) \wedge r(x)) = (p(x) \vee q(x)) \wedge (p(x) \vee r(x)).$$

- De Morgan’s laws:

Proposition 2.17: De Morgan’s laws (sets)

Given the sets $B, C \subseteq U$, we have

$$\overline{B \cap C} = \overline{B} \cup \overline{C}$$

$$\overline{B \cup C} = \overline{B} \cap \overline{C}$$

- They can be rewritten in terms of predicates by observing that the complement operation corresponds to the negation $\neg$ (“not”).

Proposition 2.18: De Morgan’s laws (predicates)

Given two predicates $q(x)$ and $r(x)$ we have:

$$\neg(q(x) \wedge r(x)) = \neg q(x) \vee \neg r(x)$$

$$\neg(q(x) \vee r(x)) = \neg q(x) \wedge \neg r(x)$$

Number sets and intervals

3.1 The five main number sets

3.1.1 The natural numbers

3.1.2 The integers

$\mathbb{Z}$ is the set of **integers** (also called signed numbers), that is:

$$0, 1, -1, 2, -2, \dots$$

3.1.3 The rational numbers

$\mathbb{Q}$ is the set of **rational numbers**, that is, of the fractions:

$$\frac{n}{m}, \quad \text{where } n, m \in \mathbb{Z}, \text{ and } m \neq 0.$$

Rational numbers can also be written in decimal form. For example:

$$\frac{3}{4} = 0,75, \quad \text{while} \quad \frac{1}{3} = 0,333\dots = 0,\overline{3}.$$

A rational number, written in decimal form, may have after the decimal point a finite number of digits (different from zero), or an infinite number of digits different from zero, which however repeat *periodically*.

Observation 3.1

$$0,\bar{9} = 1$$

There are different proofs of this remark, based on different mathematical techniques.

Proof. A simple proof follows directly from the definition of 1 divided by 3; indeed, we have:

$$\begin{aligned}\frac{1}{3} &= 0,\bar{3} \\ \frac{1}{3} \cdot 3 &= 0,\bar{3} \cdot 3 \\ 1 &= 0,\bar{9}\end{aligned}$$

□

Proof. Using algebraic arguments we can write:

$$\begin{array}{ll}x = 0.999\dots & \\10x = 9.999\dots & \text{multiplying by 10} \\10x = 9 + 0.999\dots & \text{separating the integer part from the fractional part} \\10x = 9 + x & \text{by definition of } x \\9x = 9 & \text{subtracting } x \\x = 1 & \text{dividing by 9}\end{array}$$

□

Proof. A proof by contradiction is the following:

$$\begin{aligned}
 0,\bar{9} &\neq 1 \\
 0,\bar{9} \cdot 9 &\neq 1 \cdot 9 \\
 0,\bar{9} \cdot 9 + 0,\bar{9} &\neq 1 \cdot 9 + 0,\bar{9} \\
 0,\bar{9} \cdot 9 + 0,\bar{9} &\neq 9,\bar{9} \\
 0,\bar{9} \cdot (9 + 1) &\neq 9,\bar{9} \\
 0,\bar{9} \cdot (10) &\neq 9,\bar{9} \\
 9,\bar{9} &\neq 9,\bar{9} \quad \text{contradiction!}
 \end{aligned}$$

□

Other proofs start from the assumption that two numbers are identical if and only if their difference is equal to zero, and they are based on computing the value of $1 - 0,\bar{9}$.

Proof. We write the number $0,999\dots$ with n digits after the decimal point as $0,(9)_n$, hence $0,(9)_1 = 0.9$, $0,(9)_2 = 0.99$, $0,(9)_3 = 0.999$, and so on.

Given $\frac{1}{10^n} = 0,0\dots01$, with n digits after the decimal point, the addition rules for decimal numbers imply

$$0,(9)_n + \frac{1}{10^n} = 1 \quad \text{moreover} \quad 0,(9)_n < 1, \forall n \in \mathbb{N}.$$

We must prove that 1 is the smallest number that is not less than all the $0,(9)_n$. For this it is enough to prove that, if a number x is not greater than 1 and not less than all the $0,(9)_n$, then $x = 1$.

So let x be such that

$$0,(9)_n \leq x \leq 1$$

for every positive integer n . Hence

$$1 - 1 \leq 1 - x \leq 1 - 0,(9)_n$$

which, using basic arithmetic and the first equality established above, simplifies to

$$0 \leq 1 - x \leq \frac{1}{10^n}$$

This implies that the difference between 1 and x is less than the reciprocal of any positive integer. Hence this difference must be zero, and therefore $x = 1$; which in turn implies

$$0.999\dots = 1$$

□

This proof is based on the fact that 0 is the only non-negative number less than all the reciprocals of the positive integers, or equivalently that there is no number greater than every integer.

This is the **Archimedean property**, which holds for the rational and the real numbers.

3.1.4 The real numbers

$\mathbb{R}$ is the set of **real numbers**, that is, those which, written in decimal form, have after the decimal point any sequence of digits different from zero, possibly even *infinite* and *non-periodic*.

That numbers of this last kind exist (that is, real but not rational), can be understood by reflecting on examples such as:

$$0,10110111011110\dots$$

After the decimal point, the previous number has: one digit equal to 1, then 0, then two digits equal to 1, then 0, then three digits equal to 1 ... and so on. It is clear that this rule defines a decimal number precisely. On the other hand, the string of nonzero digits after the decimal point is neither finite nor periodic: therefore this number is irrational.

3.1.5 The complex numbers

$\mathbb{C}$ is the set of **complex numbers**, that is, numbers of the form $a + i b$, where a, b are real numbers, and i is the *imaginary unit*, that is, a number whose square is -1 .

Among the number sets introduced, the following inclusions hold:

$$\mathbb{N} \subsetneq \mathbb{Z} \subsetneq \mathbb{Q} \subsetneq \mathbb{R} \subsetneq \mathbb{C}.$$

As indicated by the symbols, all the inclusions are strict:

- there exist integers that are not natural numbers (the negative numbers),
- there exist rational numbers that are not integers (the proper fractions),
- there exist real numbers that are not rational (the irrational numbers),
- there exist complex numbers that are not real (the imaginary numbers).

3.2 Intervals

Definition 3.2: of interval

Given two real numbers a, b , an **interval** with endpoints a and b is one of the following sets:

$$\begin{aligned} [a, b] &= \{x \in \mathbb{R} : a \leq x \leq b\}, & [a, b) &= \{x \in \mathbb{R} : a \leq x < b\} \\ (a, b] &= \{x \in \mathbb{R} : a < x \leq b\}, & (a, b) &= \{x \in \mathbb{R} : a < x < b\} \end{aligned}$$


- As can be seen, the square (round) bracket at one of the two endpoints indicates that this endpoint is included in (excluded from) the interval.
- The intervals $[a, b]$ are called **closed**; the intervals (a, b) are called **open**.
- All the intervals listed above are bounded; half-lines are also called (unbounded) intervals, for example:

$$(-\infty, b) = \{x \in \mathbb{R} : x < b\}$$

$$[a, +\infty) = \{x \in \mathbb{R} : x \geq a\}$$

or the whole line

$$\mathbb{R} = (-\infty, +\infty)$$




It can be proved that the intervals, bounded or unbounded, are exactly the subsets I of $\mathbb{R}$ that satisfy the following property (called **connectedness**):

$$x_1 < x_2 < x_3, \text{ if } x_1, x_3 \in I, \text{ then } x_2 \in I$$

- In what follows we will sometimes consider the Cartesian product of two (or more) intervals, which can be given the geometric meaning of a rectangle (in two dimensions) or of a parallelepiped (in three dimensions).

Example 3.1: Cartesian product of intervals

Let $A = [0, 1]$, $B = [1, 2]$; the figure



illustrates the sets $A \times B$, $B \times A$, $A \times A$, also denoted by A^2 . In general, $A \times B$ is different from $B \times A$.

Binary relations

4.1 Binary relations

Definition 4.1: of binary relation

Given two sets A and B , a **binary relation** R is a subset of the Cartesian product $A \times B$

When we say that R is a binary relation on a single set A , we mean that R is a subset of $A \times A$.

- A binary relation will simply be called a relation (however, non-binary relations also exist).

Example 4.1

- The relation “less than” (“ $<$ ”) on the natural numbers is the set:

$$R_{<} = \left\{ (a, b) : a, b \in \mathbb{N} \text{ and } a < b \right\}.$$

- The relation “less than or equal to” (“ $\leq$ ”) on the natural numbers is the set:

$$R_{\leq} = \left\{ (a, b) : a, b \in \mathbb{N} \text{ and } a \leq b \right\}.$$

- The relation “is a subset of” $R_{\subseteq}$ on the power set of the natural numbers (denoted by $2^{\mathbb{N}}$) is the set:

$$R_{\subseteq} = \left\{ (A, B) : A, B \in 2^{\mathbb{N}} \text{ and } A \subseteq B \right\}.$$

- A relation $R \subseteq A \times A$ is **reflexive** if:

$$\forall a \in A, \quad (a, a) \in R$$

- A relation $R \subseteq A \times A$ is **symmetric** if:

$$\forall a, b \in A, \quad (a, b) \in R \Rightarrow (b, a) \in R$$

- A relation $R \subseteq A \times A$ is **transitive** if:

$$\forall a, b, c \in A, \quad (a, b) \in R \text{ and } (b, c) \in R \Rightarrow (a, c) \in R$$

Example 4.2

- The relation $R_{\leq}$ is reflexive, but $R_{<}$ is not.
- The relations $R_{<}$ and $R_{\leq}$ are not symmetric.
- The relations $R_{<}$ and $R_{\leq}$ are transitive, but for example the relation:

$$R_{ab}^1 = \left\{ (a, b) : a, b \in \mathbb{N} \text{ and } a = b - 1 \right\}$$

is not, e.g., $(3, 4) \in R_{ab}^1$ and $(4, 5) \in R_{ab}^1$ but $(3, 5) \notin R_{ab}^1$.

4.2 Partial order relations

- A relation $R \subseteq A \times A$ is **antisymmetric** if:

$$\forall a, b \in A, \quad (a, b) \in R \text{ and } (b, a) \in R \Rightarrow a = b$$

Example 4.3

- The relation $R_{\leq}$ is antisymmetric, since $a \leq b$ and $b \leq a$ imply $a = b$.

Definition 4.2: of partial order relation

A reflexive, antisymmetric and transitive relation is a **partial order relation**.

Example 4.4

- The relation $R_{\leq}$ is a partial order relation, but the relation $R_{<}$ is not, since it is not reflexive.

Observation 4.3

The relation $R_{\subseteq}$ is a partial order relation

Proof. We must prove that the relation is reflexive, antisymmetric and transitive.

- For it to be reflexive we must prove that $(S, S) \in R_{\subseteq}$, which is true since $S \subseteq S$.
- For it to be antisymmetric we must prove that if $S_1 \neq S_2$ then $S_1 \not\subseteq S_2$ or $S_2 \not\subseteq S_1$ or both, the negation of the desired property.

Since $S_1 \neq S_2$:

either there exists some element that is in S_1 but not in S_2 , hence $S_1 \not\subseteq S_2$
 or there exists some element that is in S_2 but not in S_1 , hence $S_2 \not\subseteq S_1$
 or both of the previous options hold

- For it to be transitive we must prove that $(S_1, S_2) \in R_{\subseteq}$ and $(S_2, S_3) \in R_{\subseteq}$ imply $(S_1, S_3) \in R_{\subseteq}$. Clearly, since $S_1 \subseteq S_2$ and $S_2 \subseteq S_3$, we have $S_1 \subseteq S_3$.

□

Definition 4.4: of partially ordered set

A **partially ordered set** is a pair consisting of a set and a partial order relation defined on it.

Example 4.5


- The sets of natural, rational or real numbers with the relation $R_{\leq}$ are partially ordered sets.
- The relation “is a descendant of” defined on a subset of people is a partial order

relation (if we consider individuals as descendants of themselves). Consequently, the subset of people considered with the relation “is a descendant of” is a partially ordered set.

Relations can be represented by a **directed graph**, where the **vertices** are the elements of the set on which the relation R is defined and an **arc** (a, b) means that $(a, b) \in R$. If the graph is acyclic, then the relation is a partial order relation.

Example 4.6

The directed acyclic graph associated with the relation $R_{\subseteq}$ on the set $\{1, 2, 3, 4\} \subseteq \mathbb{N}$ is the following (the sets consisting of a single number and the empty set are not shown in the figure):




- In a partially ordered set there may be no single **maximum element**, that is, an element a such that:

$$\forall b \in A, \quad (b, a) \in R$$

A partially ordered set may therefore contain several maximal elements a such that, for no $b \in A$ with $b \neq a$, we have $(a, b) \in R$.

Example 4.7

The directed acyclic graph associated with the partial order relation “is a divisor of” on the set $\{2, 3, \dots, 15\} \subseteq \mathbb{N}$ is the following:

**Example 4.8**

Given a set of boxes of different sizes, the relation “one box is contained in the other” on the set of boxes may have several maximal boxes, that is, boxes that are not contained in any other box.

4.3 Total order relations**Definition 4.5: of total relation**

A relation R on a set A is a **total relation** if:

$$\forall a, b \in A, \quad (a, b) \in R \quad \text{or} \quad (b, a) \in R \quad (\text{or both})$$

Example 4.9

- The relation $R_{\leq}$ is a total relation.
- The relation $R_{\subseteq}$ is not a total relation since, taking for example $S_1 = \{1, 2\}$ and $S_2 = \{2, 3\}$, $(S_1, S_2) \notin R_{\subseteq}$ and $(S_2, S_1) \notin R_{\subseteq}$.
- The relation “is a descendant of” is not a total relation since there exist pairs of individuals (a, b) for which neither a descends from b nor b descends from a .

Definition 4.6: of total order relation

A partial order relation that is also a total relation is a **total order relation**.

Example 4.10

- The relation $R_{\leq}$ is a total order relation.

Definition 4.7: of totally ordered set

A **totally ordered set** is a pair consisting of a set and a total order relation defined on it.

Example 4.11

- The sets of natural, rational or real numbers with the relation $R_{\leq}$ are totally ordered sets.

4.4 Functions

Definition 4.8: of function

Given two sets A and B , a binary relation f on A and B is a **function** if, for each $a \in A$, there exists one and only one $b \in B$ such that $(a, b) \in f$.

Definition 4.9: of domain and codomain

The set A is called the **domain** of f , and the set B is called the **codomain** of f .

- We write:

$$f : A \rightarrow B$$

and if $(a, b) \in f$, we write:

$$b = f(a)$$

since b is uniquely determined by the choice of a .

- Intuitively, the function f assigns an element of B to each element of A . No element of A is associated with two different elements of B . The same element of B can however be assigned to different elements of A .

Example 4.12

- The binary relation:

$$f = \left\{ (a, b) : a, b \in \mathbb{N} \text{ and } b = a \pmod{2} \right\}$$

is a function $f : \mathbb{N} \rightarrow \{0, 1\}$ since for all natural numbers a , there is exactly one value $b \in \{0, 1\}$ such that $b = a \pmod{2}$. For example,

$$0 = f(0), \quad 1 = f(1), \quad 0 = f(2), \dots$$

Example 4.13

- The binary relation

$$g = \left\{ (a, b) : a, b \in \mathbb{N} \text{ and } a + b \text{ is even} \right\}$$

is not a function, since for example $(1, 3)$ and $(1, 5)$ are both in g . In other words, for $a = 1$, we do not have one and only one b such that $(a, b) \in g$.

Definition 4.10: of argument and value

Given a function $f : A \rightarrow B$, if $b = f(a)$, we say that $a \in A$ is the argument of f and that $b \in B$ is the value of f associated with a .

- We can **define a function** by *directly defining its value* for all the *elements* of its *domain*.

Example 4.14

For example, we can define $f(n) = 2n$ for $n \in \mathbb{N}$, which means:

$$f = \{(n, 2n) : n \in \mathbb{N}\} \text{ and } f : \mathbb{N} \rightarrow \mathbb{N}$$

Ordered fields, supremum/infimum and the completeness axiom

5.1 Properties R_1 and R_2

- We now begin to study more closely the **structure of the number sets** that we introduced earlier, and in particular: the set $\mathbb{Q}$ of **rational numbers** and the set $\mathbb{R}$ of **real numbers**.
- We denote by a, b, c three generic **rational** or **real** numbers.

$R_1 \rightarrow$ In $\mathbb{Q}$ and $\mathbb{R}$ an operation is defined (called **addition** or **sum**) which has the following four properties:

- 1 $\forall a, b, \quad a + b = b + a$ (commutative property)
- 2 $\forall a, b, c, \quad (a + b) + c = a + (b + c)$ (associative property)
- 3 there exists a **neutral element for the sum**, denoted by 0, such that:

$$\forall a, \quad a + 0 = a$$

- 4 for every a there exists an **inverse** element of a **with respect to the sum**, called the **opposite** of a and denoted by $-a$, such that:

$$a + (-a) = 0$$

$R_2 \rightarrow$ In $\mathbb{Q}$ and $\mathbb{R}$ an operation is defined (called **multiplication** or **product**) which has the following four properties:

- 1 $\forall a, b, \quad a \cdot b = b \cdot a$ (commutative property)

2 $\forall a, b, c, \quad (a \cdot b) \cdot c = a \cdot (b \cdot c) \quad (\text{associative property})$

3 there exists a **neutral element for the product**, denoted by 1, such that:

$$\forall a, \quad a \cdot 1 = a$$

4 for every $a \neq 0$ there exists an **inverse element of a with respect to the product**, called the **reciprocal** of a and denoted by a^{-1} , such that:

$$a \cdot a^{-1} = 1$$

- The operations of sum and product are linked by the following property:

$$\forall a, b, c, \quad (a + b) \cdot c = a \cdot c + b \cdot c \quad (\text{distributive property})$$

- From properties R_1 and R_2 follows the **possibility of performing the four fundamental operations without restrictions**:

1. **addition** and **multiplication** (defined above)

2. **subtraction**, by setting:

$$a - b = a + (-b)$$

3. **division**, by setting:

$$a/b = a \cdot b^{-1} \quad \text{provided that } b \neq 0$$

5.2 Commensurable magnitudes

In mathematics, the word **magnitude** denotes a **property** of a **geometric figure** that can be **measured**. Examples of magnitudes are: the length of a segment, the measure of an angle, the area of a plane figure or the volume of a solid.

Definition 5.1: of homogeneous magnitudes

Two magnitudes are **homogeneous** if they have the same dimension, i.e., if they can be expressed using the same unit of measurement.

- The length of a segment and the height of a solid are two homogeneous magnitudes; indeed, both can be expressed with the same unit of length.

- The area of a square and the volume of a cone are not homogeneous magnitudes; indeed, area is expressed in square meters or in other units of surface, which however cannot be used to express volume.

Definition 5.2: of commensurable and incommensurable magnitudes

Two homogeneous magnitudes a and b are called **commensurable** if their ratio is a rational number, i.e., if $a/b \in \mathbb{Q}$. They are instead called **incommensurable** if their ratio is not a rational number, i.e., if $a/b \notin \mathbb{Q}$.

Example 5.1: of commensurable magnitudes

The volume of a sphere and the volume of the cylinder circumscribed about the sphere are commensurable. In formulas, the volume V_s of the sphere as a function of its radius r is given by the formula:

$$V_s = \frac{4}{3} \pi r^3$$

The cylinder circumscribed about it has base radius equal to the radius r of the sphere and height h equal to twice the radius. The volume V_c of this cylinder is therefore given by the formula:

$$V_c = \pi r^2 h = \pi r^2 2r = 2 \pi r^3$$

Taking the ratio we obtain:

$$\frac{V_s}{V_c} = \frac{\frac{4}{3} \pi r^3}{2 \pi r^3} = \frac{2}{3} \in \mathbb{Q}$$

Example 5.2: of incommensurable magnitudes

The length of a circle and the length of its diameter are incommensurable. In formulas, denoting by d the diameter of the circle and by c the length of the circle, we have:

$$c = \pi d$$

Now, taking the ratio between the length of the circle and its diameter, we obtain:

$$\frac{c}{d} = \frac{\pi d}{d} = \pi \notin \mathbb{Q}$$


A proof that $\pi \notin \mathbb{Q}$ will be given later.

5.3 Geometric representation of $\mathbb{Q}$

A geometric representation of $\mathbb{Q}$ can be obtained by associating with each rational number a point of the **Euclidean line** (which can be imagined as a line in the plane of infinite length)

- With an arbitrarily chosen point of the Euclidean line we associate 0, and with another point, distinct from the first, we associate 1, thus identifying the oriented segment 01, which constitutes the **unit of measurement**
- At this point we have a **one-to-one correspondence** between the rational numbers and those points P of the line that are endpoints of oriented segments $0P$ **commensurable** with 01. They are clearly commensurable since:

$$\frac{P}{1} \in \mathbb{Q} \quad \text{with} \quad P \in \mathbb{Q}$$



5.4 Total order relations

Observation 5.3

The sets of rational numbers $\mathbb{Q}$ and of real numbers $\mathbb{R}$ with the relation “less than or equal to” (“ $\leq$ ”) are totally ordered sets.

- The relations “less than or equal to” (“ $\leq$ ”) for the set of rational or real numbers:

$$R_{\leq} = \left\{ (a, b) : a, b \in \mathbb{Q} \quad \text{and} \quad a \leq b \right\}, \quad R_{\leq} = \left\{ (a, b) : a, b \in \mathbb{R} \quad \text{and} \quad a \leq b \right\}$$

are **partial order relations**. That is, they satisfy the following properties:

$$\forall a, \quad a \leq a \quad (\text{reflexive property})$$

$$\forall a, b, \quad a \leq b \quad \text{and} \quad b \leq a \Rightarrow a = b \quad (\text{antisymmetric property})$$

$$\forall a, b, c, \quad a \leq b \quad \text{and} \quad b \leq c \Rightarrow a \leq c \quad (\text{transitive property})$$

Moreover, the relations “less than or equal to” (“ $\leq$ ”) for $\mathbb{Q}$ and $\mathbb{R}$ are **total relations** since:

$$\forall a, b, \quad a \leq b \quad \text{or} \quad b \leq a$$

- Consequently, the pair consisting of the set $\mathbb{Q}$ or $\mathbb{R}$ and the corresponding relation $R_{\leq}$ is a totally ordered set.

5.5 Property R_3

- We now highlight the properties of the ordering of rational or real numbers:

$R_3 \rightarrow$ In $\mathbb{Q}$ and $\mathbb{R}$ the total order relation “less than or equal to” (“ $\leq$ ”) is defined, compatible with the algebraic structure ^a, that is:

1. $\forall a, b, c, \quad a \leq b \Rightarrow a + c \leq b + c$
2. $\forall a, b, c > 0, \quad a \leq b \Rightarrow a \cdot c \leq b \cdot c$

^aan algebraic structure is a set, called the underlying set (of the structure), equipped with one or more operations

- We observe that all the rules of algebraic computation follow from the properties: R_1 , R_2 , R_3 .

Example 5.3

For example, all the usual procedures used to solve inequalities are a consequence of the axioms and of the algebraic properties of the sum and the product, expressed by R_1 and R_2 .

5.6 Ordered fields

Definition 5.4: of (ordered) field

An **ordered field** is a set on which two operations (sum and product) and a total order relation are defined, satisfying the properties R_1 , R_2 , R_3 . A set with only the properties R_1 , R_2 is called a **field**.


- Everything we have said so far about the operations of sum and product and about the total order relation “less than or equal to” (“ $\leq$ ”) holds both for the set of rational numbers and for the set of real numbers.

Observation 5.5

The sets of rational numbers $\mathbb{Q}$ and of real numbers $\mathbb{R}$ are ordered fields.

- From this point of view, therefore, $\mathbb{Q}$ and $\mathbb{R}$ appear to have similar properties. In what follows we will highlight the **property that substantially distinguishes $\mathbb{R}$ from $\mathbb{Q}$** and that makes $\mathbb{R}$ the **right setting for developing mathematical analysis**.
- We begin by observing that the **set of rational numbers is inadequate for expressing the lengths of segments** (but also areas, volumes, times, speeds, etc.)

As we have seen, there exist magnitudes that are not **commensurable** with each other. The classic example is given by the diagonal and the side of a square: if the side measures 1, the abscissa d that measures the diagonal is not a rational number.



Indeed, by the Pythagorean Theorem we have:

$$d^2 = 1^2 + 1^2 = 2$$

but, as we have seen, there is no rational number whose square is 2. Hence the point d on the line does not represent any rational number. This means that, after “occupying” the points of the line with the rational numbers, there are still some **empty places** left on it.

5.7 Bounded sets, maxima and minima

Definition 5.6: of bounded set

Let E be a set contained in $\mathbb{Q}$ or in $\mathbb{R}$. The set E is called **bounded** if there exist two numbers m and M such that:

$$\forall x \in E, \quad m \leq x \leq M$$

- E is **bounded below** if there exists a number m such that:

$$\forall x \in E, \quad m \leq x$$

- E is **bounded above** if there exists a number M such that:

$$\forall x \in E, \quad x \leq M$$

- It makes no difference whether we require m and/or M to belong to $\mathbb{Q}$ or to $\mathbb{R}$, since the condition only concerns the existence of such numbers.

Definition 5.7: of maximum and minimum of a set

An element $x_M \in E$ is called the **maximum** of E if:

$$\forall x \in E, \quad x \leq x_M$$

An element $x_m \in E$ is called the **minimum** of E if:

$$\forall x \in E, \quad x_m \leq x$$

The existence of the maximum and of the minimum of a set implies that the set is bounded, that is:

$$\text{existence of maximum and minimum} \Rightarrow \text{bounded set} \quad (5.1)$$

but the converse is not true (we will give a counterexample later):

$$\text{bounded set} \not\Rightarrow \text{existence of maximum and minimum} \quad (5.2)$$

Hence the fact that a set is bounded is a necessary but not sufficient condition for the set to have a maximum and a minimum. Moreover, the existence of a maximum and a minimum is a sufficient but not necessary condition for a set to be bounded.

From the contrapositive of (5.1) we have:

$$\text{unbounded set} \Rightarrow \text{non-existence of maximum and minimum}$$


that is, if a set is not bounded, it has no maximum or it has no minimum.

Example 5.4: of maxima and minima

Example	set E	minimum	maximum
I)	$\mathbb{N}$	0	does not exist
II)	even integers	does not exist	does not exist

Example 5.5: of maxima and minima

Example	set E	minimum	maximum
III)	$\left\{ \frac{1}{n} : n \in \mathbb{N} \setminus \{0\} \right\}$	does not exist	1




Example 5.6: of maxima and minima

Example	set E	minimum	maximum
IV)	$\left\{ \frac{n-1}{n+1} : n \in \mathbb{N} \right\}$	-1	does not exist

We have:

$$\frac{n-1}{n+1} = \frac{n+1-1-1}{n+1} = 1 - \frac{2}{n+1}$$



Note that sometimes, even though the set is bounded, it may have no maximum or no minimum.

Example 5.7: of maxima and minima

Example	set E	minimum	maximum
V)	$\{x \in \mathbb{R} : 27 \leq x^3\}$	3	does not exist


We have:

$$27 \leq x^3 \iff \sqrt[3]{27} = 3 \leq x \quad \text{and} \quad 3 \in E$$

Example 5.8: of maxima and minima

Example	set E	minimum	maximum
VI)	$\{x \in \mathbb{Q} : x \geq 0, x < \sqrt{2}\}$	0	does not exist


The value $\sqrt{2}$ is irrational, hence by truncating we obtain rational numbers that get closer and closer to $\sqrt{2}$, none of which has the property of being the maximum.



It is an example of a bounded set that has no maximum.

Example	set E	minimum	maximum
VII)	$\{x \in \mathbb{Q} : x > \sqrt{2}, x \leq 4\}$	does not exist	4

The value $\sqrt{2}$ is irrational, hence by truncating and rounding up we obtain rational numbers that get closer and closer to $\sqrt{2}$, none of which has the property of being the minimum.



It is an example of a bounded set that has no minimum.

5.8 Upper/lower bounds and suprema/infima

Definition 5.8: of upper bound of a set

Let E be a set contained in $\mathbb{Q}$ or in $\mathbb{R}$; a number k belonging to $\mathbb{Q}$ or to $\mathbb{R}$ respectively is called an **upper bound** of E if:

$$\forall x \in E, \quad x \leq k$$

Definition 5.9: of lower bound of a set

Let E be a set contained in $\mathbb{Q}$ or in $\mathbb{R}$; a number k belonging to $\mathbb{Q}$ or to $\mathbb{R}$ respectively is called a **lower bound** of E if:

$$\forall x \in E, \quad k \leq x$$

We observe that a set bounded above (below) has many upper (lower) bounds. Moreover, the upper (lower) bounds do not necessarily belong to the set itself.

Definition 5.10: of supremum

The **supremum** of E (denoted by $\sup E$) is the minimum of the upper bounds of E .

Definition 5.11: of infimum

The **infimum** of E (denoted by $\inf E$) is the maximum of the lower bounds of E .

We observe that the supremum (infimum) may not exist (unbounded sets). We also observe that if the set has a maximum (minimum), it coincides with the supremum (infimum).

Example 5.9: of suprema and infima

Example	set E	$\inf E$	$\sup E$	minimum	maximum
I)	$\{x \in \mathbb{Q} : x \geq 0, x < \sqrt{2}\}$	0	does not exist	0	does not exist

We have:

$$\sqrt{2} \notin E \quad \text{and} \quad \sqrt{2} \notin \mathbb{Q}$$

hence the maximum does not exist, and neither does the sup.

Example	set E	$\inf E$	$\sup E$	minimum	maximum
II)	$\{x \in \mathbb{R} : x \geq 0, x < \sqrt{2}\}$	0	$\sqrt{2}$	0	does not exist

We have:

$$\sqrt{2} \notin E \quad \text{and} \quad \sqrt{2} \in \mathbb{R}$$

that is, the maximum does not exist, but $\sqrt{2}$ is the minimum of the upper bounds and therefore it is the sup.

5.9 Axiomatic definition of the real numbers

A totally ordered number set X is said to have the **supremum property** (least upper bound property) if:

$R_4 \rightarrow$ Every non-empty set $E \subset X$ that is bounded above has a supremum in X

- It is easy to prove that, if this property holds, then it is also true that every non-empty subset of X that is bounded below has an infimum.
- It is not required that X itself has a supremum (for example, in the case $X = \mathbb{Q}$ or $\mathbb{R}$ this is certainly false!), but that every non-empty subset of X that is bounded above has one.

Example VI shows that $\mathbb{Q}$ certainly does not have the supremum property. On the other hand, $\mathbb{R}$ has this property. ^a

^aHowever, we will not present a formal proof.

- In the **axiomatic definition** of $\mathbb{R}$, this property is part of the very definition of $\mathbb{R}$

Definition 5.12: (axiomatic) of the real numbers

We call $\mathbb{R}$ a set that satisfies the properties R_1, R_2, R_3, R_4 , that is, an ordered field that has the supremum property

- The supremum property is also called the **Dedekind axiom**, or **axiom of continuity**, or **completeness axiom**, and it can also be stated in the following equivalent form.

Let $\{A, B\}$ be a partition of $\mathbb{R}$ (i.e., A and B are non-empty disjoint sets whose union is $\mathbb{R}$); the partition is called a **cut** (section) if:

$$\forall a \in A \text{ and } \forall b \in B \text{ we have } a < b.$$

Then one proves that:

$R'_4 \rightarrow$ For every cut $\{A, B\}$ of $\mathbb{R}$ there exists a unique real number s (called the **separating element**) such that:


$$\forall a \in A \text{ and } \forall b \in B, \quad a \leq s \leq b$$

This separating element is $\sup A$ and $\inf B$, hence $\sup A = \inf B$

5.10 Geometric representation of $\mathbb{R}$

Can the set of real numbers $\mathbb{R}$ be put in one-to-one correspondence with the points of the Euclidean line?

- Let us now return to the problem of the incommensurability of the side and the diagonal of a square. When, with straightedge and compass, having fixed the unit side of the square, we construct its diagonal and transfer it onto the starting line, the arc drawn by the compass "splits the line in two", generating what we have called a cut; the separating element, which exists by axiom R_4 , geometrically represents the point of intersection, and therefore that number measures the diagonal.



Hence the geometric interpretation of property R_4 shows us how the set $\mathbb{R}$ is an adequate representation of our intuitive idea of a line, so adequate that **in mathematics the expression “the real line” is often used to denote $\mathbb{R}$** , identifying the number set with its natural geometric representation.

Roots, powers, logarithms and modular arithmetic

6.1 Roots, powers, logarithms

- As a consequence of property R_4 we can perform, in the real field, operations that are only occasionally possible in the rational field, such as extracting roots or raising to a power.

6.1.1 Arithmetic n -th roots

Theorem 6.1

For every $y \in \mathbb{R}$, $y > 0$ and $n \in \mathbb{N}$, $n \geq 1$, there exists one and only one $x \in \mathbb{R}$, $x > 0$, such that $x^n = y$.

- This number is called the arithmetic n -th root of y and is denoted by one of the symbols

$$\sqrt[n]{y} \text{ or } y^{\frac{1}{n}}.$$

Proof. The proof of this Theorem will be given using the properties of continuous functions. □

The arithmetic n -th root is non-negative.

Example 6.1: arithmetic n -th root

For example:

$$\sqrt{4} = 2, \quad \sqrt{9} = 3.$$

Moreover, we have:

$$\sqrt{x^2} = |x| \quad (6.1)$$

Construction of the decimal representation of the n -th root.

- We look for the decimal expansion of

$$\sqrt{2} \approx 1.41421356237$$

this number, not being rational, will be represented by an infinite (non-periodic) expansion.

- We proceed as follows: we build a class of rational numbers of the form:

$$\begin{aligned} 0 &< a_0 \\ &< a_0, a_1 \\ &< a_0, a_1 a_2 \\ &< a_0, a_1 a_2 a_3 \\ &< \dots \\ &< a_0, a_1 a_2 a_3 \dots a_n \\ &< \dots \end{aligned}$$

The rule is: each of these numbers is the largest among those with the same number of digits after the decimal point whose square is less than 2. The first of these numbers are:

$$\begin{array}{lll} 1 & 1^2 & = 1 \\ 1, 4 & (1, 4)^2 & = 1, 96 \\ 1, 41 & (1, 41)^2 & = 1, 9881 \\ 1, 414 & (1, 414)^2 & = 1, 999396 \\ 1, 4142 & (1, 4142)^2 & = 1, 99996164 \\ \dots & & \end{array}$$

- This set of numbers, which we call E_- , is bounded above (each one is < 2); by property R_4 it has a supremum, and this is used to define $\sqrt{2}$.

The number $\sqrt{2}$ is defined precisely as $\sup E_-$

- We could also have built a class of numbers E_+ like the previous one, with the rule that each of them is the smallest among those with the same number of digits after the decimal point whose square is greater than 2; we would have obtained:

$$\begin{array}{rcl}
 2 & 2^2 & = 4 \\
 1,5 & (1,5)^2 & = 2,25 \\
 1,42 & (1,42)^2 & = 2,0164 \\
 1,415 & (1,415)^2 & = 2,002225 \\
 1,4143 & (1,4143)^2 & = 2,00024449 \\
 \dots & &
 \end{array}$$

This set E_+ is bounded below (every element is > 1), therefore it has an infimum; one proves that:

$$\sqrt{2} = \inf E_+ = \sup E_-$$

The numbers of the class E_- approximate $\sqrt{2}$ from below, those of the class E_+ from above.

6.1.2 Powers with real exponent

Extracting the n -th root is the inverse operation of raising to an integer power.

Rational exponent

- The operation of raising to a power can be extended to every rational exponent if the base is positive (using the previous theorem).

$$\text{If } r = \frac{m}{n} \quad \text{and} \quad a > 0 \quad \text{then} \quad a^r = (a^m)^{\frac{1}{n}} = \sqrt[n]{a^m}$$

(we assume $m \in \mathbb{Z}$ and $n \in \mathbb{Z}$ positive)

Real exponent

- If the exponent is real, $b = b_0, b_1 b_2 b_3 \dots b_n \dots$, the number a^b ($a > 0$) will be determined by the class of numbers

$$a^{b_0} \quad a^{b_0, b_1} \quad a^{b_0, b_1 b_2} \quad \dots$$

in a way similar to the case of the root.

If the base a is negative, the operation of raising to a power a^b is defined only in certain cases:

1. if the exponent b is an integer, or
2. if the exponent $b = \frac{n}{m}$ is rational, provided that it is not the case that n is odd and m is even.

If $c < 0$ and m is odd, we define

$$\sqrt[m]{c} = -\sqrt[m]{-c}$$

Example 6.2: negative base

For example:

$$(-2)^{\frac{3}{5}} = \sqrt[5]{(-2)^3} = \sqrt[5]{-8} = -\sqrt[5]{8}$$

$$(-2)^{\frac{2}{5}} = \sqrt[5]{(-2)^2} = \sqrt[5]{4}$$

- When we say “does not exist in $\mathbb{R}$ ” we mean that it is not possible to define this operation in such a way that the usual rules of computation remain valid.

Example 6.3: non-existence of a power with real exponent

For example:

$$(-2)^{\frac{3}{4}}$$

does not exist in the real field!

Let

$$a, b \in \mathbb{R}, \quad a > 0 \quad \text{and} \quad b > 0 \quad (\text{positive reals})$$

and

$$c, d \in \mathbb{R} \quad (\text{arbitrary reals}),$$

the main properties of raising to a power are:

$$a^0 = 1 \quad \forall a \neq 0 \quad (6.2a)$$

$$1^c = 1 \quad \forall c \quad (6.2b)$$

$$a^c > 0 \quad \forall c \quad (6.2c)$$

$$a^c > 1 \quad \forall a > 1 \text{ and } c > 0 \quad (6.2d)$$

$$a^c < 1 \quad \forall a < 1 \text{ and } c > 0 \quad (6.2e)$$

$$a^{c+d} = a^c \cdot a^d \quad (6.3a)$$

$$(a \cdot b)^c = a^c \cdot b^c \quad (6.3b)$$

$$(a^b)^c = a^{b \cdot c} \quad (6.3c)$$

$$c < d \Rightarrow a^c < a^d \text{ if } a > 1 \quad (6.4a)$$

$$c < d \Rightarrow a^c > a^d \text{ if } a < 1 \quad (6.4b)$$

$$0 < a \leq b \Rightarrow a^c \leq b^c \quad \forall c > 0 \quad (6.4c)$$

6.1.3 Logarithms

- Consider the equation

$$a^x = y, \quad a > 0$$

with y given and x unknown. First of all, if $a = 1$, it is solvable only if $y = 1$ (and in that case every real number x is a solution). So let $a \neq 1$.

If $y \leq 0$ it has no solution, see property (6.2c).

The following theorem tells us that it has exactly one solution for every $y > 0$:

Theorem 6.2

Let $a > 0$, $a \neq 1$, $y > 0$. There exists a unique real number x such that $a^x = y$.

- This number is called the logarithm to base a of y and is denoted by the symbol:

$$\log_a y$$

- The logarithm is the inverse operation of raising to a power.
- the logarithm of a number to a given base is the exponent to which the base must be raised to obtain the number itself.

Let

$$x, y, a \in \mathbb{R}, \quad x > 0, y > 0, a > 0 \quad (\text{positive reals}) \quad \text{and} \quad a \neq 1,$$

the main properties of logarithms are (they follow from those of exponentials):

$$\log_a(x \cdot y) = \log_a x + \log_a y \quad (6.5a)$$

$$\log_a\left(\frac{x}{y}\right) = \log_a x - \log_a y \quad (6.5b)$$

$$\log_a\left(\frac{1}{y}\right) = -\log_a y \quad (6.5c)$$

$$\log_a x^\alpha = \alpha \log_a x \quad (\forall \alpha \in \mathbb{R}) \quad (6.6a)$$

$$\log_b x = \frac{\log_a x}{\log_a b} \quad (\forall b > 0, b \neq 1) \quad (6.6b)$$

$$\log_a x = \frac{1}{\log_x a} = -\log_{\frac{1}{a}} x \quad (x \neq 1) \quad (6.6c)$$

Notation:

- $\lg n = \log_2 n \rightarrow$ **binary logarithm**
- $\ln n = \log_e n \rightarrow$ **natural logarithm** (where $e \approx 2.718$ is **Euler's number** (Napier's constant)) (sometimes the natural logarithm $\ln$ is also denoted by $\log$)
- $\lg^k n = (\lg n)^k \rightarrow$ **power**
- $\lg \lg n = \lg(\lg n) \rightarrow$ **composition**

6.1.4 Approximations

- a rational number can always be expressed with absolute precision, using either the fractional or the decimal notation (possibly with repeating digits).

- On the other hand, it is of course not possible to write all the decimal digits of an irrational number, since there are infinitely many of them and they follow one another without periodicity.
- What does it mean, then, to “know” or “specify” an irrational number? It means knowing some algorithm that allows us (at least in theory) to write as many exact decimal digits as we wish.

Example 6.4: irrational numbers

- In the (already considered) example of the irrational number:

$$0,101001000100001\dots$$

built according to the rule: write one digit 1, one digit 0, one digit 1, two digits 0, one digit 1, three digits 0, and so on) it is clear that we could write as many digits as we wish.

- In other cases, such as

$$\sqrt{2} \text{ or } \log_2 3,$$

things are more laborious and require iterative computations to determine each successive decimal digit; nevertheless, these computations can actually be carried out.

- At other times, in Mathematical Analysis, a number is specified in a non-constructive way, by denoting it as the unique number that solves a given problem (once it has been proved that such a solution exists and is unique). This is a less satisfactory approach from an operational point of view, but theoretically flawless.

Practical aspects of computing irrational numbers

- Whenever we perform computations with a **pocket calculator**, it will only display numbers with a fixed number of decimal digits after the decimal point (typically 9). This means that we are working only with rational numbers, indeed with a finite subset of the set $\mathbb{Q}$.
- Working with a **computer** things improve somewhat, but we still remain within finite subsets of $\mathbb{Q}$. Of course, one needs to be aware of this.
- Besides the approximation produced by computing tools, sometimes we ourselves are not interested in too many decimal digits; we then deliberately introduce approximations, writing for example

$$\sqrt{2} \approx 1,414$$

Rounding rule: the last digit written is rounded down (up) if the first digit discarded is from 0 to 4 (respectively, from 5 to 9).

Example 6.5: Rounding

For example:

$$2,4138 \approx 2,41$$

$$2,4152 \approx 2,42$$

6.2 Modular arithmetic**Definition 6.3: of integer part**

Given a real number $a \in \mathbb{R}$, we denote by $[a]$ or $\lfloor a \rfloor$ the **integer part** (or “**floor**”) of a :

$$[a] = \text{integer } n \text{ such that } n \leq a < n + 1$$

Example 6.6: Integer part

$$[2,38] = 2; \quad [3] = 3; \quad [-1,8] = -2.$$

- While for positive numbers the integer part is obtained simply by “throwing away the digits after the decimal point”, for negative numbers one must take the largest integer $\leq a$, which is different from what is obtained by throwing away the digits after the decimal point (except when a is already an integer)

Definition 6.4: of fractional part

Given a real number $a \in \mathbb{R}$, we denote by (a) the **fractional part** (or decimal part, in Italian “*mantissa*”) of a :

$$(a) = a - [a]$$

Example 6.7: Fractional part

$$(2,38) = 0.38; \quad (3) = 0; \quad (-1,8) = 0.2.$$

- The fractional part is therefore not an integer but a real number, lying in $[0, 1)$.

- For positive numbers, it is obtained simply by “throwing away the digits before the decimal point”; for negative numbers the fractional part is the **one’s complement** of the number obtained by throwing away the digits before the decimal point.

Definition 6.5: of upper integer part

Given a real number $a \in \mathbb{R}$, we denote by $\lceil a \rceil$ the **upper integer part** (or “**ceil**”) of a :

$$\lceil a \rceil = \text{integer } n \text{ such that } n - 1 < a \leq n$$

Example 6.8: of ceil

$$\lceil 2,38 \rceil = 3; \quad \lceil 3 \rceil = 3; \quad \lceil -1,8 \rceil = -1.$$

Some properties:

Given a real number $a \in \mathbb{R}$, we have

$$a - 1 < \lfloor a \rfloor \leq a \leq \lceil a \rceil < a + 1$$

Given an integer $n \in \mathbb{Z}$, we have

$$\left\lfloor \frac{n}{2} \right\rfloor + \left\lceil \frac{n}{2} \right\rceil = n$$

Given two positive integers $r, s \in \mathbb{Z}$, $r, s > 0$, we have

$$\left\lceil \frac{r}{s} \right\rceil \leq \frac{r + (s - 1)}{s} \quad (6.7a)$$

$$\left\lfloor \frac{r}{s} \right\rfloor \geq \frac{r - (s - 1)}{s} \quad (6.7b)$$

Given, in addition, a non-negative real number $p \in \mathbb{R}$, $p \geq 0$, we have

$$\left\lceil \frac{\left\lfloor \frac{p}{r} \right\rfloor}{s} \right\rceil = \left\lceil \frac{p}{r s} \right\rceil \quad (6.8a)$$

$$\left\lfloor \frac{\left\lceil \frac{p}{r} \right\rceil}{s} \right\rfloor = \left\lfloor \frac{p}{r s} \right\rfloor \quad (6.8b)$$

Definition 6.6: Divisor

A **divisor** of an integer $n \in \mathbb{Z}$, also called a **factor** of n , is an integer $m \in \mathbb{Z}$ that can be multiplied by some integer $q \in \mathbb{Z}$ to obtain n , i.e., if $n = q \cdot m$.

- If m is a divisor of n , n is a **multiple** of m .
- An integer n is **divisible** by another integer m if m is a *divisor* of n .
- For an integer $a \in \mathbb{Z}$ and a positive integer $n \in \mathbb{Z}, n > 0$, the value $a \bmod n$ is the remainder of the division $\frac{a}{n}$.

Definition 6.7: of modulo (remainder)

Given $a \in \mathbb{Z}$ and $n \in \mathbb{Z}, n > 0$,

$$a \bmod n = a - n \left\lfloor \frac{a}{n} \right\rfloor$$

- It follows that $0 \leq a \bmod n < n$

If $(a \bmod n) = (b \bmod n)$, we write

$$a \equiv b \pmod{n}$$

and we say that a is **equivalent** to b , modulo n .

Example 6.9

For example, 23 and 13 are equivalent modulo 5 and we write $23 \equiv 13 \pmod{5}$

- Equivalently, $a \equiv b \pmod{n}$ if a and b have the same remainder when divided by n .
- Equivalently, $a \equiv b \pmod{n}$ if and only if n is a divisor of $|b - a|$.

Summations and geometric progressions

7.1 Summations

Definition 7.1: summation

Let $a_1, a_2, \dots, a_n$ be n real numbers. Their sum

$$a_1 + a_2 + \dots + a_n$$

can be written in compact form with the summation symbol:

$$\sum_{k=1}^n a_k$$

which is read: “sum for k from 1 to n of a_k ”. The symbol k is called the summation index.

- The summation symbol is therefore just a shorthand, which is nevertheless very useful when the terms a_k are defined explicitly as a function of the index k .

Example 7.1: summations

$$\sum_{k=1}^{10} \frac{1}{k} = 1 + \frac{1}{2} + \frac{1}{3} + \frac{1}{4} + \frac{1}{5} + \frac{1}{6} + \frac{1}{7} + \frac{1}{8} + \frac{1}{9} + \frac{1}{10}$$

$$\sum_{k=3}^n k^2 = 3^2 + 4^2 + 5^2 + \dots + n^2$$

- The summation index is a **dummy index**. This means that if we replace k with i , j or any other index (in all its occurrences), the value of the summation does not change.

Example 7.2: dummy index

We have:

$$\sum_{k=1}^n k^2 = \sum_{i=1}^n i^2$$

since both symbols denote the sum of the squares of the first n natural numbers (without zero).

On the other hand, we have:

$$\sum_{k=1}^n k^2 \neq \sum_{k=1}^m k^2$$

since the two symbols denote the sum of, respectively, the first n or the first m squares (if $n \neq m$ the result will be different).

7.1.1 Main properties of summations

Observation 7.2

Given $c \in \mathbb{R}$, we have:

$$\sum_{k=1}^n (c \cdot a_k) = c \sum_{k=1}^n a_k \quad (\text{product by a constant}) \quad (7.1)$$

$$\sum_{k=1}^n c = c \cdot n \quad (\text{summation with a constant term}) \quad (7.2)$$

Proof. First property (7.1). By the distributive property we have:

$$\underbrace{c a_1 + c a_2 + \cdots + c a_n}_{=\sum_{k=1}^n (c \cdot a_k)} = \underbrace{c (a_1 + a_2 + \cdots + a_n)}_{=c \sum_{k=1}^n a_k}$$

Second property (7.2):

$$\underbrace{c + c + \cdots + c}_{=\sum_{k=1}^n c \quad \text{that is, } c \text{ added } n \text{ times}} = c n$$

□

Observation 7.3

Given two summations $\sum_{k=1}^n a_k$ and $\sum_{k=1}^n b_k$, we have:

$$\sum_{k=1}^n a_k + \sum_{k=1}^n b_k = \sum_{k=1}^n (a_k + b_k) \quad (7.3)$$

Proof. We have:

$$\underbrace{a_1 + a_2 + \cdots + a_n + b_1 + b_2 + \cdots + b_n}_{=\sum_{k=1}^n a_k + \sum_{k=1}^n b_k} = \underbrace{a_1 + b_1 + a_2 + b_2 + \cdots + a_n + b_n}_{=\sum_{k=1}^n (a_k + b_k)}$$

□

Observation 7.4

Given two natural numbers $n, m \in \mathbb{N}$, we have:

$$\sum_{k=1}^{n+m} a_k = \sum_{k=1}^n a_k + \sum_{k=n+1}^{n+m} a_k \quad (\text{splitting}) \quad (7.4)$$

$$\sum_{k=1}^n a_k = \sum_{k=1+m}^{n+m} a_{k-m} = \sum_{k=1-m}^{n-m} a_{k+m} \quad (\text{index shift}) \quad (7.5)$$

$$\sum_{k=1}^n a_k = \sum_{k=1}^n a_{n-k+1} = \sum_{k=0}^{n-1} a_{n-k} \quad (\text{index reflection}) \quad (7.6)$$

Proof. The three properties are simply different ways of writing and/or ordering the terms of the summations. □

7.1.2 Some important summations

Observation 7.5: sum of the first n natural numbers (without zero)

For every natural number $n \geq 1$, we have:

$$\sum_{k=1}^n k = \frac{n(n+1)}{2}$$

Proof.

$$\begin{aligned} \sum_{k=1}^n k &= \frac{1}{2} \left(\sum_{k=1}^n k + \sum_{k=1}^n k \right) = \frac{1}{2} \left(\sum_{k=1}^n k + \sum_{k=1}^n (n - k + 1) \right) \\ &= \frac{1}{2} \sum_{k=1}^n (k + n - k + 1) = \frac{1}{2} \sum_{k=1}^n (n + 1) = \frac{n(n+1)}{2} \end{aligned}$$

□

Observation 7.6: sum of the first n odd numbers

For every natural number $n \geq 1$, we have:

$$\sum_{k=1}^n (2k-1) = n^2 \quad \text{or equivalently} \quad \sum_{k=0}^{n-1} (2k+1) = n^2$$

Proof. Using the properties of summations we have:

$$\begin{aligned} \sum_{k=1}^n (2k-1) &= 2 \sum_{k=1}^n k - \sum_{k=1}^n 1 \\ &= 2 \left(\frac{n(n+1)}{2} \right) - n = n^2 + n - n = n^2 \\ \sum_{k=0}^{n-1} (2k+1) &= 2 \sum_{k=0}^{n-1} k + \sum_{k=0}^{n-1} 1 \\ &= 2 \sum_{k=1}^n (k-1) + n = 2 \left(\sum_{k=1}^n k - \sum_{k=1}^n 1 \right) + n \\ &= 2 \left(\frac{n(n+1)}{2} - n \right) + n = n^2 + n - 2n + n = n^2 \end{aligned}$$

□

Observation 7.7: sum of the first n even numbers (without zero)

For every natural number $n \geq 1$, we have:

$$\sum_{k=1}^n 2k = n(n+1)$$

Proof.

$$\sum_{k=1}^n 2k = 2 \sum_{k=1}^n k = 2 \left(\frac{n(n+1)}{2} \right) = n(n+1)$$

□

7.2 Geometric progressions

Definition 7.8: geometric progression

A sequence of real numbers is a **geometric progression** if the ratio between each term (starting from the second one) and the previous one is constant. This constant is called the common ratio of the progression.

Given the first term $a \in \mathbb{R}$ and the common ratio $q \in \mathbb{R}$, the associated geometric progression is:

$$a, \quad aq, \quad aq^2, \quad aq^3, \quad aq^4, \quad \dots$$


Each term (starting from the second one) is obtained from the previous one by multiplying it by q . The k -th term ($k \in \mathbb{N}, k \geq 1$) can be written as aq^{k-1} and we have:

$aq^{1-1} = aq^0 = a$	first term, in position $k = 1$
$aq^{2-1} = aq^1 = aq$	second term, in position $k = 2$
$aq^{3-1} = aq^2$	third term, in position $k = 3$
$aq^{4-1} = aq^3$	fourth term, in position $k = 4$
$\dots$	$\dots$

Example 7.3: geometric progressions

- With $a = 1$ and $q = \frac{1}{2}$, the first 4 terms are:

$$1, \quad \frac{1}{2}, \quad \frac{1}{4}, \quad \frac{1}{8}$$



- With $a = 1$ and $q = 2$, the first 4 terms are:

$$1, \quad 2, \quad 4, \quad 8$$

7.2.1 Sums of the terms of geometric progressions
Observation 7.9: sum of the first n terms of the geometric progression ($a = 1$)

Given $q \in \mathbb{R}_+$, for every natural number $n \geq 1$ we have:

$$\sum_{k=1}^n q^{k-1} = \begin{cases} \frac{q^n - 1}{q - 1} & \text{if } q \neq 1 \\ n & \text{otherwise} \end{cases} \quad (7.7)$$

Proof. If $q \neq 1$, we prove the equation in the following equivalent form:

$$(q - 1) \sum_{k=1}^n q^{k-1} = q^n - 1$$

Applying the properties of summations, we get:

$$\begin{aligned}
 (q-1) \sum_{k=1}^n q^{k-1} &= q \sum_{k=1}^n q^{k-1} - \sum_{k=1}^n q^{k-1} = \\
 &= \sum_{k=1}^n q^k - \sum_{k=1}^n q^{k-1} = \sum_{k=1}^n q^k - \sum_{k=0}^{n-1} q^k = \\
 &= \sum_{k=1}^{n-1} q^k + q^n - \left(1 + \sum_{k=1}^{n-1} q^k\right) = q^n - 1
 \end{aligned}$$

If $q = 1$, we have instead:

$$\sum_{k=1}^n q^{k-1} = \sum_{k=1}^n 1 = n$$

□

- Given $q \in \mathbb{R}_+$, $n \in \mathbb{N}$, $n \geq 1$ and $a \in \mathbb{R}$, formula (7.7) extends as follows:

$$\sum_{k=1}^n a q^{k-1} = \begin{cases} a \left(\frac{q^n - 1}{q - 1} \right) & \text{if } q \neq 1 \\ a n & \text{otherwise} \end{cases} \quad \text{since} \quad \sum_{k=1}^n a q^{k-1} = a \sum_{k=1}^n q^{k-1} \quad (7.8)$$

clearly:

$$\frac{q^n - 1}{q - 1} = \frac{1 - q^n}{1 - q} \quad \text{and hence we also have} \quad \sum_{k=1}^n a q^{k-1} = \begin{cases} a \left(\frac{1 - q^n}{1 - q} \right) & \text{if } q \neq 1 \\ a n & \text{otherwise} \end{cases}$$

Example 7.4: sum of the first n terms of geometric progressions

- With $a = 1$ and $q = \frac{1}{2}$, the first 4 terms are:

$$1, \quad \frac{1}{2}, \quad \frac{1}{4}, \quad \frac{1}{8} \quad \text{and} \quad 1 + \frac{1}{2} + \frac{1}{4} + \frac{1}{8} = \frac{8 + 4 + 2 + 1}{8} = \frac{15}{8}$$

the sum of the first $n = 4$ terms is given by the formula:

$$\sum_{k=1}^4 \left(\frac{1}{2} \right)^{k-1} = \sum_{k=1}^4 \frac{1}{2^{k-1}} = \frac{1 - \frac{1}{2^4}}{1 - \frac{1}{2}} = \frac{1 - \frac{1}{16}}{\frac{1}{2}} = \frac{15}{16} \cdot 2 = \frac{15}{8}$$

- With $a = 1$ and $q = 2$, the first 4 terms are:

$$1, \quad 2, \quad 4, \quad 8 \quad \text{and} \quad 1 + 2 + 4 + 8 = 15$$

the sum of the first $n = 4$ terms is given by the formula:

$$\sum_{k=1}^4 2^{k-1} = \frac{2^4 - 1}{2 - 1} = 16 - 1 = 15$$

Principle of mathematical induction

8.1 The principle of mathematical induction

- We now present a proof technique called proof by induction. This procedure can be applied to theorems with the following structure:

“ for every $n \in \mathbb{N}$, $n \geq n_0$, property $p(n)$ holds ”

- The number n_0 is the smallest integer for which we want the property to be true; if $n_0 = 0$ the theorem simply states that the property is true for every $n \in \mathbb{N}$.

A proof by induction consists of the following two steps:

1. We prove that $p(n)$ is true for $n = n_0$ (**base case of the induction**).
2. We prove that, if n is an arbitrary natural number $\geq n_0$, the fact that $p(n)$ is true implies that $p(n + 1)$ is true (**inductive step**).

We can then conclude that for every $n \geq n_0$, $p(n)$ is true.

- The validity of this proof technique, *intuitively*, is based on the following fact:
 1. By point 1, we know that $p(n_0)$ is true. Suppose for example $n_0 = 1$: we therefore know that $p(1)$ is true (this must be proved explicitly).
 2. By point 2, since $p(1)$ is true, $p(2)$ will be true: indeed, we have proved that for any n , if $p(n)$ is true then $p(n + 1)$ is also true. But then, since $p(2)$ is true, $p(3)$ will be true; but then $p(4)$ is true, $\dots$ and so on, hence $p(n)$ is true for every $n \geq 1$.
- In practice, the proof consists of two phases.
 1. Prove $p(n_0)$ directly;

2. Assume $p(n)$ as a hypothesis (**inductive hypothesis**) and prove $p(n+1)$.

- This is the delicate point, often subject to misunderstandings. “Assuming as a hypothesis” $p(n)$ does not mean assuming the thesis as a hypothesis. What must be proved is that:

“ for every $n \geq n_0$, if $p(n)$ is true then $p(n+1)$ is also true ”

and not

“ if $p(n)$ is true for every n , then $p(n+1)$ is also true”

8.2 Proofs based on the principle of mathematical induction

8.2.1 Bernoulli's inequality

Observation 8.1: Bernoulli's inequality

For every integer $n \geq 0$, $x \in \mathbb{R}$, $x \geq -1$, we have:

$$(1+x)^n \geq 1+n x \quad (8.1)$$

Proof. By induction on n .

- **Base case of the induction**

Let $n = 0$. Then the statement becomes:

$$(1+x)^0 \geq 1+0 x \quad \text{i.e.} \quad 1 \geq 1$$

which is clearly true.

- **Inductive step**

Suppose it is true for n , and let us prove it for $(n+1)$. By the inductive hypothesis, we have:

$$(1+x)^n \geq 1+n x$$

and moreover we have

$$(1+x) \geq 0 \quad \text{since} \quad x \geq -1.$$

Then we can write:

$$\begin{aligned}
 (1+x)^{n+1} &= (1+x) \cdot (1+x)^n \\
 &\geq (1+x) \cdot (1+nx) \\
 &= 1 + (n+1)x + nx^2 \\
 &\geq 1 + (n+1)x
 \end{aligned}$$

where in the last inequality we used the fact that $nx^2 \geq 0$.



The chain of inequalities shows that, for $n+1$, we have


$$(1+x)^{n+1} \geq 1 + (n+1)x$$

which is exactly the desired statement, for $n+1$.

□

Example 8.1: Bernoulli's inequality





8.2.2 Some important summations

Observation 8.2: sum of the first n natural numbers

For every integer $n \geq 1$, we have:

$$\sum_{k=1}^n k = \frac{n(n+1)}{2}$$

Proof. By induction on n .

- **Base case of the induction**

Let $n = 1$. Then the statement becomes:

$$\sum_{k=1}^1 k = \frac{1(1+1)}{2} \quad \text{i.e.} \quad 1 = 1$$

which is clearly true.

- **Inductive step**

Suppose it is true for n , and let us prove it for $(n+1)$. By the inductive hypothesis, we have:

$$\sum_{k=1}^n k = \frac{n(n+1)}{2}.$$

Hence we can write:

$$\begin{aligned} \sum_{k=1}^{n+1} k &= \sum_{k=1}^n k + (n+1) \\ &= \frac{n(n+1)}{2} + (n+1) \\ &= \frac{n(n+1) + 2(n+1)}{2} \\ &= \frac{(n+1)(n+2)}{2} = \frac{(n+1)((n+1)+1)}{2} \end{aligned}$$

which is exactly the desired statement, for $n+1$.

□

8.2.3 Sum of the terms of the geometric progression

Observation 8.3: sum of the first n terms of the geometric progression ($a = 1$)

Given $q \in \mathbb{R}_+$, for every integer $n \geq 1$ we have:

$$\sum_{k=1}^n q^{k-1} = \begin{cases} \frac{q^n - 1}{q - 1} & \text{if } q \neq 1 \\ n & \text{otherwise} \end{cases} \quad (8.2)$$

Proof. By induction on n .

- **Base case of the induction**

Let $n = 1$. Then the statement becomes:

$$\sum_{k=1}^1 q^{k-1} = \frac{q^1 - 1}{q - 1} \quad \text{i.e.} \quad 1 = 1$$

which is clearly true.

- **Inductive step**

Suppose it is true for n , and let us prove it for $(n+1)$. By the inductive hypothesis, we have:

$$\sum_{k=1}^n q^{k-1} = \frac{q^n - 1}{q - 1}$$

Hence we can write

$$\begin{aligned}\sum_{k=1}^{n+1} q^{k-1} &= \sum_{k=1}^n q^{k-1} + q^n = \frac{q^n - 1}{q - 1} + q^n \\ &= \frac{q^n - 1 + q^{n+1} - q^n}{q - 1} = \frac{q^{n+1} - 1}{q - 1}\end{aligned}$$

which is exactly the desired statement, for $n+1$.

□

Factorials, binomial coefficients and triangle inequality

9.1 Factorials

Definition 9.1: factorial of n

The factorial of n is the product of the first n positive integers. It is denoted by $n!$ and is read “ n factorial”. In formulas:

$$n! = \prod_{k=1}^n k = 1 \cdot 2 \cdot 3 \cdot \dots \cdot (n-1) \cdot n$$

By definition, we set $0! = 1$.

- The number $n!$ grows very rapidly as n increases. The first values are:

n	0	1	2	3	4	5	6	7	8	9	10
$n!$	1	1	2	6	24	120	720	5.040	40.320	362.880	3.628.800

- Some properties of the factorial, which are immediate to verify, are:

$$n! = n \cdot (n-1)!$$

$$\frac{n!}{(n-k)!} = n \cdot (n-1) \cdot (n-2) \cdot \dots \cdot (n-k+1), \quad \text{with } k \geq 1 \quad (9.1)$$

With $k \geq 1$, it becomes the product of k factors, starting from n and decreasing by one unit at a time.

Example 9.1: Computing the factorial

$$\frac{100!}{95!} = \frac{100!}{(100-5)!} = 100 \cdot 99 \cdot 98 \cdot 97 \cdot 96 = 9.034.502.400$$

It is always convenient to simplify expressions containing the factorial as much as possible, before computing them!

9.2 Binomial coefficients**Definition 9.2: binomial coefficient**

The **binomial coefficient** is defined as the number:

$$c_{n,k} = \frac{n!}{k! (n-k)!} \quad \text{with } 0 \leq k \leq n. \quad (9.2)$$

The binomial coefficient $c_{n,k}$ is usually denoted by the symbol: $\binom{n}{k}$ which is read “ n choose k ”.

- Given rule (9.1), we have:

$$c_{n,k} = \frac{n \cdot (n-1) \cdot (n-2) \cdot \dots \cdot (n-k+1)}{k!} \quad (9.3)$$

with $k \geq 1$, an expression that is more convenient for computing the binomial coefficient.

We have:

$$\binom{n}{n-k} = \binom{n}{k} \quad (9.4)$$

Since:

$$\binom{n}{n-k} = \frac{n!}{(n-k)! (n-(n-k))!} = \frac{n!}{k! (n-k)!} = \binom{n}{k}$$

We have:

$$\binom{n-1}{k-1} + \binom{n-1}{k} = \binom{n}{k} \quad (9.5)$$

Since:

$$\begin{aligned}
 \binom{n-1}{k-1} + \binom{n-1}{k} &= \frac{(n-1)!}{(k-1)! \underbrace{(n-1-(k-1))!}_{=(n-k)! = (n-k)(n-k-1)!}} + \frac{(n-1)!}{k! (n-k-1)!} \\
 &= \frac{(n-1)!}{(k-1)! (n-k) (n-k-1)!} + \frac{(n-1)!}{k (k-1)! (n-k-1)!} \\
 &= \frac{k (n-1)! + (n-k) (n-1)!}{k (k-1)! (n-k) (n-k-1)!} = \frac{\overbrace{(n-1)!}^{=n!} n}{\underbrace{k (k-1)!}_{k!} \underbrace{(n-k) (n-k-1)!}_{(n-k)!}} \\
 &= \frac{n!}{k! (n-k)!} = \binom{n}{k}
 \end{aligned}$$

It also follows that:

$$\binom{n}{k-1} + \binom{n}{k} = \binom{n+1}{k}$$

9.2.1 Newton's formula

The n -th power of a binomial $(a+b)$ can be computed with the following formula (from which the name binomial coefficient derives):

Observation 9.3: Newton's formula

For every integer $n \geq 0$, with $a, b \in \mathbb{R}$, we have:

$$(a+b)^n = \sum_{k=0}^n \binom{n}{k} a^{n-k} b^k \quad (9.6)$$

Proof. By induction on n .

- **Base case of the induction**

Let $n = 0$. Then the statement becomes: $(a+b)^0 = \binom{0}{0} a^0 b^0$ i.e. $1 = 1$ which is clearly true.

- **Inductive step**

Suppose it is true for n , and let us prove it for $(n+1)$. By the inductive hypothesis,

we have: $(a + b)^n = \sum_{k=0}^n \binom{n}{k} a^{n-k} b^k$. Then:

$$\begin{aligned}
 (a + b)^{n+1} &= (a + b) \cdot (a + b)^n = (a + b) \sum_{k=0}^n \binom{n}{k} a^{n-k} b^k \\
 &= \sum_{k=0}^n \binom{n}{k} a^{n-k+1} b^k + \underbrace{\sum_{k=0}^n \binom{n}{k} a^{n-k} b^{k+1}}_{= \sum_{k=1}^{n+1} \binom{n}{k-1} a^{n-k+1} b^k} \\
 &= a^{n+1} + \sum_{k=1}^n \binom{n}{k} a^{n-k+1} b^k + b^{n+1} + \sum_{k=1}^n \binom{n}{k-1} a^{n-k+1} b^k \\
 &= a^{n+1} + b^{n+1} + \sum_{k=1}^n \underbrace{\left(\binom{n}{k} + \binom{n}{k-1} \right)}_{= \binom{n+1}{k}} a^{n-k+1} b^k \\
 &= \sum_{k=0}^{n+1} \binom{n+1}{k} a^{n+1-k} b^k
 \end{aligned}$$

which is exactly the desired statement, for $n + 1$.

□

Observation 9.4

For every integer $n \geq 0$ and integer k such that $0 \leq k \leq n$, we have:

$$\sum_{k=0}^n \binom{n}{k} = 2^n \tag{9.7}$$

Proof. We write:

$$2^n = (1 + 1)^n$$

We now apply Newton's formula (9.6):

$$(1 + 1)^n = \sum_{k=0}^n \binom{n}{k} 1^{n-k} 1^k = \sum_{k=0}^n \binom{n}{k}$$

□

9.2.2 Recursive computation of binomial coefficients

Relation (9.5) allows us to compute the binomial coefficients $\binom{n}{k}$ by means of the so-called **Pascal's triangle** (or Tartaglia's triangle).

The rules for building the triangle are:

1. At the top of the triangle we place the number $\binom{0}{0} = 1$ (by definition).
2. On the sides we place the numbers $\binom{n}{0} = \binom{n}{n} = 1$ for every $n \geq 1$.
3. For $0 < k < n$, the number $\binom{n}{k}$ is written at the intersection of the n -th row and the k -th column.
4. The number $\binom{n}{k}$ is the sum of the two numbers located in the previous row, the one in the same column and the one in the previous column.

Example 9.2: Pascal's triangle (or Tartaglia's triangle)

Pascal's triangle with $n \leq 10$ and $k \leq 10$ is:

	$k = 0$	$k = 1$	$k = 2$	$k = 3$	$k = 4$	$k = 5$	$k = 6$	$k = 7$	$k = 8$	$k = 9$	$k = 10$
$n = 0$	1										
$n = 1$	1	1									
$n = 2$	1	2	1								
$n = 3$	1	3	3	1							
$n = 4$	1	4	6	4	1						
$n = 5$	1	5	10	10	5	1					
$n = 6$	1	6	15	20	15	6	1				
$n = 7$	1	7	21	35	35	21	7	1			
$n = 8$	1	8	28	56	70	56	28	8	1		
$n = 9$	1	9	36	84	126	126	84	36	9	1	
$n = 10$	1	10	45	120	210	252	210	120	45	10	1

To compute the binomial coefficient $\binom{5}{3}$, corresponding to the blue cell, we can use relation (9.5) and add the two binomial coefficients $\binom{4}{2}$ and $\binom{4}{3}$, corresponding to the red cells:

$$\binom{5}{3} = \binom{4}{2} + \binom{4}{3} = 6 + 4 = 10.$$

Example 9.3: computing the power of a binomial using Pascal's triangle

Thanks to the previous Pascal's triangle and to Newton's formula (9.6) we can compute:

$$(a + b)^5 = a^5 + 5 a^4 b + 10 a^3 b^2 + 10 a^2 b^3 + 5 a b^4 + b^5$$

9.3 Absolute value

Definition 9.5: absolute value

The **absolute value** of a real number $a \in \mathbb{R}$ (or **modulus** of a) is the non-negative number defined as follows:

$$|a| = \begin{cases} a & \text{if } a \geq 0 \\ -a & \text{if } a < 0 \end{cases} \quad (9.8)$$

From the definition of absolute value it immediately follows that:

$$\forall \varepsilon \geq 0, a \in \mathbb{R}, \quad |a| \leq \varepsilon \iff -\varepsilon \leq a \leq \varepsilon \quad (9.9)$$

9.3.1 Triangle inequality in $\mathbb{R}$

Observation 9.6: triangle inequality in $\mathbb{R}$

$$|b + c| \leq |b| + |c| \quad \forall b, c \in \mathbb{R} \quad (9.10)$$

Proof. We write the two relations:

$$-|b| \leq b \leq |b|, \quad -|c| \leq c \leq |c|$$

and add them side by side:

$$-(|b| + |c|) \leq b + c \leq |b| + |c|$$

Hence, by (9.9), with $\varepsilon = |b| + |c| \geq 0$ and $a = b + c$, (9.10) follows. $\square$

- The triangle inequality is also used in the following form:

$$|d - e| \leq |d - f| + |e - f| \quad \forall d, e, f \in \mathbb{R} \quad (9.11)$$

To obtain it, it suffices to set in (9.10):

$$b = d - f, \quad c = f - e$$

we obtain:

$$|d - f + f - e| = |d - e| \leq |d - f| + |f - e| = |d - f| + |e - f|$$

since:

$$|f - e| = |e - f| \quad \forall e, f \in \mathbb{R}$$

- Moreover, the triangle inequality can also be written in the following form:

$$|g| \leq |g - h| + |h| \quad \text{i.e.} \quad |g| - |h| \leq |g - h| \quad \forall g, h \in \mathbb{R} \quad (9.12)$$

To obtain it, it suffices to set in (9.10):

$$b = g - h, \quad c = h$$

Observation 9.7: reverse triangle inequality in $\mathbb{R}$

$$||g| - |h|| \leq |g - h|, \quad \forall g, h \in \mathbb{R}. \quad (9.13)$$

Proof. From (9.12) we have

$$|g| - |h| \leq |g - h| \quad \forall g, h \in \mathbb{R}$$

Similarly, swapping g and h in (9.12) we obtain:

$$|h| - |g| \leq |h - g| = |g - h| \quad \text{that is} \quad |g| - |h| \geq -|g - h|$$

hence we have

$$-(|g - h|) \leq |g| - |h| \leq |g - h| \quad \forall g, h \in \mathbb{R}$$

Hence, by (9.9), with $\varepsilon = |g - h| \geq 0$ and $a = |g| - |h|$, the reverse triangle inequality follows. $\square$

- Inequality (9.10) can easily be extended to the case of k terms:

$$\left| \sum_{i=1}^k b_i \right| \leq \sum_{i=1}^k |b_i|. \quad (9.14)$$

- The following immediate properties also hold:

$$|bc| = |b| |c|, \quad \left| \frac{b}{c} \right| = \frac{|b|}{|c|}, \quad |-b| = |b| \quad \forall b, c \in \mathbb{R}. \quad (9.15)$$

The Fibonacci sequence

10.1 The Fibonacci sequence

Definition 10.1: Fibonacci sequence

$$\begin{aligned} F_0 &= 0, \\ F_1 &= 1, \\ F_i &= F_{i-1} + F_{i-2}, \quad \text{with } i \geq 2, i \in \mathbb{N} \end{aligned} \quad (10.1)$$

- Therefore, each Fibonacci number is the sum of the two previous ones, which gives the sequence

$$0, 1, 1, 2, 3, 5, 8, 13, 21, 34, 55, \dots$$

- The Fibonacci sequence is related to the **golden ratio** ϕ and to the **conjugate of the golden ratio** $\hat{\phi}$,

Definition 10.2: golden ratio


The **golden ratio** ϕ and the **conjugate of the golden ratio** $\hat{\phi}$ are the two roots of the equation:

$$x^2 = x + 1$$

and are given by the formulas:

$$\begin{aligned} \phi &= \frac{1 + \sqrt{5}}{2} & \hat{\phi} &= \frac{1 - \sqrt{5}}{2} \\ &\approx 1.61803 & &\approx -0.61803 \end{aligned}$$

The following figure shows the graph of the function $f(x) = x^2 - x - 1$ on the interval $[-3, 3]$; the red dots correspond to the roots of the equation $x^2 = x + 1$.



The golden ratio ϕ and the conjugate of the golden ratio $\hat{\phi}$ clearly satisfy the equation $x^2 = x + 1$ since:

$$\phi^2 = \left(\frac{1 + \sqrt{5}}{2} \right)^2 = \frac{1 + 2\sqrt{5} + 5}{4} = \frac{3 + \sqrt{5}}{2} = \frac{1 + \sqrt{5}}{2} + 1 = \phi + 1$$

$$\hat{\phi}^2 = \left(\frac{1 - \sqrt{5}}{2} \right)^2 = \frac{1 - 2\sqrt{5} + 5}{4} = \frac{3 - \sqrt{5}}{2} = \frac{1 - \sqrt{5}}{2} + 1 = \hat{\phi} + 1$$

Observation 10.3

$$F_i = \frac{\phi^i - \hat{\phi}^i}{\sqrt{5}}, \quad i = 0, 1, 2, \dots$$

Proof. By induction on i .

- **Base case of the induction**

We prove that the formula holds for $i = 0$ and $i = 1$:

$$F_0 = \frac{\phi^0 - \hat{\phi}^0}{\sqrt{5}} = \frac{1 - 1}{\sqrt{5}} = 0, \quad F_1 = \frac{\phi^1 - \hat{\phi}^1}{\sqrt{5}} = \frac{\sqrt{5}}{\sqrt{5}} = 1.$$

- **Inductive step**

Suppose it is true for $i = k$ and $i = k - 1$ with $k \geq 1$, and let us prove it for $i = k + 1$. By the inductive hypothesis, we have: $F_{k+1} = F_k + F_{k-1}$, hence we can write

$$\begin{aligned} F_{k+1} &= F_k + F_{k-1} \\ &= \frac{\phi^k - \hat{\phi}^k}{\sqrt{5}} + \frac{\phi^{k-1} - \hat{\phi}^{k-1}}{\sqrt{5}} = \frac{(\phi^k - \hat{\phi}^k) + (\phi^{k-1} - \hat{\phi}^{k-1})}{\sqrt{5}} \\ &= \frac{(\phi^k + \phi^{k-1}) - (\hat{\phi}^k + \hat{\phi}^{k-1})}{\sqrt{5}} = \frac{\phi^{k-1}(\phi + 1) - \hat{\phi}^{k-1}(\hat{\phi} + 1)}{\sqrt{5}} \\ &= \frac{\phi^{k-1}(\phi^2) - \hat{\phi}^{k-1}(\hat{\phi}^2)}{\sqrt{5}} = \frac{\phi^{k+1} - \hat{\phi}^{k+1}}{\sqrt{5}} \end{aligned}$$

which is exactly the desired statement, for $i = k + 1$.

□

Observation 10.4

$$F_i = \left\lfloor \frac{\phi^i}{\sqrt{5}} + \frac{1}{2} \right\rfloor, \quad i = 0, 1, 2, \dots$$

Proof. Since $|\hat{\phi}| < 1$, we have

$$\frac{|\hat{\phi}^i|}{\sqrt{5}} \leq \frac{1}{\sqrt{5}} < \frac{1}{2} \quad \text{and} \quad -\frac{1}{2} < \frac{\hat{\phi}^i}{\sqrt{5}} < \frac{1}{2}, \quad i = 0, 1, 2, \dots$$


Since $F_i \in \mathbb{N}$ and $F_i = \frac{\phi^i}{\sqrt{5}} - \frac{\hat{\phi}^i}{\sqrt{5}}$, the i -th Fibonacci number F_i is equal to $\frac{\phi^i}{\sqrt{5}}$ rounded to the nearest integer:

$$F_i = \left\lfloor \frac{\phi^i}{\sqrt{5}} + \frac{1}{2} \right\rfloor, \quad i = 0, 1, 2, \dots$$

□


- The values of

$$\frac{\hat{\phi}^i}{\sqrt{5}} \text{ with } i = 0, 1, \dots, 10 \text{ are :}$$



- The values of


$$\frac{\phi^i}{\sqrt{5}} - \left\lfloor \frac{\phi^i}{\sqrt{5}} \right\rfloor \text{ with } i = 0, 1, \dots, 10 \text{ are :}$$



- Since $F_i \in \mathbb{N}$, we moreover have:

$$\left(\frac{\phi^i}{\sqrt{5}} - \left\lfloor \frac{\phi^i}{\sqrt{5}} \right\rfloor \right) - \frac{\hat{\phi}^i}{\sqrt{5}} \in \{0, 1\}, \quad i = 0, 1, 2, \dots$$

- The values of F_i , with $i = 0, 1, \dots, 10$, are:



Cardinality of infinite sets

11.1 Countable cardinality

Is it possible to compare the “size” of infinite sets?

- Let us approach the question starting from the important number sets that we have introduced: $\mathbb{N}$, $\mathbb{Z}$, $\mathbb{Q}$, $\mathbb{R}$, and ask ourselves: how many elements does each of these sets have?
- Intuitively, the answer seems obvious: each of these sets has infinitely many elements; however, the rational numbers are more numerous than the integers, since $\mathbb{Z} \subset \mathbb{Q}$, and for the same reason the real numbers are more numerous than the rational numbers, since $\mathbb{Q} \subset \mathbb{R}$.
- How can we state, at the same time, that two sets are both infinite, but one is more numerous than the other?

How is it possible, in general, to compare the size of infinite sets?

- To give meaning to these questions, even before being able to answer them, we must define what we mean by two sets having the same size.
- Abstracting from the experience of counting the elements of a finite set, the idea of **same size** has been identified with that of **one-to-one correspondence** (bijection):

Definition 11.1: equal cardinality of two sets

Two sets A, B are said to have **equal cardinality** (or **power**) if they can be put in one-to-one correspondence with each other, i.e., if there exists a rule that associates with each element of A one and only one element of B , and vice versa.

- The cardinality (or power) of a set expresses the intuitive idea of size

Observation 11.2

The set of integers $\mathbb{Z}$ and the set of natural numbers $\mathbb{N}$ have the same cardinality

Proof. The following rule puts the integers and the natural numbers in one-to-one correspondence:

$$\begin{array}{cccccccccc}
 \mathbb{Z} & 0 & 1 & -1 & 2 & -2 & \dots & n & -n & \dots \\
 & \updownarrow & \updownarrow & \updownarrow & \updownarrow & \updownarrow & & \updownarrow & \updownarrow & \\
 \mathbb{N} & 0 & 1 & 2 & 3 & 4 & \dots & 2n-1 & 2n & \dots
 \end{array}$$

□

- Therefore, even though from the point of view of inclusion $\mathbb{Z}$ has “more elements” than $\mathbb{N}$ (in the sense that it has all the elements of $\mathbb{N}$ plus others), the two sets have the same cardinality.

Two sets that have the same cardinality are also called **equipotent** and should be thought of as equally numerous.

Definition 11.3: countable set

A set that has the same cardinality as $\mathbb{N}$ is called **countable**.

- The term countable indicates that the elements of the set can be enumerated, that is, arranged in a numbered list (place 1, place 2, place 3, ...)

The cardinality of $\mathbb{N}$ is called the **cardinality of the countable** (countable infinity).

Theorem 11.4

The set $\mathbb{Q}$ is countable

Proof. • We begin by proving that the set of positive rational numbers is countable.

- To do this, we represent the positive rational numbers as fractions n/m , with n, m positive integers, and we arrange these fractions in an **infinite triangular table**, in the following way:

$\frac{n}{m}$, with $n + m = \dots$				
2	$\frac{1}{1}$			
3	$\frac{1}{2}$	$\frac{2}{1}$		
4	$\frac{1}{3}$	$\frac{2}{2}$	$\frac{3}{1}$	
5	$\frac{1}{4}$	$\frac{2}{3}$	$\frac{3}{2}$	$\frac{4}{1}$
...	...			

Note that: all the fractions representing positive rational numbers appear in this table at least once; some numbers are repeated several times (e.g., $\frac{1}{1} = \frac{2}{2}$). Each row has finite length.

- Hence we can put $\mathbb{N}$ in one-to-one correspondence with the set of positive rationals, by going through the table row by row (and skipping an element when it is equal to one already encountered). For example:

1	2	3	4	5	6	7	8	9	...
$\frac{1}{1}$	$\frac{1}{2}$	$\frac{2}{1}$	$\frac{1}{3}$	$\frac{3}{1}$	$\frac{1}{4}$	$\frac{2}{3}$	$\frac{3}{2}$	$\frac{4}{1}$	...

(note that we skipped $\frac{2}{2}$ because it is equal to $\frac{1}{1}$, already encountered).

- This proves that the set of positive rationals is countable.
- Then $\mathbb{Q}$ is also countable, and this can be proved with an argument similar to the one we used to prove that $\mathbb{Z}$ is countable: denoting by $q_1, q_2, q_3, \dots$ the positive rationals, we set:

$\mathbb{N}$	0	1	2	3	4	5	6	...
	$\updownarrow$	$\updownarrow$	$\updownarrow$	$\updownarrow$	$\updownarrow$	$\updownarrow$	$\updownarrow$	
$\mathbb{Q}$	0	q_1	$-q_1$	q_2	$-q_2$	q_3	$-q_3$	...

which gives a one-to-one correspondence between $\mathbb{N}$ and $\mathbb{Q}$.

□

- Having discovered that several infinite sets, each properly contained in the next ($\mathbb{N}, \mathbb{Z}, \mathbb{Q}$), have the same cardinality, might suggest that this is true for all infinite sets. This is not true, as shown in the next section.

11.2 Cardinality of the continuum

Theorem 11.5

The set of real numbers $\mathbb{R}$ is not countable

- The proof technique is called **Cantor's diagonal argument**

Proof. • We will prove that the interval $[0, 1]$ has uncountable cardinality, from which the uncountability of $\mathbb{R}$ obviously follows

- Suppose therefore, **by contradiction**, that $[0, 1]$ is countable, and arrange **all** the real numbers of the interval $[0, 1]$ in a list $r_1, r_2, r_3, \dots$
- We write each number r_i in decimal form:

$$0.a_1a_2a_3\dots$$

where the a_i are digits from 0 to 9; if the digits are all zero we have $r_i = 0$ and if they are all nine we have $r_i = 0.\overline{9} = 1$):

$$r_1 = 0.a_{11}a_{12}a_{13}\dots$$

$$r_2 = 0.a_{21}a_{22}a_{23}\dots$$

$$r_3 = 0.a_{31}a_{32}a_{33}\dots$$

...

- We now define the following decimal number:

$$r = 0.b_1b_2b_3\dots$$

where the digits b_i are defined by the following rule:

$$b_i = \begin{cases} 5 & \text{if } a_{ii} \text{ is one of the digits } 0, 1, 2, 3 \text{ or } 4 \\ 4 & \text{if } a_{ii} \text{ is one of the digits } 5, 6, 7, 8 \text{ or } 9 \end{cases}$$

With this definition we have $b_i \neq a_{ii}$ for every i . Note that the number r has been constructed by reasoning on the diagonal of the infinite table whose rows are the numbers r_i , hence the name of the argument. □


Proof. • We now observe that r is a real number belonging to the interval $[0, 1]$ (because it is of the form $0.b_1b_2b_3\dots$ with all digits equal to 4 or 5) and on the other hand it is not equal to any of the numbers r_i in the list, since:

→ $r \neq r_1$: the first digit of r is different from the first digit of r_1 ($b_1 \neq a_{11}$);
 → $r \neq r_2$: the second digit of r is different from the second digit of r_2 ($b_2 \neq a_{22}$);
 → ...

- But this leads to a **contradiction**, since we had assumed that the r_i completely exhausted the set of real numbers of the interval $[0, 1]$. Therefore $[0, 1]$ is not countable. □

Since $\mathbb{N}$ cannot be put in one-to-one correspondence with $\mathbb{R}$, but it can be put in one-to-one correspondence with a proper subset of $\mathbb{R}$ ($\mathbb{N}$ itself!), we say that $\mathbb{N}$ has **smaller cardinality** than $\mathbb{R}$.

- We also note that the interval $[0, 1]$, and any interval of $\mathbb{R}$ (open or closed, bounded or unbounded), has the same cardinality as $\mathbb{R}$.
- This fact can be proved, for example, with an elementary geometric construction that gives a one-to-one correspondence between the points of the line and those of a segment $[A, B]$; as shown in the following figure, by drawing suitable segments from the two fixed points P_1 , P_2 to the line itself:



- This implies that, for example, the points of a line have the same cardinality as the points of a segment.

The cardinality of $\mathbb{R}$ is called the **cardinality of the continuum**.

- It is not only every interval of $\mathbb{R}$ that has this same cardinality. It can be proved that the same holds for the plane, for three-dimensional space and for their “continuous” subsets: for example, a plane and a sphere both have the cardinality of the continuum.
- So far we have encountered only two hierarchical levels of infinity: the cardinality of the countable and that of the continuum. One should not believe that only these two exist! For example, the set of all subsets of $\mathbb{R}$ (its power set) is even more numerous than $\mathbb{R}$; and with this procedure we can always construct a set more numerous than a given set.
- Therefore the hierarchical levels of infinite cardinalities are themselves infinite.

Complex numbers


12.1 Definition of $\mathbb{C}$ and field structure

- We have denoted by $\mathbb{R}^2$ (short for $\mathbb{R} \times \mathbb{R}$) the set of ordered pairs (a, b) of real numbers.
- On these pairs we directly define the operations of sum and product with the following rules:


$$(a, b) + (c, d) = (a + c, b + d) \quad (12.1)$$

$$(a, b) \cdot (c, d) = (ac - bd, ad + bc) \quad (12.2)$$

Example 12.1: sum



Example 12.2: product



- This “sum” and this “product” satisfy the **commutative**, **associative** and **distributive** properties
- We also observe that, $\forall (a, b) \in \mathbb{R}^2$:

$$(a, b) + (0, 0) = (0, 0) + (a, b) = (a, b)$$

hence the pair $(0, 0)$ is the **identity element for the sum**. Moreover:

$$(a, b) \cdot (1, 0) = (1, 0) \cdot (a, b) = (a, b)$$

hence the pair $(1, 0)$ is the **identity element for the product**. We have:

$$(a, b) + (-a, -b) = (0, 0)$$

hence $(-a, -b)$ is the **opposite** of (a, b) . Moreover, if $(a, b) \neq (0, 0)$ then:

$$(a, b) \cdot \left(\frac{a}{a^2 + b^2}, \frac{-b}{a^2 + b^2} \right) = (1, 0)$$

hence the pair $(a/(a^2 + b^2), -b/(a^2 + b^2))$ is the **reciprocal** of (a, b) .

Definition 12.1: field of complex numbers

Properties R_1, R_2 are satisfied by the sum and the product defined above, and therefore the set $\mathbb{R}^2$ with this structure is a field, which we will call the **field of complex numbers** and denote by $\mathbb{C}$

- We now observe that $\mathbb{C}$ contains the subset $\mathbb{C}_0$ of pairs of the form $(a, 0)$; it is a subfield of $\mathbb{C}$, since the sum and product of pairs of this form are again pairs of the same form; indeed we have:

$$(a, 0) + (b, 0) = (a + b, 0) \quad \text{and} \quad (a, 0) \cdot (b, 0) = (a \cdot b, 0)$$

Moreover, $\mathbb{C}_0$ can be ordered by setting $(a, 0) < (b, 0)$ if $a < b$.

- If we then put the set of real numbers $\mathbb{R}$ in one-to-one correspondence with $\mathbb{C}_0$, by setting

$$(a, 0) \longleftrightarrow a$$

we can identify the real numbers $a \in \mathbb{R}$ with the complex numbers of the form $(a, 0) \in \mathbb{R}^2$. In this sense the field of complex numbers $\mathbb{C}$ is an **extension** of the field of real numbers $\mathbb{R}$.

Let us now consider the number $(0, 1) \in \mathbb{C}$. It has the remarkable property that:

$$(0, 1) \cdot (0, 1) = (-1, 0)$$

i.e., its square coincides with the real number -1

Definition 12.2: imaginary unit

The pair $(0, 1) \in \mathbb{C}$ is denoted by the letter “ i ” and is called the **imaginary unit**

12.2 Algebraic form of complex numbers

We observe that, if we write any complex number $(c, 0)$ simply as c , we have:

$$(a, b) = (a, 0) + \underbrace{(0, 1)}_{=i} \cdot (b, 0) = a + ib$$

With this notation, rules (12.1) and (12.2) are the ordinary rules of algebraic calculation, keeping in mind that $i^2 = -1$:

$$(a + ib) + (c + id) = (a + c) + i(b + d) \quad (12.3)$$

$$(a + ib) \cdot (c + id) = (ac - bd) + i(ad + bc) \quad (12.4)$$

Definition 12.3: algebraic form, real part and imaginary part


The expression:

$$z = a + ib \quad (12.5)$$

is called the **algebraic form of complex numbers**; a is called the **real part** of z and is denoted by $\Re(z)$ (or $\text{Re}(z)$) while b is called the **imaginary part** and is denoted by $\Im(z)$ (or $\text{Im}(z)$).

Complex plane

- In a Cartesian plane, the complex numbers $a + ib$ can be represented as points with coordinates (a, b) . In this context:
 - the plane is called the **complex plane** or **Gauss plane**
 - the x, y axes are called the **real axis** and the **imaginary axis**
 - the points on the real axis are the **real numbers**
 - the points on the imaginary axis are the **purely imaginary numbers** (i.e., of the form ib)
- The **sum of two complex numbers** is the complex number whose coordinates are the sums of the coordinates: the geometric meaning of this fact is that the point $z + t$ is constructed from the points z, t according to the “**parallelogram rule**”, illustrated in the following figure:

**Observation 12.4**

The set of complex numbers $\mathbb{C}$ is not an ordered field

Proof. • We have seen that $\mathbb{C}$ satisfies the field axioms; however, it does not satisfy those of an ordered field, that is, it is not possible to define a relation $\leq$ between complex numbers in such a way that properties $R3$ hold

- It can be proved that properties $R3$ imply that the square of any number is never

negative, and on the other hand, if a number is positive its opposite is negative. Now, in $\mathbb{C}$ we have:

$$1^2 = 1 \quad \text{and} \quad i^2 = -1$$

We therefore have two squares, each the opposite of the other. Neither of them, however, can be negative (because they are squares), and this is absurd (because between a and $-a$ one must be negative, if $a \neq 0$). We conclude that $\mathbb{C}$ is not an ordered field. □

Definition 12.5: conjugate

The complex number $a - ib$ is called the **complex conjugate** of $z = a + ib$ and is denoted by $\bar{z}$.

We have:

$$z + \bar{z} = 2a = (a + ib) + (a - ib) = 2a = 2 \Re(z) \quad (12.6)$$

$$z - \bar{z} = 2ib = (a + ib) - (a - ib) = 2ib = 2i \Im(z) \quad (12.7)$$

$$z \cdot \bar{z} = (a + ib) \cdot (a - ib) = a^2 - aib + iba - \underbrace{i^2}_{=-1} b^2 = a^2 + b^2 \geq 0 \quad (12.8)$$

- The conjugation operation has the following elementary properties with respect to sum and product:

$$\overline{(z_1 + z_2)} = \bar{z}_1 + \bar{z}_2 \quad (12.9)$$

$$\overline{(z_1 \cdot z_2)} = \bar{z}_1 \cdot \bar{z}_2 \quad (12.10)$$

$$\overline{\left(\frac{1}{z}\right)} = \frac{1}{\bar{z}} \quad (12.11)$$

Definition 12.6: modulus

The **modulus** of $z = a + ib$ is the non-negative real number $\sqrt{a^2 + b^2}$; it is denoted by $|z|$.

- If $z = a$ is real, its modulus is called the absolute value and is still denoted by $|a|$. The following properties hold:

- a) $|z| = 0 \iff z = 0$, moreover $|z| \geq 0$
 - b) $|z| = |\bar{z}|$
 - c) $\Re(z) \leq |z| \quad \Im(z) \leq |z| \quad |z| \leq |\Re(z)| + |\Im(z)|$
 - d) $|z_1 + z_2| \leq |z_1| + |z_2|$ **triangle inequality**
 - e) $|z_1 + z_2| \geq ||z_1| - |z_2||$
- Properties a), b), c) are immediately verified.

Proof. • Let us prove properties d) and e). They are equivalent to the following:

$$(|z_1| - |z_2|)^2 \leq |z_1 + z_2|^2 \leq (|z_1| + |z_2|)^2$$

- Setting $z_1 = a + ib$, $z_2 = c + id$ we obtain:

$$\begin{aligned} |z_1 + z_2|^2 &= |(a + ib) + (c + id)|^2 \\ &= |(a + c) + i(b + d)|^2 \\ &= \left(\sqrt{(a + c)^2 + (b + d)^2} \right)^2 = (a + c)^2 + (b + d)^2 \end{aligned}$$

and hence we have:

$$\begin{aligned} \underbrace{\left(\sqrt{a^2 + b^2} - \sqrt{c^2 + d^2} \right)^2}_{=(a^2+b^2)+(c^2+d^2)-2\sqrt{a^2+b^2}\cdot\sqrt{c^2+d^2}} &\leq \underbrace{(a + c)^2 + (b + d)^2}_{=a^2+2ac+c^2+b^2+2bd+d^2} \\ &\leq \underbrace{\left(\sqrt{a^2 + b^2} + \sqrt{c^2 + d^2} \right)^2}_{=(a^2+b^2)+(c^2+d^2)+2\sqrt{a^2+b^2}\cdot\sqrt{c^2+d^2}} \end{aligned}$$

After simplification, this double inequality reduces to:

$$-\sqrt{a^2 + b^2} \cdot \sqrt{c^2 + d^2} \leq ac + bd \leq \sqrt{a^2 + b^2} \cdot \sqrt{c^2 + d^2}$$

which is equivalent to the following:

$$|ac + db| \leq \sqrt{a^2 + b^2} \cdot \sqrt{c^2 + d^2}$$

Squaring both sides we arrive at:

$$\underbrace{(ac + db)^2}_{=a^2c^2+d^2b^2+2acbd} \leq \underbrace{(a^2 + b^2) \cdot (c^2 + d^2)}_{=a^2c^2+a^2d^2+b^2c^2+b^2d^2}$$


that is,

$$0 \leq -2acbd + a^2d^2 + b^2c^2 = (ad - bc)^2$$

which is true for every $a, b, c, d \in \mathbb{R}$.

□

- Geometrically, $|z|$ represents the distance of the point (or complex number) z from the origin; $|z_1 - z_2|$ represents the distance between the two points z_1 and z_2 ; inequalities d) and e) express the well-known theorem on the **lengths of the sides of a triangle**¹:



- Using the concepts just introduced, we can write the quotient of two complex numbers in algebraic form:

$$\frac{a + ib}{c + id}$$

it suffices to multiply the numerator and the denominator by $c - id$, and we obtain:

$$\frac{a + ib}{c + id} = \frac{(a + ib)(c - id)}{\underbrace{(c + id)(c - id)}_{=c^2+d^2}} = \frac{ac - aid + ibc - i^2bd}{c^2 + d^2} = \frac{(ac + bd)}{c^2 + d^2} + i \frac{(bc - ad)}{c^2 + d^2}$$

12.2.1 Equations in the complex field

- Let us see how to solve an equation in the complex field, when it involves the unknown $z = x + iy$ also through $\Re(z)$, $\Im(z)$, z , $|z|$.
- We illustrate, with the following example, the procedure of transforming the equation in one complex unknown into a system of two equations in two real unknowns. The method consists in taking the real and imaginary parts of the equation.

¹In a non-degenerate triangle, the sum of the lengths of two sides is greater than the length of the third one.

Example 12.3: equations in the complex field (algebraic method)

- We want to solve:

$$z^2 + i \Im z + 2 \bar{z} = 0$$

We set $z = x + iy$, with x, y real unknowns, and rewrite the equation:

$$\begin{cases} z^2 = (x + iy)^2 = x^2 - y^2 + 2ixy \\ i \Im z = iy \\ 2\bar{z} = 2(x - iy) = 2x - 2iy \end{cases}$$

Substituting, we therefore obtain:

$$(x^2 - y^2 + 2ixy) + (iy) + (2x - 2iy) = 0$$

- Now, a complex number is zero if and only if its real part and imaginary part are zero. Therefore we separate the real part and the imaginary part of the left-hand side:

$$(x^2 - y^2 + 2x) + i(2xy + y - 2y) = 0$$

and set both equal to zero:

$$\begin{cases} x^2 - y^2 + 2x = 0 \\ 2xy - y = 0 \end{cases}$$

We have thus transformed the equation in one complex unknown into a system of two equations in two real unknowns.

- We solve the system. The second equation gives:

$$y = 0 \quad \text{or} \quad x = \frac{1}{2}$$

1. For $y = 0$ the first equation becomes

$$x^2 + 2x = 0 \quad \text{which gives} \quad x = 0 \quad \text{or} \quad x = -2$$

2. For $x = \frac{1}{2}$ the first equation becomes

$$-y^2 + \frac{5}{4} = 0 \quad \text{which gives} \quad y = \pm \frac{\sqrt{5}}{2}$$


- Hence the equation has the following 4 solutions:

$$z = 0, \quad z = -2, \quad z = \frac{1}{2} + i \frac{\sqrt{5}}{2}, \quad z = \frac{1}{2} - i \frac{\sqrt{5}}{2}$$

The method seen in this example can in principle be applied to any equation in $\mathbb{C}$, but a generic system of two equations in two unknowns is almost always impossible to solve algebraically.

12.3 Trigonometric form of complex numbers

- As is known from Geometry, the points of the plane can be identified not only by their Cartesian coordinates, but also by their **polar coordinates**:
 - $\varrho \rightarrow$ **polar radius**, i.e., the distance of the point from the origin
 - $\vartheta \rightarrow$ **polar angle**, i.e., the angle that the line joining the point to the origin forms with the positive x -axis, measured counterclockwise.
- Clearly, a pair ϱ, ϑ , with $\varrho > 0$, identifies a well-defined point of the plane; conversely, a point of the plane uniquely determines the coordinate ϱ , but the angle ϑ , measured in radians, is determined only up to multiples of 2π .



- Given a complex number z , its **modulus** $|z|$ coincides with the **polar radius** ϱ of the point representing it in the complex plane.
- We call **argument** of z , and denote by $\arg(z)$, any of the angles ϑ associated with the point z . In this way the argument of z is not uniquely determined. Often this indeterminacy does not cause any problem. At other times, however, it is preferable to assign a well-defined argument to a complex number. This can be done in infinitely many ways, by fixing any interval of length 2π within which the angle ϑ is allowed to vary
- The intervals most commonly used for this purpose are $[0, 2\pi)$ and $(-\pi, \pi]$; the argument of z is then called the **principal argument**.

Example 12.4: arguments of complex numbers

- the number $-i$ has argument $-\pi/2$ or $3\pi/2$ or any other value of the form $-\pi/2 + 2k\pi$ with $k \in \mathbb{Z}$. Its principal argument will be $3\pi/2$ if we adopt the convention $\vartheta \in [0, 2\pi)$, and $-\pi/2$ with the convention $\vartheta \in (-\pi, \pi]$.
- Positive real numbers have principal argument 0 and negative ones π , with both conventions.

Given the number $z = a + ib$, from trigonometry we immediately obtain the relations between the Cartesian coordinates a, b and the polar ones ϱ, ϑ :

$$a = \varrho \cos \vartheta \quad \text{and} \quad b = \varrho \sin \vartheta \quad (12.12)$$

The inverse relations are:

$$\varrho = \sqrt{a^2 + b^2}, \quad \cos \vartheta = \frac{a}{\sqrt{a^2 + b^2}} \quad \text{and} \quad \sin \vartheta = \frac{b}{\sqrt{a^2 + b^2}} \quad (12.13)$$

Definition 12.7: trigonometric form

A complex number $z = a + ib$ can also be written in the form

$$z = \varrho (\cos \vartheta + i \sin \vartheta) \quad (12.14)$$

which is called the **trigonometric form** of complex numbers.

Example 12.5: trigonometric form

Let us write the following complex number in trigonometric form:

$$z = \sqrt{3} + i$$

We have $\varrho = \sqrt{a^2 + b^2} = \sqrt{3 + 1} = 2$, hence:

$$\sqrt{3} + i = 2 \left(\frac{\sqrt{3}}{2} + i \frac{1}{2} \right) = 2 \left(\cos \frac{\pi}{6} + i \sin \frac{\pi}{6} \right)$$

12.3.1 De Moivre's formulas

- The trigonometric form is convenient for expressing products and quotients of complex numbers. Indeed, if we have:

$$z_1 = \varrho_1 (\cos \vartheta_1 + i \sin \vartheta_1) \quad z_2 = \varrho_2 (\cos \vartheta_2 + i \sin \vartheta_2)$$

for the **product** we obtain:

$$\begin{aligned} z_1 z_2 &= \varrho_1 \varrho_2 \cdot \left\{ \cos \vartheta_1 \cos \vartheta_2 - \sin \vartheta_1 \sin \vartheta_2 + i (\sin \vartheta_1 \cos \vartheta_2 + \cos \vartheta_1 \sin \vartheta_2) \right\} \\ &= \varrho_1 \varrho_2 \cdot \left\{ \cos (\vartheta_1 + \vartheta_2) + i \sin (\vartheta_1 + \vartheta_2) \right\} \end{aligned} \quad (12.15)$$

for the **quotient**, if $z_2 \neq 0$, we have:

$$\frac{z_1}{z_2} = \frac{\varrho_1}{\varrho_2} \cdot \frac{\cos \vartheta_1 + i \sin \vartheta_1}{\cos \vartheta_2 + i \sin \vartheta_2},$$

multiplying numerator and denominator by $(\cos \vartheta_2 - i \sin \vartheta_2)$ and, taking into account that $(\cos \vartheta_2)^2 + (\sin \vartheta_2)^2 = 1$, we obtain:

$$\begin{aligned} \frac{z_1}{z_2} &= \frac{\varrho_1}{\varrho_2} \cdot \left\{ (\cos \vartheta_1 + i \sin \vartheta_1) \cdot (\cos \vartheta_2 - i \sin \vartheta_2) \right\} \\ &= \frac{\varrho_1}{\varrho_2} \cdot \left\{ \cos(\vartheta_1 - \vartheta_2) + i \sin(\vartheta_1 - \vartheta_2) \right\} \end{aligned} \quad (12.16)$$

- Therefore the modulus of the product and of the quotient of two complex numbers is, respectively, the product and the quotient of the moduli; the argument is, respectively, the sum and the difference of the arguments:

$$|z_1 \cdot z_2| = |z_1| \cdot |z_2| \quad \text{and} \quad \arg(z_1 \cdot z_2) = \arg(z_1) + \arg(z_2) \quad (12.17)$$

$$\left| \frac{z_1}{z_2} \right| = \frac{|z_1|}{|z_2|} \quad \text{and} \quad \arg\left(\frac{z_1}{z_2}\right) = \arg(z_1) - \arg(z_2) \quad (12.18)$$

- Formula (12.15) generalizes to the case of any number of factors $z_1, z_2, \dots, z_n$:

$$z_1 z_2 \dots z_n = \varrho_1 \varrho_2 \dots \varrho_n \cdot \left\{ \cos(\vartheta_1 + \vartheta_2 + \dots + \vartheta_n) + i \sin(\vartheta_1 + \vartheta_2 + \dots + \vartheta_n) \right\} \quad (12.19)$$

If, moreover, all the factors are equal, we obtain:

$$z^n = \varrho^n \cdot \left\{ \cos(n \vartheta) + i \sin(n \vartheta) \right\} \quad (12.20)$$

- These relations on products and quotients of complex numbers are known as De Moivre's formulas.

Example 12.6: powers of complex numbers with De Moivre's formulas

Write in algebraic form:

$$z = (1 + i)^7$$

We determine the modulus and argument of $(1 + i)$ and then apply De Moivre's formula.

$$|1 + i| = \sqrt{2} \quad \text{and} \quad \arg(1 + i) = \frac{\pi}{4} \quad \left(\cos \vartheta = \frac{1}{\sqrt{2}}, \quad \sin \vartheta = \frac{1}{\sqrt{2}} \right)$$

Hence we have:

$$|(1 + i)^7| = (\sqrt{2})^7 = 2^{\frac{7}{2}} = 2^{3+\frac{1}{2}} = 8\sqrt{2} \quad \text{and} \quad \arg(1 + i)^7 = \frac{7}{4} \pi$$

Consequently:

$$(1+i)^7 = 8\sqrt{2} \left(\underbrace{\cos \frac{7}{4}\pi}_{=\frac{1}{\sqrt{2}}} + i \underbrace{\sin \frac{7}{4}\pi}_{=-\frac{1}{\sqrt{2}}} \right) = 8\sqrt{2} \left(\frac{1}{\sqrt{2}} - i \frac{1}{\sqrt{2}} \right) = 8 - 8i$$

- De Moivre's formulas allow us to give a geometric interpretation of the product of complex numbers.
- Let z , to begin with, be a complex number with modulus 1, hence of the form $(\cos \vartheta + i \sin \vartheta)$. Then, multiplying a number by z means adding ϑ to its argument, i.e., performing a **rotation by the angle ϑ** .
- If z has modulus ϱ instead of 1, in addition to a rotation we perform a **dilation by the factor ϱ** .

Example 12.7: geometric interpretation of the product of complex numbers

- multiplying by i means performing a rotation by $\frac{\pi}{2}$;
- multiplying by -1 means performing a rotation by π ;
- multiplying by $(1+i)$ means performing a dilation by the factor $\sqrt{2}$ and a rotation by $\frac{\pi}{4}$

Example 12.8: equations in the complex field (trigonometric method)

- We want to solve:

$$z^3 - |z| = 0$$

We rewrite the equation in the form

$$z^3 = |z|$$

and we set $z = \varrho (\cos \vartheta + i \sin \vartheta)$ (trigonometric form) and rewrite the equation:

$$\begin{cases} z^3 = \varrho^3 (\cos 3\vartheta + i \sin 3\vartheta) \\ |z| = \varrho \end{cases}$$

Substituting, we therefore obtain:

$$\varrho^3 (\cos 3\vartheta + i \sin 3\vartheta) = \varrho$$

The equation is satisfied if and only if the two sides have equal moduli and arguments that differ by multiples of 2π (the right-hand side has argument 0), that is:

$$\begin{cases} \varrho^3 = \varrho \\ 3\vartheta = 2k\pi \quad \text{with } k \in \mathbb{Z} \end{cases}$$

The first equation gives $\varrho = 0$ and $\varrho = 1$ (careful: ϱ must be ≥ 0 because it is the modulus of the complex number; therefore $\varrho = -1$ is not acceptable); the second one gives $\vartheta = \frac{2k\pi}{3}, k \in \mathbb{Z}$. Hence:

$$z = 0, \quad z = \cos \frac{2k\pi}{3} + i \sin \frac{2k\pi}{3} \quad \text{with } k \in \mathbb{Z}$$

Explicitly:

$$z = 0, \quad z = 1, \quad z = -\frac{1}{2} + i \frac{\sqrt{3}}{2}, \quad z = -\frac{1}{2} - i \frac{\sqrt{3}}{2}$$

12.3.2 n -th roots of complex numbers

Definition 12.8: n -th root of a complex number

Given a complex number w , we say that z is an n -th (complex) root of w if $z^n = w$.

Theorem 12.9

Let $w \in \mathbb{C}$, $w \neq 0$, and n an integer ≥ 1 . There exist exactly n complex n -th roots $z_0, z_1, \dots, z_{n-1}$ of w ; setting

$$w = r (\cos \varphi + i \sin \varphi) \quad \text{and} \quad z_k = \varrho_k (\cos \vartheta_k + i \sin \vartheta_k)$$

we have

$$\begin{cases} \varrho_k = r^{1/n} \\ \vartheta_k = \frac{\varphi + 2k\pi}{n} \end{cases} \quad k = 0, 1, 2, \dots, n-1 \quad (12.21)$$

Proof. • The numbers z_k are clearly roots of w , as can be seen by computing z_k^n with De Moivre's formula. Let us show that there are no others.

- If a number $R(\cos \psi + i \sin \psi)$ is an n -th root of w , we must have

$$R^n = r \quad \text{and} \quad n \psi = \varphi + 2h\pi \quad \text{with } h \in \mathbb{Z}$$

or equivalently:

$$R = r^{1/n} \quad \text{and} \quad \psi = \varphi/n + 2h\pi/n \quad \text{with} \quad h \in \mathbb{Z}$$

- Giving h the values $0, 1, \dots, n-1$ we find exactly the numbers z_k .
- Giving h any other value $\bar{h}$ different from the previous ones, it can be written in the form $\bar{h} = k + mn$ ($m \in \mathbb{Z}$ is the quotient and k is the remainder of the division of $\bar{h}$ by n), so that we would have

$$\psi = \frac{\varphi}{n} + \frac{2k\pi}{n} + 2m\pi = \vartheta_k + 2m\pi$$

and we would find again the same z_k as before. □

Example 12.9: fifth root of a complex number

- Let us compute $\sqrt[5]{1+i}$: the complex number $(1+i)$ has modulus $\sqrt{2}$ and argument $\pi/4$. The fifth roots will therefore be:

$$\begin{aligned} \sqrt[5]{1+i} &= \sqrt[5]{\sqrt{2}} \left[\cos \left(\frac{\frac{\pi}{4} + 2k\pi}{5} \right) + i \sin \left(\frac{\frac{\pi}{4} + 2k\pi}{5} \right) \right] = \\ &= \sqrt[10]{2} \left[\cos \left(\frac{\pi}{20} + \frac{2}{5}k\pi \right) + i \sin \left(\frac{\pi}{20} + \frac{2}{5}k\pi \right) \right] \quad \text{with} \quad k = 0, 1, 2, 3, 4 \end{aligned}$$

- The angles found are not standard angles, but if desired, their sines and cosines can be computed approximately using a calculator.

- Unfortunately, a somewhat ambiguous notation is used for complex roots, the same one used to denote the arithmetic root; that is, $\sqrt[n]{z}$ or $z^{1/n}$ denotes the set of the n complex roots of z .
- This can create confusion when z is real. Indeed, the symbol $\sqrt{4}$, understood as the arithmetic root of 4, is 2; understood as the complex root of 4, it is the set of the two numbers $+2$ and -2 .

Example 12.10: cube root of a complex number

- Let us compute $\sqrt[3]{-1}$: the complex number (-1) has modulus 1 and argument π . The cube roots will therefore be:

$$\sqrt[3]{-1} = \underbrace{\sqrt[3]{1}}_{=1} \left[\cos \left(\frac{\pi}{3} + \frac{2}{3}k\pi \right) + i \sin \left(\frac{\pi}{3} + \frac{2}{3}k\pi \right) \right] \quad \text{with} \quad k = 0, 1, 2$$

that is, the numbers z_k of the form:

$$\cos \vartheta_k + i \sin \vartheta_k \quad \text{with} \quad \vartheta_k = \frac{\pi}{3} + \frac{2}{3}k\pi \quad \text{and} \quad k = 0, 1, 2$$


- Explicitly we have:

$$\begin{cases} z_0 = \cos\left(\frac{\pi}{3}\right) + i \sin\left(\frac{\pi}{3}\right) = \frac{1}{2} + i \frac{\sqrt{3}}{2} = \frac{1}{2}(1 + i\sqrt{3}) \\ z_1 = \cos\left(\frac{\pi}{3} + \frac{2\pi}{3}\right) + i \sin\left(\frac{\pi}{3} + \frac{2\pi}{3}\right) = \cos(\pi) + i \sin(\pi) = -1 \\ z_2 = \cos\left(\frac{\pi}{3} + \frac{4\pi}{3}\right) + i \sin\left(\frac{\pi}{3} + \frac{4\pi}{3}\right) \\ = \cos\left(\frac{5\pi}{3}\right) + i \sin\left(\frac{5\pi}{3}\right) = \frac{1}{2} - i \frac{\sqrt{3}}{2} = \frac{1}{2}(1 - i\sqrt{3}) \end{cases}$$

- The arrangement of the roots of complex numbers in the Gauss plane is not random.
- Indeed, if $w = r(\cos \varphi + i \sin \varphi)$, the n -th roots $z_0, z_1, \dots, z_{n-1}$ of w lie at the vertices of the regular polygon with n sides inscribed in the circle with center 0 and radius $r^{1/n}$, with the vertex z_0 located at the point with argument $\vartheta = \varphi/n$.

Example 12.11: roots in the Gauss plane

- The figure shows the cube roots of -1 : z_0, z_1, z_2 from the previous exercise:




- The figure shows the sixth roots of i : $z_0, z_1, z_2, z_3, z_4, z_5$,

$$\sqrt[6]{i} = \underbrace{\sqrt[6]{1}}_{=1} \left[\cos\left(\frac{\pi}{2 \cdot 6} + \frac{2}{6}k\pi\right) + i \sin\left(\frac{\pi}{2 \cdot 6} + \frac{2}{6}k\pi\right) \right] \quad \text{with } k = 0, 1, 2, 3, 4, 5$$

that is, the numbers z_k of the form:

$$\cos \vartheta_k + i \sin \vartheta_k \quad \text{with} \quad \vartheta_k = \frac{\pi}{12} + \frac{1}{3}k\pi \quad \text{and} \quad k = 0, 1, 2, 3, 4, 5$$




12.3.3 Exponential form of complex numbers

- The following notation is useful:

$$e^{i\varphi} = \cos \varphi + i \sin \varphi \quad \text{with } \varphi \in \mathbb{R}$$

where e is Euler's number (Napier's constant).



With this notation, the *trigonometric form* can be equivalently rewritten as:

$$z = |z| (\cos \varphi + i \sin \varphi) = |z| e^{i\varphi}, \quad \varphi = \arg(z) \quad \text{with } z \neq 0$$

This notation is called the **exponential form** of complex numbers.

- This form is useful since we have:

$$e^{i\varphi_1} \cdot e^{i\varphi_2} = e^{i(\varphi_1+\varphi_2)}$$

that is, it is very easy to compute products of complex numbers.

De Moivre's formulas become:

$$(e^{i\varphi})^n = e^{in\varphi}$$

The exponential notation is widely used in applications because it greatly simplifies manipulations involving trigonometric quantities.

We have:

$$e^{2k\pi i} = 1 \quad \text{and} \quad e^{i(\varphi+2k\pi)} = e^{i\varphi} \quad \forall k \in \mathbb{Z}, \quad \text{moreover} \quad |e^{i\varphi}| = 1$$

We also have:

$$e^{i\pi} = -1, \quad e^{i\frac{\pi}{2}} = i, \quad e^{i\frac{3\pi}{2}} = -i, \quad e^{i\frac{3\pi}{4}} = \frac{-1+i}{\sqrt{2}}$$

Rewriting, we obtain **Euler's formula**:

$$e^{i\pi} + 1 = 0$$

which links in a simple way five of the most important constants: 0, 1, i , π , e .

The exponential notation can be generalized consistently with the properties of powers, by defining the **complex exponential**:

$$e^z = e^{x+iy} = e^x e^{iy} = e^x (\cos y + i \sin y), \quad \forall z = x + iy \in \mathbb{C}$$

- From the definition of exponential form it follows that

$$e^{-i\varphi} = \cos(-\varphi) + i \sin(-\varphi) = \cos \varphi - i \sin \varphi$$

and we easily obtain:

$$\cos \varphi = \frac{e^{i\varphi} + e^{-i\varphi}}{2}, \quad \sin \varphi = \frac{e^{i\varphi} - e^{-i\varphi}}{2i} \quad \text{with } \varphi \in \mathbb{R}$$

These formulas allow us to express sine and cosine as combinations of complex exponentials.

12.4 Quadratic equations

- A **second-degree equation** or **quadratic equation** in one **real** unknown x is an algebraic equation in which the highest power of the unknown is 2, and it can always be written in the form:

$$a x^2 + b x + c = 0 \quad (a \neq 0) \quad (12.22)$$

where a, b and c are real numbers.

The solutions are also called **roots** or **zeros** of the equation.

Observation 12.10: quadratic formula

Given a quadratic equation $a x^2 + b x + c = 0$ ($a \neq 0$), its zeros or roots are:

$$x = \frac{-b \pm \sqrt{b^2 - 4 a c}}{2 a} \quad (12.23)$$

- A quadratic equation is called a “complete quadratic equation” when all its coefficients are different from 0. It is solved with the so-called **method of completing the square**, so called because the equation is modified until its left-hand side becomes the square of a binomial.

Proof. Isolating the constant term, we obtain:

$$\begin{aligned} a x^2 + b x &= -c \\ \underbrace{4 a^2 x^2}_{=(2 a x)^2} + \underbrace{4 a b x}_{=2 (2 a x) b} &= -4 a c \\ (2 a x)^2 + 2 (2 a x) b + b^2 &= b^2 - 4 a c \\ (2 a x + b)^2 &= \underbrace{b^2 - 4 a c}_{:=\Delta \text{ (discriminant)}} \\ 2 a x + b &= \pm \sqrt{b^2 - 4 a c} \\ x &= \frac{-b \pm \sqrt{b^2 - 4 a c}}{2 a} \end{aligned}$$

□

- If the discriminant Δ is negative there are no real solutions.
- If $\Delta = 0$, the quadratic formula becomes:

$$x = -\frac{b}{2a}$$

therefore there is only one root, with *multiplicity two*.

- Finally, if $\Delta < 0$, the equation has no real solutions. In particular, there are always two solutions, but they belong to the field of complex numbers: they are **two complex conjugate numbers** and are computed with the two formulas:

$$x_+ = \frac{-b}{2a} + i \left(\frac{\sqrt{4ac - b^2}}{2a} \right)$$

$$x_- = \frac{-b}{2a} - i \left(\frac{\sqrt{4ac - b^2}}{2a} \right)$$

where i is the **imaginary unit** ($i^2 = -1$).

A quadratic equation in one **complex** unknown z has the following form

$$a z^2 + b z + c = 0 \quad (a \neq 0) \quad (12.24)$$

where a, b and c are complex numbers. It is solved with the same formula as in the case of a real unknown:

$$z = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a} \quad (12.25)$$

where the square root is understood in the complex sense (the $\pm$ sign is superfluous, because in the complex field the root denotes two numbers, each the opposite of the other)

Example 12.12: quadratic equations in the complex field

- We want to solve:

$$z^2 + 2iz - \sqrt{3}i = 0 \quad a = 1, b = 2i, c = -\sqrt{3}i$$

we have (switching to the trigonometric form to compute the root):

$$\begin{aligned} z &= \frac{-2i \pm \sqrt{2^2 i^2 + 2^2 \sqrt{3}i}}{2} = \frac{-2i \pm 2\sqrt{-1 + \sqrt{3}i}}{2} = -i \pm \sqrt{-1 + \sqrt{3}i} \\ &= -i \pm \sqrt{2 \left(-\frac{1}{2} + i \frac{\sqrt{3}}{2} \right)} = -i \pm \sqrt{2 \left(\cos \frac{2}{3}\pi + i \sin \frac{2}{3}\pi \right)} \\ &= -i \pm \sqrt{2} \left(\underbrace{\cos \frac{\pi}{3}}_{=\frac{1}{2}} + i \underbrace{\sin \frac{\pi}{3}}_{=\frac{\sqrt{3}}{2}} \right) = \pm \frac{\sqrt{2}}{2} + i \left(-1 \pm \frac{\sqrt{6}}{2} \right) \end{aligned}$$

- The previous theorem tells us that a polynomial of the form $z^n + a$ (with a complex) has exactly n roots in $\mathbb{C}$; in the real field, instead, the equation $x^n + a = 0$ may have two, one, or no roots (examples: $x^2 - 1 = 0, x^3 - 1 = 0, x^2 + 1 = 0$); we now know that such an equation always has n roots in $\mathbb{C}$, but only occasionally one or two of them lie in $\mathbb{R}$.

- The result is much more general, as stated by the following theorem, whose proof we do not report.


Theorem 12.11: fundamental theorem of algebra

A polynomial equation of the form

$$a_0 + a_1 z + \cdots + a_n z^n = 0 \quad (a_n \neq 0)$$

with arbitrary complex coefficients has exactly n roots in $\mathbb{C}$, if each of them is counted with its multiplicity ^a.

^aIf $P(z)$ is a polynomial in z of degree n and z_0 is one of its roots, we say that z_0 has multiplicity k (k integer, ≥ 1) if the formula $P(z) = (z - z_0)^k Q(z)$ holds, where Q is a polynomial such that $Q(z_0) \neq 0$.

Cosines and sines of the main angles


Exercises

The sheets of worked exercises on the topics of this volume, one per section, with the solution right after each statement. The sheets are by Fabio Furini and Gianluca Priori; many exercises are taken from M. Bramanti, *Esercitazioni di Analisi matematica 1*, Esculapio.

13.1 Maxima, minima, suprema and infima

Exercise 13.1

Determine whether the set

$$A = [0, \sqrt{2}] \cap \mathbb{Q}$$

has a maximum, a minimum, a supremum, an infimum and, if so, determine these elements.

Solution

The minimum of A is 0. A has no maximum; its supremum is $\sqrt{2}$.

Exercise 13.2

Determine whether the set

$$A = [\sqrt{2}, \sqrt{3}] \cap \mathbb{Q}$$

has a maximum, a minimum, a supremum, an infimum and, if so, determine these elements.

Solution

A has neither a minimum nor a maximum. The infimum is $\sqrt{2}$, the supremum is $\sqrt{3}$.

Exercise 13.3

Determine whether the set

$$A = [0, \sqrt{2}] \cap (\mathbb{R} - \mathbb{Q})$$

has a maximum, a minimum, a supremum, an infimum and, if so, determine these elements.

Solution

A has no minimum; the infimum is 0. The maximum of A is $\sqrt{2}$.

Exercise 13.4

In each of the following cases, say whether the set

$$A = (-\infty, \sqrt{2}] \cap \mathbb{Q}$$

has a maximum, a minimum, a supremum, an infimum and, if so, determine these elements.

Solution

A is not bounded below, hence it has no infimum in $\mathbb{R}$. A has no maximum; the supremum is $\sqrt{2}$.

Exercise 13.5

Say whether the following set $A \subset \mathbb{R}$ has a maximum, a minimum, a supremum, an infimum, and whether it is bounded:

$$A = \left\{ \frac{2n+1}{n} : n \in \mathbb{N} \setminus \{0\} \right\}$$

Solution

From $(2n+1)/n = 2 + (1/n)$ we have that the maximum of A is attained for $n = 1$ and is equal to 3. A has no minimum; the infimum is 2. In particular, A is bounded.

Exercise 13.6

Say whether the following set $A \subset \mathbb{R}$ has a maximum, a minimum, a supremum, an infimum, and whether it is bounded:

$$A = \left\{ \frac{n-1}{n} : n \in \mathbb{N} \setminus \{0\} \right\}$$

Solution

From $(n-1)/n = 1 - (1/n)$ we have that the minimum of A is attained for $n = 1$ and is equal to 0. A has no maximum; the supremum is 1. In particular, A is bounded.

Exercise 13.7

Say whether the following set $A \subset \mathbb{R}$ has a maximum, a minimum, a supremum, an infimum, and whether it is bounded:

$$A = \left\{ \frac{3n^2 + 1}{n^2} : n \in \mathbb{N} \setminus \{0\} \right\}$$

Solution

From $(3n^2 + 1)/n^2 = 3 + (1/n^2)$ we have that the maximum of A is attained for $n = 1$ and is equal to 4. A has no minimum; the infimum is 3. In particular, A is bounded.

Exercise 13.8

Say whether the following set $A \subset \mathbb{R}$ has a maximum, a minimum, a supremum, an infimum, and whether it is bounded:

$$A = \left\{ \frac{1}{n^2 + 1} : n \in \mathbb{N} \setminus \{0\} \right\}$$

Solution

The maximum value of A is attained for $n = 1$ and is equal to $1/2$. A has no minimum; the infimum is 0. In particular, A is bounded.

Exercise 13.9

Let

$$A = \left\{ \left| \frac{3x}{x+1} \right| : -\frac{1}{2} < x \leq 2 \right\}.$$

Choose the correct statements among the following:

- (a) $\sup A \notin A$ (b) $\inf A = \min A = 2$
- (c) $\max A = 3$ (d) none of the other answers is correct

Solution


A is the set of values of $|f(x)|$ on the interval $(-1/2, 2]$, with $f(x) = \frac{3x}{x+1}$. From

$$\frac{3x}{x+1} = 3 - \frac{3}{x+1}$$

we have that the function f is strictly increasing on the given interval $(-1/2, 2]$, negative on $(-1/2, 0)$, zero for $x = 0$, and positive on $(0, 2]$. It follows that

$$|f(x)| = \left| \frac{3x}{x+1} \right|$$

is strictly decreasing on $(-1/2, 0]$, where it takes all the values in $[0, 3)$, and strictly increasing on $[0, 2]$, where it takes all the values in $[0, 2]$. The minimum value is therefore attained for $x = 0$ and is equal to 0. The supremum is 3; there is no maximum value. The correct answer is (a).



Exercise 13.10

For $I = [1/3, +\infty)$ consider the function

$$f : I \rightarrow \mathbb{R}, f(x) = \exp\left(\left|\frac{2x-1}{x}\right|\right).$$

Determine, among the following intervals, the set $J = f(I)$ of the values taken by f .

- (a) $[e, e^2]$ (b) $(e^{-2}, e]$ (c) $[1, e^2]$
 (d) $(e, +\infty)$ (e) $(0, e^2)$ (f) another interval



Solution

The function f has the same monotonicity behavior as

$$|g(x)| = \left|\frac{2x-1}{x}\right|$$

with

$$g(x) = \frac{2x-1}{x} = 2 - \frac{1}{x}.$$




Solution

The function g is strictly increasing on the interval $I = [1/3, +\infty)$, negative on $[1/3, 1/2)$, zero for $x = 1/2$, and positive on $(1/2, +\infty)$. It follows that $|g(x)|$ is strictly decreasing on $[1/3, 1/2]$, where it takes all the values in $[0, 1]$, and strictly increasing on $[1/2, +\infty)$, where it takes all the values in $[0, 2)$. The set of values $|g|(I)$ is $[0, 2)$, hence for $f(x) = e^{|g(x)|}$ we have

$$f(I) = [1, e^2).$$

The correct answer is (c).



More precisely, f is strictly decreasing on $[1/3, 1/2]$, where it takes all the values in $[1, e]$, and strictly increasing on $[1/2, +\infty)$, where it takes all the values in $[1, e^2)$.

Exercise 13.11

Let $A = A_+ \cup A_-$ with

$$A_+ = \left\{ x + \frac{2}{x} : x > 0 \right\}, \quad A_- = \left\{ x + \frac{2}{x} : x < 0 \right\}.$$

Choose the correct statements among the following:

- (a) $\sup A = +\infty$ (b) $\inf A_+ = 2\sqrt{2}$ (c) A_+ has no minimum
 (d) A_- has no maximum (e) none of the other answers is correct

Solution

Clearly, A_+ is not bounded above, hence statement (a) is correct (and statement (e) is

false). For $x \neq 0$, we consider the equation


$$x + \frac{2}{x} = y,$$

$$x^2 - yx + 2 = 0.$$

There are solutions $x \in \mathbb{R}$ if and only if $y \in (-\infty, -2\sqrt{2}] \cup [2\sqrt{2}, +\infty)$. Moreover, for every $y \in (-\infty, -2\sqrt{2}]$ the solutions x are negative, while for every $y \in [2\sqrt{2}, +\infty)$ the solutions x are positive. This means

$$A_+ = [2\sqrt{2}, +\infty), \quad A_- = (-\infty, -2\sqrt{2}].$$

Therefore (b) is true, (c) is false, (d) is false.



13.2 Equations and inequalities in one real variable

Exercise 13.12

Solve the following inequality in the real variable x :

$$\frac{x^2 - 2x}{x^2 - 4x + 3} > 0$$

Solution

The numerator $x^2 - 2x$ is positive for $x < 0$ or for $x > 2$, zero for $x = 0$, $x = 2$, and negative for $0 < x < 2$. The denominator $x^2 - 4x + 3$ is positive for $x < 1$ or for $x > 3$, and negative

for $1 < x < 3$. The quotient is defined and has the required positive sign for

$$x \in (-\infty, 0) \cup (1, 2) \cup (3, +\infty).$$

Exercise 13.13

Solve the following inequality in the real variable x :

$$\log(x-1)^2 - \log(x-2) > 0$$

Solution

The function $f(x) = \log(x-1)^2 - \log(x-2)$ is defined for $x > 2$, since for such x , and only for such x , we have $x-2 > 0$, $(x-1)^2 > 0$. In its domain, we have, as required, $f(x) > 0$ if and only if

$$\log(x-1)^2 > \log(x-2), \quad x > 2$$

hence if and only if

$$\begin{cases} (x-1)^2 > x-2 \\ x > 2 \end{cases}$$

We obtain

$$\begin{cases} x^2 - 3x + 3 > 0 \\ x > 2 \end{cases}$$

which is equivalent to

$$x > 2$$

since $x^2 - 3x + 3 > 0$ is satisfied for every x . On its whole domain $(2, +\infty)$ the function $f(x)$ takes positive values.

Exercise 13.14

Solve the following inequality in the real variable x :

$$e^x + e^{-x} < \frac{10}{3}$$

Solution

For every $x \in \mathbb{R}$

$$e^x + e^{-x} < \frac{10}{3}$$

is equivalent to

$$3e^{2x} - 10e^x + 3 < 0$$

hence to

$$\frac{1}{3} < e^x < 3.$$

It follows that the inequality is solved by

$$-\log 3 < x < \log 3.$$

Exercise 13.15

Solve the following inequality in the real variable x :

$$8^{x+1} \geq 2^{x^2}$$

Solution

For every $x \in \mathbb{R}$

$$8^{x+1} \geq 2^{x^2}$$

is equivalent to

$$2^{3x+3} \geq 2^{x^2}$$

hence to

$$3x + 3 \geq x^2$$

since 2^x is a strictly increasing function on $\mathbb{R}$. It follows that the inequality is solved by

$$\frac{3 - \sqrt{21}}{2} \leq x \leq \frac{3 + \sqrt{21}}{2}.$$

Exercise 13.16

Solve the following inequality in the real variable x :

$$\log(1 - \sin x) \geq 0$$

Solution

The function $f(x) = \log(1 - \sin x)$ is defined for $\sin x < 1$, hence for

$$x \neq \frac{\pi}{2} + 2k\pi, \quad k \in \mathbf{Z}.$$

For such x we have $f(x) \geq 0$ if and only if

$$1 - \sin x \geq 1$$

hence for

$$\sin x \leq 0.$$

The given inequality is solved for

$$\pi + 2k\pi \leq x \leq 2\pi + 2k\pi, \quad k \in \mathbf{Z}$$

i.e., for

$$x \in \bigcup_{k \in \mathbf{Z}} [\pi + 2k\pi, 2\pi + 2k\pi].$$

Exercise 13.17

Solve the following inequality in the real variable x :

$$|x|\sqrt{1-2x^2} > 2x^2 - 1$$

Solution

$\sqrt{1-2x^2}$ is defined for $1-2x^2 \geq 0$. When $1-2x^2 = 0$ the inequality is not satisfied, since it reduces to $0 > 0$. When $1-2x^2 > 0$ the inequality is satisfied, since the left-hand side $|x|\sqrt{1-2x^2}$ is the product of two positive numbers while the right-hand side $2x^2 - 1 = -(1-2x^2)$ is negative. Therefore the inequality is solved for

$$-\sqrt{\frac{1}{2}} < x < \sqrt{\frac{1}{2}}.$$

Exercise 13.18

Solve the following equation in the real variable x :

$$\sin^2 x = 2 \cos^2 x - \frac{1}{2}$$

Solution

The equation

$$\sin^2 x = 2 \cos^2 x - \frac{1}{2}$$

is equivalent to

$$2 \sin^2 x = 4(1 - \sin^2 x) - 1$$

$$6 \sin^2 x = 3$$

$$\sin x = \pm \sqrt{\frac{1}{2}}.$$

The solutions are given by

$$x = \frac{\pi}{4} + \frac{k\pi}{2}, \quad k \in \mathbf{Z}.$$

Exercise 13.19

Solve the following inequality in the real variable x :

$$e^{\sin^2 x - \sin x} \leq 1$$

Solution

For every $x \in \mathbb{R}$, $e^{\sin^2 x - \sin x} \leq 1$ is equivalent to

$$\sin^2 x - \sin x \leq 0$$

hence to

$$0 \leq \sin x \leq 1.$$

The solutions are given by

$$2k\pi \leq x \leq \pi + 2k\pi, \quad k \in \mathbf{Z}$$

i.e., by

$$x \in \bigcup_{k \in \mathbf{Z}} [2k\pi, \pi + 2k\pi].$$

Exercise 13.20

Solve the following inequality in the real variable x :

$$2|x^2 - x| > |x|$$

Solution

The inequality $2|x^2 - x| > |x|$ is not satisfied for $x = 0$. For $x \neq 0$, dividing by the positive term $|x|$, it is equivalent to

$$2|x - 1| > 1$$

hence to

$$|x - 1| > 1/2.$$

The solutions are given by

$$(x - 1 < -1/2, x \neq 0) \vee (x - 1 > 1/2),$$

$$(x < 1/2, x \neq 0) \vee x > 3/2$$

hence by

$$x \in (-\infty, 0) \cup (0, 1/2) \cup (3/2, +\infty).$$

Exercise 13.21

Solve the following inequality in the real variable x :

$$(x + 1)^{x^2 - 1} > 1$$

Solution

The function $f(x) = (x + 1)^{x^2 - 1}$ is defined for $x > -1$. For such x , $f(x) > 1$ is equivalent to

$$e^{(x^2 - 1) \log(x + 1)} > 1$$

hence to

$$(x^2 - 1) \log(x + 1) > 0, \quad x > -1.$$

Dividing by the positive term $x + 1$ we obtain

$$(x - 1) \log(x + 1) > 0, \quad x > -1.$$

The factor $x - 1$ is positive for $x > 1$, zero for $x = 1$, and negative for $-1 < x < 1$. The factor $\log(x + 1)$ is positive for $x > 0$, zero for $x = 0$, and negative for $-1 < x < 0$. By the rule of signs, the solutions are given by

$$x \in (-1, 0) \cup (1, +\infty).$$

Exercise 13.22

Solve the following equation in the real variable x :

$$3^{|x^2 - 4|} = 0$$

Solution

$3^{|x^2-4|} = 0$ has no solutions, since $3^y > 0$ for every $y \in \mathbb{R}$.

Exercise 13.23

Solve the following inequality in the real variable x :

$$\frac{\log_a(4x-3)}{\log_a(2x-1)} > 1 \quad 0 < a < 1$$

Solution

Because of the existence conditions of the logarithm and of the denominator, we must impose the system

$$\begin{cases} 4x-3 > 0 \\ 2x-1 > 0 \\ \log_a(2x-1) \neq 0 \end{cases}$$

which gives $\frac{3}{4} < x < 1, x > 1$. Now moving everything to the left-hand side, finding the least common denominator and using the properties of logarithms, we obtain

$$\frac{\log_a\left(\frac{4x-3}{2x-1}\right)}{\log_a(2x-1)} > 0$$

Studying the sign of the numerator and of the denominator, and keeping in mind that $a < 1$, for the numerator we obtain

$$\log_a\left(\frac{4x-3}{2x-1}\right) > 0 \iff \frac{4x-3}{2x-1} < 1 \iff \frac{1}{2} < x < 1$$

while for the denominator we obtain

$$\log_a(2x-1) > 0 \iff x < 1$$

Combining the signs, the inequality is satisfied for $\frac{1}{2} < x < 1, x > 1$. Intersecting this set of solutions with the existence conditions studied above, we obtain that the inequality is satisfied for every x belonging to the set $(\frac{3}{4}; 1) \cup (1, +\infty)$.

Exercise 13.24

Solve the following inequality in the real variable x :

$$\sqrt{x^2 - 3x + 5} \leq x + 3$$

Solution

This irrational inequality has the form $\sqrt{A(x)} \leq B(x)$, with $A(x)$ and $B(x)$ polynomials in x . First of all, we must impose the existence condition of the root, i.e., $x^2 - 3x + 5 \geq 0$. Moreover, since a square root is always positive or zero in $\mathbb{R}$, we impose $x + 3 \geq 0$. We therefore solve the system:

$$\begin{cases} x^2 - 3x + 5 \geq 0 \\ x + 3 \geq 0 \\ x^2 - 3x + 5 \leq (x + 3)^2 \end{cases}$$

where, in the last inequality, we were allowed to square both sides, since they are positive. The system reduces to:

$$\begin{cases} \forall x \in \mathbb{R} \\ x \geq -3 \\ x \geq -\frac{4}{9} \end{cases}$$

whose set of solutions is the interval $[-\frac{4}{9}, +\infty)$.

Exercise 13.25

Solve the following rational inequality in the real variable x :

$$\frac{x^3 - 3x^2 + 2x - 6}{x^2 - 2x - 3} > \frac{2}{x + 1}$$

Solution

First of all, we must write the inequality in canonical form (0 on the right-hand side). The existence conditions (denominators different from zero) will be automatically included in the sign study. We obtain

$$\frac{x^3 - 3x^2 + 2x - 6}{x^2 - 2x - 3} - \frac{2}{x + 1} > 0$$

We factor the numerator and the denominator of the first fraction, obtaining:

$$\frac{(x^2 + 2)(x - 3)}{(x - 3)(x + 1)} - \frac{2}{x + 1} > 0$$

Since we can cancel $x - 3$ and obtain an equivalent fraction, provided that $x \neq 3$ by the existence condition, we have

$$\frac{x^2 + 2}{x + 1} - \frac{2}{x + 1} > 0$$

that is,

$$\frac{x^2}{x + 1} > 0$$

The numerator is positive $\forall x \in \mathbb{R}, x \neq 0$, and the denominator is positive for $x > -1$. Combining the signs, and taking into account the condition $x \neq 3$, we obtain the solution set $(-1, 0) \cup (0, 3) \cup (3, +\infty)$.

13.3 Summations

Exercise 13.26

Prove, using the properties of summations, that for every integer $n \geq 1$ we have:

$$\sum_{k=1}^n k^2 = \frac{n(n+1)(2n+1)}{6}$$

(sum of the squares of the first n natural numbers, excluding zero).

Solution

We start by writing the sum $\sum_{k=0}^n (k+1)^3$ in two different ways:

$$\begin{aligned} 1) \quad \sum_{k=0}^n (k+1)^3 &= \sum_{k=0}^n (k^3 + 3k^2 + 3k + 1) \\ &= \left(\sum_{k=1}^n k^3 + 3 \sum_{k=1}^n k^2 + 3 \sum_{k=1}^n k + \sum_{k=1}^n 1 \right) + 1 \\ &= \sum_{k=1}^n k^3 + 3 \sum_{k=1}^n k^2 + \frac{3n(n+1)}{2} + n + 1 \end{aligned}$$

$$2) \quad \sum_{k=0}^n (k+1)^3 = \sum_{k=1}^{n+1} k^3 = \sum_{k=1}^n k^3 + (n+1)^3$$

Equating the two expressions and canceling the sum with k^3 , we obtain:

$$3 \sum_{k=1}^n k^2 + \frac{3n(n+1)}{2} + n + 1 = (n+1)^3$$

Isolating the sum with k^2 , we obtain:

$$\begin{aligned} 3 \sum_{k=1}^n k^2 &= (n+1)^3 - \frac{3n(n+1)}{2} - n - 1 \\ &= n^3 + 3n^2 + 3n + 1 - \frac{3n^2 + 3n}{2} - n - 1 \\ &= \frac{2n^3 + 3n^2 + n}{2} \end{aligned}$$

Hence:

$$\sum_{k=1}^n k^2 = \frac{2n^3 + 3n^2 + n}{6} = \frac{n(n+1)(2n+1)}{6}$$

Exercise 13.27

Prove, using the properties of summations, that for every integer $n \geq 1$ we have:

$$\sum_{k=1}^n k^3 = \frac{n^2(n+1)^2}{4}$$

(sum of the cubes of the first n natural numbers, excluding zero). That is, we have:

$$\sum_{k=1}^n k^3 = \frac{n^2(n+1)^2}{4} = \left(\frac{n(n+1)}{2}\right)^2 = \left(\sum_{k=1}^n k\right)^2$$

and hence the sum of the cubes is equal to the square of the sum of the natural numbers.

Solution

We start by writing the following sum in two different ways: $\sum_{k=0}^n (k+1)^4$.

$$\begin{aligned} 1) \quad \sum_{k=0}^n (k+1)^4 &= \sum_{k=0}^n (k^4 + 4k^3 + 6k^2 + 4k + 1) \\ &= \left(\sum_{k=1}^n k^4 + 4 \sum_{k=1}^n k^3 + 6 \sum_{k=1}^n k^2 + 4 \sum_{k=1}^n k + \sum_{k=1}^n 1 \right) + 1 \\ &= \sum_{k=1}^n k^4 + 4 \sum_{k=1}^n k^3 + n(n+1)(2n+1) + 2n(n+1) + n + 1 \\ &= \sum_{k=1}^n k^4 + 4 \sum_{k=1}^n k^3 + 2n^3 + 5n^2 + 4n + 1 \\ 2) \quad \sum_{k=0}^n (k+1)^4 &= \sum_{k=1}^{n+1} k^4 = \sum_{k=1}^n k^4 + (n+1)^4 \end{aligned}$$

Equating the two expressions and canceling the sum with k^4 , we obtain:

$$4 \sum_{k=1}^n k^3 + 2n^3 + 5n^2 + 4n + 1 = (n+1)^4$$

Isolating the sum with k^3 , we obtain:

$$4 \sum_{k=1}^n k^3 = n^4 + 4n^3 + 6n^2 + 4n + 1 - 2n^3 - 5n^2 - 4n - 1 = n^4 + 2n^3 + n^2$$

Hence:

$$\sum_{k=1}^n k^3 = \frac{n^4 + 2n^3 + n^2}{4} = \frac{n^2 (n+1)^2}{4}$$

13.4 Principle of mathematical induction

Exercise 13.28

Prove by induction that for every integer $n \geq 1$ the following holds:

$$\sum_{k=1}^n k^2 = \frac{n(n+1)(2n+1)}{6}$$

(sum of the squares of the first n natural numbers)

Solution

By induction on n .

- **Base case**

Let $n = 1$. Then the statement becomes:

$$\sum_{k=1}^1 k^2 = \frac{1(1+1)(2 \cdot 1 + 1)}{6} \quad \text{i.e.} \quad 1 = 1$$

which is clearly true.

- **Inductive step**

Assume that it is true for n , and let us prove it for $(n+1)$. By the inductive hypothesis, we have:

$$\sum_{k=1}^n k^2 = \frac{n(n+1)(2n+1)}{6}.$$

Hence we can write:

$$\begin{aligned}
 \sum_{k=1}^{n+1} k^2 &= \sum_{k=1}^n k^2 + (n+1)^2 \\
 &= \frac{n(n+1)(2n+1)}{6} + (n+1)^2 = \frac{n(n+1)(2n+1) + 6(n+1)^2}{6} \\
 &= \frac{(n+1)(n(2n+1) + 6(n+1))}{6} = \frac{(n+1)(2n^2 + n + 6n + 6)}{6} \\
 &= \frac{(n+1)(2n^2 + 4n + 6)}{6} = \frac{(n+1)((2n)(n+2) + 3(n+2))}{6} \\
 &= \frac{(n+1)(n+2)(2n+3)}{6} = \frac{(n+1)(n+1+1)(2(n+1)+1)}{6}
 \end{aligned}$$

which is exactly the desired statement for $n+1$.

Exercise 13.29

Prove by induction that for every integer $n \geq 1$ the following holds:

$$\sum_{k=0}^{n-1} (2k+1) = n^2 \quad \text{or} \quad \sum_{k=1}^n (2k-1) = n^2$$

(sum of the first n odd numbers)

Solution

By induction on n .

- **Base case**

Let $n = 1$. Then the statement becomes:

$$\sum_{k=0}^0 (2k+1) = 1^2 \quad \text{i.e.} \quad 1 = 1$$

which is clearly true.

- **Inductive step**

Assume that it is true for n , and let us prove it for $(n+1)$. By the inductive hypothesis, we have:

$$\sum_{k=0}^{n-1} (2k+1) = n^2$$

Hence we can write:

$$\begin{aligned}\sum_{k=0}^{n-1+1} (2k+1) &= \sum_{k=0}^{n-1} (2k+1) + 2n+1 \\ &= n^2 + 2n + 1 \\ &= (n+1)^2\end{aligned}$$

which is exactly the desired statement for $n+1$.

Exercise 13.30

Prove by induction that for every integer $n \geq 1$ the following holds:

$$\sum_{k=1}^n 2k = n(n+1)$$

(sum of the first n even numbers)

Solution

By induction on n .

- **Base case**

Let $n = 1$. Then the statement becomes:

$$\sum_{k=1}^1 2k = 1 \cdot 2 \quad \text{i.e.} \quad 2 = 2$$

which is clearly true.

- **Inductive step**

Assume that it is true for n , and let us prove it for $(n+1)$. By the inductive hypothesis, we have:

$$\sum_{k=1}^n 2k = n(n+1)$$

Hence we can write:

$$\begin{aligned}
 \sum_{k=1}^{n+1} 2k &= \sum_{k=1}^n 2k + 2(n+1) \\
 &= 2 \left(\sum_{k=1}^n k + (n+1) \right) \\
 &= 2 \left(\frac{n(n+1)}{2} + (n+1) \right) \\
 &= 2 \left(\frac{n(n+1) + 2(n+1)}{2} \right) \\
 &= (n+1)(n+2) = (n+1)(n+1+1)
 \end{aligned}$$

which is exactly the desired statement for $n+1$.

Exercise 13.31

Prove by induction that for every integer $n \geq 1$ the following holds:

$$\sum_{k=1}^n \frac{k}{2^k} = 2 - \frac{n+2}{2^n}$$

Solution

By induction on n .

- **Base case**

Let $n = 1$. Then the statement becomes:

$$\sum_{k=1}^1 \frac{k}{2^k} = 2 - \frac{1+2}{2^1} \quad \text{i.e.} \quad \frac{1}{2} = \frac{1}{2}$$

which is clearly true.

- **Inductive step**

Assume that it is true for n , and let us prove it for $(n+1)$. By the inductive hypothesis, we have:

$$\sum_{k=1}^n \frac{k}{2^k} = 2 - \frac{n+2}{2^n}$$

Hence we can write:

$$\begin{aligned}
 \sum_{k=1}^{n+1} \frac{k}{2^k} &= \sum_{k=1}^n \frac{k}{2^k} + \frac{n+1}{2^{n+1}} \\
 &= 2 - \frac{n+2}{2^n} + \frac{n+1}{2^{n+1}} \\
 &= 2 - \frac{2(n+2)}{2^{n+1}} + \frac{n+1}{2^{n+1}} \\
 &= 2 - \left(\frac{2(n+2) - n - 1}{2^{n+1}} \right) \\
 &= 2 - \left(\frac{n+3}{2^{n+1}} \right) \\
 &= 2 - \frac{(n+1)+2}{2^{n+1}}
 \end{aligned}$$

which is exactly the desired statement for $n+1$.

Exercise 13.32

Given a set X with n elements, prove by induction that the power set $\mathcal{P}(X)$ has 2^n elements, that is,

$$|\mathcal{P}(X)| = 2^n \quad \text{for } n \geq 0$$

Solution

By induction on n .

- **Base case**

Let $n = 0$. A set with 0 elements is the empty set $(\emptyset)$. The power set of the empty set contains only the empty set as an element. Then the statement becomes:

$$|\mathcal{P}(\emptyset)| = 2^0 \quad \text{i.e.} \quad 1 = 1$$

which is clearly true.

- **Inductive step**

Assume the statement is true for a set of n elements, and let us prove it for a set of $(n+1)$ elements. By the inductive hypothesis, the power set of a set of n elements has 2^n elements.

A set Y of $(n+1)$ elements can be written as:

$$Y = \{a_1, a_2, \dots, a_n, a_{n+1}\}$$

hence:

$$\mathcal{P}(Y) = \{S : S \subseteq Y\} = \{S : S \subseteq \{a_1, a_2, \dots, a_n, a_{n+1}\}\}$$

The subsets of Y either contain the element a_{n+1} or do not contain it. We therefore define the set Z_a formed by the first n elements of Y :

$$Z_a = \{a_1, a_2, \dots, a_n\} \quad \text{hence} \quad |\mathcal{P}(Z_a)| = 2^n$$

We now define the following set Z_b :

$$Z_b = \{S \cup \{a_{n+1}\} : S \subseteq \underbrace{\{a_1, a_2, \dots, a_n\}}_{=Z_a}\} = \{S \cup \{a_{n+1}\} : S \subseteq Z_a\} \quad \text{and} \quad |Z_b| = 2^n$$

since Z_b has one element for each element of $\mathcal{P}(Z_a)$.

In other words, $\mathcal{P}(Z_a)$ contains all the subsets of Y in which the element a_{n+1} is not present, while Z_b contains all the subsets of Y in which the element a_{n+1} is present.

We also note that the sets $\mathcal{P}(Z_a)$ and Z_b are disjoint, that is: $\mathcal{P}(Z_a) \cap Z_b = \emptyset$. Hence $|\mathcal{P}(Z_a) \cup Z_b| = |\mathcal{P}(Z_a)| + |Z_b|$. Finally, we have:

$$|\mathcal{P}(Y)| = |\mathcal{P}(Z_a) \cup Z_b| = 2^n + 2^n = 2^{n+1}$$

which is exactly the desired statement for $n + 1$.

13.5 Complex numbers

Exercise 13.33

Write the following complex numbers z in algebraic form, identifying their real part and imaginary part. Then, for each of them, find the modulus, the conjugate and the reciprocal.

$$z = \frac{2 + 5i}{1 + i} \qquad z = \frac{2 + i}{1 - 3i}$$

Solution

$$z = \frac{2 + 5i}{1 + i} \cdot \frac{1 - i}{1 - i} = \frac{7 + 3i}{2} = \frac{7}{2} + i\frac{3}{2}$$

from which

$$\Re z = \frac{7}{2}, \quad \Im z = \frac{3}{2}, \quad |z| = \sqrt{\frac{49}{4} + \frac{9}{4}} = \frac{\sqrt{58}}{2}, \quad \bar{z} = \frac{7}{2} - i\frac{3}{2}.$$

Finally

$$z^{-1} = \frac{1}{z} \cdot \frac{\bar{z}}{\bar{z}} = \frac{\bar{z}}{|z|^2} = \frac{7 - 3i}{2} \cdot \frac{4}{58} = \frac{7}{29} - i\frac{3}{29}.$$

Solution

$$z = \frac{2+i}{1-3i} \cdot \frac{1+3i}{1+3i} = \frac{-1+7i}{10} = -\frac{1}{10} + i\frac{7}{10}$$

from which

$$\Re z = -\frac{1}{10}, \Im z = \frac{7}{10}, |z| = \sqrt{\frac{1}{100} + \frac{49}{100}} = \frac{\sqrt{2}}{2}, \bar{z} = -\frac{1}{10} - i\frac{7}{10}.$$

Finally

$$z^{-1} = \frac{1}{z} \cdot \frac{\bar{z}}{\bar{z}} = \frac{\bar{z}}{|z|^2} = -\frac{1+7i}{10} \cdot 2 = -\frac{1}{5} - i\frac{7}{5}.$$

Exercise 13.34

Determine all the complex numbers z that satisfy:

$$|z| \leq |z - 2i|$$

Solution

It is convenient to use the form $z = x + iy$, $x, y \in \mathbb{R}$.

$$|x + iy| \leq |x + i(y - 2)|$$

$$|x + iy|^2 \leq |x + i(y - 2)|^2$$

$$x^2 + y^2 \leq x^2 + (y - 2)^2$$

$$0 \leq -4y + 4$$

$$y \leq 1.$$

Representing the solutions in the Cartesian plane, we obtain a closed half-plane.

Exercise 13.35

Determine all the complex numbers z that satisfy:

$$|z - (3 + i)| \leq 2$$

Solution

This inequality is satisfied by those complex numbers $z = x + iy$ which, in the Gauss plane (complex plane), are represented by the points $P(x, y)$ whose distance from the point $(3, 1)$ is less than or equal to 2.

That is, all and only the points of the **closed disk** with center $C(3, 1)$ and radius 2 (careful: it is not the circle, but the disk, i.e., the interior together with the boundary). The circle has equation:

$$\gamma : (x - 3)^2 + (y - 1)^2 = 4$$

while the closed disk has equation:

$$(x - 3)^2 + (y - 1)^2 \leq 4$$

Alternatively, the inequality can be handled through the substitution $z = x + iy$, obtaining

$$|x + iy - 3 - i| \leq 2$$

in which, factoring out the imaginary unit,

$$|x - 3 + i(y - 1)| \leq 2$$

Applying the definition of the modulus of a complex number, we have

$$\sqrt{(x - 3)^2 + (y - 1)^2} \leq 2$$

Since we can square both non-negative sides, we finally obtain:

$$(x - 3)^2 + (y - 1)^2 \leq 4$$

Exercise 13.36

Determine all the complex numbers z that satisfy:

$$(1 - i)z - (1 + i)\bar{z} = i$$

Solution

$$(1 - i)(x + iy) - (1 + i)(x - iy) = i$$

$$x + y - ix + iy - x - y - ix + iy = i$$

$$i(2y - 2x) = i$$

$$2y - 2x = 1.$$

Representing the solutions in the Cartesian plane, we obtain a line.

Exercise 13.37

Determine all the complex numbers z that satisfy:

$$|z + \bar{z}| + |z - \bar{z}| \leq 2$$

Solution

$$|z + \bar{z}| + |z - \bar{z}| \leq 2$$

$$2|\Re z| + 2|\Im z| \leq 2$$

$$|x| + |y| \leq 1.$$

Representing the solutions in the Cartesian plane, we obtain a closed square with vertices $\pm 1, \pm i$.

Exercise 13.38

Determine all the complex numbers z that satisfy:

$$\Im(z) - |z + \bar{z}|^2 < 1$$

Solution

$$\Im(z) - |z + \bar{z}|^2 < 1$$

$$\Im(z) - 4|\Re z|^2 < 1$$

$$y < 4x^2 + 1.$$

Representing the solutions in the Cartesian plane, we obtain the open region below the parabola $y = 4x^2 + 1$.

Exercise 13.39

Determine all the complex numbers z that satisfy:

$$\begin{cases} z\bar{z} \leq 2 \\ z + \bar{z} \leq 2 \end{cases}$$

Solution

$$\begin{cases} |z|^2 \leq 2 \\ 2\Re z \leq 2 \end{cases} \quad \begin{cases} x^2 + y^2 \leq 2 \\ x \leq 1 \end{cases}$$

Representing the solutions in the Cartesian plane, we obtain the part of the plane common to the closed disk with center at the origin and radius $\sqrt{2}$ and to the closed half-plane to the left of the vertical line $x = 1$.

Exercise 13.40

The system in $\mathbb{C}$

$$\begin{cases} z^2 + \bar{z}^2 = 0 \\ |\Re(z)| + |\Im(z)| = 1 \end{cases}$$

has:

- (a) two solutions (b) three solutions
(c) four solutions (d) none of the other answers is correct

Solution

Setting $z = x + iy$, $x, y \in \mathbb{R}$, we have

$$\begin{cases} (x + iy)^2 + (x - iy)^2 = 0 \\ |x| + |y| = 1 \end{cases}$$

$$\begin{cases} x^2 - y^2 + 2ixy + x^2 - y^2 - 2ixy = 0 \\ |x| + |y| = 1 \end{cases}$$

$$\begin{cases} x^2 = y^2 \\ |x| + |y| = 1 \end{cases} \quad \begin{cases} |x| = |y| \\ |x| + |y| = 1 \end{cases} \quad \begin{cases} |x| = 1/2 \\ |y| = 1/2 \end{cases}$$

We obtain the four points $(1 + i)/2$, $(-1 + i)/2$, $(1 - i)/2$, $(-1 - i)/2$. The correct answer is (c).

Exercise 13.41

The solutions $z = x + iy$ of the complex equation

$$z^2 - z\bar{z} + iz = -3 + 2i$$

satisfy

$$\begin{array}{ll} \text{(a)} \quad \left(x + \frac{1}{6}\right)^2 + \left(y + \frac{1}{4}\right)^2 = 1 & \text{(b)} \quad y = \frac{2}{3}x + 1 \\ \text{(c)} \quad \left(x - \frac{1}{6}\right)^2 + \left(y - \frac{1}{4}\right)^2 = 1 & \text{(d)} \quad y = \frac{3}{2}x \end{array}$$

Solution

$$(x + iy)^2 - (x^2 + y^2) + ix - y = -3 + 2i$$

$$-2y^2 - y + i(x + 2xy) = -3 + 2i$$

$$\begin{cases} 2y^2 + y - 3 = 0 \\ x + 2xy = 2 \end{cases} \quad \begin{cases} y = 1 \vee y = -3/2 \\ x = 2/(1 + 2y) \end{cases}$$

We obtain the two points $(2/3) + i$, $-1 - i(3/2)$, which lie on the line $y = (3/2)x$. The correct answer is (d).

Exercise 13.42

Find the modulus and the principal argument of

$$\text{(a)} \quad z = -1 - i\sqrt{3} \quad , \quad \text{(b)} \quad z = -4i.$$

Solution

(a) We have $|z| = \sqrt{1+3} = 2$. The principal argument $\text{Arg } z$ is the unique $\vartheta \in [-\pi, \pi)$ such that

$$\cos \vartheta = \Re z / |z| = -1/2, \quad \sin \vartheta = \Im z / |z| = -\sqrt{3}/2$$

hence $\text{Arg } z = -2\pi/3$.

Solution

(b) We have $|z| = 4$. The principal argument $\text{Arg } z$ is the unique $\vartheta \in [-\pi, \pi)$ such that

$$\cos \vartheta = \Re z / |z| = 0, \quad \sin \vartheta = \Im z / |z| = -1$$

hence $\text{Arg } z = -\pi/2$.

Exercise 13.43

Let

$$w = \frac{z + 1 - i}{z + i}$$

Determine, locating them in the Gauss plane, the complex numbers z for which w is a purely imaginary number.

Solution

The complex number w is defined for $z \neq -i$. Moreover,

$$w = \frac{z + 1 - i}{z + i} \cdot \frac{z - i}{z - i} = \frac{z^2 + z - 1 + i(-2z - 1)}{z^2 + 1}$$

That is, separating the real part of w from its imaginary part:

$$w = \frac{z^2 + z - 1}{z^2 + 1} + i \frac{-2z - 1}{z^2 + 1}$$

The complex number w is purely imaginary when $\Re(w) = 0$, i.e.,

$$\Re(w) = \frac{z^2 + z - 1}{z^2 + 1} = 0$$

This rational equation, under the condition $z \neq \pm\sqrt{-1}$, i.e., $z \neq \pm i$, gives the real numbers

$$z_1 = -\frac{1}{2} + \frac{\sqrt{5}}{2}, \quad z_2 = -\frac{1}{2} - \frac{\sqrt{5}}{2}$$

which, in the Gauss plane, can be represented as two points on the real axis. In conclusion, w is purely imaginary for the two values of z found.

Exercise 13.44

Let $z = \sqrt{3} - 3i$. The number of solutions w with $\Im(w) < 0$ of the equation

$$w^6 = |z|(z + \bar{z})$$

is:

- (a) two (b) three
(c) zero (d) none of the other answers is correct

Solution

We have $|z| = \sqrt{12}$, $z + \bar{z} = 2\sqrt{3}$, hence

$$w^6 = 12.$$

The complex sixth roots of a positive real number are six, pairwise opposite, and two of them are real. Of the remaining four, two have positive imaginary part and two have negative imaginary part. The correct answer is (a). We nevertheless want to determine their values. Setting $w = r(\cos \vartheta + i \sin \vartheta)$ and taking into account that $|12| = 12$, $\arg 12 = 2k\pi$, $k \in \mathbb{Z}$, we obtain

$$\begin{cases} r^6 = 12 \\ 6\vartheta = 2k\pi \end{cases} \quad \begin{cases} r = 12^{1/6} \\ \vartheta = k\pi/3 \end{cases}$$

We therefore have the six roots

$$w_k = 12^{1/6}(\cos k\pi/3 + i \sin k\pi/3), \quad k = 0, 1, 2, 3, 4, 5,$$

i.e.,

$$w_0 = 12^{1/6}, \quad w_1 = 12^{1/6}(1/2 + i\sqrt{3}/2), \quad w_2 = 12^{1/6}(-1/2 + i\sqrt{3}/2)$$

$$w_3 = -12^{1/6}, \quad w_4 = 12^{1/6}(-1/2 - i\sqrt{3}/2), \quad w_5 = 12^{1/6}(1/2 - i\sqrt{3}/2).$$

The two with negative imaginary part are w_4 and w_5 .

Exercise 13.45

Solve the following system in the unknown $z \in \mathbb{C}$.

$$\begin{cases} z^6 = i \\ \Re(z)\Im(z) < 0 \end{cases}$$

Solution

We first solve the equation $z^6 = i$. Setting $z = r(\cos \vartheta + i \sin \vartheta)$ and taking into account that $|i| = 1$, $\arg(i) = \pi/2 + 2k\pi$, $k \in \mathbb{Z}$, we obtain

$$\begin{cases} r^6 = 1 \\ 6\vartheta = \pi/2 + 2k\pi \end{cases} \quad \begin{cases} r = 1 \\ \vartheta = \pi/12 + k\pi/3. \end{cases}$$

We therefore have the six roots

$$z_k = \cos((\pi + 4k\pi)/12) + i \sin((\pi + 4k\pi)/12), \quad k = 0, 1, 2, 3, 4, 5.$$

Those satisfying the condition $\Re(z)\Im(z) < 0$ (points in the second or fourth quadrant) are:

$$z_2 = -\sqrt{1/2} + i\sqrt{1/2}, \quad z_5 = \sqrt{1/2} - i\sqrt{1/2}.$$

Exercise 13.46

Solve the following equation in the complex variable z .

$$z^3 + 8 = 0$$

Solution

The equation to be solved is, clearly,

$$z^3 = -8$$

Setting $z = r(\cos \vartheta + i \sin \vartheta)$ and taking into account that $|-8| = 8$, $\arg(-8) = \pi + 2k\pi$, $k \in \mathbb{Z}$, we obtain

$$\begin{cases} r^3 = 8 \\ 3\vartheta = \pi + 2k\pi \end{cases} \quad \begin{cases} r = 2 \\ \vartheta = (2k+1)\pi/3. \end{cases}$$

We therefore have the three roots

$$z_k = 2 \cos((2k+1)\pi/3) + 2i \sin((2k+1)\pi/3), \quad k = 0, 1, 2,$$

i.e.,

$$z_0 = 1 + \sqrt{3}i, \quad z_1 = -2, \quad z_2 = 1 - \sqrt{3}i.$$

Exercise 13.47

Solve the following equation in the complex variable z .

$$z^3 = -27i$$

Solution

Setting $z = r(\cos \vartheta + i \sin \vartheta)$ and taking into account that $|-27i| = 27$, $\arg(-27i) = -\pi/2 + 2k\pi$, $k \in \mathbb{Z}$, we obtain

$$\begin{cases} r^3 = 27 \\ 3\vartheta = -\pi/2 + 2k\pi \end{cases} \quad \begin{cases} r = 3 \\ \vartheta = (4k-1)\pi/6. \end{cases}$$

We therefore have the three roots

$$z_k = 3 \cos((4k-1)\pi/6) + 3i \sin((4k-1)\pi/6), \quad k = 0, 1, 2,$$

i.e.,

$$z_0 = 3\sqrt{3}/2 - 3i/2, \quad z_1 = 3i, \quad z_2 = -3\sqrt{3}/2 - 3i/2.$$

Exercise 13.48

Solve the following equation in the complex variable z .

$$z^5 + (3 + i\sqrt{3})z = 0$$

Solution

We have

$$z(z^4 + 3 + i\sqrt{3}) = 0$$

hence one solution is $z = 0$. The other four are found by solving

$$z^4 = -3 - i\sqrt{3}.$$

Setting $z = r(\cos \vartheta + i \sin \vartheta)$ and taking into account that

$$|-3 - i\sqrt{3}| = 2\sqrt{3}, \quad \arg(-3 - i\sqrt{3}) = -5\pi/6 + 2k\pi, \quad k \in \mathbb{Z}$$

we obtain

$$\begin{cases} r^4 = 12^{1/2} \\ 4\vartheta = -5\pi/6 + 2k\pi \end{cases} \quad \begin{cases} r = 12^{1/8} \\ \vartheta = -5\pi/24 + k\pi/2. \end{cases}$$

Besides the solution $z = 0$, we therefore have four more roots

$$z_k = 12^{1/8} \cos((12k - 5)\pi/24) + 12^{1/8}i \sin((12k - 5)\pi/24), \quad k = 0, 1, 2, 3.$$

Exercise 13.49

Solve the following equation in the complex variable z .

$$\left(\frac{z+i}{1+i} \right)^3 = -8i$$

Solution

Setting $w = (z+i)/(1+i)$, we solve $w^3 = -8i$. Writing $w = r(\cos \vartheta + i \sin \vartheta)$ and taking into account that $|-8i| = 8$, $\arg(-8i) = -\pi/2 + 2k\pi$, $k \in \mathbb{Z}$, we obtain

$$\begin{cases} r^3 = 8 \\ 3\vartheta = -\pi/2 + 2k\pi \end{cases} \quad \begin{cases} r = 2 \\ \vartheta = (4k - 1)\pi/6. \end{cases}$$

We therefore have the three roots

$$w_k = 2 \cos((4k - 1)\pi/6) + 2i \sin((4k - 1)\pi/6), \quad k = 0, 1, 2,$$

i.e.,

$$w_0 = \sqrt{3} - i, \quad w_1 = 2i, \quad w_2 = -\sqrt{3} - i.$$

Going back to the variable z , from $z = (1+i)w - i$, the three solutions of the given equation are

$$z_0 = \sqrt{3} + 1 + i\sqrt{3} - 2i, \quad z_1 = -2 + i, \quad z_2 = 1 - \sqrt{3} - i\sqrt{3} - 2i.$$

Exercise 13.50

Solve the following equation in the complex variable z .

$$\left(\frac{z-i}{i}\right)^4 = -16$$

Solution

From $i^4 = 1$ we have $(z-i)^4 = -16$. Setting $w = z - i$, we solve $w^4 = -16$. Writing $w = r(\cos \vartheta + i \sin \vartheta)$ and taking into account that $|-16| = 16$, $\arg(-16) = -\pi + 2k\pi$, $k \in \mathbb{Z}$, we obtain

$$\begin{cases} r^4 = 16 \\ 4\vartheta = -\pi + 2k\pi \end{cases} \quad \begin{cases} r = 2 \\ \vartheta = -\pi/4 + k\pi/2. \end{cases}$$

We therefore have the four roots

$$w_k = 2 \cos((2k-1)\pi/4) + 2i \sin((2k-1)\pi/4), \quad k = 0, 1, 2, 3,$$

i.e.,

$$w_0 = \sqrt{2} - i\sqrt{2}, \quad w_1 = \sqrt{2} + i\sqrt{2}, \quad w_2 = -\sqrt{2} + i\sqrt{2}, \quad w_3 = -\sqrt{2} - i\sqrt{2}.$$

Going back to the variable z , from $z = w + i$, the four solutions of the given equation are

$$z_0 = \sqrt{2} - i\sqrt{2} + i, \quad z_1 = \sqrt{2} + i\sqrt{2} + i, \quad z_2 = -\sqrt{2} + i\sqrt{2} + i, \quad z_3 = -\sqrt{2} - i\sqrt{2} + i.$$

Exercise 13.51

Solve the following equation in the complex variable z .

$$z^5 = -\bar{z}$$

Solution

The equation can be seen as

$$z^5 = (-1)\bar{z}$$

Setting $z = r(\cos \vartheta + i \sin \vartheta)$ and taking into account that $|\bar{z}| = r$, $\arg(\bar{z}) = -\vartheta + 2k\pi$, $k \in \mathbb{Z}$, and recalling that $-1 = e^{i\pi}$, we obtain

$$\begin{cases} r^5 = r \\ 5\vartheta = \pi - \vartheta + 2k\pi \end{cases} \quad \begin{cases} r = 0, r = 1 \\ \vartheta = (2k+1)\pi/6. \end{cases}$$

At this point,

- for $r = 0$, we obtain $z_0 = 0$;
- for $r = 1$, we obtain the roots $z = \cos((2k+1)\pi/6) + i \sin((2k+1)\pi/6)$, $k = 0, 1, 2, 3, 4, 5$.

Exercise 13.52

The equation

$$z^4 - 2z^3 + 4z^2 - 2z + 3 = 0$$

has $z_1 = i$ as a solution. Denoting by z_2, z_3, z_4 the other solutions, we have:

- (a) $\Im(z_1 z_2 z_3 z_4) = 0$ (b) $\Re(z_1 + z_2 + z_3 + z_4) = 0$
 (c) $\Re((z_1 + z_2 + z_3 + z_4)^{-1}) = 1/2$ (d) none of the other answers is correct

Solution

The equation has real coefficients, hence the solutions come in conjugate pairs; in particular, another solution is $z_2 = -i$. It follows that the polynomial

$$p(z) = z^4 - 2z^3 + 4z^2 - 2z + 3$$

is divisible by $(z - i)(z + i)$, i.e., by $z^2 + 1$. Performing the division we obtain

$$p(z) = (z^2 + 1)(z^2 - 2z + 3)$$

hence among the solutions we also have the roots of

$$z^2 - 2z + 3 = 0$$

which are $z_3 = 1 - i\sqrt{2}$, $z_4 = 1 + i\sqrt{2}$.

Summarizing, the solutions of the given equation are

$$z_1 = i, \quad z_2 = -i, \quad z_3 = 1 - i\sqrt{2}, \quad z_4 = 1 + i\sqrt{2}.$$

Since $z_1 + z_2 + z_3 + z_4 = 2$ and $z_1 z_2 z_3 z_4 = 3$, the correct answers are (a) and (c).

Exercise 13.53

Let z_h , $h = 1, \dots, 6$ be the solutions of

$$z^6 - (8 - i)z^3 - 8i = 0.$$

We have:

$$(a) \sum_{h=1}^6 \Re(z_h) = -1 \quad (b) \sum_{h=1}^6 \Im(z_h) = -1$$

$$(c) \sum_{h=1}^6 \Re(z_h) = 1 \quad (d) \sum_{h=1}^6 \Im(z_h) = 0$$

$$(e) \sum_{h=1}^6 \Re(z_h) = 2 \quad (f) \text{ none of the other answers is correct}$$

Solution

It is convenient to factor it as

$$(z^3 - 8)(z^3 + i) = 0$$

and then solve the two equations

$$z^3 = 8, \quad z^3 = -i$$

with the usual method illustrated several times in the solutions of the previous exercises. We obtain the values:

$$z_1 = 2, \quad z_2 = -1 + i\sqrt{3}, \quad z_3 = -1 - i\sqrt{3},$$

$$z_4 = \sqrt{3}/2 - i/2, \quad z_5 = i, \quad z_6 = -\sqrt{3}/2 - i/2.$$

The correct answer is (d).