



SAPIENZA
UNIVERSITÀ DI ROMA

Mathematical Analysis 1

Volume 2 — Functions

The second volume: functions and their graphs. Functions between sets and real functions of a real variable, power, exponential and logarithmic functions, trigonometric functions, vibration phenomena, integer part and piecewise functions, hyperbolic functions, operations on graphs, composite and inverse functions. Definitions, results, many graphs and worked examples; at the end, the exercises of the part with solutions.

Fabio Furini

sites.google.com/view/fabiofurini

Department of Computer, Control and Management Engineering — Sapienza University of Rome

Online version and downloadable material:
fabiofurini.github.io/mathematical-analysis-1/
github.com/fabiofurini/mathematical-analysis-1

Academic year 2026–2027

Contents

1	Functions	3
1.1	The concept of function	3
1.2	Definition of function, domain, codomain and image	5
1.3	Surjections, injections and bijections	7
2	Real functions of a real variable	11
2.1	Real function of a real variable	11
2.2	Bounded functions	13
2.3	Symmetric functions	15
2.4	Monotonic functions	17
2.5	Periodic functions	18
3	Power functions	20
3.1	Power functions	20
3.2	Direct and inverse proportionality functions	26
4	Exponential and logarithmic functions	29
4.1	Exponential functions	29
4.2	Logarithmic functions	30
4.3	Change of base	30
5	Trigonometric functions	33
5.1	Trigonometric functions	33
5.2	Values of the trigonometric functions	36
5.3	Main trigonometric formulas	37
6	Vibration phenomena	40
6.1	Vibration phenomena	40
6.2	Damping and amplification effects	43
7	Integer part and fractional part functions	47
7.1	Integer part and fractional part functions	47
7.2	Piecewise-defined functions	49
8	Hyperbolic functions	51
8.1	Hyperbolic functions	51
8.2	Main hyperbolic formulas	52
9	Operations on graphs	54
9.1	Operations on graphs	54



10 Composite functions	62
10.1 Composite functions	62
11 Inverse functions	66
11.1 Invertible functions and inverse functions	66
12 Exercises	83
12.1 Domains of functions	83
12.2 Inverse functions	84

Functions

1.1 The concept of function

The existence of a **variable quantity** implies the existence of a **relation** between **two quantities**, that is, the dependence of one quantity on another.


- In each case, this relation follows a certain **law** or **formula**.

Example 1.1: Laws/formulas $\rightarrow$ functions

If a heavy object is dropped from a certain height, the distance traveled by the object varies with time t according to the formula:

$$s(t) = \frac{1}{2} g t^2 \quad \text{with } t \geq 0$$

where $g \approx 9,8$ is the gravitational acceleration constant.



Hence, the time t is associated with the distance traveled $s(t)$: $t \mapsto s(t)$.

Example 1.2: Laws/formulas $\rightarrow$ functions

If a unit capital is invested for one year at the annual rate i , with interest paid monthly at a rate $\frac{i}{12}$, we have:

- after the first month, a capital equal to:

$$1 + \frac{i}{12} \cdot 1 = \underbrace{1 + \frac{i}{12}}_{\alpha}$$

- after the second month, a capital equal to:


$$\underbrace{1 + \frac{i}{12}}_{\alpha} + \frac{i}{12} \underbrace{\left(1 + \frac{i}{12}\right)}_{\alpha} = \left(1 + \frac{i}{12}\right) \underbrace{\left(1 + \frac{i}{12}\right)}_{\beta} = \underbrace{\left(1 + \frac{i}{12}\right)^2}_{\beta}$$

- after the third month, a capital equal to:

$$\underbrace{\left(1 + \frac{i}{12}\right)^2}_{\beta} + \frac{i}{12} \underbrace{\left(1 + \frac{i}{12}\right)^2}_{\beta} = \underbrace{\left(1 + \frac{i}{12}\right)^2}_{\beta} \underbrace{\left(1 + \frac{i}{12}\right)}_{\beta} = \underbrace{\left(1 + \frac{i}{12}\right)^3}_{\beta}$$

Repeating the argument, the capital at the end of the year therefore depends on the rate i according to the formula:

$$k(i) = \left(1 + \frac{i}{12}\right)^{12} \quad \text{with } i \in [0, 1]$$



Hence, the rate i is associated with the capital $k(i)$ at the end of the year: $i \mapsto k(i)$

- In each of the previous examples, a *real number* (**input**) is associated *uniquely* with another *real number* (**output**). It is precisely the **uniqueness of the relation** that characterizes a function.
- In general, the **admissible inputs** for a given relation (function) are subject to natural restrictions, related to the very nature of the relation.

Example 1.3: Admissible inputs

- In the first example the “starting” real number has the meaning of *time*; imagining that the object is dropped at an initial time $t = 0$, it is clear that we must restrict ourselves to times $t \geq 0$.
- In the second example, clearly we must have $0 \leq i \leq 1$, where 1 corresponds to a rate of 100%.

1.2 Definition of function, domain, codomain and image

Definition 1.1: of domain

The set of admissible inputs for a given function is called the **domain**.

- The expressions **independent variable** (for a generic *input*) and **dependent variable** (for the *output*) are often used.

Example 1.4: Other types of relations

Let us consider the set of students of a given course.

- If A is the set of students and B is the set of their names, the relation that associates each student with his or her name is *unique* (single-valued).
- In this case neither A nor B is a set of numbers, but nevertheless a unique (single-valued) relation between these two sets is well defined.

On the other hand, the converse is not necessarily true: there could be two students with the same name.

Definition 1.2: of function

Given any two sets A , B , a **function** f with **domain** A and values in B (or also “with **codomain** B ”) is any law that associates with each element of A one and only one element of B .

- The notation:

$$f : A \rightarrow B$$

(which reads “ f defined from A to B ”), indicates the **domain** and the **codomain** of the function f .

- The notation

$$f : x \mapsto f(x)$$

(which reads “ f maps x to $f(x)$ ”) indicates how the function f acts on the elements.

- The **symbol** $f(x)$ **denotes the output or the value** that the function f associates with x , and it must not be confused with the symbol f , which denotes the **function itself**.

Example 1.5: Notation for functions

For the function

$$k(i) = \left(1 + \frac{i}{12}\right)^{12}$$

of the previous example we would have:

$$k : [0, 1] \rightarrow \mathbb{R}$$

$$k : i \mapsto \left(1 + \frac{i}{12}\right)^{12}$$


- The notation

$$y = f(x)$$

is also used to indicate that the variable y is a function of x . In this case y is the output or the value that f associates with the input x .

For brevity, sometimes the value of $f(x)$ is defined directly in terms of x , as for example $f(x) = x^2 + 10$.

- In general, we can think of a function as a **black box** that associates with each admissible input x (**input**) a unique output $f(x)$ (**output**):



Definition 1.3: of image and image of the domain

The output corresponding to x is called the **image** of x ; the set of possible outputs is called the **image of the domain A under f** and is denoted by the symbol $f(A)$ or $\text{Im } f$.

- Note that in the notation $f : A \rightarrow B$, the codomain B can be larger than the image $f(A)$, that is, in general we have

$$f(A) \subseteq B.$$


- If f is real-valued, we usually write $f : A \rightarrow \mathbb{R}$ without specifying the actual image of f .

1.3 Surjections, injections and bijections

Definition 1.4: of surjection (surjective function)

A function f is a **surjection** if the image of its domain coincides with its codomain.

- A surjection $f : A \rightarrow B$ is sometimes also called a mapping *onto* B (from A).
- Set representation:


Example 1.6: surjective and non-surjective functions

- The function $f(n) = \lfloor \frac{n}{2} \rfloor$ is a surjective function from $\mathbb{N}$ to $\mathbb{N}$, since every element of the codomain $\mathbb{N}$ is the image of some value of the domain.
- The function $f(n) = 2n$ is not a surjective function from $\mathbb{N}$ to $\mathbb{N}$, since no argument of f produces 3 as a value.
- The function $f(n) = 2n$ is, however, a surjective function from the natural numbers to the even numbers.

Observation 1.5

Given two sets A and B , and a function $f : A \rightarrow B$, if f is surjective then $|A| \geq |B|$.

Proof. We proceed by induction on the number n of elements in the codomain of the function.

- **Base case**

If there is only one element in the codomain ($|B| = 1$), since the function is surjective, we have $|A| \geq 1$ (every element of B is the image of at least one element of A). Hence $1 \geq 1$, which is clearly true.

- **Inductive step**

Assume then that all functions with codomain of size n satisfy $|A_n| \geq |B_n|$.

Considering codomains of size $n+1$ we have $|B_{n+1}| = |B_n| + 1$. Since the codomain has one more element and the function is surjective, we have $|A_{n+1}| \geq |A_n| + 1$. Substituting we obtain:


$$\underbrace{|A_n|}_{\leq |A_{n+1}|-1} \geq \underbrace{|B_n|}_{= |B_{n+1}|-1} \quad \text{hence } |A_{n+1}| \geq |B_{n+1}|.$$

□

Definition 1.6: of injection (injective function)

A function f is an **injection** if distinct arguments of f produce distinct values, that is, if $a \neq b$ implies $f(a) \neq f(b)$.

- set representation:



Example 1.7

- The function $f(n) = 2n$ is an injective function from $\mathbb{N}$ to $\mathbb{N}$, since each even number b is the image under f of exactly one element of the domain, namely $n = \frac{b}{2}$
- The function $f(n) = \lfloor \frac{n}{2} \rfloor$ is not an injective function, since the value 1 is obtained from two arguments: 2 and 3.

- An injection is also called a **one-to-one** function.

Observation 1.7

Given two sets A and B , and a function $f : A \rightarrow B$, if f is injective then $|A| \leq |B|$.

Proof. We proceed by induction on the number n of elements in the domain of the function.

- **Base case**

If there is only one element in the domain ($|A| = 1$), since each element of the domain is associated with one and only one element of the codomain, we have $|B| \geq 1$. Hence $1 \leq 1$, which is clearly true.

- **Inductive step**

Assume then that all functions with domain of size n satisfy $|A_n| \leq |B_n|$.

Considering domains of size $n+1$ we have $|A_{n+1}| = |A_n| + 1$ and, since the function is injective, $|B_{n+1}| \geq |B_n| + 1$. Substituting we obtain


$$\underbrace{|A_n|}_{= |A_{n+1}| - 1} \leq \underbrace{|B_n|}_{= |B_{n+1}| - 1} \quad \text{hence} \quad |A_{n+1}| \leq |B_{n+1}|.$$

□

Definition 1.8: of bijection (bijective function or one-to-one correspondence)

A function f is a **bijection** if: (i) it is injective and (ii) it is surjective.

- set representation:


Example 1.8

- The function $f(n) = (-1)^n \lceil \frac{n}{2} \rceil$ is a bijection from $\mathbb{N}$ to $\mathbb{Z}$. The values of the function are:

$$f(0) = 0, \quad f(1) = -1, \quad f(2) = 1, \quad f(3) = -2, \quad f(4) = 2 \dots$$

The function is injective, since no element of $\mathbb{Z}$ is the image of more than one element of $\mathbb{N}$. The function is surjective, since all elements of $\mathbb{Z}$ are images of some element of $\mathbb{N}$.

- A bijection is also called a **one-to-one** correspondence, since it pairs elements of the domain with elements of the codomain.
- A bijection from a set A to itself is also called a **permutation**.

Real functions of a real variable

2.1 Real function of a real variable

Definition 2.1: real function of a real variable

A function whose domain D is a subset of $\mathbb{R}$ and whose codomain is $\mathbb{R}$:


$$f : D \subseteq \mathbb{R} \rightarrow \mathbb{R}, \quad f : x \mapsto f(x)$$

is called a **real function of a real variable**.

- These are functions in which the “input” variable x and the “output” variable $f(x)$ are real numbers.
- The most common real functions of a real variable have as domain D and as image $f(D)$ an **interval** (possibly the whole of $\mathbb{R}$) or the union of a **finite number of intervals**.

The dependence of the output $f(x)$ on the input x is effectively visualized by drawing the **graph** of f , that is, the set of points of the plane with coordinates (x, y) such that $y = f(x)$ and x in the domain D .

Example of the graph of a real function of a real variable with domain $D = [a, b]$ (a closed and bounded interval):



Every line parallel to the y -axis that intersects the x -axis at a point x_0 of the domain D intersects the graph of f at one and only one point. ^a **Therefore, not all curves are graphs of functions.**


Note, instead, that nothing prevents a line parallel to the x -axis from intersecting the graph of f at several points or at no point at all.

^aIf the line did not intersect the graph, this would mean that no output corresponds to the input x . If the line intersected the graph at more than one point, this would mean that several distinct outputs correspond to the input x , and hence the function would no longer be single-valued.

Example 2.1: curve that does not correspond to the graph of a function

Consider, for example, the curve made of the points of the circle of radius r :

$$x^2 + y^2 = r^2 \iff y = \pm\sqrt{r^2 - x^2}$$



It is not possible to associate a unique output with the input $x_0 \in (-r, r)$, hence this curve does not correspond to the graph of a function!

2.2 Bounded functions

Definition 2.2: bounded functions

Given a function $f : D \subseteq \mathbb{R} \rightarrow \mathbb{R}$, the function is said to be

$$\begin{cases} \text{bounded above} & \text{if } \exists M \in \mathbb{R} : f(x) \leq M, \quad \forall x \in D \\ \text{bounded below} & \text{if } \exists M \in \mathbb{R} : f(x) \geq M, \quad \forall x \in D \\ \text{bounded} & \text{if } \exists M \in \mathbb{R}_{\geq 0} : |f(x)| \leq M, \quad \forall x \in D \end{cases}$$

• Graphically, we have:

1. a function is bounded above if its graph is contained in the lower half-plane bounded by a line parallel to the x -axis
2. a function is bounded below if its graph is contained in the upper half-plane bounded by a line parallel to the x -axis
3. a function is bounded if its graph is contained in a horizontal strip

Example 2.2: bounded function


Consider the function:

$$f : \mathbb{R} \rightarrow \mathbb{R}, \quad f : x \mapsto \frac{1}{1+x^2} + 1$$

we have

$$\frac{1}{1+x^2} + 1 = \frac{1+x^2-x^2}{1+x^2} + 1 = 2 - \frac{x^2}{1+x^2} \quad \text{and} \quad \frac{x^2}{1+x^2} \geq 0, \quad \forall x \in \mathbb{R}$$


$$\text{hence} \quad 1 < \frac{1}{1+x^2} + 1 \leq 2, \quad \forall x \in \mathbb{R}$$


Example 2.3: Unbounded function

The function

$$f : \mathbb{R} \rightarrow \mathbb{R}, \quad f : x \mapsto x^3 \quad (y = x^3)$$

is bounded neither above nor below.




Example 2.4: Function bounded below

The function

$$f : \mathbb{R} \rightarrow \mathbb{R}, \quad f : x \mapsto x^2 \quad (y = x^2)$$

is bounded below; indeed, $x^2 \geq 0, \forall x \in \mathbb{R}$



- Equivalently, we can say that a function is *bounded above* (*bounded below*, *bounded*) if, respectively, its **image** is a subset of $\mathbb{R}$ that is *bounded above* (*bounded below*, *bounded*).

2.3 Symmetric functions

Definition 2.3: even function

Functions whose graph is symmetric with respect to the y -axis are called **even**.

- They are characterized by the relation

$$f(-x) = f(x)$$

which expresses the equality of the ordinates corresponding to the points x and $-x$, which are symmetric with respect to $x = 0$.

Definition 2.4: Odd function

Functions whose graph is symmetric with respect to the origin are called **odd**.



- They are characterized by the relation

$$f(-x) = -f(x)$$

which expresses the fact that the ordinates corresponding to the points x and $-x$, which are symmetric with respect to $x = 0$, are the opposite of each other.

Example 2.5: Even and odd functions

For example, the function $x \mapsto x^2$ is even, while $x \mapsto x^3$ is odd. More generally, powers with integer exponent are even functions if the exponent is even and odd functions if the exponent is odd.

1. Example of the graph of an even function:

2. Example of the graph of an odd function:


A function *cannot* have a graph that is symmetric with respect to the x -axis, since the correspondence would no longer be single-valued.

2.4 Monotonic functions

Definition 2.5: Increasing function

A function is called **non-decreasing** if for every pair of points x_1, x_2 in the domain of f we have:

$$x_1 > x_2 \implies f(x_1) \geq f(x_2) \quad (2.1)$$

Definition 2.6: Strictly increasing function

A function is called **increasing** if for every pair of points x_1, x_2 in the domain of f we have:

$$x_1 > x_2 \implies f(x_1) > f(x_2) \quad (2.2)$$

Definition 2.7: Decreasing function

A function is called **non-increasing** if for every pair of points x_1, x_2 in the domain of f we have:

$$x_1 > x_2 \implies f(x_1) \leq f(x_2) \quad (2.3)$$

Definition 2.8: Strictly decreasing function

A function is called **decreasing** if for every pair of points x_1, x_2 in the domain of f we have:

$$x_1 > x_2 \implies f(x_1) < f(x_2) \quad (2.4)$$


- A function f is *non-decreasing* if, as x increases, the corresponding ordinate on the graph of the function does not decrease (hence it either stays the same or increases);
- A function f is *non-increasing* if, as x increases, the corresponding ordinate on the graph of the function does not increase (hence it either stays the same or decreases).

Increasing or decreasing functions are called **monotonic**. Strictly increasing or strictly decreasing functions are called **strictly monotonic**.


Example 2.6: Monotonic functions

For example, the function $x \mapsto x^3$ is strictly monotonic increasing; the constant function $x \mapsto k$ (whose graph is the line with equation $y = k$) is both non-decreasing and non-increasing.

1. Example of the graph of a non-decreasing function (horizontal segment):



2. Example of the graph of an increasing function:



- Increasing or decreasing functions (strictly or not) are called **monotonic**.

2.5 Periodic functions

Definition 2.9: Periodic function

A (non-constant) function $f : D \rightarrow \mathbb{R}$ is **periodic** with period T , $T > 0$, if T is the smallest positive real number such that

$$f(x + T) = f(x) \quad \text{for every } x \in D$$


- Every interval of length T contained in D is called a **periodicity interval**.

Example 2.7: Periodic functions

Typical examples of periodic functions are the *trigonometric functions* $x \mapsto \sin(x)$ ($T = 2\pi$), $x \mapsto \cos(x)$ ($T = 2\pi$) and $x \mapsto \tan(x)$ ($T = \pi$).

Example 2.8: Graph of periodic functions

Graph of a periodic function with period $T = 2$:



- Lines parallel to the x -axis have equation

$$y = k \quad k \in \mathbb{R}$$

Power functions

3.1 Power functions

Definition 3.1: power functions

Given $\alpha \in \mathbb{R}, \alpha \neq 0$, the function:

$$f : D \subseteq \mathbb{R} \rightarrow \mathbb{R}, \quad f : x \mapsto x^\alpha \quad (3.1)$$

is called the **power function** with exponent α . The domain D depends on the value of α :

- with rational exponent $\alpha = \frac{m}{n} \in \mathbb{Q}$, $m \in \mathbb{Z}$ and $n \in \mathbb{N}_+$ coprime, we have four cases:

$$D = \begin{cases} \mathbb{R} & \text{if } n \text{ odd and } \frac{m}{n} > 0 \\ \mathbb{R} \setminus \{0\} & \text{if } n \text{ odd and } \frac{m}{n} < 0 \\ [0, +\infty) & \text{if } n \text{ even and } \frac{m}{n} > 0 \\ (0, +\infty) & \text{if } n \text{ even and } \frac{m}{n} < 0 \end{cases}$$

- with real, non-rational exponent $\alpha \in \mathbb{R} \setminus \mathbb{Q}$, we have two cases:

$$D = \begin{cases} [0, +\infty) & \text{if } \alpha > 0 \\ (0, +\infty) & \text{if } \alpha < 0 \end{cases}$$

3.1.1 Rational exponent

With $\alpha = \frac{m}{n} \in \mathbb{Q}$, $m \in \mathbb{Z}$ and $n \in \mathbb{N}_+$ coprime, we have a power function with rational exponent:


$$f : D \subseteq \mathbb{R} \rightarrow \mathbb{R}, \quad f : x \mapsto x^{\frac{m}{n}} = \sqrt[n]{x^m}$$

We have the following three cases, depending on whether m and n are even or odd.

1. With n odd and m even, it is an **even function**, i.e., $(-x)^{\frac{m}{n}} = x^{\frac{m}{n}}$. For the sign and the monotonicity we have:

$$x^{\frac{m}{n}} > 0, \quad \forall x \in \mathbb{R} \setminus \{0\}; \quad x^{\frac{m}{n}} = 0 \iff x = 0 \quad \text{and} \quad \frac{m}{n} > 0$$


$$\begin{cases} \text{if } \frac{m}{n} > 0 & \text{then } f \text{ is decreasing in } (-\infty, 0] \text{ and increasing in } [0, +\infty) \\ \text{if } \frac{m}{n} < 0 & \text{then } f \text{ is increasing in } (-\infty, 0) \text{ and decreasing in } (0, +\infty) \end{cases}$$



2. With n odd and m odd, it is an **odd function**, i.e., $(-x)^{\frac{m}{n}} = -x^{\frac{m}{n}}$. For the sign and the monotonicity we have:


$$x^{\frac{m}{n}} < 0, \quad \forall x < 0, \quad x^{\frac{m}{n}} > 0, \quad \forall x > 0; \quad x^{\frac{m}{n}} = 0 \iff x = 0 \quad \text{and} \quad \frac{m}{n} > 0$$

$$\begin{cases} \text{if } \frac{m}{n} > 0 & \text{then } f \text{ is increasing in } \mathbb{R} \\ \text{if } \frac{m}{n} < 0 & \text{then } f \text{ is decreasing in } (-\infty, 0) \text{ and in } (0, +\infty) \end{cases}$$




Example 3.1: graphs of power functions with rational exponent $\frac{m}{n}$ and n odd

With m even:




With m odd:




3. With n even and m odd (m cannot be even since they are coprime), for the sign and the monotonicity we have:

$$x^{\frac{m}{n}} > 0, \quad \forall x > 0; \quad x^{\frac{m}{n}} = 0 \iff x = 0 \quad \text{and} \quad \frac{m}{n} > 0$$

$$\begin{cases} \text{if } \frac{m}{n} > 0 & \text{then } f \text{ is increasing in } [0, +\infty) \\ \text{if } \frac{m}{n} < 0 & \text{then } f \text{ is decreasing in } (0, +\infty) \end{cases}$$



Example 3.2: graphs of power functions with rational exponent $\frac{m}{n}$ and n even




3.1.2 Real exponent


- With $\alpha \in \mathbb{R} \setminus \mathbb{Q}$, for the sign and the monotonicity we have:

$$x^\alpha > 0, \quad \forall x > 0; \quad x^\alpha = 0 \iff x = 0 \text{ and } \alpha > 0$$

$$\begin{cases} \text{if } \alpha > 0 & \text{then } f \text{ is increasing in } [0, +\infty) \\ \text{if } \alpha < 0 & \text{then } f \text{ is decreasing in } (0, +\infty) \end{cases}$$



Example 3.3: graphs of power functions with real exponent



3.1.3 Polynomials

Definition 3.2: polynomial

Given $n + 1$ values $a_i \in \mathbb{R}$ with $i \in \{0, 1, \dots, n\}$ and $a_n \neq 0$, the function:

$$f : \mathbb{R} \rightarrow \mathbb{R}, \quad f : x \mapsto \underbrace{\sum_{i=0}^n a_i x^i}_{=P_n(x)} \quad (3.2)$$

is called a **polynomial** of degree n .


- For each **monomial** $i \in \{0, 1, \dots, n\}$:
 1. the value a_i is the **coefficient of the monomial**
 2. the power function x^i with integer exponent i is the **literal part** of the monomial

The value a_0 is the **constant term** of the polynomial since $x^0 = 1$.

Example 3.4: polynomial

For example, with $n = 4$, $a_0 = 1$, $a_1 = -2$, $a_2 = 0$, $a_3 = \frac{1}{2}$ and $a_4 = \frac{1}{9}$ we have the following polynomial of degree four:

$$P_4(x) = \sum_{i=0}^4 a_i x^i = 1 - 2x + \frac{1}{2}x^3 + \frac{1}{9}x^4$$




3.2 Direct and inverse proportionality functions

With $\alpha = 1$ and given $\lambda \in \mathbb{R}$, we have the family of **direct proportionality (or linear) functions**:


$$f : \mathbb{R} \rightarrow \mathbb{R}, \quad f : x \mapsto \lambda x \quad \text{where } \lambda \text{ is the constant of direct proportionality}$$

Moreover, $\lambda = \tan \vartheta$ and ϑ is the angle between the line $y = \lambda x$ and the x -axis.



With $\alpha = -1$ and given $\lambda \in \mathbb{R}$, we have the family of **inverse proportionality functions (or rectangular hyperbolas)**:




$$f : \mathbb{R} \setminus \{0\} \rightarrow \mathbb{R}, \quad f : x \mapsto \frac{\lambda}{x} \quad \text{where } \lambda \text{ is the constant of inverse proportionality}$$





Given $\alpha \in \mathbb{R}$ and $\lambda \in \mathbb{R}$, we have the family of **power functions multiplied by a constant**:

$$f : D \subseteq \mathbb{R} \rightarrow \mathbb{R}, \quad f : x \mapsto \lambda x^\alpha \quad \text{where } \lambda \text{ is the multiplicative constant}$$

- For example, with α equal to 2 or equal to 3 and $\lambda \in \mathbb{R}$ we have:



- For example, with $\alpha = \frac{1}{2}$ and $\lambda \in \mathbb{R}$ we have:



Exponential and logarithmic functions

4.1 Exponential functions

Definition 4.1: exponential functions


Given $b \in \mathbb{R}_+ \setminus \{1\}$, the function:

$$f : \mathbb{R} \rightarrow \mathbb{R}, \quad f : x \mapsto b^x \quad (4.1)$$

is called the **exponential function** with base b .

- Regarding positivity and monotonicity, we have:

$$b^x > 0, \quad \forall x \in \mathbb{R} \quad \text{and} \quad \begin{cases} \text{if } 0 < b < 1 & \text{then } f \text{ is decreasing on } \mathbb{R} \\ \text{if } b > 1 & \text{then } f \text{ is increasing on } \mathbb{R} \end{cases}$$



4.2 Logarithmic functions

Definition 4.2: logarithmic functions

Given $a \in \mathbb{R}_{>0} \setminus \{1\}$, the function:

$$f : (0, +\infty) \rightarrow \mathbb{R}, \quad f : x \mapsto \log_a x \quad (4.2)$$


is called the **logarithmic function** with base a .

- Regarding positivity and monotonicity, we have:

$$\left\{ \begin{array}{l} \text{if } 0 < a < 1 \quad \log_a x > 0, \quad \forall x \in (0, 1), \quad \log_a x < 0, \quad \forall x \in (1, +\infty) \\ \quad \text{and } f \text{ is decreasing on } \mathbb{R}_{>0} \\ \\ \text{if } a > 1 \quad \log_a x < 0, \quad \forall x \in (0, 1), \quad \log_a x > 0, \quad \forall x \in (1, +\infty) \\ \quad \text{and } f \text{ is increasing on } \mathbb{R}_{>0} \end{array} \right.$$

Moreover, we have:

$$\log_a x = 0 \iff x = 1, \quad \forall a \in \mathbb{R}_{>0} \setminus \{1\}$$



4.3 Change of base

Given any base $c \in \mathbb{R}_{>0} \setminus \{1\}$, all exponential and logarithmic functions can be rewritten in terms of another base $d \in \mathbb{R}_{>0} \setminus \{1\}$ as follows:


$$c^x = d^{\log_d c^x} = d^{x \log_d c} \quad \text{hence with } \lambda = \log_d c \quad \text{we get} \quad c^x = d^{\lambda x}$$

$$\log_c x = \frac{\log_d x}{\log_d c} \quad \text{hence with } \lambda = \frac{1}{\log_d c} \quad \text{we get} \quad \log_c x = \lambda \log_d x$$


The logarithmic function with base e is also written $\log x$ or $\ln x$, while the one with base 2 is also written $\lg x$.

Example 4.1: graphs of exponential and logarithmic functions

Exponential functions:



Logarithmic functions:



Trigonometric functions

5.1 Trigonometric functions

Definition 5.1: sine function and cosine function

The trigonometric functions:

$$f : \mathbb{R} \rightarrow \mathbb{R}, \quad f : x \mapsto \sin x \quad (5.1)$$

$$f : \mathbb{R} \rightarrow \mathbb{R}, \quad f : x \mapsto \cos x \quad (5.2)$$


are called **sine** and **cosine**.

- The sine is an **odd** function, while the cosine is an **even** function. Hence we have:

$$\sin(-x) = -\sin x, \quad \forall x \in \mathbb{R} \quad \text{and} \quad \cos(-x) = \cos x, \quad \forall x \in \mathbb{R}$$

- The sine and the cosine are periodic functions with period:

$$T = 2\pi \quad \text{and we have the fundamental identity} \quad \sin^2 x + \cos^2 x = 1, \quad \forall x \in \mathbb{R}$$



We have

$$\cos\left(x - \frac{\pi}{2}\right) = \sin x, \quad \forall x \in \mathbb{R}$$

- From the sign and the monotonicity of the sine function we have:

$$\sin x = 0 \iff x = k\pi, \quad \forall k \in \mathbb{Z} \quad \begin{cases} \sin x > 0 & \text{if } x \in (2k\pi, 2k\pi + \pi), \quad \forall k \in \mathbb{Z} \\ \sin x < 0 & \text{if } x \in (2k\pi - \pi, 2k\pi), \quad \forall k \in \mathbb{Z} \end{cases}$$

$$\begin{cases} \text{if } x \in (2k\pi - \frac{\pi}{2}, 2k\pi + \frac{\pi}{2}), \quad \forall k \in \mathbb{Z} & f \text{ is increasing} \\ \text{if } x \in (2k\pi + \frac{\pi}{2}, 2k\pi + \frac{3\pi}{2}), \quad \forall k \in \mathbb{Z} & f \text{ is decreasing} \end{cases}$$

- From the sign and the monotonicity of the cosine function we have:

$$\cos x = 0 \iff x = k\pi - \frac{\pi}{2}, \quad \forall k \in \mathbb{Z}$$

$$\begin{cases} \cos x > 0 & \text{if } x \in (2k\pi - \frac{\pi}{2}, 2k\pi + \frac{\pi}{2}), \quad \forall k \in \mathbb{Z} \\ \cos x < 0 & \text{if } x \in (2k\pi + \frac{\pi}{2}, 2k\pi + \frac{3\pi}{2}), \quad \forall k \in \mathbb{Z} \end{cases}$$



$$\begin{cases} \text{if } x \in (2k\pi + \pi, 2k\pi + 2\pi), \quad \forall k \in \mathbb{Z} & f \text{ is increasing} \\ \text{if } x \in (2k\pi, 2k\pi + \pi), \quad \forall k \in \mathbb{Z} & f \text{ is decreasing} \end{cases}$$

The trigonometric functions:

$$f : \mathbb{R} \setminus \{k\pi - \frac{\pi}{2}, \quad \forall k \in \mathbb{Z}\} \rightarrow \mathbb{R}, \quad f : x \mapsto \tan x = \frac{\sin x}{\cos x} \quad (5.3)$$

$$f : \mathbb{R} \setminus \{k\pi, \quad \forall k \in \mathbb{Z}\} \rightarrow \mathbb{R}, \quad f : x \mapsto \cot x = \frac{\cos x}{\sin x} \quad (5.4)$$

are called **tangent** and **cotangent**; they are both odd and periodic with period $T = \pi$.




The trigonometric functions:

$$f : \mathbb{R} \setminus \{k\pi - \frac{\pi}{2}, \forall k \in \mathbb{Z}\} \rightarrow \mathbb{R}, \quad f : x \mapsto \sec x = \frac{1}{\cos x} \quad (5.5)$$


$$f : \mathbb{R} \setminus \{k\pi, \forall k \in \mathbb{Z}\} \rightarrow \mathbb{R}, \quad f : x \mapsto \csc x = \frac{1}{\sin x} \quad (5.6)$$

are called **secant** and **cosecant** and they are periodic with period $T = 2\pi$. The secant function is even, while the cosecant function is odd.



5.2 Values of the trigonometric functions

	cos	sin	tan	cot	sec	csc
0	1	0	0	$\pm\infty$	1	$\pm\infty$
$x = \frac{\pi}{6}$ (30°)	$\frac{\sqrt{3}}{2}$	$\frac{1}{2}$	$\frac{\sqrt{3}}{3}$	$\sqrt{3}$	$\frac{2}{3}\sqrt{3}$	2
$x = \frac{\pi}{4}$ (45°)	$\frac{\sqrt{2}}{2}$	$\frac{\sqrt{2}}{2}$	1	1	$\sqrt{2}$	$\sqrt{2}$
$x = \frac{\pi}{3}$ (60°)	$\frac{1}{2}$	$\frac{\sqrt{3}}{2}$	$\sqrt{3}$	$\frac{\sqrt{3}}{3}$	2	$\frac{2}{3}\sqrt{3}$
$x = \frac{\pi}{2}$ (90°)	0	1	$\pm\infty$	0	$\pm\infty$	1
$x = \pi$ (180°)	-1	0	0	$\pm\infty$	-1	$\pm\infty$



	cos	sin	tan	cot
x	$\cos x$	$\sin x$	$\tan x$	$\cot x$
$-x$	$\cos x$	$-\sin x$	$-\tan x$	$-\cot x$
$\frac{\pi}{2} + x$	$-\sin x$	$\cos x$	$-\cot x$	$-\tan x$
$\frac{\pi}{2} - x$	$\sin x$	$\cos x$	$\cot x$	$\tan x$
$\pi + x$	$-\cos x$	$-\sin x$	$\tan x$	$\cot x$
$\pi - x$	$-\cos x$	$\sin x$	$-\tan x$	$-\cot x$

5.3 Main trigonometric formulas

Relations among the trigonometric functions

	$\sin x$	$\cos x$	$\tan x$	$\cot x$
$\sin x$	-	$\pm\sqrt{1 - \cos^2 x}$	$\pm\sqrt{\frac{\tan^2 x}{1 + \tan^2 x}}$	$\pm\sqrt{\frac{1}{1 + \cot^2 x}}$
$\cos x$	$\pm\sqrt{1 - \sin^2 x}$	-	$\pm\sqrt{\frac{1}{1 + \tan^2 x}}$	$\pm\sqrt{\frac{\cot^2 x}{1 + \cot^2 x}}$
$\tan x$	$\pm\sqrt{\frac{\sin^2 x}{1 - \sin^2 x}}$	$\pm\sqrt{\frac{1 - \cos^2 x}{\cos^2 x}}$	-	$\frac{1}{\cot x}$
$\cot x$	$\pm\sqrt{\frac{1 - \sin^2 x}{\sin^2 x}}$	$\pm\sqrt{\frac{\cos^2 x}{1 - \cos^2 x}}$	$\frac{1}{\tan x}$	-

where the symbol $\pm$ means that the sign depends on the quadrant in which x lies

Addition and subtraction

$$\sin(x_1 \pm x_2) = \sin x_1 \cos x_2 \pm \sin x_2 \cos x_1 \quad (5.7a)$$

$$\cos(x_1 \pm x_2) = \cos x_1 \cos x_2 \mp \sin x_1 \sin x_2 \quad (5.7b)$$

$$\tan(x_1 \pm x_2) = \frac{\tan x_1 \pm \tan x_2}{1 \mp \tan x_1 \tan x_2} \quad (5.8a)$$

$$\cot(x_1 \pm x_2) = \frac{\cot x_1 \cot x_2 \mp 1}{\cot x_2 \pm \cot x_1} \quad (5.8b)$$

Double-angle formulas

$$\sin(2x) = 2 \sin x \cos x \quad (5.9a)$$

$$\begin{aligned} \cos(2x) &= \cos^2 x - \sin^2 x \\ &= 2 \cos^2 x - 1 \\ &= 1 - 2 \sin^2 x \end{aligned} \quad (5.9b)$$

$$\tan(2x) = \frac{2 \tan x}{1 - \tan^2 x} \quad (5.10a)$$

$$\cot(2x) = \frac{\cot^2 x - 1}{2 \cot x} \quad (5.10b)$$

Triple-angle formulas

$$\sin(3x) = 3 \sin x - 4 \sin^3 x \quad (5.11a)$$

$$\cos(3x) = 4 \cos^3 x - 3 \cos x \quad (5.11b)$$

Half-angle formulas

$$\sin \frac{x}{2} = \pm \sqrt{\frac{1 - \cos x}{2}} \quad (5.12a)$$

$$\cos \frac{x}{2} = \pm \sqrt{\frac{1 + \cos x}{2}} \quad (5.12b)$$

$$\begin{aligned} \tan \frac{x}{2} &= \pm \sqrt{\frac{1 - \cos x}{1 + \cos x}} \\ &= \frac{\sin x}{1 + \cos x} \\ &= \frac{1 - \cos x}{\sin x} \end{aligned} \quad (5.13a)$$

$$\begin{aligned}
 \cot \frac{x}{2} &= \pm \sqrt{\frac{1 + \cos x}{1 - \cos x}} \\
 &= \frac{\sin x}{1 - \cos x} \\
 &= \frac{1 + \cos x}{\sin x}
 \end{aligned} \tag{5.14a}$$

Parametric formulas (tangent half-angle substitution)

$$\sin x = \frac{2 \tan \frac{x}{2}}{1 + \tan^2 \frac{x}{2}} \tag{5.15a}$$

$$\cos x = \frac{1 - \tan^2 \frac{x}{2}}{1 + \tan^2 \frac{x}{2}} \tag{5.15b}$$

Sum-to-product formulas (prosthaphaeresis)

$$\sin x_1 + \sin x_2 = 2 \sin \frac{x_1 + x_2}{2} \cos \frac{x_1 - x_2}{2} \tag{5.16a}$$

$$\sin x_1 - \sin x_2 = 2 \sin \frac{x_1 - x_2}{2} \cos \frac{x_1 + x_2}{2} \tag{5.16b}$$

$$\cos x_1 + \cos x_2 = 2 \cos \frac{x_1 + x_2}{2} \cos \frac{x_1 - x_2}{2} \tag{5.16c}$$

$$\cos x_1 - \cos x_2 = -2 \sin \frac{x_1 + x_2}{2} \sin \frac{x_1 - x_2}{2} \tag{5.16d}$$

Product-to-sum formulas (Werner's formulas)

$$\sin x_1 \sin x_2 = \frac{1}{2} \left(\cos(x_1 - x_2) - \cos(x_1 + x_2) \right) \tag{5.17a}$$

$$\sin x_1 \cos x_2 = \frac{1}{2} \left(\sin(x_1 - x_2) + \sin(x_1 + x_2) \right) \tag{5.17b}$$

$$\cos x_1 \cos x_2 = \frac{1}{2} \left(\cos(x_1 - x_2) + \cos(x_1 + x_2) \right) \tag{5.17c}$$

Additional formulas

$$\sin^2 x = \frac{1 - \cos(2x)}{2} \tag{5.18a}$$


$$\cos^2 x = \frac{1 + \cos(2x)}{2} \tag{5.18b}$$

$$\sin x \cos x = \frac{\sin(2x)}{2} \tag{5.18c}$$


Vibration phenomena

6.1 Vibration phenomena

- We have seen that the sine and cosine functions are periodic with **period** 2π . We will use the variable t to denote time.



- Recall that $\sin(t + \varphi)$ corresponds to a **phase shift** φ (horizontal translation).



The functions:

$$t \mapsto a \sin(\omega t), \quad t \mapsto b \cos(\omega t) \quad (6.1)$$

where a , b and ω are positive real numbers, are called **elementary vibrations**. They are periodic functions with

$$\text{period } T = \frac{2\pi}{\omega}$$

Example 6.1: period

For example, considering $t' = t + \frac{2\pi}{\omega}$:

$$\sin \left[\omega \underbrace{\left(t + \frac{2\pi}{\omega} \right)}_{t'} \right] = \sin(\omega t + 2\pi) = \sin \omega t$$

- Clearly we have:

$$|a \sin(\omega t)| \leq a \quad \text{and} \quad |b \cos(\omega t)| \leq b$$

The functions (6.1) describe elementary vibrations characterized by

amplitude a and b (respectively)

$$\text{angular frequency } \omega = \frac{2\pi}{T}$$


also called *angular velocity*, which indicates how many periods there are in an interval of length 2π . Moreover, they are characterized by the

$$\text{frequency } \nu = \frac{\omega}{2\pi}$$

which indicates how many times the function repeats itself in an interval of length 1

Example 6.2: of amplitude, angular frequency and frequency

$$t \mapsto 2 \sin \left(\frac{3}{2} t \right) \quad a = 2, \quad \omega = \frac{3}{2}, \quad T = \frac{4\pi}{3}$$



The angular frequency $\omega = \frac{3}{2}$ means that there are 1.5 periods in an interval of length 2π . The frequency ν is $\frac{3}{4\pi}$.

- The function

$$h(t) = a \sin(\omega t) + b \cos(\omega t) \quad (6.2)$$

describes the **superposition** of the two elementary vibrations with the **same angular frequency** ω . The latter is again an elementary vibration, **phase-shifted** with respect to the previous ones.

- Indeed, setting

$$A = \sqrt{a^2 + b^2}$$

we can write

$$h(t) = A \underbrace{\frac{a}{\sqrt{a^2 + b^2}}}_{\alpha} \sin(\omega t) + A \underbrace{\frac{b}{\sqrt{a^2 + b^2}}}_{\beta} \cos(\omega t) \quad (6.3)$$

- Now observe that the numbers

$$\alpha = \frac{a}{\sqrt{a^2 + b^2}} \quad \text{and} \quad \beta = \frac{b}{\sqrt{a^2 + b^2}}$$

satisfy the conditions

$$-1 \leq \alpha \leq 1 \quad -1 \leq \beta \leq 1 \quad \alpha^2 + \beta^2 = 1$$


Hence there exists a **unique angle** φ such that

$$\cos \varphi = \alpha \quad \sin \varphi = \beta$$

so that (6.3) can be rewritten in the following form

$$\begin{aligned} h(t) &= A \cos \varphi \sin(\omega t) + A \sin \varphi \cos(\omega t) \\ &= A \sin(\omega t + \varphi) \end{aligned} \quad (6.4)$$

Therefore $h(t)$ represents an elementary vibration with amplitude A , angular frequency ω , phase-shifted by an angle φ .



In summary:

$$a \sin(\omega t) + b \cos(\omega t) = A \sin(\omega t + \varphi)$$

$$A = \sqrt{a^2 + b^2} \quad \begin{cases} a = A \cos \varphi \\ b = A \sin \varphi \end{cases}$$

- Under fairly general conditions, a periodic natural phenomenon can be written as the superposition of a finite or infinite number of elementary vibrations with different frequencies (**Fourier series**).

6.2 Damping and amplification effects

- By multiplying an elementary vibration by **powers** or **exponentials**, we can model **damping** or **amplification** effects.

First example

- For example, consider the function

$$h(t) = t \sin(\omega t)$$

which models an **amplified vibration**.

- Since $-1 \leq \sin(\omega t) \leq 1$, we have

$$-t \leq t \sin(\omega t) \leq t$$

and hence the graph of $h(t)$ lies between the graphs of the lines with equations $y = -t$, $y = t$.

- At the points where

$$\sin(\omega t) = 1$$

i.e.,

$$t = \frac{\pi}{2\omega} + k \frac{2\pi}{\omega} \quad (k = 0, 1, 2, \dots)$$

the graph of $h(t)$ touches that of $y = t$.


- At the points where

$$\sin(\omega t) = -1$$


i.e.,


$$t = \frac{3\pi}{2\omega} + k \frac{2\pi}{\omega} \quad (k = 0, 1, 2, \dots)$$

the graph of $h(t)$ touches that of $y = -t$.



- From the graph we can see that multiplication by t has the effect of amplifying the vibration as t increases.





Second example

- For example, consider the function

$$k(t) = e^{-\alpha t} \sin(\omega t) \quad (\alpha > 0)$$

which models a **damped vibration**.


- Recalling that


$$e^{-\alpha t} = \left(\frac{1}{e^\alpha}\right)^t$$

is an exponential with base less than 1, considerations analogous to those made for the function $h(t)$ show that the graph of k lies between the graphs of the functions

$$y_1 = e^{-\alpha t} \quad \text{and} \quad y_2 = -e^{-\alpha t}$$

as shown in the figure:





Integer part and fractional part functions

7.1 Integer part and fractional part functions

- Two functions that are typically encountered when writing *algorithms* are the integer part (floor) function and the fractional part (or decimal part) function.

Definition 7.1: of the integer part function

The *integer part (floor) function* is:

$$f : \mathbb{R} \rightarrow \mathbb{Z}, x \mapsto [x] \quad (\text{or } x \mapsto \lfloor x \rfloor) \quad (7.1)$$

Definition 7.2: of the upper integer part function

The *upper integer part (ceiling) function* is:



$$f : \mathbb{R} \rightarrow \mathbb{Z}, x \mapsto \lceil x \rceil \quad (7.2)$$

Definition 7.3: of the fractional part function


The *fractional part function* is:

$$f : \mathbb{R} \rightarrow [0, 1), x \mapsto (x) \quad (7.3)$$

- The integer part function $f(x) = [x]$ is *monotone increasing*, like the upper integer part function $f(x) = \lceil x \rceil$.
- Graphs of the functions $f(x) = \lceil x \rceil$ and $f(x) = [x]$ on the interval $[-4, 4]$.



- Note that the fractional part is a periodic function with period 1:



7.2 Piecewise-defined functions

- Starting from the elementary functions, new ones can be built by using **different analytic definitions on different intervals**.

Definition 7.4: of piecewise-defined function


A function f whose value $f(x)$ is computed through different “instructions” depending on the interval in which x lies is called a **piecewise-defined function**.

Example 7.1: Piecewise-defined functions

Consider for example

$$f(x) = \begin{cases} \ln x & \text{if } x > 1, \\ x^2 & \text{if } 0 < x \leq 1, \\ x & \text{if } x \leq 0 \end{cases}$$

its graph is:



- From the *logical* / *algorithmic* point of view, one can say that a function of this kind has the peculiarity of being built using not only mathematical functions but also the **logical function** “if ... then”.

Hyperbolic functions

8.1 Hyperbolic functions

Definition 8.1: hyperbolic sine function and hyperbolic cosine function

The hyperbolic functions:

$$f : \mathbb{R} \rightarrow \mathbb{R}, \quad f : x \mapsto \operatorname{Sh} x = \frac{e^x - e^{-x}}{2} \quad (8.1)$$

$$f : \mathbb{R} \rightarrow \mathbb{R}, \quad f : x \mapsto \operatorname{Ch} x = \frac{e^x + e^{-x}}{2} \quad (8.2)$$


are called **hyperbolic sine** and **hyperbolic cosine**.

- The hyperbolic sine is an **odd** function, while the hyperbolic cosine is an **even** function. Hence we have:

$$\operatorname{Sh}(-x) = -\operatorname{Sh} x, \quad \forall x \in \mathbb{R} \quad \text{and} \quad \operatorname{Ch}(-x) = \operatorname{Ch} x, \quad \forall x \in \mathbb{R}$$

- We have the fundamental relations:

$$\operatorname{Ch}^2 x - \operatorname{Sh}^2 x = 1, \quad \forall x \in \mathbb{R} \quad \text{and} \quad \operatorname{Sh} x \leq \frac{e^x}{2} \leq \operatorname{Ch} x, \quad \forall x \in \mathbb{R} \quad (8.3)$$



- For the sign and monotonicity of the hyperbolic sine function we have:

$$\operatorname{Sh} x = 0 \iff x = 0 \quad \begin{cases} \operatorname{Sh} x > 0 & \text{if } x > 0 \\ \operatorname{Sh} x < 0 & \text{if } x < 0 \end{cases}$$

$$f \text{ is increasing } \quad \forall x \in \mathbb{R}$$

- For the sign and monotonicity of the hyperbolic cosine function we have:


$$\operatorname{Ch} x > 0, \quad \forall x \in \mathbb{R}$$

$$\begin{cases} \text{if } x < 0 & f \text{ is decreasing} \\ \text{if } x > 0 & f \text{ is increasing} \end{cases}$$

The hyperbolic function:

$$f : \mathbb{R} \rightarrow \mathbb{R}, \quad f : x \mapsto \operatorname{Th} x = \frac{e^x - e^{-x}}{e^x + e^{-x}} \quad (8.4)$$

is called the **hyperbolic tangent** and it is an odd function.



8.2 Main hyperbolic formulas

Addition

$$\operatorname{Sh}(x_1 + x_2) = \operatorname{Sh} x_1 \operatorname{Ch} x_2 + \operatorname{Sh} x_2 \operatorname{Ch} x_1 \quad (8.5a)$$

$$\operatorname{Ch}(x_1 + x_2) = \operatorname{Ch} x_1 \operatorname{Ch} x_2 + \operatorname{Sh} x_1 \operatorname{Sh} x_2 \quad (8.5b)$$

$$\operatorname{Th}(x_1 + x_2) = \frac{\operatorname{Th} x_1 + \operatorname{Th} x_2}{1 + \operatorname{Th} x_1 \operatorname{Th} x_2} \quad (8.5c)$$

Double-argument formulas

$$\operatorname{Sh}(2x) = 2 \operatorname{Sh} x \operatorname{Ch} x \quad (8.6a)$$

$$\operatorname{Ch}(2x) = \operatorname{Ch}^2 x + \operatorname{Sh}^2 x \quad (8.6b)$$

Operations on graphs

9.1 Operations on graphs

- Knowing the graph of a function $y = f(x)$, by means of simple geometric transformations it is possible to draw the graph of the following functions:

$$y_1 = f(x) + a, \quad a \in \mathbb{R} \quad (9.1a)$$

$$y_2 = f(x + a), \quad a \in \mathbb{R} \quad (9.1b)$$

$$y_3 = k f(x), \quad k \in \mathbb{R} \quad (9.1c)$$

$$y_4 = f(kx), \quad k \in \mathbb{R} \quad (9.1d)$$

$$y_5 = |f(x)| \quad (9.1e)$$

$$y_6 = f(|x|) \quad (9.1f)$$


- Therefore, starting from the elementary functions, it is possible to build, by means of these operations, a large variety of new functions. These operations create **translations**, **dilations** and **reflections** of the original function.

9.1.1 Operations related to $y_1 = f(x) + a$

Example 9.1: Operations on graphs related to $y_1 = f(x) + a$

Consider $y = \ln x$. Then

$$y_1 = \ln(x) + a \quad (x > 0).$$



- The graph of y_1 is obtained from that of y by a **translation** of a units **upward** if $a > 0$, **downward** if $a < 0$.


9.1.2 Operations related to $y_2 = f(x + a)$

Example 9.2: Operations on graphs related to $y_2 = f(x + a)$

Consider again $y = \ln x$. Then

$$y_2 = \ln(x + a).$$

Since $\ln x$ is defined for $x > 0$, $\ln(x + a)$ will be defined for $x + a > 0$, i.e., for $x > -a$. Since $\ln x = 0$ if $x = 1$, we have $\ln(x + a) = 0$ for $x + a = 1$, i.e., $x = 1 - a$.




- The graph of y_2 is obtained from that of y by a **translation** of a units to the **left** if $a > 0$, to the **right** if $a < 0$.


9.1.3 Operations related to $y_3 = k f(x)$

- The graph of $y_3 = k f(x)$ is obtained from that of f by multiplying all the ordinates $f(x)$ by k .
- In particular, if $k = -1$ the ordinates simply change sign, so that the graph of y_3 is symmetric to that of f with respect to the x -axis.
- Note that if $k > 1$, the graph is “**stretched**” in the **vertical direction**, dilating the **positive** ordinates **upward** and the **negative** ones **downward**.
- Conversely, if $0 < k < 1$ the graph “**shrinks**”, again in the vertical direction.
- Therefore, the operation of multiplying $f(x)$ by k has the geometric meaning of a **dilation** (if $|k| > 1$) or a **contraction** (if $|k| < 1$) along the y -axis, possibly accompanied by a **reflection with respect to the x -axis**, if $k < 0$.

Example 9.3: Operations on graphs related to $y_3 = k f(x)$

Consider $y = \sin x$, then $y_3 = k \sin(x)$.







9.1.4 Operations related to $y_4 = f(kx)$

- The graph of $y_4 = f(kx)$ is obtained from that of $f(x)$ by a **change of scale** along the x -axis.
- If $k > 1$, kx grows faster than x , and therefore the graph of y_4 will be similar to that of f but with faster oscillations, that is, it will be “**compressed**” in the **horizontal direction** by a factor $\frac{1}{k}$.
- Similarly, if $0 < k < 1$ the graph will appear “**dilated**” in the **horizontal direction**, with gentler oscillations.
- If $k < 0$, in addition to a compression (if $|k| > 1$) or a dilation (if $|k| < 1$) along the x -axis, there will be a **reflection with respect to the y -axis**.

Example 9.4: Operations on graphs related to $y_4 = f(kx)$


Consider $y = \sin x$, then $y_4 = \sin(kx)$. With $k = \frac{1}{2}$ and $k = 2$ we have:





Example 9.5: Operations on graphs related to $y_4 = f(kx)$

Consider $y = \ln x$, then $y_4 = \ln(kx)$. With $k = -1$ we have:



9.1.5 Operations related to $y_5 = |f(x)|$



Absolute value of $f(x)$:

$$|f(x)| = \begin{cases} f(x) & \text{if } f(x) \geq 0, \\ -f(x) & \text{if } f(x) < 0 \end{cases}$$


- When passing from the graph of $y = f(x)$ to that of $y_5 = |f(x)|$, the points with **non-negative ordinate remain unchanged**, while those with **negative ordinate are transformed into their symmetric points with respect to the x -axis**.
- The graph of $y_5 = |f(x)|$ is obtained from that of f by “flipping” symmetrically with respect to the x -axis the part of the graph of f that lies in the lower half-plane, and leaving the rest unchanged.

Example 9.6: Operations on graphs related to $y_5 = |f(x)|$

Consider $y = x$, then $y_5 = |x|$

**Example 9.7: Operations on graphs related to $y_5 = |f(x)|$**

Consider $y = \sin x$, then $y_5 = |\sin x|$





9.1.6 Operations related to $y_6 = f(|x|)$

- Finally, to draw the graph of $y_6 = f(|x|)$, we observe that $|x| = x$ for $x \geq 0$, and hence in the right half-plane the two graphs coincide.
- We have $|-x| = |x|$, and hence y_6 is an even function, therefore symmetric with respect to the y -axis.
- Consequently, the graph of $y_6 = f(|x|)$ is drawn by **leaving the graph of f unchanged in the right half-plane and flipping it symmetrically with respect to the y -axis.**

Example 9.8: Operations on graphs related to $y_6 = f(|x|)$

Consider $y = e^x$, then $y_6 = e^{|x|}$



Composite functions

10.1 Composite functions

Definition 10.1: of composite function


Given two functions:

$$f : E \rightarrow \mathbb{R} \quad \text{and} \quad g : F \rightarrow \mathbb{R},$$

if $f(E) \subseteq F$ (i.e., if for every $x \in E$ we have $f(x) \in F$), one can define the function $h : E \rightarrow \mathbb{R}$, the **composite** of f and g (in this order), denoted by the symbol $g \circ f$, through the formula

$$h(x) = (g \circ f)(x) = g[f(x)]$$

- The composition scheme is the following:




Example 10.1: of composite function

The function $x \mapsto |f(x)|$ is actually “composed” of two functions:

1. given x , we compute $f(x)$
2. once $f(x)$ is computed, we compute $|f(x)|$

This amounts to operating in series with two black boxes, the first corresponding to f , the second to its absolute value, according to the following scheme:



- It may happen that both compositions $(g \circ f)$ and $(f \circ g)$ are well defined, but in general

$$(f \circ g) \neq (g \circ f)$$

In other words, **the commutative property does not hold**.

Example 10.2: of composition of functions


Consider the functions:

$$f : \mathbb{R} \rightarrow \mathbb{R}, \quad f : x \mapsto x^2 \quad \text{and} \quad g : \mathbb{R} \rightarrow \mathbb{R}, \quad g : x \mapsto \cos x$$

that is, $f(x) = x^2$ and $g(x) = \cos x$.


1. Since g is defined on all of $\mathbb{R}$, $h = g \circ f$ is well defined on $\mathbb{R}$ and the following formula holds

$$h(x) = (g \circ f)(x) = g[f(x)] = \cos x^2$$



2. Since f is defined on all of $\mathbb{R}$, $k = f \circ g$ is well defined on $\mathbb{R}$ and the following formula holds

$$k(x) = (f \circ g)(x) = f[g(x)] = \cos^2 x$$



Example 10.3: of composition of functions


Consider the functions:

$$f : \mathbb{R} \rightarrow \mathbb{R}, \quad f : x \mapsto e^x \quad g : [0, +\infty) \rightarrow \mathbb{R}, \quad g : x \mapsto \sqrt{x}$$

that is, $f(x) = e^x$ and $g(x) = \sqrt{x}$.


1. Since g is defined on $[0, +\infty)$ and $f(x) > 0$ for every $x \in \mathbb{R}$, $h = g \circ f$ is well defined on $\mathbb{R}$ and the following formula holds

$$h(x) = (g \circ f)(x) = g[f(x)] = \sqrt{e^x}$$



2. Since g is defined only on $[0, +\infty)$, $k = f \circ g$ is well defined only on $\mathbb{R}_+$ and the following formula holds

$$k(x) = (f \circ g)(x) = f[g(x)] = e^{\sqrt{x}}$$



- The composition operation can be extended to three or more factors. One can verify that if the composition $(f \circ g) \circ r$ exists, then $f \circ (g \circ r)$ also exists and they are equal

$$(f \circ g) \circ r = f \circ (g \circ r)$$

In other words, **the associative property holds**.

- If a function $f : D \rightarrow \mathbb{R}$ is such that $f(D) \subseteq D$, then it can be composed with itself

$$f^2 = f \circ f \quad \text{that is} \quad f^2(x) = f[f(x)]$$

f^2 is called the **second iterate** of f . Similarly, f^n is called the **n -th iterate function** of f .

Inverse functions

11.1 Invertible functions and inverse functions

- Given a real function of a real variable $f : D \rightarrow \mathbb{R}$, for every input x in the domain D there exists a unique output $y = f(x)$ in the image of the domain $f(D)$.
- If, in addition, for every output $y = f(x) \in f(D)$ there exists a unique input $x \in D$, then f is called **invertible**, and it establishes a **one-to-one correspondence** between the domain D of f and the image of the domain $f(D)$.

Definition 11.1: of invertible function

A function $f : D \rightarrow \mathbb{R}$ is **invertible** on the domain D if one of the following equivalent conditions holds:

$$\forall x_1, x_2 \in D, \quad x_1 \neq x_2 \implies f(x_1) \neq f(x_2)$$

$$\forall x_1, x_2 \in D, \quad f(x_1) = f(x_2) \implies x_1 = x_2$$

$$\forall y \in f(D), \quad \exists! x \in D : y = f(x)$$

that is, if f is **injective**.

WARNING: the definition of invertible function given here for real functions of a real variable is different from the usual one given for functions between arbitrary sets, which also requires surjectivity. In this case *injectivity is enough* because we define the inverse on the image of f and not on its whole codomain.


Definition 11.2: of inverse function

Given an invertible function $f : D \rightarrow \mathbb{R}$, the function that associates to each output $y \in f(D)$ the unique input $x \in D$ such that $f(x) = y$ is called the **inverse function** of f and is denoted by the symbol f^{-1} .

The pair f and f^{-1} is written as:


$$f : \begin{cases} y = f(x) \\ x \in D \end{cases} \quad f^{-1} : \begin{cases} x = f^{-1}(y) \\ y \in f(D) \end{cases} \quad (11.1)$$

The black box of f^{-1} works backwards with respect to that of f , according to the following scheme:




The invertibility condition is equivalent to requiring that the graph of f be intersected in at most one point by every line parallel to the x -axis.

Example 11.1: graph of an invertible function



Graph of a function that is **invertible** on $[a, b]$, since every line parallel to the x -axis either does not intersect the graph of f or intersects it in exactly one point.

Example 11.2: graph of a non-invertible function





Graph of a function that is **not invertible** on $[a, b]$, since for the indicated value $\tilde{y}$ there are several values x (precisely x_1 , x_2 and x_3) whose image is $\tilde{y}$.

Theorem 11.3

If a function $f : D \rightarrow \mathbb{R}$ is strictly increasing (decreasing) on D , then it is invertible on D . Moreover, its inverse function is strictly increasing (decreasing).

For the strictly increasing case, graphically we have:





Proof. Consider the case of a function that is strictly increasing on D and two arbitrary values $x_1, x_2 \in D$ (the strictly decreasing case is analogous).

If $x_1 \neq x_2$, then either $x_1 < x_2$ or $x_1 > x_2$. Since f is strictly increasing, we have:

$$\text{if } x_1 < x_2, \text{ then } f(x_1) < f(x_2)$$

$$\text{if } x_1 > x_2, \text{ then } f(x_1) > f(x_2)$$

In both cases $f(x_1) \neq f(x_2)$, therefore f is invertible.

Consider the inverse function f^{-1} , that is, $x = f^{-1}(y)$, and let us prove that it is strictly increasing. Consider two arbitrary values $y_1, y_2 \in f(D)$, with $y_1 < y_2$.

If we had $f^{-1}(y_1) = x_1 \geq x_2 = f^{-1}(y_2)$, since f is strictly increasing we would have $y_1 = f(x_1) \geq f(x_2) = y_2$, that is, $y_1 \geq y_2$. This cannot happen, since it contradicts $y_1 < y_2$ (**contradiction**).


Consequently, we have $f^{-1}(y_1) = x_1 < x_2 = f^{-1}(y_2)$, that is, $f^{-1}(y_1) < f^{-1}(y_2)$, and hence f^{-1} is strictly increasing. □

However, a function can be invertible even without being strictly increasing or decreasing.

Example 11.3: invertible function that is neither strictly increasing nor strictly decreasing

For example, the following piecewise-defined function is invertible:

$$f(x) = \begin{cases} \frac{1}{2}x & \text{if } -1 \leq x < 0, \\ \frac{1}{2} - \frac{1}{2}x & \text{if } 0 \leq x \leq \frac{2}{3} \end{cases}$$



- Two classes of functions that are certainly not invertible are:

1. even (symmetric) functions, since:

$$f(-x) = f(x)$$

2. periodic functions, since:

$$f(x + T) = f(x)$$


11.1.1 Graph of the inverse function

- The relations between an invertible function f and its inverse function f^{-1} :

$$f : \begin{cases} y = f(x) \\ x \in D \end{cases} \quad f^{-1} : \begin{cases} x = f^{-1}(y) \\ y \in f(D) \end{cases}$$

indicate that if the point (x_0, y_0) lies on the graph of f , then the point (y_0, x_0) lies on the graph of f^{-1} .

- Since the points (x_0, y_0) and (y_0, x_0) are **symmetric with respect to the bisector** with equation $y = x$, the graph of f^{-1} is obtained from that of f by symmetry with respect to the bisector.



If the analytic expression of f is known and f is invertible, the **analytic expression** of f^{-1} is found by trying to solve for x the equation:

$$f(x) = y \quad \text{that is, find the input } x \text{ that produces the output } y$$

Note that it is not always possible to find it, even in cases where the inverse function exists!

Given $a, b \in \mathbb{R}, a \neq 0$, we have:

$$f : \begin{cases} y = ax + b \\ x \in \mathbb{R} \end{cases} \quad f^{-1} : \begin{cases} x = \frac{y-b}{a} \\ y \in \mathbb{R} \end{cases}$$

Example 11.4: inverse function of an affine function


The function $f : \mathbb{R} \rightarrow \mathbb{R}, x \mapsto 2x + 3$ is strictly increasing, hence invertible on $\mathbb{R}$, and the equation

$$2x + 3 = y \text{ solved for } x, \quad \text{that is } x = \frac{y-3}{2}$$


gives for f^{-1} the analytic expression:

$$f^{-1}(y) = \frac{y-3}{2}$$

that is, the function: $f^{-1} : \mathbb{R} \rightarrow \mathbb{R}, y \mapsto \frac{y-3}{2}$.



The graphs $y = 2x + 3$ and $y = \frac{x-3}{2}$ are symmetric with respect to the bisector $y = x$



We have:

$$f : \begin{cases} y = x^2 \\ x \geq 0 \end{cases} \quad f^{-1} : \begin{cases} x = \sqrt{y} \\ y \geq 0 \end{cases}$$

Example 11.5: inverse function of $f(x) = x^2$

The function $f : \mathbb{R} \rightarrow \mathbb{R}, x \mapsto x^2$ is strictly increasing on the interval $[0, +\infty)$, hence

invertible there; the equation


$$x^2 = y \text{ solved for } x, \text{ that is } x = \sqrt{y}$$

gives for f^{-1} the analytic expression

$$f^{-1}(y) = \sqrt{y}$$

that is, the function: $f^{-1} : [0, +\infty) \rightarrow \mathbb{R}, y \mapsto \sqrt{y}$.

The graphs $y = x^2$ and $y = \sqrt{x}$ are symmetric with respect to the bisector $y = x$



Given $a \in \mathbb{R}_+, a \neq 1$, we have:

$$f : \begin{cases} y = a^x \\ x \in \mathbb{R} \end{cases} \quad f^{-1} : \begin{cases} x = \log_a y \\ y > 0 \end{cases}$$

Example 11.6: inverse function of $f(x) = e^x$

The function $f : \mathbb{R} \rightarrow \mathbb{R}, x \mapsto e^x$ is strictly increasing on all of $\mathbb{R}$ and hence invertible; the equation


$$e^x = y \text{ solved for } x, \text{ that is } x = \log y$$

gives for f^{-1} the analytic expression

$$f^{-1}(y) = \log y$$

that is, the function: $f^{-1} : (0, +\infty) \rightarrow \mathbb{R}, y \mapsto \log y$.

The graphs $y = e^x$ and $y = \log x$ are symmetric with respect to the bisector $y = x$



- Now consider the function:

$$f(x) = x + e^x$$

Being the sum of two functions that are strictly increasing on all of $\mathbb{R}$, $f(x)$ is strictly increasing and hence invertible on all of $\mathbb{R}$ (see Theorem (11.3)). However, we would try in vain to solve the equation $x + e^x = y$ for x . In other words, f^{-1} exists, but we cannot write it explicitly.

11.1.2 Inverse power functions

Given $\alpha \in \mathbb{R}, \alpha \neq 0$, we have:

$$f : \begin{cases} y = x^\alpha \\ x > 0 \end{cases} \quad f^{-1} : \begin{cases} x = y^{\frac{1}{\alpha}} \\ y > 0 \end{cases}$$

If $\alpha > 0$ we have:

$$f : \begin{cases} y = x^\alpha \\ x \geq 0 \end{cases} \quad f^{-1} : \begin{cases} x = y^{\frac{1}{\alpha}} \\ y \geq 0 \end{cases}$$


If $\alpha = \frac{m}{n} \in \mathbb{Q}$, with $m \in \mathbb{Z}, n \in \mathbb{N}_+$ odd and coprime, we have:

$$f : \begin{cases} y = x^{\frac{m}{n}} \\ x \in \mathbb{R} \end{cases} \quad f^{-1} : \begin{cases} x = y^{\frac{n}{m}} \\ y \in \mathbb{R} \end{cases}$$

- The **even powers**:

$$x^{2n} \quad \text{with } n = 1, 2, \dots$$

are not invertible on the whole real line, but only on the half-line $x \geq 0$.




- The **odd powers**:

$$x^{2n+1} \quad \text{with } n = 1, 2, \dots$$

and the **powers with rational exponent**

$$x^{\frac{m}{n}} \quad \text{with } n, m \text{ positive odd integers}$$

being strictly increasing monotone functions, are invertible from $-\infty$ to $+\infty$.



11.1.3 Inverse trigonometric functions

- Being periodic, the trigonometric functions cannot be invertible. Indeed, for example, the equation

$$y = \sin x$$

has infinitely many solutions if $-1 \leq y \leq 1$ (the output y corresponds to infinitely many inputs), or has no real solutions if $|y| > 1$.

- To speak of inverse functions of sine, cosine and tangent, we will need to restrict ourselves to intervals on which these functions are strictly monotone and therefore invertible.
- An interval on which the sine function is invertible is $[-\frac{\pi}{2}, \frac{\pi}{2}]$.

Definition 11.4: of arcsine


The inverse function of the sine on the interval $[-\frac{\pi}{2}, \frac{\pi}{2}]$ is the **arcsine**:

$$f : [-1, 1] \rightarrow \left[-\frac{\pi}{2}, \frac{\pi}{2}\right], \quad f : y \mapsto \arcsin y$$

We have:

$$f : \begin{cases} y = \sin x \\ x \in [-\frac{\pi}{2}, \frac{\pi}{2}] \end{cases} \quad f^{-1} : \begin{cases} x = \arcsin y \\ y \in [-1, 1] \end{cases} \quad (11.2)$$

- The graph of the arcsine is obtained from the arc of the sine curve restricted to the interval $[-\frac{\pi}{2}, \frac{\pi}{2}]$ (strictly monotone), by symmetry with respect to the bisector $y = x$,



- An interval on which the cosine function is invertible is $[0, \pi]$.

Definition 11.5: of arccosine


The inverse function of the cosine on the interval $[0, \pi]$ is the **arccosine**:

$$f : [-1, 1] \rightarrow [0, \pi], \quad f : y \mapsto \arccos y$$

We have:

$$f : \begin{cases} y = \cos x \\ x \in [0, \pi] \end{cases} \quad f^{-1} : \begin{cases} x = \arccos y \\ y \in [-1, 1] \end{cases} \quad (11.3)$$

- The graph of the arccosine is obtained from the arc of the cosine curve restricted to the interval $[0, \pi]$ (strictly monotone), by symmetry with respect to the bisector $y = x$,



- Note that on the interval $(-\frac{\pi}{2}, \frac{\pi}{2})$ the tangent is strictly monotone and hence invertible. Its inverse function is called the arctangent ($\arctan$), and it is defined on $\mathbb{R}$.

Definition 11.6: of arctangent


The inverse function of the tangent on the interval $(-\frac{\pi}{2}, \frac{\pi}{2})$ is the **arctangent**:

$$f : \mathbb{R} \rightarrow \left(-\frac{\pi}{2}, \frac{\pi}{2}\right), \quad f : y \mapsto \arctan y$$

We have:

$$f : \begin{cases} y = \tan x \\ x \in (-\frac{\pi}{2}, \frac{\pi}{2}) \end{cases} \quad f^{-1} : \begin{cases} x = \arctan y \\ y \in \mathbb{R} \end{cases} \quad (11.4)$$

- The graph of the arctangent is obtained from that of the tangent restricted to the interval $(-\frac{\pi}{2}, \frac{\pi}{2})$ (strictly monotone), by symmetry with respect to the bisector $y = x$,



- By means of the inverse trigonometric functions, we can express the solutions of a **trigonometric equation or inequality** when it involves angles that are not standard angles.


Example 11.7: Trigonometric equations/inequalities

- The solutions of:

$$\sin x = \frac{1}{4}$$

are:

$$(i) \quad x = \arcsin \frac{1}{4} + 2k\pi; \quad (ii) \quad x = \pi - \arcsin \frac{1}{4} + 2k\pi \quad (k \in \mathbb{Z})$$




- The solutions of:

$$\cos x < \frac{1}{5}$$

are:

$$\arccos \frac{1}{5} + 2k\pi < x < 2\pi - \arccos \frac{1}{5} + 2k\pi \quad (k \in \mathbb{Z})$$



- The solutions of:

$$\tan x \geq 3$$

are:

$$\arctan 3 + k\pi \leq x < \frac{\pi}{2} + k\pi \quad (k \in \mathbb{Z})$$

11.1.4 Inverse hyperbolic functions

- Consider the **hyperbolic sine** function:

$$y = \text{Sh } x = \frac{e^x - e^{-x}}{2}$$

It is defined and strictly increasing on all of $\mathbb{R}$, therefore it is invertible.

- To solve the equation for x , multiplying both sides by e^x we obtain:

$$\begin{aligned} 0 &= e^x y - e^x \frac{e^x - e^{-x}}{2} \\ &= 2e^x y - (e^x e^x - \underbrace{e^x e^{-x}}_{=e^{x-x}=e^0}) \\ &= 2y e^x - e^{2x} + 1 \\ &= e^{2x} - 2y e^x - 1 \end{aligned}$$

which is a quadratic equation in the unknown e^x . Setting $t = e^x$ we obtain

$$0 = t^2 - 2y t - 1$$

We obtain:

$$t = y \pm \sqrt{y^2 + 1} \quad \text{and substituting back} \quad e^x = y \pm \sqrt{y^2 + 1}$$

since $e^x > 0$, the solution with the minus sign must be discarded. Hence we are left with:

$$e^x = \underbrace{y + \sqrt{y^2 + 1}}_{>0, \forall y \in \mathbb{R}} \quad \text{and hence} \quad x = \log \left(y + \sqrt{y^2 + 1} \right)$$

- This is the analytic expression of the inverse function of $\text{Sh } x$, which is called the **inverse hyperbolic sine** (area hyperbolic sine), and is also denoted by SettSh . It is defined for every real y .

Definition 11.7: of inverse hyperbolic sine


The inverse function of the hyperbolic sine is the **inverse hyperbolic sine**:

$$f : \mathbb{R} \rightarrow \mathbb{R}, \quad f : y \mapsto \text{SettSh } y$$

We have:

$$f : \begin{cases} y = \text{Sh } x \\ x \in \mathbb{R} \end{cases} \quad f^{-1} : \begin{cases} x = \text{SettSh } y \\ y \in \mathbb{R} \end{cases} \quad (11.5)$$

- The graph of the inverse hyperbolic sine is obtained from that of the hyperbolic sine on $\mathbb{R}$ (strictly monotone), by symmetry with respect to the bisector $y = x$



- Consider the **hyperbolic cosine** function:

$$y = \text{Ch } x = \frac{e^x + e^{-x}}{2}$$

It is defined on all of $\mathbb{R}$, strictly increasing for $x \geq 0$, decreasing for $x \leq 0$. Therefore it is not invertible on all of $\mathbb{R}$.

- Its restriction to $x \geq 0$, however, is invertible. We want to determine the analytic expression of the inverse function of this restriction.
- Proceeding as before, we obtain:

$$e^x = y \pm \sqrt{y^2 - 1}$$

- This time both numbers $y \pm \sqrt{y^2 - 1}$ are positive; recall, however, that we are reasoning only for $x \geq 0$, which is equivalent to choosing the plus sign. Therefore:

$$x = \log \left(y + \sqrt{y^2 - 1} \right)$$

This is the analytic expression of the inverse function of $\text{Ch } x$, which is called the inverse hyperbolic cosine (area hyperbolic cosine), and is also denoted by SettCh . Note that it is defined for $y \geq 1$.

Definition 11.8: of inverse hyperbolic cosine


The inverse function of the hyperbolic cosine on the interval $[0, +\infty)$ is the **inverse hyperbolic cosine**:

$$f : [1, +\infty) \rightarrow [0, +\infty), \quad f : y \mapsto \text{SettCh } y$$

We have:

$$f : \begin{cases} y = \text{Ch } x \\ x \in [0, +\infty) \end{cases} \quad f^{-1} : \begin{cases} x = \text{SettCh } y \\ y \in [1, +\infty) \end{cases}$$

- The graph of the inverse hyperbolic cosine is obtained from that of the hyperbolic cosine on $[0, +\infty)$ (strictly monotone), by symmetry with respect to the bisector $y = x$.



Example 11.8: Hyperbolic equations

- The equation:

$$\operatorname{Sh} x = 2$$

has the unique solution:


$$x = \operatorname{SettSh} 2 = \log(2 + \sqrt{5})$$

- The equation:

$$\operatorname{Ch} x = 3$$

has two solutions:

$$x = \pm \operatorname{SettCh} 3 = \pm \log(3 + 2\sqrt{2})$$



Exercises

The sheets of worked exercises on the topics of this volume, one per section, with the solution right after each statement. The sheets are by Fabio Furini and Gianluca Priori; many exercises are taken from M. Bramanti, *Esercitazioni di Analisi matematica 1*, Esculapio.

12.1 Domains of functions

Exercise 12.1

Determine the domain of the function:

$$f(x) = \frac{\log_2(3+x)}{\sqrt[3]{x+2}}$$

it is determined by studying the numerator and the denominator of the function

- The numerator:

The function $\log_2 t$ is defined for $t > 0$, hence the function $\log_2(3+x)$ is defined for $3+x > 0$, i.e., $x > -3$.

- The denominator:

The function $\sqrt[3]{t}$ is always defined; however, since it is in the denominator, it must not vanish. Hence we impose $\sqrt[3]{x+2} \neq 0$, i.e., $x \neq -2$.

Hence the domain is

$$(-3, -2) \cup (-2, +\infty)$$

Exercise 12.2

Determine the domain of the function:

$$f(x) = \sqrt{2^x - 8} \cdot \log|x-4|$$

Exercise 12.3

Determine the domain of the function:

$$f(x) = \tan(\log x)$$

Exercise 12.4

Determine the domain of the function:

$$f(x) = \log(\tan x)$$

12.2 Inverse functions

A function $f : D \rightarrow \mathbb{R}$ is **invertible** on the domain D if one of the following equivalent conditions holds:

$$\forall x_1, x_2 \in D, \quad x_1 \neq x_2 \implies f(x_1) \neq f(x_2)$$

$$\forall x_1, x_2 \in D, \quad f(x_1) = f(x_2) \implies x_1 = x_2$$

$$\forall y \in f(D), \quad \exists! x \in D : y = f(x)$$

that is, if f is **injective**. In the following exercises, the second condition will mostly be used to verify the injectivity of a function f .

Given an invertible function $f : D \rightarrow \mathbb{R}$, the function that assigns to each output $y \in f(D)$ the unique input $x \in D$ such that $f(x) = y$ is called the **inverse function** of f and is denoted by the symbol f^{-1} .

Recall that the invertibility condition is equivalent to requiring that the graph of f be intersected in at most one point by every line parallel to the x -axis.

If a function $f : D \rightarrow \mathbb{R}$ is strictly increasing (decreasing) on D , then it is invertible on D . Moreover, its inverse function is strictly increasing (decreasing).

In conclusion, to establish whether or not a function $f : D \rightarrow \mathbb{R}$ is invertible on D , it suffices to verify any one of the following three conditions:

- f is **injective**;
- f is bijective, since every bijective function is both injective and surjective;
- f is strictly increasing (or decreasing) on D , since this is a *sufficient* condition for injectivity.

The relations between an invertible function f and its inverse function f^{-1} :

$$f : \begin{cases} y = f(x) \\ x \in D \end{cases} \quad f^{-1} : \begin{cases} x = f^{-1}(y) \\ y \in f(D) \end{cases}$$

show that if the point (x_0, y_0) is on the graph of f , then the point (y_0, x_0) is on the graph of f^{-1} . Since the points (x_0, y_0) and (y_0, x_0) are **symmetric with respect to the bisector** of equation $y = x$, the graph of f^{-1} is obtained from that of f by reflection across the bisector.

Exercise 12.5

Prove that the function

$$f(x) = \log(2 + 3x)$$

is invertible on its domain and determine the analytic expression of the inverse f^{-1} . Specify the domain of f^{-1} .

Solution

The domain of the given function is

$$\left(-\frac{2}{3}, +\infty\right),$$

the interval where the argument of the logarithm is strictly positive. To prove that f is invertible, it suffices to verify that it is injective, i.e.,

$$\forall x_1, x_2 \in D \quad f(x_1) = f(x_2) \implies x_1 = x_2$$

Imposing $f(x_1) = f(x_2)$ is equivalent to imposing that

$$\log(2 + 3x_1) = \log(2 + 3x_2)$$

$$2 + 3x_1 = 2 + 3x_2$$

$$3x_1 = 3x_2 \iff x_1 = x_2$$

The function f is injective and, consequently, invertible. We look for the analytic expression of the inverse function f^{-1} by solving, with respect to the variable x , the equation

$$\log(2 + 3x) = y.$$

which has the unique solution

$$x = \frac{1}{3}(e^y - 2)$$


Solution

More formally, the inverse function of f is

$$f^{-1}(y) = \frac{1}{3}(e^y - 2)$$

whose domain is the image of the original function f , that is, $\mathbb{R}$. Hence $f : (-\frac{2}{3}, +\infty) \rightarrow (-\infty, +\infty)$, while $f^{-1} : (-\infty, +\infty) \rightarrow (-\frac{2}{3}, +\infty)$.

The graphs $y = \log(2 + 3x)$ and $y = \frac{1}{3}(e^x - 2)$ are symmetric with respect to the bisector $y = x$:



Exercise 12.6

Prove that the function

$$f(x) = \sqrt{1 - 2x}$$

is invertible on its domain and determine the analytic expression of the inverse f^{-1} . Specify the domain of f^{-1} .

Solution

The domain of the given function is

$$\left(-\infty, \frac{1}{2}\right],$$

the interval where the radicand is non-negative.

Solution

To prove that f is invertible, it suffices to verify that it is injective, i.e.,

$$\forall x_1, x_2 \in D \quad f(x_1) = f(x_2) \implies x_1 = x_2$$

Imposing $f(x_1) = f(x_2)$ is equivalent to imposing that

$$\begin{aligned}\sqrt{1-2x_1} &= \sqrt{1-2x_2} \\ 1-2x_1 &= 1-2x_2 \\ -2x_1 &= -2x_2 \iff x_1 = x_2\end{aligned}$$

The function f is injective and, consequently, invertible. We look for the analytic expression of the inverse function f^{-1} by solving, with respect to the variable x , the equation

$$\sqrt{1-2x} = y.$$

which, for $y \geq 0$ (a square root is never negative in $\mathbb{R}$), has the unique solution


$$x = -\frac{1}{2}y^2 + \frac{1}{2}$$

More formally, the inverse function of f is

$$f^{-1}(y) = -\frac{1}{2}y^2 + \frac{1}{2}$$

whose domain is the image of the original function f , that is, $[0, +\infty)$. Hence $f : (-\infty, \frac{1}{2}] \rightarrow [0, +\infty)$, $f^{-1} : [0, +\infty) \rightarrow (-\infty, \frac{1}{2}]$.

The graphs $y = \sqrt{1-2x}$ and $y = -\frac{1}{2}x^2 + \frac{1}{2}$ are symmetric with respect to the bisector $y = x$:



Exercise 12.7

Prove that the function

$$f(x) = e^{2x-3}$$

is invertible on its domain and determine the analytic expression of the inverse f^{-1} . Specify the domain of f^{-1} .

Solution

The domain of the given function is

$$(-\infty, +\infty),$$

since the exponential function is defined for every $x \in \mathbb{R}$. To prove that f is invertible, it suffices to verify that it is injective, i.e.,

$$\forall x_1, x_2 \in D \quad f(x_1) = f(x_2) \implies x_1 = x_2$$

Imposing $f(x_1) = f(x_2)$ is equivalent to imposing that

$$\begin{aligned} e^{2x_1-3} &= e^{2x_2-3} \\ 2x_1 - 3 &= 2x_2 - 3 \\ 2x_1 &= 2x_2 \iff x_1 = x_2 \end{aligned}$$

The function f is injective and, consequently, invertible. We look for the analytic expression of the inverse function f^{-1} by solving, with respect to the variable x , the equation

$$e^{2x-3} = y.$$

It makes sense to solve this equation only for $y > 0$, since it would have no solutions when $y \leq 0$ (the exponential never vanishes and is never negative). The equation has the unique solution

$$x = \frac{3}{2} + \frac{1}{2} \log y$$

More formally, the inverse function of f is

$$f^{-1}(y) = \frac{3}{2} + \frac{1}{2} \log y$$

whose domain is the image of the original function f , that is, $(0, +\infty)$. Hence $f : (-\infty, +\infty) \rightarrow (0, +\infty)$, $f^{-1} : (0, +\infty) \rightarrow (-\infty, +\infty)$.

Exercise 12.8

Prove that the function

$$f(x) = \arctan(2x - 1)$$

is invertible on its domain and determine the analytic expression of the inverse f^{-1} . Specify the domain of f^{-1} .

Solution

The arctangent function, which we recall is the inverse function of the restriction of $y = \tan(x)$ to the interval $(-\frac{\pi}{2}, \frac{\pi}{2})$, has the following properties:

- its *domain* is the set $\mathbb{R}$;
- its image is the interval $(-\frac{\pi}{2}, \frac{\pi}{2})$;
- it is a **strictly increasing** monotonic function.

Since it is strictly increasing, the function f is invertible (note that this is only a *sufficient* condition). We look for the analytic expression of the inverse function f^{-1} by solving, with respect to the variable x , the equation

$$\arctan(2x - 1) = y,$$

where $y \in (-\frac{\pi}{2}, \frac{\pi}{2})$. The equation becomes

$$\begin{aligned} 2x - 1 &= \tan(y) \\ x &= \frac{1}{2} + \frac{\tan(y)}{2} \end{aligned}$$

More formally, the inverse function of f is

$$f^{-1}(y) = \frac{1}{2} + \frac{\tan(y)}{2}$$

whose domain is the image of the original function f , that is, $(-\frac{\pi}{2}, \frac{\pi}{2})$. Hence $f : \mathbb{R} \rightarrow (-\frac{\pi}{2}, \frac{\pi}{2})$, while $f^{-1} : (-\frac{\pi}{2}, \frac{\pi}{2}) \rightarrow \mathbb{R}$

Exercise 12.9

Prove that the function

$$f(x) = \frac{x-1}{x+2}$$

is invertible on its domain and determine the analytic expression of the inverse f^{-1} . Specify the domain of f^{-1} .

Solution

The domain of the given function is

$$(-\infty, -2) \cup (-2, +\infty),$$

since it is defined $\forall x \in \mathbb{R}, x \neq -2$. To prove that f is invertible, it suffices to verify that it is injective, i.e.,

$$\forall x_1, x_2 \in D \quad f(x_1) = f(x_2) \implies x_1 = x_2$$

To simplify the subsequent computations, it is convenient to rewrite the function f as:

$$f(x) = \frac{x-1}{x+2} = \frac{x+2-3}{x+2} = 1 - \frac{3}{x+2}$$

Imposing $f(x_1) = f(x_2)$ is equivalent to imposing that

$$\begin{aligned} 1 - \frac{3}{x_1 + 2} &= 1 - \frac{3}{x_2 + 2} \\ \frac{3}{x_1 + 2} &= \frac{3}{x_2 + 2} \\ x_1 + 2 &= x_2 + 2 \iff x_1 = x_2 \end{aligned}$$

The function f is injective and, consequently, invertible. We look for the analytic expression of the inverse function f^{-1} by solving, with respect to the variable x , the equation

$$\frac{x - 1}{x + 2} = y.$$

Rewriting the function again as before

$$1 - \frac{3}{x + 2} = y$$

we obtain

$$x = \frac{3}{1 - y} - 2 = \frac{2y + 1}{1 - y}$$

More formally, the inverse function of f is

$$f^{-1}(y) = \frac{2y + 1}{1 - y}$$

whose domain is the image of the original function f , that is, $(-\infty, 1) \cup (1, +\infty)$. Hence $f : (-\infty, -2) \cup (-2, +\infty) \rightarrow (-\infty, 1) \cup (1, +\infty)$, $f^{-1} : (-\infty, 1) \cup (1, +\infty) \rightarrow (-\infty, -2) \cup (-2, +\infty)$.

Exercise 12.10

Prove that the function

$$f(x) = e^{1+x^2}$$

is invertible on a restriction of its domain and determine the analytic expression of the inverse f^{-1} . Specify the domain of f^{-1} .

Solution

The domain of the given function is

$$(-\infty, +\infty),$$

since the exponential function is defined for every $x \in \mathbb{R}$. It is clear that f is not invertible on all of $\mathbb{R}$, since it is an even function, i.e., symmetric with respect to the y -axis:

$$f(-x) = e^{1+(-x)^2} = e^{1+x^2} = f(x)$$

This implies that any horizontal line intersects the graph of the function in two points. The function is not injective, hence not invertible on $\mathbb{R}$. However, we can restrict the domain

by considering only the positive x -half-axis including the origin, i.e., the set

$$[0, +\infty)$$

Let us verify that, on this restricted domain, the function f is injective. To this end, we impose $f(x_1) = f(x_2)$, i.e.,

$$\begin{aligned} e^{1+x_1^2} &= e^{1+x_2^2} \\ 1 + x_1^2 &= 1 + x_2^2 \\ x_1^2 &= x_2^2 \iff x_1 = x_2 \end{aligned}$$

where the last equivalence holds since $x \in [0, +\infty)$, i.e., $x \geq 0$, so the solution $x_1 = -x_2$ is excluded. We have thus proved that the function f is injective on the restricted domain and, consequently, invertible on $[0, +\infty)$.

We look for the analytic expression of the inverse function f^{-1} by solving, with respect to the variable x , the equation

$$e^{1+x^2} = y.$$

The equation is solved by

$$x = \pm \sqrt{\log(y) - 1}$$

where, however, it is now clear that the negative solution must be excluded, since x varies in the restriction $[0, +\infty)$.

Solution

More formally, the inverse function of f is

$$f^{-1}(y) = \sqrt{\log(y) - 1}$$

whose domain is the image of the original function f , that is, $[e, +\infty)$. Hence $f : [0, +\infty) \rightarrow [e, +\infty)$, $f^{-1} : [e, +\infty) \rightarrow [0, +\infty)$.

Exercise 12.11

Prove that the function

$$f(x) = \frac{4}{5^{3x}}$$

is invertible on its domain and determine the analytic expression of the inverse f^{-1} . Specify the domain of f^{-1} .

Solution

The domain of the given function is

$$(-\infty, +\infty),$$

since the denominator is always nonzero (the exponential never vanishes). To prove that f is invertible, it suffices to verify that it is injective, i.e.,

$$\forall x_1, x_2 \in D \quad f(x_1) = f(x_2) \implies x_1 = x_2$$

Imposing $f(x_1) = f(x_2)$ is equivalent to imposing that

$$\begin{aligned}\frac{4}{5^{3x_1}} &= \frac{4}{5^{3x_2}} \\ 5^{3x_1} &= 5^{3x_2} \\ 3x_1 &= 3x_2 \iff x_1 = x_2\end{aligned}$$

The function f is injective and, consequently, invertible. We look for the analytic expression of the inverse function f^{-1} by solving, with respect to the variable x , the equation

$$\frac{4}{5^{3x}} = y,$$

with $y > 0$, which has the unique solution

$$x = \frac{1}{3} \log_5 \left(\frac{4}{y} \right) = -\frac{1}{3} \log_5 \left(\frac{y}{4} \right)$$

Solution

More formally, the inverse function of f is

$$f^{-1}(y) = -\frac{1}{3} \log_5 \left(\frac{y}{4} \right)$$

whose domain is the image of the original function f , that is, $(0, +\infty)$. Hence $f : \mathbb{R} \rightarrow (0, +\infty)$, while $f^{-1} : (0, +\infty) \rightarrow \mathbb{R}$.

Exercise 12.12

Prove that the function

$$f(x) = -1 + \sqrt[5]{1+x}$$

is invertible on its domain and determine the analytic expression of the inverse f^{-1} . Specify the domain of f^{-1} .

Solution

The domain of the given function is

$$(-\infty, +\infty),$$

since roots of odd index are always defined. To prove that f is invertible, it suffices to verify that it is injective, i.e.,

$$\forall x_1, x_2 \in D \quad f(x_1) = f(x_2) \implies x_1 = x_2$$

Imposing $f(x_1) = f(x_2)$ is equivalent to imposing that

$$\begin{aligned}-1 + \sqrt[5]{1+x_1} &= -1 + \sqrt[5]{1+x_2} \\ \sqrt[5]{1+x_1} &= \sqrt[5]{1+x_2} \\ 1+x_1 &= 1+x_2 \iff x_1 = x_2\end{aligned}$$

The function f is injective (it is also strictly increasing) and, consequently, invertible. We look for the analytic expression of the inverse function f^{-1} by solving, with respect to the variable x , the equation

$$-1 + \sqrt[5]{1+x} = y,$$

which has the unique solution

$$x = -1 + (y+1)^5$$

Solution

More formally, the inverse function of f is

$$f^{-1}(y) = -1 + (y+1)^5$$

whose domain is the image of the original function f , that is, $\mathbb{R}$. Hence $f : \mathbb{R} \rightarrow \mathbb{R}$, and also $f^{-1} : \mathbb{R} \rightarrow \mathbb{R}$.

Exercise 12.13

Prove that the function

$$f(x) = \frac{1-3x}{x+1}$$

is invertible on its domain and determine the analytic expression of the inverse f^{-1} . Specify the domain of f^{-1} .

(Hint: perform the division of $1-3x$ by $x+1$).

Exercise 12.14

Prove that the function

$$f(x) = 3^{x^2}$$

is invertible on a restriction of its domain and determine the analytic expression of the inverse f^{-1} on the restriction. Specify the domain of f^{-1} .

Exercise 12.15

Prove that the function

$$f(x) = \sqrt{\log\left(\frac{x-1}{x}\right)}$$

is invertible on its domain and determine the analytic expression of the inverse f^{-1} . Specify the domain of f^{-1} .