

Basics of logic and proof techniques



Outline

Logical symbols

Universal implications and proofs

Counterexamples

Law of contraposition

Sufficient conditions and necessary conditions

Proofs by contradiction

Logic and sets

Logical implication and set inclusion

Equality of sets and universal implications

Set operations and logical operations

Logical symbols

Introduction: Logical symbols

- ▶ In this section we introduce the **logical symbols** that we will use throughout the course
- ▶ Quantifiers, implication, membership, conjunction, disjunction and negation
- ▶ These symbols allow us to write definitions and theorems in a precise and compact way

Quantifiers and implication

- a) the symbol “ $\forall$ ” is called the *universal quantifier* and is read “for every”, “for all”, “for each”
- b) the symbol “ $\exists$ ” is called the *existential quantifier* and is read “there exists”, “there exist”
- c) the symbol “ $\Rightarrow$ ” is called *logical implication* and is read “implies” or “if . . . then”

Other symbols

- a) the symbol “ $:$ ” is read “such that”
 - b) the symbol “ $\in$ ” is read “belongs to”
 - c) the symbol “ $\notin$ ” is read “does not belong to”
-
- a) the symbol “ $\vee$ ” is called *logical disjunction* and is read “or”
 - b) the symbol “ $\wedge$ ” is called *logical conjunction* and is read “and”
 - c) the symbol “ $\neg$ ” is called *logical negation* and is read “not”.

Summary

- ▶ $\forall$: for every; $\exists$: there exists; $\Rightarrow$: implies
- ▶ $:$ such that; $\in$ belongs to; $\notin$ does not belong to
- ▶ $\vee$: or (disjunction); $\wedge$: and (conjunction); $\neg$: not (negation)

Universal implications and proofs



Introduction: Universal implications and proofs

- ▶ In this section we distinguish **predicates** from **propositions**
- ▶ We define **universal implications**, the logical structure of most theorems
- ▶ We will see how to prove a universal implication and how to disprove it with a **counterexample**

Predicates

Predicates (or properties) and **propositions** (or statements)

- Consider the following assertion:

“the natural number n is odd” (1)

and let us ask whether it is true. Obviously the answer is: “it depends on n ”. Indeed, in (1) the symbol n represents a variable that can take different values and make the assertion **true** or **false**.

A sentence of this kind is called a **predicate** (or *property*): its truth or falsity depends on the values of the **variable(s)** appearing in it.

Propositions

- Now consider the following assertion:

“for every natural number n , if n is odd then n^2 is odd”
(2)

which we can write more formally as follows:

$$\forall n \in \mathbb{N} \quad (n \text{ odd} \Rightarrow n^2 \text{ odd}) \quad (3)$$

In this case we say that the variable n is not free, but bound by the *quantifier* $\forall$. As a consequence, (3) is true or false “once and for all” and is called a **proposition** (or *statement*).

Components of a proposition

In particular, the components of (3) are the set $\mathbb{N}$, two predicates defined on $\mathbb{N}$ given by

$$p(n) : "n \text{ odd}" \quad \text{and} \quad q(n) : "n^2 \text{ odd}"$$

and the implication

$$p(n) \Rightarrow q(n).$$

Universal implication

Definition 1: of universal implication

*In general, a statement involving a set A , two **predicates** $p(x)$ and $q(x)$ whose argument x ranges over A , and the **logical structure**:*

$$\forall x \in A \ (p(x) \Rightarrow q(x)) \quad (4)$$

*is called a **universal implication**.*

Most **theorems** consist of **universal implications**, in which the predicate $p(x)$ plays the role of the **hypothesis** and the predicate $q(x)$ plays the role of the **thesis** (conclusion).

An example of proposition

- In particular, (3) is a proposition (or a statement):

Proposition 1

$$\forall n \in \mathbb{N} \quad (n \text{ odd} \Rightarrow n^2 \text{ odd}) \quad (5)$$

In this case one is easily convinced that the proposition is true, but how can we prove it rigorously?

How to prove a universal implication

- ▶ For example, is it enough to observe that 3 is odd and $3^2 = 9$ is odd to claim that the proposition is true? Certainly not, since the proposition requires the universal implication to hold for every natural number. However, there are infinitely many odd numbers: how can we prove a universal implication for infinitely many numbers?
- ▶ The key procedure is this: we consider a generic n satisfying the hypothesis (being odd) and we prove that n satisfies the thesis (its square is odd).
- ▶ Let us see how to proceed to prove the previous proposition.

Proof

Let n be odd; we prove that then n^2 is odd.

Any odd number can be written in the form $2k + 1$, for a suitable $k \in \mathbb{N}$. We also observe that $2k$ is an even number for any $k \in \mathbb{N}$.

So let $n = 2k + 1$ be an odd number ($k \in \mathbb{N}$); we therefore need to write n^2 as an even integer plus one. We have:

$$n^2 = (2k + 1)^2 = 4k^2 + 4k + 1 = 2(2k^2 + 2k) + 1.$$

Since $2(2k^2 + 2k)$ is an even integer, n^2 is odd. □

The general procedure

To prove the correctness of a universal implication such as (4), we consider a generic x satisfying the hypothesis $p(x)$ and we try to prove that the thesis $q(x)$ is true.

The case of even numbers

- Let us now prove a similar relation for even numbers:

Proposition 2

$$\forall n \in \mathbb{N} \quad (n \text{ even} \Rightarrow n^2 \text{ even}) \quad (6)$$

Proof

Let n be even; we prove that then n^2 is even.

So let $n = 2k$ be an even number ($k \in \mathbb{N}$); we therefore need to write n^2 as an even integer. We have:

$$n^2 = (2k)^2 = 4k^2 = 2(2k^2)$$

Since $2(2k^2)$ is an even integer, n^2 is even. □

Counterexamples

Counterexamples are an important technique for proving the **falsity** of a universal implication.

- ▶ Let us ask, for example, whether the following universal implication is true or false:

$$\forall n \in \mathbb{N} \ (n \text{ prime} \Rightarrow n \text{ odd}) \quad (7)$$

A moment's reflection shows that **this proposition is false**.
Indeed, the number 2 is prime but it is even.

To claim that a universal implication is true, a proof is needed (an example is not enough), whereas to prove that a universal implication is false, one counterexample is enough.

Counterexample

- ▶ The universal implication requires that every x satisfying the hypothesis also satisfies the thesis: therefore, if we find even a single example of x that satisfies the hypothesis but not the thesis, this means that the universal implication is false. Not “false in one case”, but simply “false”, because the universal implication is true or false once and for all.

Definition 2: of counterexample

*In general, an example that satisfies the hypothesis but not the thesis of a universal implication, and that therefore proves its falsity, is called a **counterexample**.*

Proof

- ▶ The formal proof that the previous universal implication is false is the following.

The number 2 is a counterexample to the universal implication:
“For every natural number n , if n is prime then n is odd”.



Negation of a universal implication

The negation of the proposition

“for every $x \in A$, if $p(x)$ holds then $q(x)$ holds”

$$\forall x \in A \ (p(x) \Rightarrow q(x))$$

is the proposition

“there exists $x \in A$ for which $p(x)$ holds and $q(x)$ does not hold”

$$\exists x \in A \ (p(x) \wedge \neg q(x))$$

This particular x is a **counterexample**.

Summary

- ▶ A **predicate** depends on free variables; a **proposition** is true or false once and for all
- ▶ **Universal implication**: $\forall x \in A (p(x) \Rightarrow q(x))$, with hypothesis p and thesis q
- ▶ To prove it: take a generic x satisfying $p(x)$ and prove $q(x)$
- ▶ n odd $\Rightarrow n^2$ odd; n even $\Rightarrow n^2$ even
- ▶ To disprove it, one **counterexample** is enough:
 $\exists x \in A (p(x) \wedge \neg q(x))$

Law of contraposition

Introduction: Law of contraposition

- ▶ It is an indirect proof technique
- ▶ Instead of proving that the hypothesis implies the thesis, we prove that the negation of the thesis implies the negation of the hypothesis
- ▶ To use it we must be able to construct the **correct negation** of a proposition: we will see the fundamental rules

The contrapositive

The universal implication

$$\forall x \in A \quad (p(x) \Rightarrow q(x)) \quad (8)$$

is logically equivalent to

$$\forall x \in A \quad (\text{not } q(x) \Rightarrow \text{not } p(x)) \quad (9)$$

The second implication is called the **contrapositive** of the first.

An example

- For example: since we know that the universal implication (proposition (5)) holds

$$\forall n \in \mathbb{N} \ (n \text{ odd} \Rightarrow n^2 \text{ odd})$$

the following proposition holds:

Proposition 3

$$\forall n \in \mathbb{N} \ (n^2 \text{ even} \Rightarrow n \text{ even}) \quad (10)$$

Proof

Let n^2 be even; we prove that then n is even. We prove (10) relying on the truth of (5).

If n is not even then it is odd and therefore n^2 is odd by (5). This case contradicts the hypothesis that n^2 is even (hence it cannot occur).

Consequently n is even and (10) is proved. □

Law of contraposition

- ▶ The reasoning used in the previous proof has general validity, and shows precisely that if (8) is true then (9) is true; moreover, if the second is true then the first is true (because “not not $p(x)$ ” is logically equivalent to $p(x)$), so the two are logically equivalent.

The equivalence between (8) and (9) is called the **law of contraposition**. It is a method of *indirect proof* that consists in proving (9) to show that (8) is true (it requires proving that the negation of the thesis implies the negation of the hypothesis).

Negation rules

- ▶ When using the law of contraposition, one must be able to construct the **correct negation** of a given proposition or property.
- ▶ Given any two predicates or properties $p(x)$ and $q(x)$, we list schematically some **rules** for constructing the **negation** of a proposition or property.

Negation of quantifiers (1/2)

The negation of

“for every $x \in A$, $p(x)$ holds”

$$\forall x \in A \ (p(x))$$

is

“there exists $x \in A$ for which $p(x)$ does not hold”

$$\exists x \in A \ (\neg p(x))$$

Negation of quantifiers (2/2)

The negation of

“there exists $x \in A$ for which $p(x)$ holds”

$$\exists x \in A \ (p(x))$$

is

“for every $x \in A$, $p(x)$ does not hold”

$$\forall x \in A \ (\neg p(x))$$

Negation of conjunction and disjunction (1/2)

The negation of

“ $p(x)$ holds and $q(x)$ holds”

$$(p(x) \wedge q(x))$$

is

“ $p(x)$ does not hold or $q(x)$ does not hold”

$$(\neg p(x) \vee \neg q(x))$$

Negation of conjunction and disjunction (2/2)

The negation of

“ $p(x)$ holds or $q(x)$ holds”

$$(p(x) \vee q(x))$$

is

“ $p(x)$ does not hold and $q(x)$ does not hold”

$$(\neg p(x) \wedge \neg q(x))$$

Summary

- ▶ $\forall x \in A (p(x) \Rightarrow q(x))$ is equivalent to $\forall x \in A (\neg q(x) \Rightarrow \neg p(x))$
- ▶ Example: from “ n odd $\Rightarrow n^2$ odd” it follows that “ n^2 even $\Rightarrow n$ even”
- ▶ Negation swaps $\forall$ with $\exists$ and $\wedge$ with $\vee$

Sufficient conditions and necessary conditions



Introduction: Sufficient and necessary conditions

- ▶ In this section we define **sufficient conditions** and **necessary conditions**
- ▶ We will see how to read an implication in these terms and when a condition is necessary and sufficient
- ▶ Through several examples we will learn to distinguish conditions that are necessary but not sufficient and vice versa

Sufficient and necessary conditions

Definition 3: of sufficient condition

*A **sufficient condition** is one that, if satisfied, guarantees the truth of the proposition.*

Definition 4: of necessary condition

*A **necessary condition** is one that must be satisfied for the proposition to be true.*

Implications and conditions

Let $p(x)$ and $q(x)$ be any two predicates or properties. If $p(x)$ implies $q(x)$, formally:

$$p(x) \Rightarrow q(x)$$

then:

- ▶ $p(x)$ is a **sufficient condition** for $q(x)$
- ▶ $q(x)$ is a **necessary condition** for $p(x)$

Odd numbers

- ▶ We have seen that the following universal implication is true (proposition (5)):

$$\forall n \in \mathbb{N} \ (n \text{ odd} \Rightarrow n^2 \text{ odd})$$

Hence “ n odd” is a sufficient condition for “ n^2 odd” and “ n^2 odd” is a necessary condition for “ n odd”.

- ▶ Let us now prove that the following proposition is also true:

Proposition 4

$$\forall n \in \mathbb{N} \ (n^2 \text{ odd} \Rightarrow n \text{ odd}) \quad (11)$$

Proof

Let n^2 be odd; we prove that then n is odd.

So let $n^2 = 2(2k^2 + 2k) + 1$ be an odd number ($k \in \mathbb{N}$), since $2(2k^2 + 2k)$ is an even number). We therefore need to write n as an even integer plus one. We have:

$$\begin{aligned} n &= \sqrt{2(2k^2 + 2k) + 1} = \sqrt{4k^2 + 4k + 1} \\ &= \sqrt{(2k + 1)^2} = 2k + 1. \end{aligned}$$

Since $2k + 1$ is an odd integer, n is odd. □

Necessary and sufficient condition

In this way we have proved that “ n^2 odd” is a necessary and sufficient condition for “ n odd” and also that “ n odd” is a necessary and sufficient condition for “ n^2 odd”:

Proposition 5

$$\forall n \in \mathbb{N} \quad (n \text{ odd} \iff n^2 \text{ odd}) \quad (12)$$

Even numbers

- ▶ We have seen that the following universal implication is true (proposition (6)):

$$\forall n \in \mathbb{N} \quad (n \text{ even} \Rightarrow n^2 \text{ even})$$

Hence “ n even” is a sufficient condition for “ n^2 even” and “ n^2 even” is a necessary condition for “ n even”.

- ▶ Let us now prove that the following proposition is also true (direct proof, without using the law of contraposition as seen previously in (10)):

$$\forall n \in \mathbb{N} \quad (n^2 \text{ even} \Rightarrow n \text{ even})$$

Proof

Let n^2 be even; we prove that then n is even.

So let $n^2 = 2 (2 k^2)$ be an even number ($k \in \mathbb{N}$); we therefore need to write n as an even integer. We have:

$$n = \sqrt{2 (2 k^2)} = \sqrt{4 k^2} = 2 k.$$

Since $2 k$ is an even integer, n is even. □

Necessary and sufficient condition

In this way we have proved that “ n^2 even” is a necessary and sufficient condition for “ n even” and also that “ n even” is a necessary and sufficient condition for “ n^2 even”:

Proposition 6

$$\forall n \in \mathbb{N} \quad (n \text{ even} \iff n^2 \text{ even}) \quad (13)$$

Example

Necessary and sufficient conditions.

For example, for a square matrix of real numbers, the fact that its determinant is different from zero is a necessary and sufficient condition for it to be invertible.

Prime numbers greater than 2

Proposition 7

$$\forall n \in \mathbb{N} \quad (n \text{ prime} > 2 \Rightarrow n \text{ odd}) \quad (14)$$

Proof

Let $n > 2$ be a prime number; we prove that then n is odd.

If n is not odd, it is even. But no even number greater than two is prime, a fact that contradicts the hypothesis that n is prime and greater than two (hence this case cannot occur).

Consequently n is odd.



Necessary but not sufficient

- ▶ Hence “ n odd” is a necessary condition for “ n prime > 2 ” and “ n prime > 2 ” is a sufficient condition for “ n odd”.
- ▶ However, “ n odd” does not imply “ n prime > 2 ”, since for example the number 9 is not prime (counterexample). That is:

$$\forall n \in \mathbb{N} \quad (n \text{ odd} \not\Rightarrow n \text{ prime greater than } 2)$$

Hence “ n odd” is a necessary but not sufficient condition for “ n prime > 2 ” and “ n prime > 2 ” is a sufficient but not necessary condition for “ n odd”.

Numbers divisible by 6

Proposition 8

$$\forall n \in \mathbb{N} \quad (n \text{ divisible by } 6 \Rightarrow n \text{ even}) \quad (15)$$

Proof

Let n be a number divisible by six; we prove that then n is even.

If n is not even, it is odd. But no odd number is divisible by six, a fact that contradicts the hypothesis that n is divisible by six (hence this case cannot occur).

Consequently n is even.



Necessary but not sufficient

- ▶ Hence “ n even” is a necessary condition for “ n divisible by 6” and “ n divisible by 6” is a sufficient condition for “ n even”.
- ▶ However, “ n even” does not imply “ n divisible by 6”, since for example the number 2 is even but not divisible by six (counterexample). That is:

$$\forall n \in \mathbb{N} \quad (n \text{ even} \not\Rightarrow n \text{ divisible by } 6)$$

Hence “ n even” is a necessary but not sufficient condition for “ n divisible by 6” and “ n divisible by 6” is a sufficient but not necessary condition for “ n even”.

Example (1/2)

Necessary/sufficient but not sufficient/necessary conditions.

Being a square implies being a rectangle:

$$\text{being a square} \Rightarrow \text{being a rectangle}$$

since all squares are rectangles.

Hence “being a rectangle” is a necessary condition for “being a square” and “being a square” is a sufficient condition for “being a rectangle”.

Example (2/2)

But being a rectangle does not imply being a square

being a rectangle $\nRightarrow$ being a square

because there are rectangles that are not squares.

Hence “being a rectangle” is not a sufficient condition (but it is a necessary one) for “being a square” and “being a square” is not a necessary condition (but it is a sufficient one) for “being a rectangle”.

Summary

- ▶ If $p(x) \Rightarrow q(x)$: p is **sufficient** for q , q is **necessary** for p
- ▶ n odd $\Leftrightarrow n^2$ odd; n even $\Leftrightarrow n^2$ even (necessary and sufficient conditions)
- ▶ “ n odd” is necessary but not sufficient for “ n prime > 2 ”
- ▶ “ n even” is necessary but not sufficient for “ n divisible by 6”
- ▶ A counterexample shows that a condition is not sufficient

Proofs by contradiction



Introduction: Proofs by contradiction

- ▶ It is an indirect proof technique
- ▶ We assume the hypothesis true and the thesis false, and we reach a contradiction
- ▶ We will apply it to a classical result: $\sqrt{2}$ is not rational

Proof by contradiction

Definition 5: of proof by contradiction

*In general, a **proof by contradiction** consists in assuming that the hypothesis of the theorem and the negation of the thesis are true, and deducing from these facts a contradiction of any kind.*

- ▶ We illustrate **proof by contradiction** with the following theorem.

Theorem 1

There is no rational number whose square is 2.

Proof (1/2)

Suppose, by contradiction, that there exists a number $r \in \mathbb{Q}$ such that $r^2 = 2$.

We can write $r = \frac{n}{m}$ with $n, m \in \mathbb{Z}$, $m \neq 0$. Moreover, we can assume that the fraction $\frac{n}{m}$ is already reduced to lowest terms, that is, “simplified” (in other words: n , m have no common factors).

We then have the chain of implications:

$$\left(\frac{n}{m}\right)^2 = 2 \quad \Rightarrow \quad n^2 = 2 m^2$$

so n^2 is even; but then by (13) n is also even and we can write $n = 2k$ for some $k \in \mathbb{Z}$.

Proof (2/2)

Hence the relation $n^2 = 2 m^2$ can be rewritten as:

$$(2 k)^2 = 2 m^2$$

$$4 k^2 = 2 m^2$$

$$m^2 = 2 k^2.$$

So m^2 is even. But then by (13) m is also even.

Therefore both n and m are even, and this is a **contradiction**, because we had assumed that the fraction $\frac{n}{m}$ had already been simplified. □

The proof can be found in Euclid's Elements (around 300 BC).

Summary

- ▶ **Proof by contradiction:** assume the hypothesis and the negation of the thesis and deduce a contradiction
- ▶ Classical example (Euclid): there is no $r \in \mathbb{Q}$ with $r^2 = 2$
- ▶ The proof uses the fact that n^2 even $\Rightarrow n$ even

Logic and sets

Introduction: Logic and sets

- ▶ The language of logic and the language of sets are two sides of the same coin.
- ▶ We will see the parallel between implication and inclusion, and between equality of sets and universal implications
- ▶ and between set operations and logical operations (distributive properties and De Morgan's laws)

Logical implication and set inclusion (1/2)

- There is a parallel between the relation of set inclusion and logical implication. To explain it, consider the universal implication:

$$\forall n \in \mathbb{N} \quad (n \text{ divisible by } 4 \Rightarrow n \text{ divisible by } 2) \quad (16)$$

If we denote by:

$$D_4 = \{ n \in \mathbb{N} : n \text{ is divisible by } 4 \}$$

$$D_2 = \{ n \in \mathbb{N} : n \text{ is divisible by } 2 \}$$

Logical implication and set inclusion (2/2)

we can observe that the universal implication written above is equivalent to the assertion:

$$D_4 \subseteq D_2$$

Indeed, this inclusion means that every element belonging to D_4 also belongs to D_2 , that is, that every natural number divisible by 4 is also divisible by 2.

Implication and inclusion

The universal implication:

“for every $x \in A$, if $p(x)$ holds then $q(x)$ holds”

is equivalent to the set inclusion:

$$\{ x \in A : p(x) \text{ is true} \} \subseteq \{ x \in A : q(x) \text{ is true} \}.$$

Equality of sets and universal implications

- ▶ Proving the equality of two sets, i.e., $A = B$, requires proving two universal implications. Formally:

$$\forall x \ (x \in A \Rightarrow x \in B) \quad \text{and} \quad \forall x \ (x \in B \Rightarrow x \in A). \quad (17)$$

- ▶ Stating that $A \subsetneq B$ means stating that “Every element that belongs to A also belongs to B , and there exists an element of B that does not belong to A ”. Formally:

$$\forall x \ (x \in A \Rightarrow x \in B) \quad \text{and} \quad \exists x \in B : x \notin A. \quad (18)$$

Set operations and logical operations

There is a relation between operations on sets and logical operations. Precisely:

1. **Set intersection** is defined by means of “and” (**logical conjunction**).
2. **Set union** is defined by means of “or” (**logical disjunction**).
3. **Set difference** and the **complement operation** are defined by means of “not” (**logical negation**).

Distributive properties

- ▶ The distributive properties of union and intersection of sets:

Observation 1: distributive properties (sets)

Given three sets A , B and C we have:

$$A \cap (B \cup C) = (A \cap B) \cup (A \cap C)$$

$$A \cup (B \cap C) = (A \cup B) \cap (A \cup C).$$

- ▶ They can be rewritten in terms of predicates by observing that the intersection symbol $\cap$ corresponds to the conjunction $\wedge$ ("and") and that the union symbol $\cup$ corresponds to the disjunction $\vee$ ("or").

Distributive properties

Observation 2: distributive properties (predicates)

Given three predicates $p(x)$, $q(x)$ and $r(x)$ we have:

$$p(x) \wedge (q(x) \vee r(x)) = (p(x) \wedge q(x)) \vee (p(x) \wedge r(x))$$

$$p(x) \vee (q(x) \wedge r(x)) = (p(x) \vee q(x)) \wedge (p(x) \vee r(x)).$$

De Morgan's laws

- De Morgan's laws:

Proposition 9: De Morgan's laws (sets)

Given the sets $B, C \subseteq U$, we have

$$\overline{B \cap C} = \overline{B} \cup \overline{C} \qquad \overline{B \cup C} = \overline{B} \cap \overline{C}$$

De Morgan's laws

- ▶ They can be rewritten in terms of predicates by observing that the complement operation corresponds to the negation $\neg$ ("not").

Proposition 10: De Morgan's laws (predicates)

Given two predicates $q(x)$ and $r(x)$ we have:

$$\neg(q(x) \wedge r(x)) = \neg q(x) \vee \neg r(x)$$

$$\neg(q(x) \vee r(x)) = \neg q(x) \wedge \neg r(x)$$

Summary

- ▶ $\forall x \in A (p(x) \Rightarrow q(x))$ is equivalent to $\{x \in A : p(x)\} \subseteq \{x \in A : q(x)\}$
- ▶ $A = B$ requires two universal implications; $A \subsetneq B$ also requires an element of B outside A
- ▶ $\cap \leftrightarrow \wedge$, $\cup \leftrightarrow \vee$, complement $\leftrightarrow \neg$
- ▶ Distributive properties and De Morgan's laws hold for sets and for predicates

Summary (1/2)

Language of logic:

- ▶ Symbols: $\forall, \exists, \Rightarrow, \in, \vee, \wedge, \neg$
- ▶ Predicates and propositions; universal implications (hypothesis and thesis)

Proof techniques:

- ▶ Direct proof: a generic x satisfying the hypothesis
- ▶ Counterexample to show falsity
- ▶ Law of contraposition and negation rules
- ▶ Proof by contradiction ($\sqrt{2} \notin \mathbb{Q}$)

Summary (2/2)

Necessary and sufficient conditions:

- ▶ $p \Rightarrow q$: p sufficient for q , q necessary for p
- ▶ $p \Leftrightarrow q$: necessary and sufficient condition

Logic and sets:

- ▶ Implication $\Leftrightarrow$ inclusion; $\wedge, \vee, \neg \Leftrightarrow \cap, \cup$, complement
- ▶ Distributive properties and De Morgan's laws