



SAPIENZA
UNIVERSITÀ DI ROMA

Mathematical Analysis 1

Volume 1 — Numbers and logic

The first volume: the language and the numbers on which analysis is built. Sets, logic and proof techniques, number sets and binary relations, ordered fields, supremum and the completeness axiom, roots, powers and logarithms, summations and geometric progressions, mathematical induction, factorials and binomial coefficients, the Fibonacci sequence, infinite sets, complex numbers. Definitions, theorems with proofs, worked examples; at the end, the exercises of the part with solutions.

Fabio Furini

fabiofurini.github.io

Department of Computer, Control and Management Engineering — Sapienza University of Rome

Online version and downloadable material:
fabiofurini.github.io/mathematical-analysis-1/
github.com/fabiofurini/mathematical-analysis-1

Academic year 2026–2027

Contents

1	Sets	3
1.1	Informal introduction to set theory	3
1.2	Number sets	6
1.3	Relations between sets	11
1.4	Operations on sets	16
1.5	Further topics	33
2	Basics of logic and proof techniques	37
2.1	Logical symbols	37
2.2	Universal implications and proofs	37
2.3	Law of contraposition	41
2.4	Sufficient conditions and necessary conditions	42
2.5	Proofs by contradiction	46
2.6	Logic and sets	47
3	Number sets and intervals	50
3.1	The five main number sets	50
3.2	Intervals	53
4	Binary relations	55
4.1	Binary relations	55
4.2	Properties of relations	59
4.3	Representation with directed graphs	61
4.4	Equivalence relations	65
4.5	Partial order relations	70
4.6	Total order relations	75
4.7	Functions	77
5	Ordered fields, supremum/infimum and the completeness axiom	79
5.1	Properties R_1 and R_2	79
5.2	Commensurable magnitudes	80
5.3	Geometric representation of $\mathbb{Q}$	81
5.4	Total order relations	82
5.5	Property R_3	83
5.6	Ordered fields	84
5.7	Bounded sets, maxima and minima	85
5.8	Upper/lower bounds and suprema/infima	90
5.9	Axiomatic definition of the real numbers	92
5.10	Geometric representation of $\mathbb{R}$	94

6	Roots, powers, logarithms and modular arithmetic	96
6.1	Roots, powers, logarithms	96
6.2	Modular arithmetic	103
7	Summations, geometric and arithmetic progressions	108
7.1	Summations	108
7.2	Geometric progressions	113
7.3	Arithmetic progressions	118
8	Principle of mathematical induction	123
8.1	The principle of mathematical induction	123
8.2	Proofs based on the principle of mathematical induction	124
9	Factorials, binomial coefficients and triangle inequality	134
9.1	Factorials	134
9.2	Binomial coefficients	135
9.3	Absolute value	139
10	The Fibonacci sequence	141
10.1	The Fibonacci sequence	141
11	Cardinality of infinite sets	146
11.1	Countable cardinality	146
11.2	Cardinality of the continuum	149
12	Complex numbers	152
12.1	Definition of $\mathbb{C}$ and field structure	152
12.2	Algebraic form of complex numbers	154
12.3	Trigonometric form of complex numbers	160
12.4	Quadratic equations	168
13	Exercises	172
13.1	Maxima, minima, suprema and infima	172
13.2	Equations and inequalities in one real variable	178
13.3	Summations	185
13.4	Principle of mathematical induction	187
13.5	Complex numbers	199

Sets

1.1 Informal introduction to set theory

- Set theory is based on the following three key concepts:

1. Sets

The notion of set is generally taken as **primitive** (that is, not reducible to more elementary concepts). The words *collection*, *class*, *aggregate*, *family* are used as synonyms of set.

Example 1.1: sets

For example, we have the set of points of a plane, the set of students of a university, or the set of stars of a galaxy.

2. Elements

A set is determined by its *elements*, in the sense that a set is defined when we have a **criterion** to establish whether a given element is or is not an element of this set.

There are sets with a *finite number of elements* (for example the set of students of a university) and sets with an *infinite number of elements* (for example the set of points of the plane).

Sets are usually denoted by capital letters. The elements of a set are usually denoted by lowercase letters.

3. Membership

The concept of *membership* links elements to sets. When an object is an element of a set, we say that the element *belongs* to the set. To indicate that an element x belongs to a set A we write:

$$x \in A$$

To indicate that an element x does *not* belong to a set A we write:

$$x \notin A$$

1.1.1 Informal definition of sets

1. A first way to define sets is **definition by listing** (roster notation)

A set can be defined **by listing**, that is, by listing the elements that belong to it between curly brackets. This technique assumes that the set has a finite number of elements.

Example 1.2: definition by listing

- The notation:

$$A = \{a, b, c\}$$

means that the set A has as elements the three letters a , b and c . For example, a belongs to A , that is, $a \in A$, while $d \notin A$.

- The set of vowels of the Latin alphabet can be defined by listing:

$$V = \{a, e, i, o, u\}$$

- When listing the elements of a set, each element is written **only once**: a set does not contain the same object more than once.
- The elements of a set are **not ordered**: $\{1, 2, 3\}$ and $\{3, 1, 2\}$ describe the same set, because the order in which we list the elements is irrelevant.

2. A second way to define sets is **definition by a property**:

A set can be defined **by a property** as follows:

$$A = \{ x \in U : p(x) \text{ is true} \}$$

where $p(x)$ is the property that the element x of the set U must have in order to belong to the set A . This technique can be used to define sets with a finite or even an infinite number of elements.

Example 1.3: definition by a property

Consider for example the set of letters of the Latin alphabet:

$$U = \{a, b, c, d, e, f, g, h, i, j, k, l, m, n, o, p, q, r, s, t, u, v, w, x, y, z\}$$

Using for example the property $p(x)$ defined as “ x is a vowel” we can define the following set of vowels:

$$A = \underbrace{\{x \in U : x \text{ is a vowel}\}}_{\{a, e, i, o, u\}}$$

Note that to define a set A by a property we need a set U to which all the elements of the set A we want to define belong. The set U plays the role of the **universal set**.

Definition 1.1: of universal set

A **universal set** (or simply **universe**), denoted by U , is a set fixed in advance that contains all the objects under consideration in a given context: all the sets considered in that context have elements that belong to U .

It is important that the property $p(x)$ we use makes sense for every x of the set U (universal set), and therefore is true or false (without ambiguity of meaning) for every particular $x \in U$; the set A will then consist of exactly those x belonging to U for which the property $p(x)$ is true.

Fixing a single universal set for the whole discussion is convenient, but it is not necessary. We can start from **any set already known** B , which may change from one definition to the next:

Given a set B already known, the notation

$$A = \{ x \in B : p(x) \}$$

defines the set A made of exactly those elements of B for which the property $p(x)$ is true ("is true" is usually not written). The universal set is the special case in which $B = U$ is the same for all the definitions in a given context.

- This is the form used everywhere in mathematics. The property can be written in words or with a formula, and it can consist of several conditions, separated by commas or by "and".
- When the starting set is clear from the context, it is often left implicit and we simply write $\{ x : p(x) \}$. For example, when discussing real numbers, $\{ x : x > 0 \}$ denotes the set of positive real numbers. The starting set is still fixed: it is only omitted from the notation.

Example 1.4: properties with different starting sets

- Starting from $B = \{a, b, c, d, e\}$ and from the same property " x is a vowel" we obtain a set different from the one of the previous example:

$$\{x \in B : x \text{ is a vowel}\} = \{a, e\}$$

- Starting from $C = \{1, 2, 3, 4, 5, 6\}$ we can define:

$$\{x \in C : x \text{ is even}\} = \{2, 4, 6\}$$

- The conditions can be more than one:

$$\{x \in C : x \text{ is even, } x > 2\} = \{4, 6\}$$

In all cases the starting set is chosen each time and is not the universe of the whole discussion.

3. A third way to define sets is **definition by construction**: instead of choosing the elements among those of a set that already contains them, we **build** them from the elements of a set already known.

Given a set B already known, a set can be defined **by construction** as follows:

$$A = \{ \text{expression built from } x : x \in B \}$$

that is, A is made of all the objects obtained by applying the same rule to each element x of B . There is no need to know in advance a set containing the elements of A .

Example 1.5: definition by construction

- The set of the squares of the elements of $\{1, 2, 3\}$:

$$\{ x^2 : x \in \{1, 2, 3\} \} = \{1, 4, 9\}$$

- The set of the doubles of the elements of $\{1, 2, 3, 4\}$:

$$\{ 2x : x \in \{1, 2, 3, 4\} \} = \{2, 4, 6, 8\}$$

If the rule produces the same object from different elements, the object is counted only once: for example $\{ x^2 : x \in \{-1, 0, 1\} \} = \{0, 1\}$.

One must be careful when defining sets, since contradictions may arise. There is a formal definition of the concept of set, developed to avoid contradictions, but it is beyond the scope of this course.

- In all the ways seen so far we always start from a set already known. The notation $\{ x : p(x) \}$ understood as “all the objects for which $p(x)$ is true”, with no starting set at all (not even an implicit one), does **not** in general define a set.
- For example, with the property $p(x)$ defined as “ $x \notin x$ ” we would obtain the set of all sets that do not contain themselves, which however is not a set in the formal definition of sets. Admitting this set would generate the contradiction: “the set of all sets that do not belong to themselves belongs to itself if and only if it does not belong to itself” (**Russell’s paradox**, discussed in the further topics at the end of the chapter).

1.2 Number sets

Definition 1.2: numeral system

A **positional numeral system** is a way of encoding numbers using a **sequence of digits**, where each digit contributes differently to the number depending on its position. The number of distinct digits is the **base** of the system.

The **decimal expansion** of a number is a numerical representation in base 10; it expresses a number as a sum of powers of 10, and it can be finite or infinite depending on the type of number.

- Informal definition of the main number sets:

1. We denote by $\mathbb{N}$ the set of **natural numbers**, that is, the set of numbers that can be written as **decimal expansions without decimal point and without sign**. We will use the informal notation:

$$\mathbb{N} = \{0, 1, 2, 3, 4, \dots\}$$

We adopt the convention that **zero is a natural number**, that is $0 \in \mathbb{N}$ (other textbooks exclude zero from $\mathbb{N}$: it is only a convention, but it must be stated once and for all). We denote by $\mathbb{N}_{>0}$ the set of **positive natural numbers**:

$$\mathbb{N}_{>0} = \mathbb{N} \setminus \{0\} = \{1, 2, 3, 4, \dots\}$$

Definition 1.3: of even number

A natural number n is said to be **even** if there exists a natural number m such that:

$$n = 2m$$

Definition 1.4: of odd number

A natural number n is said to be **odd** if there exists a natural number m such that:

$$n = 2m + 1$$

Example 1.6: even and odd numbers

- 0 is even, because $0 = 2 \cdot 0$, and 6 is even, because $6 = 2 \cdot 3$.
- 1 is odd, because $1 = 2 \cdot 0 + 1$, and 7 is odd, because $7 = 2 \cdot 3 + 1$.

The same two definitions are used for the integers, by requiring m to be an integer: for example $-4 = 2 \cdot (-2)$ is even and $-3 = 2 \cdot (-2) + 1$ is odd.

The **well-ordering axiom** states that every non-empty subset of $\mathbb{N}$ has a minimum.

- It is a property that is **assumed as an axiom** and that characterizes $\mathbb{N}$: it does not hold in the other number sets, since $\mathbb{Z}$, $\mathbb{Q}$ and $\mathbb{R}$ are themselves non-empty subsets without a minimum.
- It is the axiom that justifies the **principle of mathematical induction**, which we will deal with later on, and it guarantees the existence of a smallest element whenever a property is satisfied by at least one natural number.

Observation 1.5

Every natural number is either even or odd, and no natural number is at the same time even and odd.

Proof. We prove the two statements separately.

Every natural number is either even or odd. Consider the set

$$S = \{ n \in \mathbb{N} : n \text{ is neither even nor odd} \}$$

and suppose by contradiction that S is not empty. By the well-ordering axiom S has a minimum, which we denote by n_0 . We have that:

- $n_0 \neq 0$, because $0 = 2 \cdot 0$ is even;
- $n_0 \neq 1$, because $1 = 2 \cdot 0 + 1$ is odd.

Hence $n_0 \geq 2$ and therefore $n_0 - 2 \in \mathbb{N}$. Since $n_0 - 2 < n_0$ and n_0 is the minimum of S , the number $n_0 - 2$ does not belong to S , that is, it is even or odd:

- if $n_0 - 2 = 2m$ with $m \in \mathbb{N}$, then $n_0 = 2m + 2 = 2(m + 1)$ and hence n_0 is even;
- if $n_0 - 2 = 2m + 1$ with $m \in \mathbb{N}$, then $n_0 = 2m + 3 = 2(m + 1) + 1$ and hence n_0 is odd.

In both cases $n_0 \notin S$, contradicting the fact that n_0 is the minimum of S . Hence S is empty.

No natural number is at the same time even and odd. If a natural number n were even and odd, there would exist $m, k \in \mathbb{N}$ with

$$n = 2m \quad \text{and} \quad n = 2k + 1 \quad \implies \quad 2(m - k) = 1$$

But $m - k$ is an integer: if $m \leq k$ the left-hand side is less than or equal to 0, while if $m \geq k + 1$ the left-hand side is greater than or equal to 2. In no case can the left-hand side be equal to 1, and therefore such an n does not exist. □

2. We denote by $\mathbb{Z}$ the set of **integers**, that is, the set of numbers that can be written as **decimal expansions without decimal point and with a sign**. We will use the informal notation:

$$\mathbb{Z} = \{0, \pm 1, \pm 2, \pm 3, \pm 4, \dots\}$$

3. We denote by $\mathbb{Q}$ the set of **rational numbers**, that is, the set of numbers that can be written as **finite or infinite periodic decimal expansions**. In other words, it is the set of numbers that can be written as a fraction $\frac{p}{q}$ where p is an integer and q is a natural number different from zero.

Example 1.7: rational numbers

- For example, with $p = 2$ and $q = 5$ we have the fraction $\frac{2}{5}$, whose decimal expansion is 0.4.
- For example, with $p = 4$ and $q = 10$ we have the fraction $\frac{4}{10}$, whose decimal expansion is again 0.4. Note that a rational number can be written with more than one fraction.
- For example, with $p = 13$ and $q = 30$ we have the fraction $\frac{13}{30}$, whose decimal expansion is $0.4\bar{3} = 0.4333\dots$

However, we can represent every rational number different from 0 by a single fraction $\frac{p}{q}$ by choosing $p \in \mathbb{Z}$ and $q \in \mathbb{N}$ coprime (that is, relatively prime: p and q are not both divisible by the same integer greater than 1).

Observation 1.6

$$0.\bar{9} = 1$$

Proof. There are different proofs of this remark, based on different mathematical techniques.

- (a) A simple proof follows directly from the definition of 1 divided by 3; indeed, we have:

$$\begin{aligned}\frac{1}{3} &= 0.\bar{3} \\ \frac{1}{3} \cdot 3 &= 0.\bar{3} \cdot 3 \\ 1 &= 0.\bar{9}\end{aligned}$$

- (b) Using algebraic arguments we can write:

$x = 0.999\dots$	
$10x = 9.999\dots$	multiplying by 10
$10x = 9 + 0.999\dots$	separating the integer part
	from the fractional part
$10x = 9 + x$	by definition of x
$9x = 9$	subtracting x
$x = 1$	dividing by 9

(c) A proof by contradiction is the following:

$$\begin{aligned}
 0.\bar{9} &\neq 1 \\
 0.\bar{9} \cdot 9 &\neq 1 \cdot 9 \\
 0.\bar{9} \cdot 9 + 0.\bar{9} &\neq 1 \cdot 9 + 0.\bar{9} \\
 0.\bar{9} \cdot 9 + 0.\bar{9} &\neq 9.\bar{9} \\
 0.\bar{9} \cdot (9 + 1) &\neq 9.\bar{9} \\
 0.\bar{9} \cdot (10) &\neq 9.\bar{9} \\
 9.\bar{9} &\neq 9.\bar{9} \quad \text{contradiction!}
 \end{aligned}$$

□

4. We denote by $\mathbb{R}$ the set of **real numbers**, that is, the set of numbers identified with finite or infinite decimal expansions, periodic or non-periodic.

Example 1.8: real numbers

- Consider for example the number

$$0.10110111011110\dots$$

obtained by putting after the decimal point one digit equal to 1, then 0, then two digits equal to 1, then 0, then three digits equal to 1 ... and so on. The string of digits after the decimal point is neither finite nor periodic: therefore this number is real but not rational.

- Other examples of real but not rational numbers are $\sqrt{2}$ and $\sqrt{3}$, or π and Euler's number e (Napier's constant), which have infinite non-periodic decimal expansions and are therefore real but not rational numbers.

Example 1.9: number sets defined by listing and by a property

- The set of the first five prime numbers can be defined by listing:

$$P = \{2, 3, 5, 7, 11\}$$

For example $2 \in P$, while $4 \notin P$.

- The set of **even numbers** can be defined by a property, with universal set $\mathbb{Z}$:

$$E = \{x \in \mathbb{Z} : \text{there exists } k \in \mathbb{Z} \text{ such that } x = 2k\}$$

For example $-4 \in E$, because $-4 = 2 \cdot (-2)$, while $3 \notin E$, because $3 = 2k$ would give $k = \frac{3}{2} \notin \mathbb{Z}$.

- The set of **positive real numbers** can be defined by a property, with universal set $\mathbb{R}$:

$$\mathbb{R}_{>0} = \{x \in \mathbb{R} : x > 0\}$$

It is a set with an infinite number of elements, which cannot be defined by listing.

- There are formal definitions of the sets of natural, integer and rational numbers that are beyond the scope of this course. Later we will give a formal definition of the real numbers.

1.3 Relations between sets

Definition 1.7: of equal sets

Two sets A and B are **equal** when they have the same elements. We write

$$A = B$$

and this means that every element that belongs to A also belongs to B and every element that belongs to B also belongs to A . If A and B are not equal we write $A \neq B$.

Example 1.10: order and multiplicity of the elements

- The concept of *order* among the elements is foreign to sets:

$$\{1, 2, 5\} = \{1, 5, 2\}$$

because the two sets have the same elements, the numbers 1, 2 and 5: the order in which we list the elements is irrelevant.

- The concept of *multiplicity of the elements* is also foreign to sets. For example, the set of solutions of the equation

$$x - 1 = 0$$

(whose only element is the number 1) is equal to the set of solutions of the equation

$$(x - 1)^2 = 0$$

The fact that the second equation has the solution $x = 1$ with algebraic multiplicity 2 does not change the set of its solutions: both sets are equal to $\{1\}$.

It may happen that only one of the two requirements expressed by the equality relation holds. For example, if we only know that every element of A is also an element of B , we can say that A is contained in B .

Definition 1.8: of subset

Given two sets A and B , we say that A is a **subset** of B , and we write

$$A \subseteq B,$$

if every element of A is also an element of B , that is, if for every x the implication $x \in A \implies x \in B$ holds. We read “ A is contained in B ” or “ A is a subset of B ”.

If we state that $A \subseteq B$, we do not exclude that $A = B$. If instead we want to state precisely that A is contained in B but does not coincide with B , we say that A is *strictly contained* in B .

Definition 1.9: of proper subset

Given two sets A and B , we say that A is a **proper subset** of B , or that A is **strictly contained** in B , and we write

$$A \subsetneq B,$$

if $A \subseteq B$ and $A \neq B$. In this case we speak of **strict inclusion**.

- In some texts the symbol $\subset$ denotes strict inclusion, in others it is a synonym of $\subseteq$. To avoid ambiguity we will use $\subseteq$ for inclusion and $\subsetneq$ for strict inclusion; when it appears, the symbol $\subset$ has the same meaning as $\subseteq$.

Example 1.11: inclusion and strict inclusion

- $\{1, 4\} \subseteq \{1, 3, 4\}$, because 1 and 4 belong to $\{1, 3, 4\}$. The inclusion is strict, $\{1, 4\} \subsetneq \{1, 3, 4\}$, because $3 \in \{1, 3, 4\}$ but $3 \notin \{1, 4\}$, so the two sets are not equal.
- $\{1, 3, 4\} \subseteq \{1, 3, 4\}$, but the inclusion is not strict, because the two sets are equal.
- $\mathbb{N} \subsetneq \mathbb{Z}$: every natural number is an integer, and $-1 \in \mathbb{Z}$ but $-1 \notin \mathbb{N}$.

Be careful not to confuse “belongs to” and “is contained in” (in symbols, $\in$ and $\subseteq$). In a sense, they are two ways of indicating that “something is inside something else”, but they have a fundamental logical difference:

- a set is contained in another set;
- an element belongs to a set.

Example 1.12: “belongs to” vs “is contained in”

For example:

$$3 \in \{1, 3, 4\};$$

$$\{3\} \subseteq \{1, 3, 4\};$$

$$\{1, 4\} \subseteq \{1, 3, 4\}.$$

In particular, do not confuse 3 (which is a number) with $\{3\}$, which is the set that contains the number 3 as its only element.

Sometimes we consider sets whose elements are other sets. In this case too, the symbols $\in$ and $\subseteq$ are not interchangeable, but must be used correctly.

Example 1.13: “belongs to” vs “is contained in”

For example, if we define the set

$$A = \left\{ \{1\}, \{2\}, \{1, 2\} \right\},$$

then it is correct to state that $\{2\} \in A$, because $\{2\}$ is a set that plays the role of an element of A , whereas it would be incorrect to say that $\{2\} \subseteq A$, because this would mean that $2 \in A$, while A has $\{2\}$ as an element but not 2 . Instead, the subset of A containing only the element $\{2\}$ is denoted by $\{\{2\}\}$ and we have $\{\{2\}\} \subseteq A$.

1.3.1 Properties of inclusion**Proposition 1.10: reflexive property of inclusion**

For every set A we have

$$A \subseteq A.$$

Proof. By the definition of subset we must check that every element of A is also an element of A , that is, that for every x the implication $x \in A \implies x \in A$ holds. The implication is true because the conclusion coincides with the hypothesis: if $x \in A$, then $x \in A$. Hence $A \subseteq A$. $\square$

Proposition 1.11: antisymmetric property of inclusion

Given two sets A and B , we have

$$A = B \iff A \subseteq B \text{ and } B \subseteq A.$$

Proof. ($\implies$) Suppose $A = B$. By the definition of equal sets every element of A belongs to B , hence $A \subseteq B$ by the definition of subset. In the same way every element of B belongs to A , hence $B \subseteq A$.

($\impliedby$) Suppose $A \subseteq B$ and $B \subseteq A$. From $A \subseteq B$ it follows that every element of A belongs to B , and from $B \subseteq A$ it follows that every element of B belongs to A . Hence A and B have the same elements, that is, $A = B$. $\square$

- The antisymmetric property gives the most common method to prove that two sets A and B are equal, called **double inclusion**: we take an arbitrary element of A and show that it belongs to B (that is, $A \subseteq B$), then we take an arbitrary element of B and show that it belongs to A (that is, $B \subseteq A$).

Proposition 1.12: transitive property of inclusion

Given three sets A , B and C , we have

$$A \subseteq B \text{ and } B \subseteq C \implies A \subseteq C.$$

Proof. Suppose $A \subseteq B$ and $B \subseteq C$, and let x be an arbitrary element of A . Since $A \subseteq B$, we have $x \in B$. Since $B \subseteq C$, from $x \in B$ it follows that $x \in C$. Hence every element of A belongs to C , that is, $A \subseteq C$. $\square$

1.3.2 Empty set, cardinality and power set**Definition 1.13: of empty set**

The **empty set** is the set that contains no elements. It is denoted by $\emptyset$.

Observation 1.14

Given a set A , we have:

$$\emptyset \subseteq A$$

Proof. By the definition of subset we must prove that every element that belongs to $\emptyset$ also belongs to A , that is, that for every x the implication

$$x \in \emptyset \implies x \in A$$

holds. The hypothesis $x \in \emptyset$ is false for every x , because no element belongs to $\emptyset$. An implication with a false hypothesis is true whatever the conclusion is: we say that the implication is **vacuously true**. The same can be seen by contradiction: if $\emptyset \subseteq A$ did not hold, there would exist an element of $\emptyset$ that does not belong to A ; but $\emptyset$ has no elements, so such an element does not exist. Hence the thesis holds. $\square$

Definition 1.15: of cardinality of a set

The number of elements of a set A is the **cardinality** of the set. It is denoted by $|A|$. If the cardinality of A is a natural number, the set A is called **finite**; otherwise it is called **infinite**.

- The empty set is finite and has cardinality $|\emptyset| = 0$.
- A set with only one element is called a **singleton**: for example $\{3\}$ is a singleton and $|\{3\}| = 1$.
- A subset with k elements of a set A is called a **k -subset** of A .

Example 1.14: cardinality and subsets with k elements

Given the set $A = \{1, 2, 3\}$ we have $|A| = 3$. The 2-subsets of A are $\{1, 2\}$, $\{1, 3\}$ and $\{2, 3\}$; the 1-subsets of A are the singletons $\{1\}$, $\{2\}$ and $\{3\}$; the only 0-subset of A is $\emptyset$ and the only 3-subset of A is A itself.

Definition 1.16: of power set

Given a set A , the set whose elements are all the subsets of A is called the **power set** of A and is denoted by the symbol $\mathcal{P}(A)$. The notation 2^A is also used.

- Every set A has two trivial subsets, namely A itself (by the reflexive property of inclusion) and the empty set $\emptyset$ (by the remark on the empty set); the two subsets coincide if A is empty.

Example 1.15: power set

For example, given

$$A = \{1, 2, 3\},$$

then

$$\mathcal{P}(A) = \left\{ \emptyset, \{1\}, \{2\}, \{3\}, \{1, 2\}, \{2, 3\}, \{1, 3\}, \{1, 2, 3\} \right\}$$

Observation 1.17

Given a set A with n elements, the power set $\mathcal{P}(A)$ has 2^n elements:

$$|\mathcal{P}(A)| = 2^n = 2^{|A|}$$

Proof. If $n = 0$, the set A is empty and its only subset is $\emptyset$, hence $|\mathcal{P}(A)| = 1 = 2^0$. If $n \geq 1$, let $a_1, a_2, \dots, a_n$ be the elements of A . To build a subset S of A we must decide, for each of the elements $a_1, a_2, \dots, a_n$, whether or not S contains that element: hence we must make a choice between two options n times, and each choice is independent of the others.

- Every sequence of n choices determines a subset of A and, conversely, every subset S of A is obtained from a sequence of choices: for each i we choose to include a_i if and only if $a_i \in S$.
- Different sequences of choices give different subsets: if two sequences differ in the choice about the element a_i , then a_i belongs to one of the two subsets but not to the other.

The total number of possible subsets is therefore equal to the number of sequences of n


choices between two options, that is,

$$\underbrace{2 \cdot 2 \cdots 2}_{n \text{ times}} = 2^n.$$

□

Example 1.16: construction of the power set with a binary tree

Given the set $A = \{1, 2, 3\}$ with $n = 3$ elements, the cardinality of its power set is $2^3 = 8$. The construction of the power set $\mathcal{P}(A)$ can be visualized through the following **binary tree** (an undirected, connected and acyclic graph) in which at each level we decide whether or not to include the object in the subset:



1.4 Operations on sets

Definition 1.18: of intersection of sets

The *intersection* of two sets $A, B \subseteq U$ is the set defined by:

$$A \cap B = \{x \in U : x \in A \text{ and } x \in B\}$$

It is the set of elements that belong both to the first and to the second set.

Definition 1.19: of union of sets

The **union** of two sets $A, B \subseteq U$ is the set defined by:

$$A \cup B = \{x \in U : x \in A \text{ or } x \in B\}$$

It is the set of elements that belong to the first or to the second set, where “or” is meant in the non-exclusive sense (the set of elements that belong to A or to B or to both).

Definition 1.20: of difference of sets

The **difference** of two sets $A, B \subseteq U$ is the set defined by:

$$A \setminus B = \{x \in A : x \notin B\}$$

It is the set of elements that belong to the first but not to the second set. The symbol “ $\setminus$ ” can also be written “ $-$ ” by analogy with arithmetic subtraction.

Example 1.17: intersection, union and difference

Given the sets $A = \{1, 2, 3, 4\}$ and $B = \{3, 4, 5\}$, we have

$$A \cap B = \{3, 4\}, \quad A \cup B = \{1, 2, 3, 4, 5\},$$

$$A \setminus B = \{1, 2\}, \quad B \setminus A = \{5\}.$$

In particular $A \setminus B \neq B \setminus A$: in the difference the order of the two sets matters.

1.4.1 Complementary sets and disjoint sets

Example 1.18: universal sets

For example, in questions of arithmetic we could have $U = \mathbb{N}$; if we consider sets made up only of integers it is natural to choose $U = \mathbb{Z}$, while in questions of analysis we could have $U = \mathbb{R}$.

Definition 1.21: of set complementation and complementary sets

The **complementation** of a set $A \subseteq U$ is the set defined by:

$$\overline{A} = \{x \in U : x \notin A\}$$

This set is called the **complement** (complementary set) of A with respect to U .

- By the definition of difference, $U \setminus A = \{x \in U : x \notin A\}$, hence the complement of A is the difference between the universal set and A :

$$\overline{A} = U \setminus A.$$

Proposition 1.22: properties of the complement

For every set $A \subseteq U$, where U is the universal set, we have the following relations:

$$\begin{aligned}\overline{\overline{A}} &= A \\ A \cap \overline{A} &= \emptyset \\ A \cup \overline{A} &= U \\ \overline{\overline{U}} &= \emptyset \\ \overline{\emptyset} &= U\end{aligned}$$

Proof. All the sets involved are subsets of U : hence it is enough to consider the elements $x \in U$. We prove the five relations.

- Let $x \in U$. We have $x \in \overline{\overline{A}}$ if and only if $x \notin \overline{A}$, that is, if and only if it is not true that $x \notin A$, that is, if and only if $x \in A$. Hence $\overline{\overline{A}} = A$.
- If there existed $x \in A \cap \overline{A}$, we would have $x \in A$ and $x \in \overline{A}$, that is, $x \notin A$: this is impossible. Hence $A \cap \overline{A}$ has no elements, that is, $A \cap \overline{A} = \emptyset$.
- By the definition of union, $A \cup \overline{A}$ is made up of elements of U , hence $A \cup \overline{A} \subseteq U$. Conversely, let $x \in U$: if $x \in A$, then $x \in A \cup \overline{A}$; if $x \notin A$, then $x \in \overline{A}$ and hence $x \in A \cup \overline{A}$. Hence $U \subseteq A \cup \overline{A}$ and, by double inclusion, $A \cup \overline{A} = U$.
- $\overline{\overline{U}} = \{x \in U : x \notin U\}$ has no elements, because no element of U can fail to belong to U . Hence $\overline{\overline{U}} = \emptyset$.
- $\overline{\emptyset} = \{x \in U : x \notin \emptyset\}$, and every $x \in U$ satisfies $x \notin \emptyset$. Hence $\overline{\emptyset} = U$.

□

Proposition 1.23: identity laws

For every set $A \subseteq U$, where U is the universal set, we have:

$$\begin{aligned}A \cap \emptyset &= \emptyset \\ A \cap U &= A \\ A \cup \emptyset &= A \\ A \cup U &= U\end{aligned}$$

Proof. We prove the four relations.

- No x can belong both to A and to $\emptyset$, because $\emptyset$ has no elements. Hence $A \cap \emptyset = \emptyset$.
- If $x \in A \cap U$, then $x \in A$. Conversely, if $x \in A$, then $x \in U$ because $A \subseteq U$, and hence $x \in A \cap U$. By double inclusion $A \cap U = A$.
- We have $x \in A \cup \emptyset$ if and only if $x \in A$ or $x \in \emptyset$. The second possibility never occurs, hence $x \in A \cup \emptyset$ if and only if $x \in A$, that is, $A \cup \emptyset = A$.

- By the definition of union, $A \cup U$ is made up of elements of U , hence $A \cup U \subseteq U$. Conversely, if $x \in U$, then $x \in A \cup U$. By double inclusion $A \cup U = U$.

□

Definition 1.24: of disjoint sets

Two sets A and B are **disjoint** if they have no elements in common:

$$A \cap B = \emptyset$$

Example 1.19: disjoint sets

The sets $\{1, 2\}$ and $\{3, 4\}$ are disjoint, while $\{1, 2\}$ and $\{2, 3\}$ are not, because $\{1, 2\} \cap \{2, 3\} = \{2\}$. By the properties of the complement, every set $A \subseteq U$ and its complement $\bar{A}$ are disjoint.

Disjoint sets allow us to “split” a set into pieces that do not overlap. Given a family $\mathcal{F}$ of sets, the **union of the family** is the set of elements that belong to at least one of the sets of the family; it is denoted by

$$\bigcup_{B \in \mathcal{F}} B.$$

For a family made up of two sets B and C we recover the union $B \cup C$.

Definition 1.25: of partition

Given a set A , a family $\mathcal{F}$ of subsets of A is a **partition** of A if:

- every set of the family is nonempty: $B \neq \emptyset$ for every $B \in \mathcal{F}$;
- the sets of the family are pairwise disjoint: if $B, C \in \mathcal{F}$ and $B \neq C$, then $B \cap C = \emptyset$;
- their union is A :

$$A = \bigcup_{B \in \mathcal{F}} B,$$

that is, every element of A belongs to at least one set of the family.

The sets of the family $\mathcal{F}$ are called the **blocks** of the partition.

Observation 1.26

A family $\mathcal{F}$ of nonempty subsets of a set A is a partition of A if and only if every element of A belongs to **exactly one** set of the family.

Proof. ($\implies$) Let $\mathcal{F}$ be a partition of A and let $x \in A$. Since the union of the family is A , the element x belongs to at least one set of the family. If it belonged to two different sets $B, C \in \mathcal{F}$, we would have $x \in B \cap C$, against the fact that $B \cap C = \emptyset$. Hence x belongs to exactly one set of the family.

($\impliedby$) Suppose that every element of A belongs to exactly one set of the family. The sets of the family are nonempty by hypothesis. Every element of A belongs to at least one set of the family, hence $A \subseteq \bigcup_{B \in \mathcal{F}} B$; conversely, every element of the union belongs to a subset of A and hence to A : by double inclusion the union is A . Finally, if two different sets $B, C \in \mathcal{F}$ had a common element x , this element of A would belong to two sets of the family, against the hypothesis: hence $B \cap C = \emptyset$. $\square$


Example 1.20: partitions

- The family $\{\{1, 2\}, \{3\}, \{4, 5, 6\}\}$ is a partition of $A = \{1, 2, 3, 4, 5, 6\}$: the three sets are nonempty and every element of A belongs to exactly one of them.
- The family $\{\{1, 2\}, \{2, 3\}\}$ is not a partition of $\{1, 2, 3\}$: the element 2 belongs to two sets of the family.
- The family $\{\{1\}, \{2\}\}$ is not a partition of $\{1, 2, 3\}$: the element 3 does not belong to any set of the family.
- With universal set $\mathbb{Z}$, the set E of even numbers and its complement $\overline{E}$, that is, the set of odd numbers, form a partition of $\mathbb{Z}$: the two sets are nonempty ($0 \in E$ and $1 \in \overline{E}$) and, by the properties of the complement, $E \cap \overline{E} = \emptyset$ and $E \cup \overline{E} = \mathbb{Z}$.


1.4.2 Venn diagrams

Venn diagrams are graphical representations in which sets are represented as regions of the plane

- Venn diagrams of intersection, union and difference:



- Venn diagram of the complement of a set:



1.4.3 Properties of operations on sets

Observation 1.27: properties of intersection

Given three sets A , B and C , intersection has the following properties:

- **Commutative:**

$$A \cap B = B \cap A$$

- **Associative:**

$$A \cap (B \cap C) = (A \cap B) \cap C$$


- **Idempotence:**


$$A \cap A = A$$

Proof. We prove the three properties by comparing the elements of the two sides.

- **Commutative:** $x \in A \cap B$ if and only if $x \in A$ and $x \in B$, that is, if and only if $x \in B$ and $x \in A$, that is, if and only if $x \in B \cap A$. Hence $A \cap B = B \cap A$.
- **Associative:** $x \in A \cap (B \cap C)$ if and only if $x \in A$ and $x \in B \cap C$, that is, if and only if x belongs to all three sets A , B and C , that is, if and only if $x \in A \cap B$ and $x \in C$, that is, if and only if $x \in (A \cap B) \cap C$.
- **Idempotence:** $x \in A \cap A$ if and only if $x \in A$ and $x \in A$, that is, if and only if $x \in A$. Hence $A \cap A = A$.

The graphical proof of the associative property of intersection is the following: in the first row we build $(A \cap B) \cap C$, in the second row $A \cap (B \cap C)$, and the two sets obtained on the right coincide.




Observation 1.28: properties of union

Given three sets A , B and C , union has the following properties:

- **Commutative:**

$$A \cup B = B \cup A$$

- **Associative:**

$$A \cup (B \cup C) = (A \cup B) \cup C$$


- **Idempotence:**


$$A \cup A = A$$

Proof. We prove the three properties by comparing the elements of the two sides.

- **Commutative:** $x \in A \cup B$ if and only if $x \in A$ or $x \in B$, that is, if and only if $x \in B$ or $x \in A$, that is, if and only if $x \in B \cup A$. Hence $A \cup B = B \cup A$.
- **Associative:** $x \in A \cup (B \cup C)$ if and only if $x \in A$ or $x \in B \cup C$, that is, if and only if x belongs to at least one of the three sets A , B and C , that is, if and only if $x \in A \cup B$ or $x \in C$, that is, if and only if $x \in (A \cup B) \cup C$.
- **Idempotence:** $x \in A \cup A$ if and only if $x \in A$ or $x \in A$, that is, if and only if $x \in A$. Hence $A \cup A = A$.

The graphical proof of the associative property of union is the following: in the first row we build $(A \cup B) \cup C$, in the second row $A \cup (B \cup C)$, and the two sets obtained on the right coincide.




Observation 1.29: distributive properties (linking union and intersection)


Given three sets A , B and C we have:

$$A \cap (B \cup C) = (A \cap B) \cup (A \cap C)$$


$$A \cup (B \cap C) = (A \cup B) \cap (A \cup C).$$

The distributive properties link union and intersection to each other.

Proof. The graphical proof of the first property is the following: in the first row we build the left-hand side $A \cap (B \cup C)$, in the second row the right-hand side $(A \cap B) \cup (A \cap C)$, and the two sets obtained on the right coincide.



The graphical proof of the second property is the following: in the first row we build the left-hand side $A \cup (B \cap C)$, in the second row the right-hand side $(A \cup B) \cap (A \cup C)$.

**Proposition 1.30: absorption laws**

Given two sets A and B we have:

$$A \cap (A \cup B) = A$$

$$A \cup (A \cap B) = A$$

Proof. We prove the two laws by double inclusion.

- If $x \in A \cap (A \cup B)$, then $x \in A$. Conversely, if $x \in A$, then $x \in A \cup B$, and hence $x \in A \cap (A \cup B)$. Hence $A \cap (A \cup B) = A$.
- If $x \in A$, then $x \in A \cup (A \cap B)$. Conversely, if $x \in A \cup (A \cap B)$, then $x \in A$ or $x \in A \cap B$; in the second case, again, $x \in A$. Hence $A \cup (A \cap B) = A$.

□

Proposition 1.31: inclusion, union and intersection

Given two sets A and B , the following statements are equivalent:

$$A \subseteq B, \quad A \cap B = A, \quad A \cup B = B.$$

Proof. We prove that $A \subseteq B$ is equivalent to each of the other two statements.

- $(A \subseteq B \implies A \cap B = A)$ Every element of $A \cap B$ belongs to A . Conversely, if $x \in A$, then $x \in B$ because $A \subseteq B$, and hence $x \in A \cap B$. By double inclusion

$$A \cap B = A.$$

- $(A \cap B = A \implies A \subseteq B)$ If $x \in A$, then $x \in A \cap B$, because $A \cap B = A$, and hence $x \in B$.
- $(A \subseteq B \implies A \cup B = B)$ Every element of B belongs to $A \cup B$. Conversely, if $x \in A \cup B$, then $x \in A$ or $x \in B$; in the first case $x \in B$ because $A \subseteq B$. In every case $x \in B$, and by double inclusion $A \cup B = B$.
- $(A \cup B = B \implies A \subseteq B)$ If $x \in A$, then $x \in A \cup B$, and hence $x \in B$ because $A \cup B = B$.

□



Proposition 1.32: De Morgan's laws (first version)

Given three sets A, B and C we have:


$$A \setminus (B \cap C) = (A \setminus B) \cup (A \setminus C)$$

$$A \setminus (B \cup C) = (A \setminus B) \cap (A \setminus C)$$

Proof. The graphical proof of the first property is the following: in the first row we build the left-hand side $A \setminus (B \cap C)$, in the second row the right-hand side $(A \setminus B) \cup (A \setminus C)$, and the two sets obtained on the right coincide.



The graphical proof of the second property is the following: in the first row we build the left-hand side $A \setminus (B \cup C)$, in the second row the right-hand side $(A \setminus B) \cap (A \setminus C)$.



Proposition 1.33: De Morgan's laws (second version)

Given the sets $B, C \subseteq U$, we have

$$\overline{B \cap C} = \overline{B} \cup \overline{C}$$


$$\overline{B \cup C} = \overline{B} \cap \overline{C}$$

Proof. These laws can be derived by setting A equal to U in the previous De Morgan's laws and recalling that $\overline{B} = U \setminus B$ and $\overline{C} = U \setminus C$, as follows:

$$\underbrace{U \setminus (B \cap C)}_{=\overline{B \cap C}} = \underbrace{(U \setminus B)}_{=\overline{B}} \cup \underbrace{(U \setminus C)}_{=\overline{C}}$$

$$\underbrace{U \setminus (B \cup C)}_{=\overline{B \cup C}} = \underbrace{(U \setminus B)}_{=\overline{B}} \cap \underbrace{(U \setminus C)}_{=\overline{C}}$$

Proof. The graphical proof of the first property is the following: in the first row we build the left-hand side $\overline{B \cap C}$ starting from $B \cap C$, in the second row the right-hand side $\overline{B} \cup \overline{C}$, and the two sets obtained on the right coincide.



- The set of elements that belong to a set B or to a set C but not to both (meaning “or” in the exclusive sense) is called the **symmetric difference** of B and C and is denoted by $B \triangle C$:


$$B \triangle C = (B \setminus C) \cup (C \setminus B).$$

It is also obtained by setting $U = B \cup C$ and taking the complement of the intersection

with respect to this universal set:

$$B \Delta C = (B \cup C) \setminus (B \cap C).$$

Proof. We prove that the two expressions of the symmetric difference coincide. We have $x \in (B \setminus C) \cup (C \setminus B)$ if and only if $(x \in B \text{ and } x \notin C)$ or $(x \in C \text{ and } x \notin B)$, that is, if and only if x belongs to only one of the two sets B and C . On the other hand, $x \in (B \cup C) \setminus (B \cap C)$ if and only if x belongs to at least one of the two sets but not to both, that is, again, if and only if x belongs to only one of the two sets. Hence the two sets have the same elements. $\square$



1.4.4 Properties of the cardinality of sets

Proposition 1.34: cardinality of the union of disjoint sets

If A and B are two disjoint finite sets, that is, $|A \cap B| = 0$, then

$$|A \cup B| = |A| + |B|.$$

More generally, if $A_1, A_2, \dots, A_m$ are pairwise disjoint finite sets, then

$$|A_1 \cup A_2 \cup \dots \cup A_m| = |A_1| + |A_2| + \dots + |A_m|.$$

Proof. Every element of $A \cup B$ belongs to A or to B , and no element belongs to both, because $A \cap B = \emptyset$. Counting first the $|A|$ elements of A and then the $|B|$ elements of B we therefore count every element of $A \cup B$ exactly once: $|A \cup B| = |A| + |B|$.

In the same way, every element of $A_1 \cup A_2 \cup \dots \cup A_m$ belongs to at least one of the sets $A_1, A_2, \dots, A_m$ and, since the sets are pairwise disjoint, to only one of them. Counting one after the other the elements of $A_1, A_2, \dots, A_m$ we therefore count every element of the union exactly once. $\square$

Proposition 1.35: cardinality of the union

For any two finite sets A and B , we have

$$|A \cup B| = |A| + |B| - |A \cap B|.$$

Proof. Every element of $A \cup B$ belongs to exactly one of the three sets

$$A \setminus B, \quad A \cap B, \quad B \setminus A :$$

if it belongs both to A and to B it lies in $A \cap B$, if it belongs only to A it lies in $A \setminus B$, if it belongs only to B it lies in $B \setminus A$. The three sets are therefore pairwise disjoint and their union is $A \cup B$, and by the cardinality of the union of disjoint sets

$$|A \cup B| = |A \setminus B| + |A \cap B| + |B \setminus A|.$$

In the same way, every element of A belongs to exactly one of the two disjoint sets $A \setminus B$ and $A \cap B$, and every element of B belongs to exactly one of the two disjoint sets $B \setminus A$ and $A \cap B$, hence

$$|A| = |A \setminus B| + |A \cap B|, \quad |B| = |B \setminus A| + |A \cap B|.$$

Adding the last two equalities we obtain

$$|A| + |B| = |A \setminus B| + |A \cap B| + |B \setminus A| + |A \cap B| = |A \cup B| + |A \cap B|,$$

that is, $|A \cup B| = |A| + |B| - |A \cap B|$. □

Proposition 1.36: inequality for the cardinality of the union

For any two finite sets A and B , we have

$$|A \cup B| \leq |A| + |B|.$$

Proof. By the cardinality of the union we have $|A \cup B| = |A| + |B| - |A \cap B|$. Since $|A \cap B| \geq 0$, we obtain

$$|A \cup B| = |A| + |B| - |A \cap B| \leq |A| + |B|.$$

□

Proposition 1.37: cardinality of a subset

Given two finite sets A and B with $A \subseteq B$, we have

$$|A| \leq |B|.$$

If moreover $A \subsetneq B$, then $|A| < |B|$.

Proof. Since $A \subseteq B$, every element of B belongs to exactly one of the two disjoint sets A and $B \setminus A$, and every element of these two sets belongs to B : hence $B = A \cup (B \setminus A)$ and, by the cardinality of the union of disjoint sets,

$$|B| = |A| + |B \setminus A| \geq |A|.$$

If moreover $A \subsetneq B$, then $A \neq B$ and hence, by the antisymmetric property of inclusion, $B \subseteq A$ cannot hold: there exists an element $b \in B$ with $b \notin A$, that is, $b \in B \setminus A$. Then $|B \setminus A| \geq 1$ and

$$|B| = |A| + |B \setminus A| \geq |A| + 1 > |A|.$$

□

1.4.5 Ordered pairs and Cartesian product

There is another operation on sets, which can be performed on any two sets (i.e., two sets not necessarily contained in the same universal set). To introduce it we need the concept of *ordered pair*: unlike the set $\{a, b\}$, in which the order of the elements is irrelevant, in an ordered pair it matters which element comes first.

Definition 1.38: of ordered pair

Given two elements a and b , the **ordered pair** (a, b) is the set

$$(a, b) = \{\{a\}, \{a, b\}\}.$$

The element a is called the **first component** and the element b the **second component** of the pair.

This definition, due to Kuratowski, expresses the ordered pair using only sets. What matters about the ordered pair is the following property.

Proposition 1.39: characteristic property of ordered pairs

Given the elements a, b, c and d , we have

$$(a, b) = (c, d) \iff a = c \text{ and } b = d.$$

Proof. ($\Leftarrow$) If $a = c$ and $b = d$, the sets $\{\{a\}, \{a, b\}\}$ and $\{\{c\}, \{c, d\}\}$ are written with the same elements, hence they are equal.

($\Rightarrow$) Suppose $\{\{a\}, \{a, b\}\} = \{\{c\}, \{c, d\}\}$ and distinguish two cases.

- If $a = b$, then $\{a, b\} = \{a\}$ and the pair $(a, b) = \{\{a\}\}$ has only one element. Hence $\{\{c\}, \{c, d\}\}$ also has $\{a\}$ as its only element: from $\{c\} = \{a\}$ it follows that $c = a$, and from $\{c, d\} = \{a\}$ it follows that $d = a$. Hence $a = c$ and $b = a = d$.
- If $a \neq b$, the pair (a, b) has two distinct elements: $\{a\}$, which has one element, and $\{a, b\}$, which has two. Then also $c \neq d$, otherwise (c, d) would have only one

element. Since $\{a\} \in \{\{c\}, \{c, d\}\}$ and $\{a\}$ has only one element, while $\{c, d\}$ has two, we must have $\{a\} = \{c\}$, that is, $a = c$. Similarly $\{a, b\}$, which has two elements, cannot be equal to $\{c\}$, hence $\{a, b\} = \{c, d\} = \{a, d\}$. Since $b \in \{a, d\}$ and $b \neq a$, we obtain $b = d$.

□

- In particular, if $a \neq b$ then $(a, b) \neq (b, a)$: if $(a, b) = (b, a)$ held, the characteristic property would give $a = b$. Instead the sets $\{a, b\}$ and $\{b, a\}$ are always equal. If $a = b$, the two pairs (a, b) and (b, a) are the same pair.

Definition 1.40: of Cartesian product

Given two (not necessarily distinct) sets A and B , the set consisting of all ordered pairs (a, b) , with $a \in A$ and $b \in B$, is called the **Cartesian product** of A and B and is denoted by the symbol $A \times B$:

$$A \times B = \{(a, b) : a \in A \text{ and } b \in B\}.$$

Proposition 1.41: cardinality of the Cartesian product

When A and B are two finite sets, the cardinality of their Cartesian product is:

$$|A \times B| = |A| \cdot |B|.$$

Proof. If $A = \emptyset$ there is no pair with first component in A , hence $A \times B = \emptyset$ and $|A \times B| = 0 = 0 \cdot |B|$. Otherwise let $a_1, a_2, \dots, a_m$ be the elements of A , with $m = |A|$, and for every $i \in \{1, 2, \dots, m\}$ consider the set of pairs with first component a_i :

$$R_i = \{(a_i, b) : b \in B\}.$$

- Every pair of $A \times B$ belongs to exactly one of the sets $R_1, R_2, \dots, R_m$, the one corresponding to its first component: by the characteristic property, pairs with different first components are different. Hence the sets $R_1, R_2, \dots, R_m$ are pairwise disjoint and their union is $A \times B$.
- Every set R_i has $|B|$ elements: by the characteristic property, $(a_i, b) = (a_i, b')$ if and only if $b = b'$, hence different elements of B give different pairs.

By the cardinality of the union of disjoint sets

$$|A \times B| = |R_1| + |R_2| + \dots + |R_m| = \underbrace{|B| + |B| + \dots + |B|}_{m \text{ times}} = |A| \cdot |B|.$$

□

Example 1.21: Cartesian product

$$\begin{aligned}\{a, b\} \times \{a, b, c\} &= \{(a, a), (a, b), (a, c), (b, a), (b, b), (b, c)\} \\ |\{a, b\}| &= 2, \quad |\{a, b, c\}| = 3, \quad |\{a, b\} \times \{a, b, c\}| = 2 \cdot 3 = 6\end{aligned}$$

The Cartesian product is not commutative: the pair (c, a) belongs to $\{a, b, c\} \times \{a, b\}$, but it does not belong to $\{a, b\} \times \{a, b, c\}$, because $c \notin \{a, b\}$. Hence $\{a, b, c\} \times \{a, b\} \neq \{a, b\} \times \{a, b, c\}$.

The Cartesian product extends to more than two sets.

Definition 1.42: of ordered n -tuple and of Cartesian product of n sets

Given $n \geq 3$ elements $a_1, a_2, \dots, a_n$, the **ordered n -tuple** $(a_1, a_2, \dots, a_n)$ is the ordered pair

$$(a_1, a_2, \dots, a_n) = ((a_1, a_2, \dots, a_{n-1}), a_n).$$

Given $n \geq 2$ sets $A_1, A_2, \dots, A_n$, their **Cartesian product** is the set of n -tuples

$$A_1 \times A_2 \times \dots \times A_n = \{(a_1, a_2, \dots, a_n) : a_i \in A_i \text{ for every } i \in \{1, 2, \dots, n\}\}.$$

- Applying repeatedly the characteristic property of ordered pairs we obtain

$$(a_1, a_2, \dots, a_n) = (b_1, b_2, \dots, b_n) \iff a_i = b_i \text{ for every } i \in \{1, 2, \dots, n\}.$$

Proposition 1.43: cardinality of the Cartesian product of n sets

If $A_1, A_2, \dots, A_n$ are finite sets, then

$$|A_1 \times A_2 \times \dots \times A_n| = |A_1| \cdot |A_2| \cdots |A_n|.$$

Proof. For $n = 2$ it is the cardinality of the Cartesian product of two sets. For $n \geq 3$, by the definition of n -tuple the n -tuples $(a_1, a_2, \dots, a_n)$ are exactly the pairs $((a_1, \dots, a_{n-1}), a_n)$ with $(a_1, \dots, a_{n-1}) \in A_1 \times \dots \times A_{n-1}$ and $a_n \in A_n$, that is,

$$A_1 \times A_2 \times \dots \times A_n = (A_1 \times A_2 \times \dots \times A_{n-1}) \times A_n.$$

By the cardinality of the Cartesian product of two sets

$$|A_1 \times A_2 \times \dots \times A_n| = |A_1 \times A_2 \times \dots \times A_{n-1}| \cdot |A_n|.$$

Repeating the same argument on $A_1 \times A_2 \times \dots \times A_{n-1}$, then on $A_1 \times A_2 \times \dots \times A_{n-2}$, and so on down to $A_1 \times A_2$, we obtain $|A_1| \cdot |A_2| \cdots |A_n|$. $\square$

- The Cartesian product of n sets equal to A is denoted by

$$A^n = \underbrace{A \times A \times \dots \times A}_{n \text{ times}}$$

and, if A is finite, by the previous proposition it has cardinality $|A^n| = |A|^n$. For example $\{0, 1\}^3$ has $2^3 = 8$ elements: the triples of 0 and 1 correspond to the “in” or “out” choices of the binary tree with which we built the power set of a set with 3 elements.

- A typical use of the Cartesian product is $\mathbb{R} \times \mathbb{R}$, which is abbreviated by the symbol $\mathbb{R}^2$ and denotes the set of ordered pairs of real numbers.
- Similarly, $\mathbb{R}^n$ (abbreviation of the Cartesian product of n sets equal to $\mathbb{R}$) is the set of *ordered n -tuples of real numbers*

$$\mathbb{R}^n = \{ (x_1, x_2, \dots, x_n) : x_i \in \mathbb{R}, i \in \{1, 2, \dots, n\} \}$$

Example 1.22: subsets of the plane

When we study subsets of the plane we choose $U = \mathbb{R}^2$ as universal set. For example, the set of points of the plane with both coordinates positive (the *first quadrant*) is

$$Q_1 = \{ (x_1, x_2) \in \mathbb{R}^2 : x_1 > 0 \text{ and } x_2 > 0 \}.$$

Since $(x_1, x_2) \in Q_1$ if and only if $x_1 \in \mathbb{R}_{>0}$ and $x_2 \in \mathbb{R}_{>0}$, we have $Q_1 = \mathbb{R}_{>0} \times \mathbb{R}_{>0}$.

1.5 Further topics

1.5.1 Russell’s paradox

“can a set be an element of itself or not?”

- For example, the set of all books in a library is not an element of itself (a set of books is not a book). On the other hand, reasoning naively, the collection of all sets with more than 20 elements would seem to be an element of itself, since it certainly has more than 20 elements. We will see at the end of this section that in the formal definition of sets this collection is not a set.
- Following this reasoning, two categories of sets can be defined:
 1. sets that are not elements of themselves
 2. sets that are elements of themselves

If we consider the set of all sets that are not elements of themselves, is it an element of itself or not?

Let us call this set S ; two hypotheses can be made:

- 1 If we suppose $S \in S$, then S contains itself as an element and therefore does not belong to S (since by definition a set belongs to S only if it does not contain itself as an element). Hence $S \notin S$, and we have a contradiction. We conclude that the hypothesis must be wrong.
- 2 If we suppose $S \notin S$, then S does not contain itself as an element and therefore belongs to S (since by definition a set belongs to S if it does not contain itself as an element). Hence $S \in S$ and we have another contradiction. We conclude that this hypothesis must be wrong too!

Russell's paradox: The set of all sets that do not belong to themselves belongs to itself if and only if it does not belong to itself.

- The formal definition of the concept of set is based on the **Zermelo-Fraenkel system of axioms**, abbreviated as **ZF**. This system of axioms includes the standard axioms of axiomatic set theory on which, together with the *axiom of choice*, all of ordinary mathematics is based.
- The **axiom of pairing** states that “Given two objects, there exists a set whose elements are the two objects”.
- The **axiom of regularity** states that “Every non-empty set A contains an element disjoint from A ”.

Observation 1.44

No set is an element of itself.

Proof. Let A be a set. By the axiom of pairing, applied to the two objects A and A , there exists the set $\{A, A\}$, which coincides with $\{A\}$ since sets do not contain repeated objects: hence $\{A\}$ is a set (it is a special case of a pair), and it is non-empty.

We now apply the axiom of regularity to the set $\{A\}$: there must exist an element of $\{A\}$ disjoint from $\{A\}$. Since the only element of $\{A\}$ is A , A is disjoint from $\{A\}$, that is, $A \cap \{A\} = \emptyset$.

If $A \in A$ held, since also $A \in \{A\}$, we would have $A \in A \cap \{A\}$, against the fact that $A \cap \{A\} = \emptyset$ (definition of disjoint sets). Hence $A \notin A$. $\square$

- In particular, the collection of all sets with more than 20 elements is not a set in ZF: if it were, it would have more than 20 elements and would therefore be an element of itself, against the remark just proved.
- The other ZF axioms are not part of the basic course in Mathematical Analysis.

1.5.2 A further proof via the Archimedean property

- A further proof of

$$0.\overline{9} = 1$$

starts from the assumption that two numbers are equal if and only if their difference is equal to zero, and it is based on computing the value of $1 - 0.\overline{9}$.

- This proof is based on the fact that 0 is the only non-negative number less than all the reciprocals of the positive integers, or equivalently that there is no number greater than every natural number. This is the **Archimedean property**, which holds for the rational and the real numbers.

The **Archimedean property** states that for every real number x there exists a natural number n such that:

$$n > x$$

- Equivalently: for every real number $\varepsilon > 0$ there exists a natural number $n > 0$ such that $\frac{1}{n} < \varepsilon$ (it is enough to take n greater than $\frac{1}{\varepsilon}$). As a consequence, the only non-negative number that is less than or equal to $\frac{1}{n}$ for every n is zero: this is the form in which the property is used in the proof below.
- It is not an additional axiom: in the chapter *Ordered fields, supremum/infimum and the completeness axiom* we will prove it starting from the supremum property of the real numbers.

Proof. We write the number $0.999\dots$ with n digits after the decimal point as $0.(9)_n$, hence $0.(9)_1 = 0.9$, $0.(9)_2 = 0.99$, $0.(9)_3 = 0.999$, and so on.

Given $\frac{1}{10^n} = 0.0\dots 01$, with n digits after the decimal point, the addition rules for decimal numbers imply

$$0.(9)_n + \frac{1}{10^n} = 1 \quad \text{moreover} \quad 0.(9)_n < 1, \forall n \in \mathbb{N}.$$

We must prove that 1 is the smallest number that is not less than all the $0.(9)_n$. For this it is enough to prove that, if a number x is not greater than 1 and not less than all the $0.(9)_n$, then $x = 1$.

So let x be such that

$$0.(9)_n \leq x \leq 1$$

for every positive integer n . Hence

$$1 - 1 \leq 1 - x \leq 1 - 0.(9)_n$$

which, using basic arithmetic and the first equality established above, simplifies to

$$0 \leq 1 - x \leq \frac{1}{10^n}$$

This implies that the difference between 1 and x is less than the reciprocal of any positive integer. Hence this difference must be zero, and therefore $x = 1$; which in turn implies

$$0.999 \dots = 1$$

□

Basics of logic and proof techniques

2.1 Logical symbols

- a) the symbol “ $\forall$ ” is called the *universal quantifier* and is read “for every”, “for all”, “for each”
- b) the symbol “ $\exists$ ” is called the *existential quantifier* and is read “there exists”, “there exist”
- c) the symbol “ $\Rightarrow$ ” is called *logical implication* and is read “implies” or “if ... then”

- a) the symbol “ $:$ ” is read “such that”
- b) the symbol “ $\in$ ” is read “belongs to”
- c) the symbol “ $\notin$ ” is read “does not belong to”

- a) the symbol “ $\vee$ ” is called *logical disjunction* and is read “or”
- b) the symbol “ $\wedge$ ” is called *logical conjunction* and is read “and”
- c) the symbol “ $\neg$ ” is called *logical negation* and is read “not”.

2.2 Universal implications and proofs

Predicates (or properties) and **propositions** (or statements)

- Consider the following assertion:

“the natural number n is odd” (2.1)

and let us ask whether it is true. Obviously the answer is: “it depends on n ”. Indeed, in (2.1) the symbol n represents a variable that can take different values and make the assertion **true** or **false**.

A sentence of this kind is called a **predicate** (or *property*): its truth or falsity depends on the values of the **variable(s)** appearing in it.

- Now consider the following assertion:

$$\text{“for every natural number } n, \text{ if } n \text{ is odd then } n^2 \text{ is odd”} \quad (2.2)$$

which we can write more formally as follows:

$$\forall n \in \mathbb{N} \ (n \text{ odd} \Rightarrow n^2 \text{ odd}) \quad (2.3)$$

In this case we say that the variable n is not free, but bound by the *quantifier* $\forall$. As a consequence, (2.3) is true or false “once and for all” and is called a **proposition** (or *statement*).

In particular, the components of (2.3) are the set $\mathbb{N}$, two predicates defined on $\mathbb{N}$ given by

$$p(n) : \text{“}n \text{ odd”} \quad \text{and} \quad q(n) : \text{“}n^2 \text{ odd”}$$

and the implication

$$p(n) \Rightarrow q(n).$$

Definition 2.1: of universal implication

In general, a statement involving a **set** A , **two predicates** $p(x)$ and $q(x)$ whose argument x ranges over A , and the **logical structure**:

$$\forall x \in A \ (p(x) \Rightarrow q(x)) \quad (2.4)$$

is called a **universal implication**.

Most **theorems** consist of **universal implications**, in which the predicate $p(x)$ plays the role of the **hypothesis** and the predicate $q(x)$ plays the role of the **thesis** (conclusion).

- In particular, (2.3) is a proposition (or a statement):

Proposition 2.2

$$\forall n \in \mathbb{N} \ (n \text{ odd} \Rightarrow n^2 \text{ odd}) \quad (2.5)$$

In this case one is easily convinced that the proposition is true, but how can we prove it rigorously?

- For example, is it enough to observe that 3 is odd and $3^2 = 9$ is odd to claim that the proposition is true? Certainly not, since the proposition requires the universal implication to hold for every natural number. However, there are infinitely many odd numbers: how can we prove a universal implication for infinitely many numbers?
- The key procedure is this: we consider a generic n satisfying the hypothesis (being odd) and we prove that n satisfies the thesis (its square is odd).

Let us see how to proceed to prove the previous proposition:

Proof. Let n be odd; we prove that then n^2 is odd.

Any odd number can be written in the form $2k + 1$, for a suitable $k \in \mathbb{N}$. We also observe that $2k$ is an even number for any $k \in \mathbb{N}$.

So let $n = 2k + 1$ be an odd number ($k \in \mathbb{N}$); we therefore need to write n^2 as an even integer plus one. We have:

$$n^2 = (2k + 1)^2 = 4k^2 + 4k + 1 = 2(2k^2 + 2k) + 1.$$

Since $2(2k^2 + 2k)$ is an even integer, n^2 is odd. □

To prove the correctness of a universal implication such as (2.4), we consider a generic x satisfying the hypothesis $p(x)$ and we try to prove that the thesis $q(x)$ is true.

- Let us now prove a similar relation for even numbers:

Proposition 2.3

$$\forall n \in \mathbb{N} \quad (n \text{ even} \Rightarrow n^2 \text{ even}) \quad (2.6)$$

Proof. Let n be even; we prove that then n^2 is even.

So let $n = 2k$ be an even number ($k \in \mathbb{N}$); we therefore need to write n^2 as an even integer. We have:

$$n^2 = (2k)^2 = 4k^2 = 2(2k^2)$$

Since $2(2k^2)$ is an even integer, n^2 is even. □

2.2.1 Counterexamples

Counterexamples are an important technique for proving the **falsity** of a universal implication.

- Let us ask, for example, whether the following universal implication is true or false:

$$\forall n \in \mathbb{N} \ (n \text{ prime} \Rightarrow n \text{ odd}) \quad (2.7)$$

A moment's reflection shows that **this proposition is false**. Indeed, the number 2 is prime but it is even.

To claim that a universal implication is true, a proof is needed (an example is not enough), whereas to prove that a universal implication is false, one counterexample is enough.

- The universal implication requires that every x satisfying the hypothesis also satisfies the thesis: therefore, if we find even a single example of x that satisfies the hypothesis but not the thesis, this means that the universal implication is false. Not “false in one case”, but simply “false”, because the universal implication is true or false once and for all.

Definition 2.4: of counterexample

*In general, an example that satisfies the hypothesis but not the thesis of a universal implication, and that therefore proves its falsity, is called a **counterexample**.*

- The formal proof that the previous universal implication is false is the following:

Proof. The number 2 is a counterexample to the universal implication: “For every natural number n , if n is prime then n is odd”.

□

The negation of the proposition

“for every $x \in A$, if $p(x)$ holds then $q(x)$ holds”

$$\forall x \in A \ (p(x) \Rightarrow q(x))$$

is the proposition

“there exists $x \in A$ for which $p(x)$ holds and $q(x)$ does not hold”

$$\exists x \in A \ (p(x) \wedge \neg q(x))$$

This particular x is a **counterexample**.

2.3 Law of contraposition

- It is an indirect proof technique

The universal implication

$$\forall x \in A \quad (p(x) \Rightarrow q(x)) \quad (2.8)$$

is logically equivalent to

$$\forall x \in A \quad (\text{not } q(x) \Rightarrow \text{not } p(x)) \quad (2.9)$$

The second implication is called the **contrapositive** of the first.

- For example: since we know that the universal implication (proposition 2.5) holds

$$\forall n \in \mathbb{N} \quad (n \text{ odd} \Rightarrow n^2 \text{ odd})$$

the following proposition holds:

Proposition 2.5

$$\forall n \in \mathbb{N} \quad (n^2 \text{ even} \Rightarrow n \text{ even}) \quad (2.10)$$

Proof. Let n^2 be even; we prove that then n is even. We prove (2.10) relying on the truth of (2.5).

If n is not even then it is odd and therefore n^2 is odd by (2.5). This case contradicts the hypothesis that n^2 is even (hence it cannot occur).

Consequently n is even and (2.10) is proved. □

- The reasoning used in the previous proof has general validity, and shows precisely that if (2.8) is true then (2.9) is true; moreover, if the second is true then the first is true (because “not not $p(x)$ ” is logically equivalent to $p(x)$), so the two are logically equivalent.

The equivalence between (2.8) and (2.9) is called the **law of contraposition**. It is a method of *indirect proof* that consists in proving (2.9) to show that (2.8) is true (it requires proving that the negation of the thesis implies the negation of the hypothesis).

When using the law of contraposition, one must be able to construct the **correct negation** of a given proposition or property.

- Given any two predicates or properties $p(x)$ and $q(x)$, we list schematically some **rules** for constructing the **negation** of a proposition or property.

The negation of

“for every $x \in A$, $p(x)$ holds”

$$\forall x \in A \ (p(x))$$

is

“there exists $x \in A$ for which $p(x)$ does not hold”

$$\exists x \in A \ (\neg p(x))$$

The negation of

“there exists $x \in A$ for which $p(x)$ holds”

$$\exists x \in A \ (p(x))$$

is

“for every $x \in A$, $p(x)$ does not hold”

$$\forall x \in A \ (\neg p(x))$$

The negation of

“ $p(x)$ holds and $q(x)$ holds”

$$(p(x) \wedge q(x))$$

is

“ $p(x)$ does not hold or $q(x)$ does not hold”

$$(\neg p(x) \vee \neg q(x))$$

The negation of

“ $p(x)$ holds or $q(x)$ holds”

$$(p(x) \vee q(x))$$

is

“ $p(x)$ does not hold and $q(x)$ does not hold”

$$(\neg p(x) \wedge \neg q(x))$$

2.4 Sufficient conditions and necessary conditions

Definition 2.6: of sufficient condition

A **sufficient condition** is one that, if satisfied, guarantees the truth of the proposition.

Definition 2.7: of necessary condition

A **necessary condition** is one that must be satisfied for the proposition to be true.

Let $p(x)$ and $q(x)$ be any two predicates or properties. If $p(x)$ implies $q(x)$, formally:

$$p(x) \Rightarrow q(x)$$

then:

- $p(x)$ is a **sufficient condition** for $q(x)$
- $q(x)$ is a **necessary condition** for $p(x)$

- We have seen that the following universal implication is true (proposition (2.5)):

$$\forall n \in \mathbb{N} \ (n \text{ odd} \Rightarrow n^2 \text{ odd})$$

Hence “ n odd” is a sufficient condition for “ n^2 odd” and “ n^2 odd” is a necessary condition for “ n odd”.

- Let us now prove that the following proposition is also true:

Proposition 2.8

$$\forall n \in \mathbb{N} \ (n^2 \text{ odd} \Rightarrow n \text{ odd}) \quad (2.11)$$

Proof. Let n^2 be odd; we prove that then n is odd.

So let $n^2 = 2(2k^2 + 2k) + 1$ be an odd number ($k \in \mathbb{N}$, since $2(2k^2 + 2k)$ is an even number). We therefore need to write n as an even integer plus one. We have:

$$n = \sqrt{2(2k^2 + 2k) + 1} = \sqrt{4k^2 + 4k + 1} = \sqrt{(2k + 1)^2} = 2k + 1.$$

Since $2k + 1$ is an odd integer, n is odd.

□

In this way we have proved that “ n^2 odd” is a necessary and sufficient condition for “ n odd” and also that “ n odd” is a necessary and sufficient condition for “ n^2 odd”:

Proposition 2.9

$$\forall n \in \mathbb{N} \ (n \text{ odd} \iff n^2 \text{ odd}) \quad (2.12)$$

- We have seen that the following universal implication is true (proposition (2.6)):

$$\forall n \in \mathbb{N} \ (n \text{ even} \Rightarrow n^2 \text{ even})$$

Hence “ n even” is a sufficient condition for “ n^2 even” and “ n^2 even” is a necessary condition for “ n even”.

- Let us now prove that the following proposition is also true (direct proof, without using the law of contraposition as seen previously in (2.10)):

$$\forall n \in \mathbb{N} \ (n^2 \text{ even} \Rightarrow n \text{ even})$$

Proof. Let n^2 be even; we prove that then n is even.

So let $n^2 = 2 (2 k^2)$ be an even number ($k \in \mathbb{N}$); we therefore need to write n as an even integer. We have:

$$n = \sqrt{2 (2 k^2)} = \sqrt{4 k^2} = 2 k.$$

Since $2 k$ is an even integer, n is even. □

In this way we have proved that “ n^2 even” is a necessary and sufficient condition for “ n even” and also that “ n even” is a necessary and sufficient condition for “ n^2 even”:

Proposition 2.10

$$\forall n \in \mathbb{N} \ (n \text{ even} \iff n^2 \text{ even}) \quad (2.13)$$

Example 2.1: necessary and sufficient conditions

For example, for a square matrix of real numbers, the fact that its determinant is different from zero is a necessary and sufficient condition for it to be invertible.

Proposition 2.11

$$\forall n \in \mathbb{N} \ (n \text{ prime} > 2 \Rightarrow n \text{ odd}) \quad (2.14)$$

Proof. Let $n > 2$ be a prime number; we prove that then n is odd.

If n is not odd, it is even. But no even number greater than two is prime, a fact that contradicts the hypothesis that n is prime and greater than two (hence this case cannot

occur).

Consequently n is odd. □

- Hence “ n odd” is a necessary condition for “ n prime > 2 ” and “ n prime > 2 ” is a sufficient condition for “ n odd”.
- However, “ n odd” does not imply “ n prime > 2 ”, since for example the number 9 is not prime (counterexample). That is:

$$\forall n \in \mathbb{N} \quad (n \text{ odd} \not\Rightarrow n \text{ prime number greater than } 2)$$

Hence “ n odd” is a necessary but not sufficient condition for “ n prime > 2 ” and “ n prime > 2 ” is a sufficient but not necessary condition for “ n odd”.

Proposition 2.12

$$\forall n \in \mathbb{N} \quad (n \text{ divisible by } 6 \Rightarrow n \text{ even}) \quad (2.15)$$

Proof. Let n be a number divisible by six; we prove that then n is even.

If n is not even, it is odd. But no odd number is divisible by six, a fact that contradicts the hypothesis that n is divisible by six (hence this case cannot occur).

Consequently n is even. □

- Hence “ n even” is a necessary condition for “ n divisible by 6” and “ n divisible by 6” is a sufficient condition for “ n even”.
- However, “ n even” does not imply “ n divisible by 6”, since for example the number 2 is even but not divisible by six (counterexample). That is:

$$\forall n \in \mathbb{N} \quad (n \text{ even} \not\Rightarrow n \text{ divisible by } 6)$$

Hence “ n even” is a necessary but not sufficient condition for “ n divisible by 6” and “ n divisible by 6” is a sufficient but not necessary condition for “ n even”.

Example 2.2: necessary/sufficient but not sufficient/necessary conditions

Being a square implies being a rectangle:

$$\text{being a square} \Rightarrow \text{being a rectangle}$$

since all squares are rectangles.

Hence “being a rectangle” is a necessary condition for “being a square” and “being a square” is a sufficient condition for “being a rectangle”.

But being a rectangle does not imply being a square

$$\text{being a rectangle} \not\Rightarrow \text{being a square}$$

because there are rectangles that are not squares.

Hence “being a rectangle” is not a sufficient condition (but it is a necessary one) for “being a square” and “being a square” is not a necessary condition (but it is a sufficient one) for “being a rectangle”.

2.5 Proofs by contradiction

- It is an indirect proof technique

Definition 2.13: of proof by contradiction

*In general, a **proof by contradiction** consists in assuming that the hypothesis of the theorem and the negation of the thesis are true, and deducing from these facts a contradiction of any kind.*

- We illustrate **proof by contradiction** with the following theorem.

Theorem 2.14

There is no rational number whose square is 2.

Proof. Suppose, by contradiction, that there exists a number $r \in \mathbb{Q}$ such that $r^2 = 2$.

We can write $r = \frac{n}{m}$ with $n, m \in \mathbb{Z}$, $m \neq 0$.

Moreover, we can assume that the fraction $\frac{n}{m}$ is already reduced to lowest terms, that is, “simplified” (in other words: n , m have no common factors).

We then have the chain of implications:

$$\left(\frac{n}{m}\right)^2 = 2$$

$$n^2 = 2m^2$$

so n^2 is even; but then by (2.13) n is also even and we can write $n = 2k$ for some $k \in \mathbb{Z}$.

Hence the relation $n^2 = 2m^2$ can be rewritten as:

$$(2k)^2 = 2m^2$$

$$4k^2 = 2m^2$$

$$m^2 = 2k^2.$$

So m^2 is even. But then by (2.13) m is also even.

Therefore both n and m are even, and this is a **contradiction**, because we had assumed that the fraction $\frac{n}{m}$ had already been simplified.

The proof can be found in Euclid's Elements (around 300 BC).

□

2.6 Logic and sets

The language of logic and the language of sets are two sides of the same coin.

2.6.1 Logical implication and set inclusion

- There is a parallel between the relation of set inclusion and logical implication. To explain it, consider the universal implication:

$$\forall n \in \mathbb{N} \quad (n \text{ divisible by } 4 \Rightarrow n \text{ divisible by } 2) \quad (2.16)$$

If we denote by:

$$D_4 = \{ n \in \mathbb{N} : n \text{ is divisible by } 4 \} \text{ and } D_2 = \{ n \in \mathbb{N} : n \text{ is divisible by } 2 \}$$

we can observe that the universal implication written above is equivalent to the assertion:

$$D_4 \subseteq D_2$$

Indeed, this inclusion means that every element belonging to D_4 also belongs to D_2 , that is, that every natural number divisible by 4 is also divisible by 2.

The universal implication:

“for every $x \in A$, if $p(x)$ holds then $q(x)$ holds”

is equivalent to the set inclusion:

$$\{ x \in A : p(x) \text{ is true} \} \subseteq \{ x \in A : q(x) \text{ is true} \}.$$

2.6.2 Equality of sets and universal implications

- Proving the equality of two sets, i.e., $A = B$, requires proving two universal implications. Formally:

$$\forall x (x \in A \Rightarrow x \in B) \quad \text{and} \quad \forall x (x \in B \Rightarrow x \in A). \quad (2.17)$$

- Stating that $A \subsetneq B$ means stating that “Every element that belongs to A also belongs to B , and there exists an element of B that does not belong to A ”. Formally:

$$\forall x (x \in A \Rightarrow x \in B) \quad \text{and} \quad \exists x \in B : x \notin A. \quad (2.18)$$

2.6.3 Set operations and logical operations

There is a relation between operations on sets and logical operations. Precisely:

1. **Set intersection** is defined by means of “and” (**logical conjunction**).
2. **Set union** is defined by means of “or” (**logical disjunction**).
3. **Set difference** and the **complement operation** are defined by means of “not” (**logical negation**).

- The distributive properties of union and intersection of sets:

Observation 2.15: distributive properties (sets)

Given three sets A , B and C we have:

$$A \cap (B \cup C) = (A \cap B) \cup (A \cap C)$$

$$A \cup (B \cap C) = (A \cup B) \cap (A \cup C).$$

- They can be rewritten in terms of predicates by observing that the intersection symbol $\cap$ corresponds to the conjunction $\wedge$ (“and”) and that the union symbol $\cup$ corresponds to the disjunction $\vee$ (“or”).

Observation 2.16: distributive properties (predicates)

Given three predicates $p(x)$, $q(x)$ and $r(x)$ we have:

$$p(x) \wedge (q(x) \vee r(x)) = (p(x) \wedge q(x)) \vee (p(x) \wedge r(x))$$

$$p(x) \vee (q(x) \wedge r(x)) = (p(x) \vee q(x)) \wedge (p(x) \vee r(x)).$$

- De Morgan's laws:

Proposition 2.17: De Morgan's laws (sets)

Given the sets $B, C \subseteq U$, we have

$$\overline{B \cap C} = \overline{B} \cup \overline{C}$$

$$\overline{B \cup C} = \overline{B} \cap \overline{C}$$

- They can be rewritten in terms of predicates by observing that the complement operation corresponds to the negation $\neg$ ("not").

Proposition 2.18: De Morgan's laws (predicates)

Given two predicates $q(x)$ and $r(x)$ we have:

$$\neg(q(x) \wedge r(x)) = \neg q(x) \vee \neg r(x)$$

$$\neg(q(x) \vee r(x)) = \neg q(x) \wedge \neg r(x)$$

Number sets and intervals

3.1 The five main number sets

3.1.1 The natural numbers

$\mathbb{N}$ is the set of **natural numbers**, that is, the numbers used for counting:

$$\mathbb{N} = \{0, 1, 2, 3, 4, \dots\}$$

- We recall the convention adopted in these notes: **zero is a natural number**, that is $0 \in \mathbb{N}$. Other textbooks exclude zero from $\mathbb{N}$: it is only a convention, but it must be stated once and for all.
- We denote by $\mathbb{N}_{>0}$ the set of **positive natural numbers**:

$$\mathbb{N}_{>0} = \mathbb{N} \setminus \{0\} = \{1, 2, 3, 4, \dots\}$$

In the chapter *Sets* we have already introduced, for the natural numbers:

- the definitions of **even** number ($n = 2m$) and **odd** number ($n = 2m + 1$), with $m \in \mathbb{N}$, and the fact that every natural number is either even or odd, but never both;
- the **well-ordering axiom**, stating that every non-empty subset of $\mathbb{N}$ has a minimum.

3.1.2 The integers

$\mathbb{Z}$ is the set of **integers** (also called signed numbers), that is:

$$0, 1, -1, 2, -2, \dots$$

3.1.3 The rational numbers

$\mathbb{Q}$ is the set of **rational numbers**, that is, of the fractions:

$$\frac{n}{m}, \quad \text{where } n, m \in \mathbb{Z}, \text{ and } m \neq 0.$$

Rational numbers can also be written in decimal form. For example:

$$\frac{3}{4} = 0.75, \quad \text{while} \quad \frac{1}{3} = 0.333\dots = 0.\bar{3}.$$

A rational number, written in decimal form, has a **finite or infinite periodic decimal expansion**: after the decimal point there is a finite number of digits, or an infinite number of digits which, from a certain point on, repeat *periodically*.

The same rational number can have two different decimal expansions. The classical example is the equality

$$0.\bar{9} = 1$$

for which we gave four different proofs in the chapter *Sets*, and which we therefore do not repeat here.

3.1.4 The real numbers

$\mathbb{R}$ is the set of **real numbers**, that is, those which are identified with **finite or infinite decimal expansions, periodic or non-periodic**: after the decimal point there may be any string of digits, possibly *infinite* and *non-periodic*.

That numbers of this last kind exist (that is, real but not rational), can be understood by reflecting on examples such as:

$$0.10110111011110\dots$$

After the decimal point, the previous number has: one digit equal to 1, then 0, then two digits equal to 1, then 0, then three digits equal to 1 ... and so on. It is clear that this rule defines a decimal number precisely. On the other hand, the string of digits after the decimal point is neither finite nor periodic: therefore this number is irrational.

3.1.5 The complex numbers


$\mathbb{C}$ is the set of **complex numbers**, that is, numbers of the form $a + ib$, where a, b are real numbers, and i is the *imaginary unit*, that is, a number whose square is -1 .

Among the number sets introduced, the following inclusions hold:

$$\mathbb{N} \subsetneq \mathbb{Z} \subsetneq \mathbb{Q} \subsetneq \mathbb{R} \subsetneq \mathbb{C}.$$

As indicated by the symbols, all the inclusions are strict:

- there exist integers that are not natural numbers (the negative numbers),
- there exist rational numbers that are not integers (the proper fractions),
- there exist real numbers that are not rational (the irrational numbers),
- there exist complex numbers that are not real (the imaginary numbers).



As for $\mathbb{N}_{>0}$, we denote by

$$\mathbb{Z}_{>0}, \quad \mathbb{Q}_{>0}, \quad \mathbb{R}_{>0}$$

the sets of **positive** integer, rational and real numbers, and by

$$\mathbb{Z}_{\geq 0}, \quad \mathbb{Q}_{\geq 0}, \quad \mathbb{R}_{\geq 0}$$

the sets of **non-negative** integer, rational and real numbers (that is, positive or zero). For example:

$$\mathbb{R}_{>0} = \{x \in \mathbb{R} : x > 0\} \quad \text{and} \quad \mathbb{Z}_{\geq 0} = \{x \in \mathbb{Z} : x \geq 0\} = \mathbb{N}$$

3.2 Intervals

Definition 3.1: of bounded interval

Given two real numbers a, b , a **bounded interval** with endpoints a and b is one of the following sets:

$$\begin{aligned} [a, b] &= \{x \in \mathbb{R} : a \leq x \leq b\}, & [a, b) &= \{x \in \mathbb{R} : a \leq x < b\} \\ (a, b] &= \{x \in \mathbb{R} : a < x \leq b\}, & (a, b) &= \{x \in \mathbb{R} : a < x < b\} \end{aligned}$$

- As can be seen, the square (round) bracket at one of the two endpoints indicates that this endpoint is included in (excluded from) the interval.
- The intervals $[a, b]$ are called **closed**; the intervals (a, b) are called **open**.


Definition 3.2: of unbounded interval

Given a real number a , an **unbounded interval** (or **half-line**) with endpoint a is one of the following sets:

$$\begin{aligned} [a, +\infty) &= \{x \in \mathbb{R} : x \geq a\}, & (a, +\infty) &= \{x \in \mathbb{R} : x > a\} \\ (-\infty, a] &= \{x \in \mathbb{R} : x \leq a\}, & (-\infty, a) &= \{x \in \mathbb{R} : x < a\} \end{aligned}$$


- The symbols $-\infty$ and $+\infty$ **are not real numbers**: they are not endpoints that can belong to the set, and therefore next to them one always writes a round bracket.
- The whole line is an unbounded interval as well:

$$\mathbb{R} = (-\infty, +\infty)$$



Example 3.1: of intervals

The graphical representation of the closed bounded interval $[-3, -2]$ and of the open unbounded interval $(2, +\infty)$ is the following:



The first one contains both of its endpoints, the second one does not contain its endpoint 2 and is not bounded above.


It can be proved that the intervals, bounded or unbounded, are exactly the subsets I of $\mathbb{R}$ that satisfy the following property (called **connectedness**):

$$\forall x_1, x_2, x_3 \in \mathbb{R} \text{ such that } x_1 < x_2 < x_3, \text{ if } x_1, x_3 \in I, \text{ then } x_2 \in I$$

- In what follows we will sometimes consider the Cartesian product of two (or more) intervals, which can be given the geometric meaning of a rectangle (in two dimensions) or of a parallelepiped (in three dimensions).

Example 3.2: Cartesian product of intervals

Let $A = [0, 1]$, $B = [1, 2]$; the figure



illustrates the sets $A \times B$, $B \times A$, $A \times A$, also denoted by A^2 . In general, $A \times B$ is different from $B \times A$.

Binary relations

4.1 Binary relations

Definition 4.1: of binary relation

Given two sets A and B , a **binary relation** R is a subset of the Cartesian product $A \times B$

When we say that R is a binary relation on a single set A , we mean that R is a subset of $A \times A$.

- A binary relation will simply be called a relation (however, non-binary relations also exist).
- If $(a, b) \in R$ we also write $a R b$ and we say that a is **related** to b .

Example 4.1

- The relation “less than” (“ $<$ ”) on the natural numbers is the set:

$$R_{<} = \left\{ (a, b) : a, b \in \mathbb{N} \text{ and } a < b \right\}.$$

- The relation “less than or equal to” (“ $\leq$ ”) on the natural numbers is the set:

$$R_{\leq} = \left\{ (a, b) : a, b \in \mathbb{N} \text{ and } a \leq b \right\}.$$

- The relation “equal to” (“ $=$ ”) on the natural numbers is the set:

$$R_{=} = \left\{ (a, b) : a, b \in \mathbb{N} \text{ and } a = b \right\}.$$

- The relation “is a subset of” $R_{\subseteq}$ on the power set of the natural numbers (denoted by $2^{\mathbb{N}}$) is the set:

$$R_{\subseteq} = \left\{ (A, B) : A, B \in 2^{\mathbb{N}} \text{ and } A \subseteq B \right\}.$$

A relation R on a set of natural numbers can be visualized with a **table**: in row a and column b we write the pair (a, b) if $(a, b) \in R$, and the cell is left empty if $(a, b) \notin R$.

Example 4.2: tables of relations

- The table of the relation $R_{<}$ (rows and columns from 0 to 4):

	0	1	2	3	4	...
0		(0, 1)	(0, 2)	(0, 3)	(0, 4)	
1			(1, 2)	(1, 3)	(1, 4)	
2				(2, 3)	(2, 4)	
3					(3, 4)	
4						
...						

- The table of the relation $R_{=}$:

	0	1	2	3	4	...
0	(0, 0)					
1		(1, 1)				
2			(2, 2)			
3				(3, 3)		
4					(4, 4)	
...						

- The table of the relation $R_{\leq}$, which contains the pairs of $R_{<}$ and those of $R_{=}$:

	0	1	2	3	4	...
0	(0, 0)	(0, 1)	(0, 2)	(0, 3)	(0, 4)	
1		(1, 1)	(1, 2)	(1, 3)	(1, 4)	
2			(2, 2)	(2, 3)	(2, 4)	
3				(3, 3)	(3, 4)	
4					(4, 4)	
...						

Definition 4.2: of congruence modulo n

Given a natural number $n \geq 1$, two natural numbers a and b are **congruent modulo n** , written $a \equiv b \pmod{n}$, if their difference is an integer multiple of n :

$$a \equiv b \pmod{n} \iff \exists q \in \mathbb{Z} \text{ such that } b - a = qn.$$

Definition 4.3: of multiple and divisor

Given two natural numbers a and b , we say that a is a **multiple** of b if there exists $k \in \mathbb{N}$ such that $a = kb$. We say that a is a **divisor** of b if there exists $k \in \mathbb{N}$ such that $b = ka$.

- Hence a is a divisor of b if and only if b is a multiple of a .
- With this definition 0 is a multiple of every natural number ($0 = 0 \cdot b$) and 0 is a divisor only of itself (if $b = k \cdot 0$ then $b = 0$).

Example 4.3: other relations on the natural numbers

- The relation “congruent modulo n ” on the natural numbers is the set:

$$R_{\equiv_n} = \left\{ (a, b) : a, b \in \mathbb{N} \text{ and } a \equiv b \pmod{n} \right\}.$$

With $n = 3$ the table is:

	0	1	2	3	4	5	6	...
0	(0, 0)			(0, 3)			(0, 6)	
1		(1, 1)			(1, 4)			
2			(2, 2)			(2, 5)		
3	(3, 0)			(3, 3)			(3, 6)	
4		(4, 1)			(4, 4)			
5			(5, 2)			(5, 5)		
6	(6, 0)			(6, 3)			(6, 6)	
...								

- The relation “is a multiple of” on the natural numbers is the set:

$$R_{\text{mul}} = \left\{ (a, b) : a, b \in \mathbb{N} \text{ and } \exists k \in \mathbb{N} \text{ such that } a = kb \right\}.$$

	0	1	2	3	4	5	6	...
0	(0, 0)	(0, 1)	(0, 2)	(0, 3)	(0, 4)	(0, 5)	(0, 6)	
1		(1, 1)						
2		(2, 1)	(2, 2)					
3		(3, 1)		(3, 3)				
4		(4, 1)	(4, 2)		(4, 4)			
5		(5, 1)				(5, 5)		
6		(6, 1)	(6, 2)	(6, 3)			(6, 6)	
...								

Example 4.4: other relations on the natural numbers

- The relation “is a divisor of” on the natural numbers is the set:

$$R_{\text{div}} = \left\{ (a, b) : a, b \in \mathbb{N} \text{ and } \exists k \in \mathbb{N} \text{ such that } b = k a \right\}.$$

Since $(a, b) \in R_{\text{div}}$ if and only if $(b, a) \in R_{\text{mul}}$, its table is obtained from that of R_{mul} by swapping the two elements of each pair:

	0	1	2	3	4	5	6	...
0	(0, 0)							
1	(1, 0)	(1, 1)	(1, 2)	(1, 3)	(1, 4)	(1, 5)	(1, 6)	
2	(2, 0)		(2, 2)		(2, 4)		(2, 6)	
3	(3, 0)			(3, 3)			(3, 6)	
4	(4, 0)				(4, 4)			
5	(5, 0)					(5, 5)		
6	(6, 0)						(6, 6)	
...								

Example 4.5: other relations on the natural numbers

- The relation “difference of one” on the natural numbers is the set:

$$R_{ab}^1 = \left\{ (a, b) : a, b \in \mathbb{N} \text{ and } a = b - 1 \right\}.$$

	0	1	2	3	4	5	6	...
0		(0, 1)						
1			(1, 2)					
2				(2, 3)				
3					(3, 4)			
4						(4, 5)		
5							(5, 6)	
6								
...								

- The relation “the sum is even” on the natural numbers is the set:

$$R_{\text{even}} = \left\{ (a, b) : a, b \in \mathbb{N} \text{ and } \exists m \in \mathbb{N} \text{ such that } a + b = 2m \right\}.$$

	0	1	2	3	4	5	6	...
0	(0, 0)		(0, 2)		(0, 4)		(0, 6)	
1		(1, 1)		(1, 3)		(1, 5)		
2	(2, 0)		(2, 2)		(2, 4)		(2, 6)	
3		(3, 1)		(3, 3)		(3, 5)		
4	(4, 0)		(4, 2)		(4, 4)		(4, 6)	
5		(5, 1)		(5, 3)		(5, 5)		
6	(6, 0)		(6, 2)		(6, 4)		(6, 6)	
...								

- The table of the relation $R_{\subseteq}$ restricted to the subsets of $\{0, 1, 2\}$ (rows and columns are the sets A and B):

	$\emptyset$	$\{0\}$	$\{1\}$	$\{2\}$	$\{0, 1\}$	$\{0, 2\}$	$\{1, 2\}$	$\{0, 1, 2\}$
$\emptyset$	$(\emptyset, \emptyset)$	$(\emptyset, \{0\})$	$(\emptyset, \{1\})$	$(\emptyset, \{2\})$	$(\emptyset, \{0, 1\})$	$(\emptyset, \{0, 2\})$	$(\emptyset, \{1, 2\})$	$(\emptyset, \{0, 1, 2\})$
$\{0\}$		$(\{0\}, \{0\})$			$(\{0\}, \{0, 1\})$	$(\{0\}, \{0, 2\})$		$(\{0\}, \{0, 1, 2\})$
$\{1\}$			$(\{1\}, \{1\})$		$(\{1\}, \{0, 1\})$		$(\{1\}, \{1, 2\})$	$(\{1\}, \{0, 1, 2\})$
$\{2\}$				$(\{2\}, \{2\})$		$(\{2\}, \{0, 2\})$	$(\{2\}, \{1, 2\})$	$(\{2\}, \{0, 1, 2\})$
$\{0, 1\}$					$(\{0, 1\}, \{0, 1\})$			$(\{0, 1\}, \{0, 1, 2\})$
$\{0, 2\}$						$(\{0, 2\}, \{0, 2\})$		$(\{0, 2\}, \{0, 1, 2\})$
$\{1, 2\}$							$(\{1, 2\}, \{1, 2\})$	$(\{1, 2\}, \{0, 1, 2\})$
$\{0, 1, 2\}$								$(\{0, 1, 2\}, \{0, 1, 2\})$

Definition 4.4: of n -ary relation

Given n sets $A_1, A_2, \dots, A_n$, an n -ary relation R is a subset of the Cartesian product $A_1 \times A_2 \times \dots \times A_n$.

- Binary relations are the n -ary relations with $n = 2$.

4.2 Properties of relations

- A relation $R \subseteq A \times A$ is **reflexive** if:

$$\forall a \in A, \quad (a, a) \in R$$

- A relation $R \subseteq A \times A$ is **symmetric** if:

$$\forall a, b \in A, \quad (a, b) \in R \Rightarrow (b, a) \in R$$

- A relation $R \subseteq A \times A$ is **transitive** if:

$$\forall a, b, c \in A, \quad (a, b) \in R \text{ and } (b, c) \in R \Rightarrow (a, c) \in R$$

- A relation $R \subseteq A \times A$ is **antisymmetric** if:

$$\forall a, b \in A, \quad (a, b) \in R \text{ and } (b, a) \in R \Rightarrow a = b$$

Observation 4.5: equivalent form of antisymmetry

A relation $R \subseteq A \times A$ is antisymmetric if and only if:

$$\forall a, b \in A, \quad (a, b) \in R \text{ and } a \neq b \Rightarrow (b, a) \notin R$$

Proof. Fix $a, b \in \mathbb{A}$. An implication is false only when the antecedent is true and the consequent is false.

- The implication “ $(a, b) \in R$ and $(b, a) \in R \Rightarrow a = b$ ” is false only when $(a, b) \in R$, $(b, a) \in R$ and $a \neq b$.
- The implication “ $(a, b) \in R$ and $a \neq b \Rightarrow (b, a) \notin R$ ” is false only when $(a, b) \in R$, $a \neq b$ and $(b, a) \in R$.

The two implications are therefore false in exactly the same cases: for every pair a, b one is true if and only if the other is, and hence one holds for all $a, b \in \mathbb{A}$ if and only if the other does. $\square$

Example 4.6

- The relation $R_{\leq}$ is reflexive, but $R_{<}$ is not.
- The relations $R_{<}$ and $R_{\leq}$ are not symmetric.
- The relations $R_{<}$ and $R_{\leq}$ are transitive, but for example the relation:

$$R_{ab}^1 = \left\{ (a, b) : a, b \in \mathbb{N} \text{ and } a = b - 1 \right\}$$

is not, e.g., $(3, 4) \in R_{ab}^1$ and $(4, 5) \in R_{ab}^1$ but $(3, 5) \notin R_{ab}^1$.

Let us now see in detail, for each property, how to prove that a relation has it and how to show, with a counterexample, that it does not.

Example 4.7: reflexive property

- The relation $R_{=}$ is reflexive, since

$$a = a, \quad \forall a \in \mathbb{N}.$$

- The relation $R_{\leq}$ is reflexive, since

$$a \leq a, \quad \forall a \in \mathbb{N}.$$

- The relation $R_{<}$ is not reflexive: a counterexample is $a = 1$, since $1 < 1$ is false and hence $(1, 1) \notin R_{<}$.

Example 4.8: symmetric property

- The relation $R_{=}$ is symmetric, since

$$a = b \Rightarrow b = a, \quad \forall a, b \in \mathbb{N}.$$

- The relations $R_{<}$ and $R_{\leq}$ are not symmetric: a counterexample is the pair $(1, 2)$, since $1 < 2$ and $1 \leq 2$, but $2 < 1$ and $2 \leq 1$ are false. Hence $(1, 2) \in R_{<}$ but $(2, 1) \notin R_{<}$, and $(1, 2) \in R_{\leq}$ but $(2, 1) \notin R_{\leq}$.

Example 4.9: transitive property

- The relation $R_=_$ is transitive, since

$$a = b \text{ and } b = c \Rightarrow a = c, \quad \forall a, b, c \in \mathbb{N}.$$

The same holds for $R_<$ and $R_\leq$: if $a < b$ and $b < c$ then $a < c$, and if $a \leq b$ and $b \leq c$ then $a \leq c$.

- The relation R_{ab}^1 is not transitive: a counterexample is given by $(3, 4) \in R_{ab}^1$ and $(4, 5) \in R_{ab}^1$, while $(3, 5) \notin R_{ab}^1$ because $3 \neq 5 - 1$.

Example 4.10: antisymmetric property

- The relation $R_\leq$ is antisymmetric, since $a \leq b$ and $b \leq a$ imply $a = b$.
- The relation $R_=_$ is antisymmetric, since

$$a = b \text{ and } b = a \Rightarrow a = b, \quad \forall a, b \in \mathbb{N}.$$

- The relation $R_<$ is antisymmetric. Indeed, there are no two natural numbers a and b with $a < b$ and $b < a$ (otherwise we would have $a < a$): the antecedent of the implication is never true, and hence the implication is true for all $a, b \in \mathbb{N}$. With the equivalent form of antisymmetry this can be seen directly:

$$a < b \text{ and } a \neq b \Rightarrow b \not< a, \quad \forall a, b \in \mathbb{N}.$$

- The relation $R_{\equiv_3}$ is not antisymmetric: a counterexample is given by $(0, 3) \in R_{\equiv_3}$ (because $3 - 0 = 1 \cdot 3$) and $(3, 0) \in R_{\equiv_3}$ (because $0 - 3 = (-1) \cdot 3$), while $0 \neq 3$.


4.3 Representation with directed graphs

Relations can be represented by a **directed graph**, where the **vertices** are the elements of the set on which the relation R is defined and an **arc** (a, b) , drawn as an arrow from a to b , means that $(a, b) \in R$. An arc (a, a) is called a **loop**.

Example 4.11: graphs of relations

- The directed graph of the relation “is a multiple of” on the set $\{0, 1, \dots, 5\}$:


$$R = \left\{ (a, b) : a, b \in \{0, 1, \dots, 5\} \text{ and } \exists k \in \mathbb{N} \text{ such that } a = kb \right\}.$$



- The directed graph of the relation “is a divisor of” on the set $\{0, 1, \dots, 5\}$:

$$R = \left\{ (a, b) : a, b \in \{0, 1, \dots, 5\} \text{ and } \exists k \in \mathbb{N} \text{ such that } b = k a \right\}.$$


It has the same arcs as the previous graph, traversed in the opposite direction.



Example 4.12: graphs of relations

- The directed graph of the relation “less than or equal to” on the set $\{0, 1, \dots, 5\}$:


$$R = \left\{ (a, b) : a, b \in \{0, 1, \dots, 5\} \text{ and } a \leq b \right\}.$$



- The directed graph of the relation “less than” on the set $\{0, 1, \dots, 5\}$, obtained from


the previous one by removing the loops:

$$R = \left\{ (a, b) : a, b \in \{0, 1, \dots, 5\} \text{ and } a < b \right\}.$$



- The directed graph of the relation “equal to” on the set $\{0, 1, \dots, 5\}$, which has only the loops:


$$R = \left\{ (a, b) : a, b \in \{0, 1, \dots, 5\} \text{ and } a = b \right\}.$$



Example 4.13: graph of the inclusion relation

The directed graph of the relation “is a subset of” on the subsets of $\{0, 1\}$:

$$R = \left\{ (A, B) : A, B \in 2^{\{0,1\}} \text{ and } A \subseteq B \right\}.$$



The main properties of a relation can be read on its directed graph:

- the relation is **reflexive** if and only if every vertex a has the loop (a, a) ;
- the relation is **symmetric** if and only if, for every pair of vertices a and b , if there is the arc from a to b then there is also the arc from b to a ;
- the relation is **transitive** if and only if, for every triple of vertices a , b and c , if there


are the arc from a to b and the arc from b to c then there is also the arc from a to c ;

- the relation is **antisymmetric** if and only if, for every pair of distinct vertices a and b , if there is the arc from a to b then there is no arc from b to a (equivalent form of antisymmetry).


- For example, in the graphs of “is a multiple of”, “is a divisor of”, “less than or equal to” and “equal to” every vertex has the loop (reflexive relations), while in the graph of “less than” no vertex has it. In none of these graphs are there two arcs in opposite directions between two distinct vertices (antisymmetric relations).

Example 4.14: graphs satisfying only two properties


- The relation $\{(a, a), (b, b), (c, c), (a, b), (b, a), (b, c), (c, b)\}$ on the set $\{a, b, c\}$ is reflexive and symmetric, but not transitive: there are the arcs (a, b) and (b, c) , but not the arc (a, c) .



- The relation $\{(a, a), (b, b), (c, c), (a, b), (a, c), (b, c)\}$ on the set $\{a, b, c\}$ is reflexive and transitive, but not symmetric: there is the arc (a, b) , but not the arc (b, a) .



- The relation $\{(a, a), (a, b), (b, a), (b, b)\}$ on the set $\{a, b, c\}$ is symmetric and transitive, but not reflexive: the loop (c, c) is missing. For transitivity it suffices to check the pairs of consecutive arcs: from (a, b) and (b, a) follows (a, a) , from (b, a) and (a, b) follows (b, b) , and the pairs containing a loop give an arc already present.



Observation 4.6: symmetric and transitive relations

Let $R \subseteq A \times A$ be a symmetric and transitive relation. If an element $a \in A$ appears in at least one pair of R , then $(a, a) \in R$.

Proof. If a appears in a pair of R , there exists $b \in A$ such that $(a, b) \in R$ or $(b, a) \in R$. By symmetry, in both cases we have both $(a, b) \in R$ and $(b, a) \in R$. By transitivity, from $(a, b) \in R$ and $(b, a) \in R$ follows $(a, a) \in R$. $\square$

- Hence a symmetric and transitive relation fails to be reflexive only if some element of A appears in no pair of the relation, like the element c of the last graph.

4.4 Equivalence relations

Definition 4.7: of equivalence relation

A reflexive, symmetric and transitive relation is an **equivalence relation**.

Example 4.15: equivalence relations

- The relation $R_=$ is an equivalence relation, because it is reflexive, symmetric and transitive.
- The relation $R_<$ is not an equivalence relation, because it is not reflexive (nor symmetric).
- The relation $R_{\leq}$ is not an equivalence relation, because it is not symmetric.

Example 4.16: congruence modulo n

Given a natural number $n \geq 1$, the relation

$$R_{\equiv_n} = \left\{ (a, b) : a, b \in \mathbb{N} \text{ and } a \equiv b \pmod{n} \right\}$$

is an equivalence relation. Recall that $(a, b) \in R_{\equiv_n}$ if and only if there exists $q \in \mathbb{Z}$ such that $b - a = qn$.

- Reflexive: for every $a \in \mathbb{N}$ we have $a - a = 0 \cdot n$, with $0 \in \mathbb{Z}$, hence $(a, a) \in R_{\equiv_n}$.
- Symmetric: for all $a, b \in \mathbb{N}$, if there exists $q \in \mathbb{Z}$ such that $b - a = qn$, then

$$a - b = (-q)n, \quad \text{with } -q \in \mathbb{Z},$$

hence $(b, a) \in R_{\equiv_n}$.

- Transitive: for all $a, b, c \in \mathbb{N}$, if there exist $q, p \in \mathbb{Z}$ such that $b - a = qn$ and $c - b = pn$, then

$$c - a = (c - b) + (b - a) = pn + qn = (q + p)n, \quad \text{with } q + p \in \mathbb{Z},$$

hence $(a, c) \in R_{\equiv_n}$.

Example 4.17: the sum is even

The relation

$$R_{\text{even}} = \left\{ (a, b) : a, b \in \mathbb{N} \text{ and } \exists m \in \mathbb{N} \text{ such that } a + b = 2m \right\}$$

is an equivalence relation.

- Reflexive: for every $a \in \mathbb{N}$ we have $a + a = 2a$, with $a \in \mathbb{N}$, hence $(a, a) \in R_{\text{even}}$.
- Symmetric: for all $a, b \in \mathbb{N}$, if $a + b = 2m$ with $m \in \mathbb{N}$, then also $b + a = a + b = 2m$, hence $(b, a) \in R_{\text{even}}$.
- Transitive: for all $a, b, c \in \mathbb{N}$, if there exist $p, q \in \mathbb{N}$ such that $a + b = 2p$ and $b + c = 2q$, then

$$(a + b) + (b + c) = 2p + 2q \Rightarrow a + c = 2(p + q - b).$$

The number $p + q - b$ is an integer, and it is not negative because $2(p + q - b) = a + c \geq 0$; hence $p + q - b \in \mathbb{N}$ and $(a, c) \in R_{\text{even}}$.

Definition 4.8: of equivalence class

Given an equivalence relation R on a set A , the **equivalence class** of an element $a \in A$ is the set

$$[a] = \left\{ b \in A : (a, b) \in R \right\}.$$

Lemma 4.9: related elements have the same class

Let R be an equivalence relation on a set A . For all $a, b \in A$:

$$(a, b) \in R \Rightarrow [a] = [b].$$

Proof. Let $(a, b) \in R$. We show the two inclusions.


- $[b] \subseteq [a]$: if $c \in [b]$, then $(b, c) \in R$. From $(a, b) \in R$ and $(b, c) \in R$, by transitivity, follows $(a, c) \in R$, that is $c \in [a]$.
- $[a] \subseteq [b]$: by symmetry $(b, a) \in R$. If $c \in [a]$, then $(a, c) \in R$, and from $(b, a) \in R$ and $(a, c) \in R$, by transitivity, follows $(b, c) \in R$, that is $c \in [b]$.

Hence $[a] = [b]$. □

- Any two elements of the same class are related to each other: if $b, c \in [a]$, then $(a, b) \in R$ and $(a, c) \in R$; by symmetry $(b, a) \in R$ and by transitivity $(b, c) \in R$. An equivalence class therefore gathers elements that are all related to each other.

Example 4.18: equivalence classes

The directed graph of an equivalence relation on the set $\{a, b, c, d, e, f\}$ with two equivalence classes, $\{a, b, c\}$ (in blue) and $\{d, e, f\}$ (in red). A double-headed arrow between x and y represents the two arcs (x, y) and (y, x) .



Definition 4.10: of partition

A **partition** of a set A is a family $\mathcal{S}$ of subsets of A such that:

- every set of the family is nonempty: $S \neq \emptyset$ for every $S \in \mathcal{S}$;
- two distinct sets of the family are disjoint: $S \cap T = \emptyset$ for all $S, T \in \mathcal{S}$ with $S \neq T$;
- the union of the sets of the family is A : $\bigcup_{S \in \mathcal{S}} S = A$, that is, every $a \in A$ belongs to at least one $S \in \mathcal{S}$.

- The family can be finite, $\mathcal{S} = \{S_1, S_2, \dots, S_n\}$, or infinite.
- From the second and the third condition it follows that every element of A belongs to **one and only one** set of the partition: to at least one by the third condition, and not to two distinct sets S and T , which are disjoint.

Proposition 4.11: equivalence classes form a partition

Let R be an equivalence relation on a set A . Then the family of the equivalence classes

$$\mathcal{S} = \left\{ [a] : a \in A \right\}$$

is a partition of A .

- The proof checks the three conditions of the definition of partition; the second one follows from the fact that two classes are either equal or disjoint.

Proof. Every class $[a]$ is a subset of A by definition.

- Nonempty classes: by reflexivity $(a, a) \in R$, hence $a \in [a]$ and $[a] \neq \emptyset$.
- Union equal to A : every $a \in A$ belongs to the class $[a]$, hence $A \subseteq \bigcup_{a \in A} [a]$; the other inclusion holds because every class is contained in A .
- Distinct classes are disjoint: we show that if two classes $[a]$ and $[b]$ have an element in common, then they are equal. Let $c \in [a] \cap [b]$. Then $(a, c) \in R$ and $(b, c) \in R$. By symmetry $(c, b) \in R$, and by transitivity, from $(a, c) \in R$ and $(c, b) \in R$, follows $(a, b) \in R$. By Lemma 4.9 we then have $[a] = [b]$. Hence two distinct classes have no element in common.

□

Proposition 4.12: every partition defines an equivalence relation

Let $\mathcal{S}$ be a partition of a set A and let

$$R = \left\{ (a, b) : a, b \in A \text{ and } \exists S \in \mathcal{S} \text{ such that } a \in S \text{ and } b \in S \right\}.$$

Then R is an equivalence relation on A , and its equivalence classes are exactly the sets of the partition $\mathcal{S}$.

Proof. We first show that R is an equivalence relation.

- Reflexive: for every $a \in A$ there exists $S \in \mathcal{S}$ with $a \in S$ (the union of the sets of the partition is A). Then a and a belong to the same set S , that is $(a, a) \in R$.
- Symmetric: if $(a, b) \in R$, there exists $S \in \mathcal{S}$ with $a \in S$ and $b \in S$; the same S contains b and a , hence $(b, a) \in R$.
- Transitive: if $(a, b) \in R$ and $(b, c) \in R$, there exist $S, T \in \mathcal{S}$ with $a, b \in S$ and $b, c \in T$. Then $b \in S \cap T$, hence $S \cap T \neq \emptyset$ and, since two distinct sets of the partition are disjoint, $S = T$. Hence a and c belong to the same set S , that is $(a, c) \in R$.

We now show that the classes are the sets of the partition. Given $a \in A$, let S_a be the only set of $\mathcal{S}$ containing a . If $b \in S_a$, then $(a, b) \in R$, that is $b \in [a]$. Conversely, if $b \in [a]$, there exists $S \in \mathcal{S}$ with $a, b \in S$; since a belongs to only one set of the partition, $S = S_a$ and hence $b \in S_a$. Therefore $[a] = S_a$: every class is a set of the partition. Finally, every $S \in \mathcal{S}$ is nonempty: taking $a \in S$, we have $S = S_a = [a]$, hence every set of the partition is a class. □

Example 4.19: classes modulo 3

The equivalence relation $R_{\equiv 3}$ divides the natural numbers into 3 equivalence classes:

$$[0] = \{0, 3, 6, 9, \dots\}, \quad [1] = \{1, 4, 7, 10, \dots\}, \quad [2] = \{2, 5, 8, 11, \dots\}.$$


Each class contains the natural numbers that have the same remainder in the division by 3.

Indeed, write $a = 3q_a + r_a$ and $b = 3q_b + r_b$, with $q_a, q_b \in \mathbb{N}$ and remainders $r_a, r_b \in \{0, 1, 2\}$. Then

$$b - a = 3(q_b - q_a) + (r_b - r_a).$$

- If $r_a = r_b$, then $b - a = 3(q_b - q_a)$ and $(a, b) \in R_{\equiv_3}$.
- If $(a, b) \in R_{\equiv_3}$, that is $b - a = 3q$ with $q \in \mathbb{Z}$, then $r_b - r_a = 3(q - q_b + q_a)$ is a multiple of 3 between -2 and 2 , hence $r_b - r_a = 0$.

Here is the directed graph of the relation on the natural numbers from 0 to 8, with one colour per class. A double-headed arrow represents the two arcs (a, b) and (b, a) ; the loops (a, a) , present for every a , are not drawn.



Example 4.20: even and odd numbers


The equivalence relation R_{even} divides the natural numbers into 2 equivalence classes, the even numbers and the odd numbers:

$$[0] = \{0, 2, 4, 6, \dots\}, \quad [1] = \{1, 3, 5, 7, \dots\}.$$

Indeed, write $a = 2q_a + r_a$ and $b = 2q_b + r_b$, with $q_a, q_b \in \mathbb{N}$ and remainders $r_a, r_b \in \{0, 1\}$. Then $a + b = 2(q_a + q_b) + (r_a + r_b)$, with $r_a + r_b \in \{0, 1, 2\}$.

- If $r_a = r_b$, then $r_a + r_b = 2r_a$ and $a + b = 2(q_a + q_b + r_a)$ is even.
- If $r_a \neq r_b$, then $r_a + r_b = 1$ and $a + b = 2(q_a + q_b) + 1$ is not even.

Hence $(a, b) \in R_{\text{even}}$ if and only if a and b are both even or both odd. Here is the directed graph of the relation on the natural numbers from 0 to 6 (even in red, odd in blue; double-headed arrows and loops omitted as in the previous example):



Observation 4.13: equivalence and antisymmetry

If an equivalence relation R on a set A is also antisymmetric, then every equivalence class has a single element: $[a] = \{a\}$ for every $a \in A$.

Proof. Let $a \in A$. By reflexivity $a \in [a]$. If $b \in [a]$, then $(a, b) \in R$; by symmetry also $(b, a) \in R$, and by antisymmetry $a = b$. Hence $[a] = \{a\}$. $\square$

- In this case $(a, b) \in R$ if and only if $b \in [a] = \{a\}$, that is if and only if $a = b$: the only equivalence relation on A that is also antisymmetric is the relation “equal to”.

4.5 Partial order relations

Definition 4.14: of partial order relation

A reflexive, antisymmetric and transitive relation is a **partial order relation**.

Example 4.21

- The relation $R_{\leq}$ is a partial order relation, but the relation $R_{<}$ is not, since it is not reflexive.
- The relation $R_{=}$ is a partial order relation, because it is reflexive, antisymmetric and transitive.

Observation 4.15

The relation $R_{\subseteq}$ is a partial order relation

Proof. We must prove that the relation is reflexive, antisymmetric and transitive.

- For it to be reflexive we must prove that $(S, S) \in R_{\subseteq}$, which is true since $S \subseteq S$.
- For it to be antisymmetric we must prove that if $S_1 \neq S_2$ then $S_1 \not\subseteq S_2$ or $S_2 \not\subseteq S_1$ or both, the negation of the desired property.

Since $S_1 \neq S_2$:

either there exists some element that is in S_1 but not in S_2 , hence $S_1 \not\subseteq S_2$
 or there exists some element that is in S_2 but not in S_1 , hence $S_2 \not\subseteq S_1$
 or both of the previous options hold

- For it to be transitive we must prove that $(S_1, S_2) \in R_{\subseteq}$ and $(S_2, S_3) \in R_{\subseteq}$ imply $(S_1, S_3) \in R_{\subseteq}$. Clearly, since $S_1 \subseteq S_2$ and $S_2 \subseteq S_3$, we have $S_1 \subseteq S_3$.



Example 4.22: multiples and divisors

The relation R_{mul} (“is a multiple of”) is a partial order relation.

- Reflexive: for every $a \in \mathbb{N}$ we have $a = 1 \cdot a$, hence $(a, a) \in R_{\text{mul}}$.
- Transitive: for all $a, b, c \in \mathbb{N}$, if there exist $q, p \in \mathbb{N}$ such that $a = qb$ and $b = pc$, then

$$a = qb = q(pc) = (qp)c, \quad \text{with } qp \in \mathbb{N},$$

hence $(a, c) \in R_{\text{mul}}$.

- Antisymmetric: let $a, b \in \mathbb{N}$ with $(a, b) \in R_{\text{mul}}$ and $(b, a) \in R_{\text{mul}}$, that is $a = qb$ and $b = pa$ with $q, p \in \mathbb{N}$. Then

$$a = qb = qpa.$$

If $a = 0$, then $b = p \cdot 0 = 0 = a$. If $a \neq 0$, dividing by a we obtain $qp = 1$. Neither factor is 0 (otherwise the product would be 0), hence $q \geq 1$ and $p \geq 1$; if $q \geq 2$ we would have $qp \geq 2p \geq 2$, hence $q = 1$, and in the same way $p = 1$. Therefore $a = 1 \cdot b = b$.

Also R_{div} (“is a divisor of”) is a partial order relation, because $(a, b) \in R_{\text{div}}$ if and only if $(b, a) \in R_{\text{mul}}$:

- reflexive: $(a, a) \in R_{\text{mul}}$, hence $(a, a) \in R_{\text{div}}$;
- transitive: if $(a, b), (b, c) \in R_{\text{div}}$, then $(c, b), (b, a) \in R_{\text{mul}}$, hence $(c, a) \in R_{\text{mul}}$ by transitivity of R_{mul} , that is $(a, c) \in R_{\text{div}}$;
- antisymmetric: if $(a, b), (b, a) \in R_{\text{div}}$, then $(b, a), (a, b) \in R_{\text{mul}}$, hence $a = b$ by antisymmetry of R_{mul} .

Definition 4.16: of partially ordered set

A **partially ordered set** is a pair consisting of a set and a partial order relation defined on it.

Example 4.23

- The sets of natural, rational or real numbers with the relation $R_{\leq}$ are partially ordered sets.
- The relation “is a descendant of” defined on a subset of people is a partial order relation (if we consider individuals as descendants of themselves). Consequently, the subset of people considered with the relation “is a descendant of” is a partially ordered set.
- The set of natural numbers with the relation R_{mul} , or with the relation R_{div} , is a partially ordered set.

Observation 4.17: graph of a partial order

The directed graph of a partial order relation contains no **cycles**, that is, sequences of arcs

$$(a_1, a_2), (a_2, a_3), \dots, (a_{k-1}, a_k), (a_k, a_1)$$


with $k \geq 2$ distinct vertices $a_1, a_2, \dots, a_k$. In other words, once the loops are removed, the graph is **acyclic**.

Proof. Suppose by contradiction that the graph contains a cycle as above. If $k = 2$, the cycle consists of the arcs (a_1, a_2) and (a_2, a_1) . If $k \geq 3$, applying transitivity to the arcs (a_2, a_3) and (a_3, a_4) we obtain $(a_2, a_4) \in R$, then from (a_2, a_4) and (a_4, a_5) we obtain $(a_2, a_5) \in R$, and so on up to the arc (a_k, a_1) , from which we obtain $(a_2, a_1) \in R$. In both cases $(a_1, a_2) \in R$ and $(a_2, a_1) \in R$, and by antisymmetry $a_1 = a_2$, against the assumption that the vertices of the cycle are distinct. $\square$

- The converse does not hold: an acyclic graph does not necessarily represent a partial order relation. For example, the relation $\{(1, 2), (2, 3)\}$ on the set $\{1, 2, 3\}$ has a graph without cycles, but it is neither reflexive ($(1, 1)$ does not belong to the relation) nor transitive ($(1, 3)$ does not belong to the relation).

Example 4.24

The directed graph of the relation $R_{\subseteq}$ on the subsets of the set $\{1, 2, 3, 4\}$ is the following. To keep the figure readable, the empty set, the sets consisting of a single number (and hence the arcs leaving them) and the loops (S, S) are not shown; all the other arcs are drawn.



4.5.1 Maximum, maximal, minimum and minimal elements

Definition 4.18: of maximum, maximal, minimum and minimal element

Let (A, R) be a partially ordered set.

- An element $\tilde{a} \in A$ is the **maximum** of A if:

$$(b, \tilde{a}) \in R, \quad \forall b \in A.$$

- An element $\bar{a} \in A$ is a **maximal element** of A if there is no $b \in A$ such that:

$$(\bar{a}, b) \in R \quad \text{and} \quad b \neq \bar{a}.$$

- An element $\tilde{a} \in A$ is the **minimum** of A if:

$$(\tilde{a}, b) \in R, \quad \forall b \in A.$$

- An element $\bar{a} \in A$ is a **minimal element** of A if there is no $b \in A$ such that:

$$(b, \bar{a}) \in R \quad \text{and} \quad b \neq \bar{a}.$$

- On the directed graph: the maximum is the vertex that receives an arc from every vertex; a maximal element is a vertex from which no arc leaves, apart from the loop. For the minimum and the minimal elements, incoming and outgoing arcs are swapped.

Proposition 4.19: uniqueness of the maximum and of the minimum

In a partially ordered set (A, R) the maximum, if it exists, is unique. The same holds for the minimum.

Proof. Let $\tilde{a}_1$ and $\tilde{a}_2$ be two maxima of A . Since $\tilde{a}_2$ is a maximum, $(\tilde{a}_1, \tilde{a}_2) \in R$; since $\tilde{a}_1$ is a maximum, $(\tilde{a}_2, \tilde{a}_1) \in R$. By antisymmetry $\tilde{a}_1 = \tilde{a}_2$. For the minimum: if $\tilde{a}_1$ and $\tilde{a}_2$ are two minima, then $(\tilde{a}_1, \tilde{a}_2) \in R$ because $\tilde{a}_1$ is a minimum and $(\tilde{a}_2, \tilde{a}_1) \in R$ because $\tilde{a}_2$ is a minimum, and again $\tilde{a}_1 = \tilde{a}_2$. $\square$

Observation 4.20: the maximum is the only maximal element

If a partially ordered set (A, R) has the maximum $\tilde{a}$, then $\tilde{a}$ is a maximal element, and it is the only maximal element of A .

Proof.

- $\tilde{a}$ is maximal: if $b \in A$ and $(\tilde{a}, b) \in R$, since $\tilde{a}$ is the maximum also $(b, \tilde{a}) \in R$, and by antisymmetry $b = \tilde{a}$. Hence there is no $b \neq \tilde{a}$ with $(\tilde{a}, b) \in R$.
- $\tilde{a}$ is the only maximal element: let $\bar{a}$ be a maximal element. Since $\tilde{a}$ is the maximum,

$(\bar{a}, \tilde{a}) \in R$; since $\bar{a}$ is maximal, $\tilde{a} \neq \bar{a}$ is impossible. Hence $\bar{a} = \tilde{a}$. □


- In a partially ordered set the **maximum element** may not exist, that is, there may be no element a such that:

$$\forall b \in A, \quad (b, a) \in R$$

A partially ordered set may therefore contain several maximal elements a such that, for no $b \in A$ with $b \neq a$, we have $(a, b) \in R$.


Example 4.25: maximal elements

The directed graph of a partial order relation on the set $\{a, b, c, d, e\}$ with three maximal elements, c, d and e , in red. The maximum does not exist: for example (d, c) does not belong to the relation. The element a is the minimum.



Example 4.26

The directed graph associated with the partial order relation “is a divisor of” on the set $\{2, 3, \dots, 15\}$ is the following (the loops (a, a) , present for every a , are not drawn). The maximal elements are 8, 9, 10, 11, 12, 13, 14 and 15, which divide no other number of the set, and the maximum does not exist. The minimal elements are 2, 3, 5, 7, 11 and 13, and the minimum does not exist.



Example 4.27

Given a set of boxes of different sizes, the relation “one box is contained in the other” on the set of boxes may have several maximal boxes, that is, boxes that are not contained in any other box.

4.6 Total order relations**Definition 4.21: of total relation**

A relation R on a set A is a **total relation** if:

$$\forall a, b \in A, \quad (a, b) \in R \quad \text{or} \quad (b, a) \in R \quad (\text{or both})$$

Observation 4.22: a total relation is reflexive

Every total relation R on a set A is reflexive.

Proof. Let $a \in A$. Choosing $b = a$ in the definition of total relation we obtain that $(a, a) \in R$ or $(a, a) \in R$, that is $(a, a) \in R$. $\square$

Example 4.28

- The relation $R_{\leq}$ is a total relation.
- The relation $R_{\subseteq}$ is not a total relation since, taking for example $S_1 = \{1, 2\}$ and $S_2 = \{2, 3\}$, $(S_1, S_2) \notin R_{\subseteq}$ and $(S_2, S_1) \notin R_{\subseteq}$.
- The relation “is a descendant of” is not a total relation since there exist pairs of individuals (a, b) for which neither a descends from b nor b descends from a .
- The relation $R_{<}$ is not a total relation, because it is not reflexive: for $a = b = 1$ the condition of the definition requires $(1, 1) \in R_{<}$, which is false.
- The relation $R_{=}$ is not a total relation: for example $(3, 4) \notin R_{=}$ and $(4, 3) \notin R_{=}$.

Definition 4.23: of total order relation

A partial order relation that is also a total relation is a **total order relation**.

Example 4.29

- The relation $R_{\leq}$ is a total order relation.
- The relation $R_{=}$ is a partial order relation but not a total order relation, because it

is not a total relation.

- The relation $R_{<}$ is not a total order relation, because it is not a partial order relation.
- The relation $R_{\subseteq}$ is a partial order relation but not a total order relation: as we have seen, the sets $S_1 = \{1, 2\}$ and $S_2 = \{2, 3\}$ are not comparable.
- The relation R_{mul} is a partial order relation but not a total order relation: for example $(6, 9) \notin R_{\text{mul}}$, because there is no $k \in \mathbb{N}$ with $6 = k \cdot 9$, and $(9, 6) \notin R_{\text{mul}}$, because there is no $k \in \mathbb{N}$ with $9 = k \cdot 6$.

Example 4.30: divisibility among powers of 2

The relation

$$R = \left\{ (a, b) : a, b \in A \text{ and } \exists k \in \mathbb{N} \text{ such that } b = k a \right\}, \quad \text{with } A = \left\{ 2^n : n \in \mathbb{N} \right\},$$

that is, the relation “is a divisor of” on the powers of 2, is a total order relation.

- It is a partial order relation: reflexivity, antisymmetry and transitivity hold for R_{div} on all the natural numbers, hence they hold in particular for the elements of A .
- It is a total relation: let $a = 2^m$ and $b = 2^n$ be two elements of A , with $m, n \in \mathbb{N}$. If $m \leq n$, then

$$b = 2^n = 2^{n-m} \cdot 2^m = 2^{n-m} a, \quad \text{with } 2^{n-m} \in \mathbb{N},$$

hence $(a, b) \in R$. If instead $n < m$, in the same way $a = 2^{m-n} b$ and $(b, a) \in R$.

Definition 4.24: of totally ordered set

A **totally ordered set** is a pair consisting of a set and a total order relation defined on it.

Example 4.31

- The sets of natural, rational or real numbers with the relation $R_{\leq}$ are totally ordered sets.

Proposition 4.25: maximal element and maximum in a totally ordered set

In a totally ordered set (A, R) an element is maximal if and only if it is the maximum. Hence a totally ordered set has at most one maximal element.


Proof. If $\tilde{a}$ is the maximum, then it is maximal, as we have already proved for every partially ordered set. Conversely, let $\bar{a}$ be a maximal element and let $b \in A$. Since the relation is total, $(b, \bar{a}) \in R$ or $(\bar{a}, b) \in R$. In the second case, since $\bar{a}$ is maximal, $b = \bar{a}$, and hence $(b, \bar{a}) = (\bar{a}, \bar{a}) \in R$ by reflexivity. In both cases $(b, \bar{a}) \in R$, hence $\bar{a}$ is the

maximum. Since the maximum, if it exists, is unique, there is at most one maximal element. $\square$

- “At most” cannot be replaced by “exactly”: the totally ordered set $(\mathbb{N}, R_{\leq})$ has no maximal elements, because for every $a \in \mathbb{N}$ we have $(a, a+1) \in R_{\leq}$ with $a+1 \neq a$.

Example 4.32: maximum

The directed graph of a total order relation on the set $\{a, b, c, d, e\}$, with the maximum e in red:



4.7 Functions

Definition 4.26: of function

Given two sets A and B , a binary relation f on A and B is a **function** if, for each $a \in A$, there exists one and only one $b \in B$ such that $(a, b) \in f$.

Definition 4.27: of domain and codomain

The set A is called the **domain** of f , and the set B is called the **codomain** of f .

- We write:

$$f : A \rightarrow B$$

and if $(a, b) \in f$, we write:

$$b = f(a)$$

since b is uniquely determined by the choice of a .

- Intuitively, the function f assigns an element of B to each element of A . No element of A is associated with two different elements of B . The same element of B can however be assigned to different elements of A .

Example 4.33

- The binary relation on $\mathbb{N}$ and $\{0, 1\}$:

$$f = \left\{ (a, b) : a \in \mathbb{N}, b \in \{0, 1\} \text{ and } b = a \bmod 2 \right\},$$

where $a \bmod 2$ is the remainder of the division of a by 2, is a function $f : \mathbb{N} \rightarrow \{0, 1\}$ since for all natural numbers a , there is exactly one value $b \in \{0, 1\}$ such that $b = a \bmod 2$. For example,

$$0 = f(0), \quad 1 = f(1), \quad 0 = f(2), \dots$$

Example 4.34

- The binary relation

$$g = \left\{ (a, b) : a, b \in \mathbb{N} \text{ and } a + b \text{ is even} \right\}$$

is not a function, since for example $(1, 3)$ and $(1, 5)$ are both in g . In other words, for $a = 1$, we do not have one and only one b such that $(a, b) \in g$.

Definition 4.28: of argument and value

Given a function $f : A \rightarrow B$, if $b = f(a)$, we say that $a \in A$ is the argument of f and that $b \in B$ is the value of f associated with a .

- We can **define a function** by *directly defining its value* for all the elements of its domain.

Example 4.35

For example, we can define $f(n) = 2n$ for $n \in \mathbb{N}$, which means:

$$f = \{(n, 2n) : n \in \mathbb{N}\} \text{ and } f : \mathbb{N} \rightarrow \mathbb{N}$$

Ordered fields, supremum/infimum and the completeness axiom

5.1 Properties R_1 and R_2

- We now begin to study more closely the **structure of the number sets** that we introduced earlier, and in particular: the set $\mathbb{Q}$ of **rational numbers** and the set $\mathbb{R}$ of **real numbers**.
- We denote by a, b, c three generic **rational** or **real** numbers.

$R_1 \rightarrow$ In $\mathbb{Q}$ and $\mathbb{R}$ an operation is defined (called **addition** or **sum**) which has the following four properties:

- 1 $\forall a, b, \quad a + b = b + a$ (**commutative property**)
- 2 $\forall a, b, c, \quad (a + b) + c = a + (b + c)$ (**associative property**)
- 3 there exists a **neutral element for the sum**, denoted by 0, such that:

$$\forall a, \quad a + 0 = a$$

- 4 for every a there exists an **inverse** element of a **with respect to the sum**, called the **opposite** of a and denoted by $-a$, such that:

$$a + (-a) = 0$$

$R_2 \rightarrow$ In $\mathbb{Q}$ and $\mathbb{R}$ an operation is defined (called **multiplication** or **product**) which has the following four properties:

- 1 $\forall a, b, \quad a \cdot b = b \cdot a$ (**commutative property**)
- 2 $\forall a, b, c, \quad (a \cdot b) \cdot c = a \cdot (b \cdot c)$ (**associative property**)

3 there exists a **neutral element for the product**, denoted by 1, such that:

$$\forall a, \quad a \cdot 1 = a$$

4 for every $a \neq 0$ there exists an **inverse element of a with respect to the product**, called the **reciprocal** of a and denoted by a^{-1} , such that:

$$a \cdot a^{-1} = 1$$

- The operations of sum and product are linked by the following property:

$$\forall a, b, c, \quad (a + b) \cdot c = a \cdot c + b \cdot c \quad (\text{distributive property})$$

- From properties R_1 and R_2 follows the **possibility of performing the four fundamental operations without restrictions**:

1. **addition** and **multiplication** (defined above)

2. **subtraction**, by setting:

$$a - b = a + (-b)$$

3. **division**, by setting:

$$a/b = a \cdot b^{-1} \quad \text{provided that } b \neq 0$$

5.2 Commensurable magnitudes

In mathematics, the word **magnitude** denotes a **property** of a **geometric figure** that can be **measured**. Examples of magnitudes are: the length of a segment, the measure of an angle, the area of a plane figure or the volume of a solid.

Definition 5.1: of homogeneous magnitudes

Two magnitudes are **homogeneous** if they have the same dimension, i.e., if they can be expressed using the same unit of measurement.

- The length of a segment and the height of a solid are two homogeneous magnitudes; indeed, both can be expressed with the same unit of length.
- The area of a square and the volume of a cone are not homogeneous magnitudes; indeed, area is expressed in square meters or in other units of surface, which however cannot be used to express volume.

Definition 5.2: of commensurable and incommensurable magnitudes

Two homogeneous magnitudes a and b are called **commensurable** if their ratio is a rational number, i.e., if $a/b \in \mathbb{Q}$. They are instead called **incommensurable** if their ratio is not a rational number, i.e., if $a/b \notin \mathbb{Q}$.

Example 5.1: of commensurable magnitudes

The volume of a sphere and the volume of the cylinder circumscribed about the sphere are commensurable. In formulas, the volume V_s of the sphere as a function of its radius r is given by the formula:

$$V_s = \frac{4}{3} \pi r^3$$

The cylinder circumscribed about it has base radius equal to the radius r of the sphere and height h equal to twice the radius. The volume V_c of this cylinder is therefore given by the formula:

$$V_c = \pi r^2 h = \pi r^2 2r = 2\pi r^3$$

Taking the ratio we obtain:

$$\frac{V_s}{V_c} = \frac{\frac{4}{3} \pi r^3}{2 \pi r^3} = \frac{2}{3} \in \mathbb{Q}$$

Example 5.2: of incommensurable magnitudes

The length of a circle and the length of its diameter are incommensurable. In formulas, denoting by d the diameter of the circle and by c the length of the circle, we have:

$$c = \pi d$$

Now, taking the ratio between the length of the circle and its diameter, we obtain:

$$\frac{c}{d} = \frac{\pi d}{d} = \pi \notin \mathbb{Q}$$

A proof that $\pi \notin \mathbb{Q}$ will be given later.


5.3 Geometric representation of $\mathbb{Q}$

A geometric representation of $\mathbb{Q}$ can be obtained by associating with each rational number a point of the **Euclidean line** (which can be imagined as a line in the plane of infinite length)

- With an arbitrarily chosen point of the Euclidean line we associate 0, and with another point, distinct from the first, we associate 1, thus identifying the oriented segment 01, which constitutes the **unit of measurement**

- At this point we have a **one-to-one correspondence** between the rational numbers and those points P of the line that are endpoints of oriented segments $0P$ **commensurable** with 01 . Indeed, if the rational number p (its **abscissa**) is associated with the point P , the segment $0P$ measures p times the unit segment 01 , and therefore the ratio of the two lengths is:

$$\frac{0P}{01} = \frac{p}{1} = p \in \mathbb{Q}$$



5.4 Total order relations

Observation 5.3

The sets of rational numbers $\mathbb{Q}$ and of real numbers $\mathbb{R}$ with the relation “less than or equal to” (“ $\leq$ ”) are totally ordered sets.

- The relations “less than or equal to” (“ $\leq$ ”) for the set of rational or real numbers:

$$R_{\leq} = \left\{ (a, b) : a, b \in \mathbb{Q} \quad \text{and} \quad a \leq b \right\}, \quad R_{\leq} = \left\{ (a, b) : a, b \in \mathbb{R} \quad \text{and} \quad a \leq b \right\}$$

are **partial order relations**. That is, they satisfy the following properties:

$$\forall a, \quad a \leq a \quad (\text{reflexive property})$$

$$\forall a, b, \quad a \leq b \quad \text{and} \quad b \leq a \Rightarrow a = b \quad (\text{antisymmetric property})$$

$$\forall a, b, c, \quad a \leq b \quad \text{and} \quad b \leq c \Rightarrow a \leq c \quad (\text{transitive property})$$

Moreover, the relations “less than or equal to” (“ $\leq$ ”) for $\mathbb{Q}$ and $\mathbb{R}$ are **total relations** since:

$$\forall a, b, \quad a \leq b \quad \text{or} \quad b \leq a$$

- Consequently, the pair consisting of the set $\mathbb{Q}$ or $\mathbb{R}$ and the corresponding relation $R_{\leq}$ is a totally ordered set.

Observation 5.4: trichotomy property

For every pair of numbers a, b (rational or real) **exactly one** of the three relations holds:

$$a < b, \quad a = b, \quad a > b$$

- This is not a new property: it is a consequence of the fact that “ $\leq$ ” is a **total** order relation, as the proof below shows. For this reason we do not count it among the axioms R_1, R_2, R_3 and R_4 .

Proof. Recall that $a < b$ means $a \leq b$ and $a \neq b$.

At least one of the three relations holds. Since the relation is total, we have $a \leq b$ or $b \leq a$:

- if $a \leq b$ and $a \neq b$, then $a < b$;
- if $b \leq a$ and $a \neq b$, then $a > b$;
- if $a = b$ the second relation holds.

At most one of the three relations holds. The relations $a < b$ and $a = b$ cannot hold together, because the first one requires $a \neq b$; for the same reason $a > b$ and $a = b$ cannot hold together. Finally, if $a < b$ and $a > b$ held together, we would have $a \leq b$ and $b \leq a$, and hence $a = b$ by the antisymmetric property, contradicting $a \neq b$. $\square$

5.5 Property R_3

- We now highlight the properties of the ordering of rational or real numbers:

$R_3 \rightarrow$ In $\mathbb{Q}$ and $\mathbb{R}$ the total order relation “less than or equal to” (“ $\leq$ ”) is defined, compatible with the algebraic structure ^a, that is:

1. $\forall a, b, c, \quad a \leq b \Rightarrow a + c \leq b + c$
2. $\forall a, b \text{ and } \forall c > 0, \quad a \leq b \Rightarrow a \cdot c \leq b \cdot c$

^aan algebraic structure is a set, called the underlying set (of the structure), equipped with one or more operations

- We observe that all the rules of algebraic computation follow from the properties: R_1, R_2, R_3 .

Example 5.3

For example, all the usual procedures used to solve inequalities are a consequence of the axioms and of the algebraic properties of the sum and the product, expressed by R_1 and R_2 , **and of the compatibility of the order relation with these operations**, expressed by R_3 . It is indeed R_3 that allows one to add the same number to both sides of an inequality and to multiply them by the same positive number without changing the direction of the inequality.

5.6 Ordered fields**Definition 5.5: of (ordered) field**

A **field** is a set on which two operations (sum and product) are defined, satisfying the properties R_1 and R_2 and the **distributive property**.

An **ordered field** is a field on which a total order relation satisfying the property R_3 is also defined.


- The distributive property is part of the definition of a field: it is the only property that links the sum and the product to each other, and without it the two operations would remain independent.
- Everything we have said so far about the operations of sum and product and about the total order relation “less than or equal to” (“ $\leq$ ”) holds both for the set of rational numbers and for the set of real numbers.

Observation 5.6

The sets of rational numbers $\mathbb{Q}$ and of real numbers $\mathbb{R}$ are ordered fields.

- From this point of view, therefore, $\mathbb{Q}$ and $\mathbb{R}$ appear to have similar properties. In what follows we will highlight the **property that substantially distinguishes $\mathbb{R}$ from $\mathbb{Q}$** and that makes $\mathbb{R}$ the **right setting for developing mathematical analysis**.
- We begin by observing that the **set of rational numbers is inadequate for expressing the lengths of segments** (but also areas, volumes, times, speeds, etc.)

As we have seen, there exist magnitudes that are not **commensurable** with each other. The classic example is given by the diagonal and the side of a square: if the side measures 1, the abscissa d that measures the diagonal is not a rational number.



Indeed, by the Pythagorean Theorem we have:

$$d^2 = 1^2 + 1^2 = 2$$

but, as we have seen, there is no rational number whose square is 2. Hence the point d on the line does not represent any rational number. This means that, after “occupying” the points of the line with the rational numbers, there are still some **empty places** left on it.

5.7 Bounded sets, maxima and minima

Definition 5.7: of bounded set

Let E be a set contained in $\mathbb{Q}$ or in $\mathbb{R}$. The set E is called **bounded** if there exist two numbers m and M such that:

$$\forall x \in E, \quad m \leq x \leq M$$

- E is **bounded below** if there exists a number m such that:

$$\forall x \in E, \quad m \leq x$$

- E is **bounded above** if there exists a number M such that:

$$\forall x \in E, \quad x \leq M$$

- It makes no difference whether we require m and/or M to belong to $\mathbb{Q}$ or to $\mathbb{R}$, since the condition only concerns the existence of such numbers. This is not an immediate consequence of the definition, though, because $\mathbb{Q}$ is a proper subset of $\mathbb{R}$: it follows from the **Archimedean property** (recalled in the chapter *Sets* and proved later in this chapter). Indeed, if $M \in \mathbb{R}$ satisfies $x \leq M$ for every $x \in E$, there exists a natural number $n > M$, and hence $x \leq M < n$ for every $x \in E$, with $n \in \mathbb{Q}$; in the same way, if $m \in \mathbb{R}$ satisfies $m \leq x$ for every $x \in E$, there exists a natural number $n > -m$, that is $-n < m$, and hence $-n < x$ for every $x \in E$, with $-n \in \mathbb{Q}$. Conversely, every rational number is also a real number, and therefore the two requirements single out the same bounded sets.

Definition 5.8: of maximum and minimum of a set

An element $x_M \in E$ is called the **maximum** of E if:

$$\forall x \in E, \quad x \leq x_M$$

An element $x_m \in E$ is called the **minimum** of E if:

$$\forall x \in E, \quad x_m \leq x$$

- Note that the maximum and the minimum, by definition, **belong** to the set E .

Observation 5.9

If a set E has a maximum, this is **unique**; in the same way, if it has a minimum, this is unique. We then write $\max E$ and $\min E$.

Proof. Let x_M and x'_M be two maxima of E . Since x'_M belongs to E and x_M is a maximum of E , we have $x'_M \leq x_M$. Exchanging the roles, since x_M belongs to E and x'_M is a maximum of E , we have $x_M \leq x'_M$. From the antisymmetric property of the order relation it follows that:

$$x'_M \leq x_M \quad \text{and} \quad x_M \leq x'_M \quad \implies \quad x_M = x'_M$$

The proof for the minimum is identical: if x_m and x'_m are two minima of E , then $x_m \leq x'_m$ and $x'_m \leq x_m$, and hence $x_m = x'_m$. $\square$

The existence of the maximum and of the minimum of a set implies that the set is bounded, that is:

$$\text{existence of maximum and minimum} \implies \text{bounded set} \quad (5.1)$$

but the converse is not true (we will give a counterexample later):

$$\text{bounded set} \not\Rightarrow \text{existence of maximum and minimum} \quad (5.2)$$

Hence the fact that a set is bounded is a necessary but not sufficient condition for the set to have a maximum and a minimum. Moreover, the existence of a maximum and a minimum is a sufficient but not necessary condition for a set to be bounded.

From the **contrapositive** of (5.1) we have:

$$\text{unbounded set} \implies \text{the maximum does not exist or the minimum does not exist}$$


that is, if a set is not bounded, it has no maximum or it has no minimum (it may well have one of the two: $\mathbb{N}$ is unbounded and nevertheless has minimum 0).

Example 5.4: of maxima and minima

Example	set E	minimum	maximum
I)	$\mathbb{N}$	0	does not exist
II)	even integers	does not exist	does not exist

Example 5.5: of maxima and minima

Example	set E	minimum	maximum
III)	$\left\{ \frac{1}{n} : n \in \mathbb{N}_{>0} \right\}$	does not exist	1




Example 5.6: of maxima and minima

Example	set E	minimum	maximum
IV)	$\left\{ \frac{n-1}{n+1} : n \in \mathbb{N} \right\}$	-1	does not exist

We have:

$$\frac{n-1}{n+1} = \frac{n+1-1-1}{n+1} = 1 - \frac{2}{n+1}$$



Note that sometimes, even though the set is bounded, it may have no maximum or no minimum.

Example 5.7: of maxima and minima

Example	set E	minimum	maximum
V)	$\{x \in \mathbb{R} : 27 \leq x^3\}$	3	does not exist


We have:

$$27 \leq x^3 \iff \sqrt[3]{27} = 3 \leq x \quad \text{and} \quad 3 \in E$$

Example 5.8: of maxima and minima

Example	set E	minimum	maximum
VI)	$\{x \in \mathbb{Q} : x \geq 0, x < \sqrt{2}\}$	0	does not exist


The value $\sqrt{2}$ is irrational, hence by truncating we obtain rational numbers that get closer and closer to $\sqrt{2}$, none of which has the property of being the maximum.



It is an example of a bounded set that has no maximum.

Example	set E	minimum	maximum
VII)	$\{x \in \mathbb{Q} : x > \sqrt{2}, x \leq 4\}$	does not exist	4

The value $\sqrt{2}$ is irrational, hence by truncating and rounding up we obtain rational numbers that get closer and closer to $\sqrt{2}$, none of which has the property of being the minimum.



It is an example of a bounded set that has no minimum.

Example 5.9: of maxima and minima of intervals

Example	set E	minimum	maximum
VIII)	$[-2, 1]$	-2	1
IX)	$(-\infty, 3)$	does not exist	does not exist

- In example VIII) the interval contains both of its endpoints: $-2 \in [-2, 1]$ and $-2 \leq x$ for every $x \in [-2, 1]$, hence -2 is the minimum; $1 \in [-2, 1]$ and $x \leq 1$ for every $x \in [-2, 1]$, hence 1 is the maximum.
- In example IX) the interval is not bounded below and therefore it has no minimum. It has no maximum either: given any $x \in (-\infty, 3)$, the number

$$y = \frac{x+3}{2} \quad \text{satisfies} \quad x < y < 3$$

indeed $y - x = \frac{3-x}{2} > 0$ and $y < 3 \Leftrightarrow x < 3$. Hence $y \in (-\infty, 3)$ and it is strictly greater than x : no element of the set can be the maximum.

Example 5.10: of the maximum and the minimum of a set defined by an inequality

Example	set E	minimum	maximum
X)	$\{x \in \mathbb{R} : -5 \leq 3x < 4\}$	$-\frac{5}{3}$	does not exist

It is convenient to **solve the inequality** first: dividing by $3 > 0$, and using the property

R_3 which preserves the direction of the inequality, we obtain

$$-5 \leq 3x < 4 \quad \Longleftrightarrow \quad -\frac{5}{3} \leq x < \frac{4}{3} \quad \text{that is} \quad E = \left[-\frac{5}{3}, \frac{4}{3} \right)$$

- $-\frac{5}{3} \in E$ and it is less than or equal to every element of E : it is therefore the minimum.
- The maximum does not exist: as in example IX), given $x \in E$, the number $y = \frac{1}{2}(x + \frac{4}{3})$ satisfies $x < y < \frac{4}{3}$ and is still an element of E strictly greater than x .

5.8 Upper/lower bounds and suprema/infima

Definition 5.10: of upper bound of a set

Let E be a set contained in $\mathbb{Q}$ or in $\mathbb{R}$; a number k belonging to $\mathbb{Q}$ or to $\mathbb{R}$ respectively is called an **upper bound** of E if:

$$\forall x \in E, \quad x \leq k$$

Definition 5.11: of lower bound of a set

Let E be a set contained in $\mathbb{Q}$ or in $\mathbb{R}$; a number k belonging to $\mathbb{Q}$ or to $\mathbb{R}$ respectively is called a **lower bound** of E if:

$$\forall x \in E, \quad k \leq x$$

We observe that a set bounded above (below) has many upper (lower) bounds. Moreover, the upper (lower) bounds do not necessarily belong to the set itself.

Observation 5.12

If k is an upper bound of E , then every number $h \geq k$ is still an upper bound of E ; if k is a lower bound of E , then every number $h \leq k$ is still a lower bound of E . Hence the set of the upper (lower) bounds of E , if it is not empty, is a **half-line**.

Proof. Let k be an upper bound of E and let $h \geq k$. For every $x \in E$ we have $x \leq k$ by definition of upper bound, and hence, by the transitive property,

$$x \leq k \leq h \quad \Longrightarrow \quad x \leq h$$

that is, h is an upper bound of E . The set of the upper bounds therefore contains, together with each of its elements k , all the numbers greater than k : it is thus a half-line unbounded above.

The proof for the lower bounds is analogous: if k is a lower bound of E and $h \leq k$, then for every $x \in E$ we have $h \leq k \leq x$ and hence $h \leq x$. $\square$

Observation 5.13

An upper bound of E that **belongs** to E is the maximum of E ; a lower bound of E that belongs to E is the minimum of E .

Proof. Let k be an upper bound of E with $k \in E$. By definition of upper bound we have $x \leq k$ for every $x \in E$: these are exactly the two conditions that define the maximum of E (belonging to E and being greater than or equal to all its elements), and hence $k = \max E$.

In the same way, if k is a lower bound of E and $k \in E$, then $k \leq x$ for every $x \in E$ and $k \in E$, that is $k = \min E$. $\square$

Definition 5.14: of supremum

The **supremum** of E (denoted by $\sup E$) is the minimum of the upper bounds of E .

Definition 5.15: of infimum

The **infimum** of E (denoted by $\inf E$) is the maximum of the lower bounds of E .

We observe that the supremum (infimum) may not exist (unbounded sets). We also observe that if the set has a maximum (minimum), it coincides with the supremum (infimum).

For unbounded sets the following **convention** is adopted: if E is not bounded above one writes

$$\sup E = +\infty$$

and if E is not bounded below one writes

$$\inf E = -\infty$$

The symbols $+\infty$ and $-\infty$ **are not numbers**: this notation is only a compact way of saying that E has no upper (lower) bounds.

Example 5.11: of suprema and infima of intervals

Example	set E	$\inf E$	$\sup E$	minimum	maximum
VIII)	$[-2, 1]$	-2	1	-2	1
IX)	$(-\infty, 3)$	$-\infty$	3	does not exist	does not exist
X)	$\{x \in \mathbb{R} : -5 \leq 3x < 4\}$	$-\frac{5}{3}$	$\frac{4}{3}$	$-\frac{5}{3}$	does not exist

- In examples VIII) and X) the minimum exists and therefore coincides with the infimum; in example VIII) the maximum exists as well and coincides with the supremum.
- In example IX) the set is not bounded below and therefore, by the convention just introduced, $\inf E = -\infty$. The number 3 is an upper bound, because $x < 3$ for every

$x \in E$; no number $k < 3$ is an upper bound, because the number $y = \frac{k+3}{2}$ satisfies $k < y < 3$ and therefore belongs to E . Hence 3 is the minimum of the upper bounds, that is $\sup E = 3$ (and it is not a maximum, because $3 \notin E$).

- In example X), with the same argument applied to $E = [-\frac{5}{3}, \frac{4}{3})$, we obtain $\sup E = \frac{4}{3} \notin E$.
- Finally, going back to example III), for $E = \{\frac{1}{n} : n \in \mathbb{N}_{>0}\}$ we have $\inf E = 0$ and $\sup E = \max E = 1$. Indeed 0 is a lower bound, because $\frac{1}{n} > 0$ for every $n \in \mathbb{N}_{>0}$; and no number $k > 0$ is a lower bound, because by the **Archimedean property** there exists $n \in \mathbb{N}_{>0}$ with $n > \frac{1}{k}$, and hence $\frac{1}{n} < k$. Hence 0 is the maximum of the lower bounds, but it is not the minimum because $0 \notin E$.

Example 5.12: of suprema and infima

Example	set E	$\inf E$	$\sup E$	minimum	maximum
I)	$\{x \in \mathbb{Q} : x \geq 0, x < \sqrt{2}\}$	0	does not exist in $\mathbb{Q}$	0	does not exist

Here the upper bounds are looked for **inside** $\mathbb{Q}$. We have:

$$\sqrt{2} \notin E \quad \text{and} \quad \sqrt{2} \notin \mathbb{Q}$$

hence the maximum does not exist; and the sup does not exist **in** $\mathbb{Q}$ either, because the rational upper bounds of E are the rational numbers greater than $\sqrt{2}$ and among them there is no smallest one (by truncating $\sqrt{2}$ from above one obtains smaller and smaller rational upper bounds). As the next example shows, in $\mathbb{R}$ the supremum does exist and is equal to $\sqrt{2}$.

Example	set E	$\inf E$	$\sup E$	minimum	maximum
II)	$\{x \in \mathbb{R} : x \geq 0, x < \sqrt{2}\}$	0	$\sqrt{2}$	0	does not exist

We have:

$$\sqrt{2} \notin E \quad \text{and} \quad \sqrt{2} \in \mathbb{R}$$

that is, the maximum does not exist, but $\sqrt{2}$ is the minimum of the upper bounds and therefore it is the sup.

5.9 Axiomatic definition of the real numbers

A totally ordered number set X is said to have the **supremum property** (least upper bound property) if:

$R_4 \rightarrow$ Every non-empty set $E \subset X$ that is bounded above has a supremum in X

- It is easy to prove that, if this property holds, then it is also true that every non-empty subset of X that is bounded below has an infimum.
- It is not required that X itself has a supremum (for example, in the case $X = \mathbb{Q}$ or $\mathbb{R}$ this is certainly false!), but that every non-empty subset of X that is bounded above has one.

Example VI shows that $\mathbb{Q}$ certainly does not have the supremum property. On the other hand, $\mathbb{R}$ has this property. ^a

^aHowever, we will not present a formal proof.

- In the **axiomatic definition** of $\mathbb{R}$, this property is part of the very definition of $\mathbb{R}$

Definition 5.16: (axiomatic) of the real numbers

We call $\mathbb{R}$ a set that satisfies the properties R_1, R_2 (together with the distributive property), R_3 and R_4 , that is, an ordered field that has the supremum property

Proposition 5.17: Archimedean property

For every real number x there exists a natural number n such that:

$$n > x$$

- This is the property that we recalled in the chapter *Sets* and used above. **It is not an additional axiom**: as the proof shows, it follows from the property R_4 .

Proof. Suppose by contradiction that there exists a real number x such that $n \leq x$ for every $n \in \mathbb{N}$. Then the set $\mathbb{N}$ is a non-empty subset of $\mathbb{R}$ bounded above (by x), and therefore, by the property R_4 , it has a supremum in $\mathbb{R}$:

$$s = \sup \mathbb{N}$$

Since s is the **minimum** of the upper bounds of $\mathbb{N}$, the number $s - 1$, which is smaller than s , is not an upper bound of $\mathbb{N}$: there exists therefore $n \in \mathbb{N}$ such that

$$n > s - 1 \quad \implies \quad n + 1 > s$$

But $n + 1$ is still a natural number, and this contradicts the fact that s is an upper bound of $\mathbb{N}$. The initial assumption is therefore absurd, and for every $x \in \mathbb{R}$ there exists $n \in \mathbb{N}$ with $n > x$. $\square$

- Equivalently: for every real number $\varepsilon > 0$ there exists $n \in \mathbb{N}_{>0}$ such that $\frac{1}{n} < \varepsilon$. Indeed it is enough to choose $n > \frac{1}{\varepsilon}$, and then $\frac{1}{n} < \varepsilon$. As a consequence, the only non-negative number that is less than or equal to $\frac{1}{n}$ for every $n \in \mathbb{N}_{>0}$ is zero.
- The supremum property is also called the **Dedekind axiom, or axiom of continuity**, or **completeness axiom**, and it can also be stated in the following equivalent form.

Let $\{A, B\}$ be a partition of $\mathbb{R}$ (i.e., A and B are non-empty disjoint sets whose union is $\mathbb{R}$); the partition is called a **cut** (section) if:

$$\forall a \in A \text{ and } \forall b \in B \text{ we have } a < b.$$

Then one proves that:

$R'_4 \rightarrow$ For every cut $\{A, B\}$ of $\mathbb{R}$ there exists a unique real number s (called the **separating element**) such that:


$$\forall a \in A \text{ and } \forall b \in B, \quad a \leq s \leq b$$

This separating element is $\sup A$ and $\inf B$, hence $\sup A = \inf B$

5.10 Geometric representation of $\mathbb{R}$

Can the set of real numbers $\mathbb{R}$ be put in one-to-one correspondence with the points of the Euclidean line?

- Let us now return to the problem of the incommensurability of the side and the diagonal of a square. When, with straightedge and compass, having fixed the unit side of the square, we construct its diagonal and transfer it onto the starting line, the arc drawn by the compass "splits the line in two", generating what we have called a cut; the separating element, which exists by axiom R_4 , geometrically represents the point of intersection, and therefore that number measures the diagonal.



Hence the geometric interpretation of property R_4 shows us how the set $\mathbb{R}$ is an adequate representation of our intuitive idea of a line, so adequate that **in mathematics the expression “the real line” is often used to denote $\mathbb{R}$** , identifying the number set with its natural geometric representation.

Roots, powers, logarithms and modular arithmetic

6.1 Roots, powers, logarithms

- As a consequence of property R_4 we can perform, in the real field, operations that are only occasionally possible in the rational field, such as extracting roots or raising to a power.

6.1.1 Arithmetic n -th roots

Theorem 6.1

For every $y \in \mathbb{R}$, $y > 0$ and $n \in \mathbb{N}$, $n \geq 1$, there exists one and only one $x \in \mathbb{R}$, $x > 0$, such that $x^n = y$.

- This number is called the arithmetic n -th root of y and is denoted by one of the symbols

$$\sqrt[n]{y} \text{ or } y^{\frac{1}{n}}.$$

Proof. The proof of this Theorem will be given using the properties of continuous functions. □

The arithmetic n -th root is non-negative.

Example 6.1: arithmetic n -th root

For example:

$$\sqrt{4} = 2, \quad \sqrt{9} = 3.$$

Moreover, we have:

$$\sqrt{x^2} = |x| \quad (6.1)$$

Construction of the decimal representation of the n -th root.

- We look for the decimal expansion of

$$\sqrt{2} \approx 1.41421356237$$

this number, not being rational, will be represented by an infinite (non-periodic) expansion.

- We proceed as follows: we build a class of rational numbers of the form:

$$\begin{aligned} 0 &< a_0 \\ &< a_0, a_1 \\ &< a_0, a_1 a_2 \\ &< a_0, a_1 a_2 a_3 \\ &< \dots \\ &< a_0, a_1 a_2 a_3 \dots a_n \\ &< \dots \end{aligned}$$

The rule is: each of these numbers is the largest among those with the same number of digits after the decimal point whose square is less than 2. The first of these numbers are:

$$\begin{array}{lll} 1 & 1^2 & = 1 \\ 1.4 & (1.4)^2 & = 1.96 \\ 1.41 & (1.41)^2 & = 1.9881 \\ 1.414 & (1.414)^2 & = 1.999396 \\ 1.4142 & (1.4142)^2 & = 1.99996164 \\ \dots & & \end{array}$$

- This set of numbers, which we call E_- , is bounded above (each one is < 2); by property R_4 it has a supremum, and this is used to define $\sqrt{2}$.

The number $\sqrt{2}$ is defined precisely as $\sup E_-$

- We could also have built a class of numbers E_+ like the previous one, with the rule that each of them is the smallest among those with the same number of digits after the decimal point whose square is greater than 2; we would have obtained:

$$\begin{array}{rcl}
 2 & 2^2 & = 4 \\
 1.5 & (1.5)^2 & = 2.25 \\
 1.42 & (1.42)^2 & = 2.0164 \\
 1.415 & (1.415)^2 & = 2.002225 \\
 1.4143 & (1.4143)^2 & = 2.00024449 \\
 \dots & &
 \end{array}$$

This set E_+ is bounded below (every element is > 1), therefore it has an infimum; one proves that:

$$\sqrt{2} = \inf E_+ = \sup E_-$$

The numbers of the class E_- approximate $\sqrt{2}$ from below, those of the class E_+ from above.

6.1.2 Powers with real exponent

Extracting the n -th root is the inverse operation of raising to an integer power.

Rational exponent

- The operation of raising to a power can be extended to every rational exponent if the base is positive (using the previous theorem).

$$\text{If } r = \frac{m}{n} \quad \text{and} \quad a > 0 \quad \text{then} \quad a^r = (a^m)^{\frac{1}{n}} = \sqrt[n]{a^m}$$

(we assume $m \in \mathbb{Z}$ and $n \in \mathbb{Z}$ positive)

Real exponent

- If the exponent is real, $b = b_0, b_1 b_2 b_3 \dots b_n \dots$, the number a^b ($a > 0$) will be determined by the class of numbers

$$a^{b_0} \quad a^{b_0, b_1} \quad a^{b_0, b_1 b_2} \quad \dots$$

in a way similar to the case of the root.

If the base a is negative, the operation of raising to a power a^b is defined only in certain cases:

- if the exponent b is an integer, or
- if the exponent $b = \frac{n}{m}$ is rational, *written in lowest terms*, provided that it is not the case that n is odd and m is even.

If $c < 0$ and m is odd, we define

$$\sqrt[m]{c} = -\sqrt[m]{-c}$$

Example 6.2: negative base

For example:

$$(-2)^{\frac{3}{5}} = \sqrt[5]{(-2)^3} = \sqrt[5]{-8} = -\sqrt[5]{8}$$

$$(-2)^{\frac{2}{5}} = \sqrt[5]{(-2)^2} = \sqrt[5]{4}$$

The assumption that the fraction is written in lowest terms is essential: $\frac{1}{3}$ and $\frac{2}{6}$ are the same rational number, but

$$\sqrt[3]{(-8)^1} = \sqrt[3]{-8} = -2 \quad \text{while} \quad \sqrt[6]{(-8)^2} = \sqrt[6]{64} = 2.$$

Only the first expression, with $\frac{n}{m} = \frac{1}{3}$ in lowest terms, defines $(-8)^{\frac{1}{3}}$.

- When we say “does not exist in $\mathbb{R}$ ” we mean that it is not possible to define this operation in such a way that the usual rules of computation remain valid.

Example 6.3: non-existence of a power with real exponent

For example:

$$(-2)^{\frac{3}{4}}$$

does not exist in the real field!

Let

$$a, b \in \mathbb{R}, \quad a > 0 \quad \text{and} \quad b > 0 \quad (\text{positive reals})$$

and

$$c, d \in \mathbb{R} \quad (\text{arbitrary reals}),$$

the main properties of raising to a power are:

$$a^0 = 1 \quad \forall a \neq 0 \quad (6.2a)$$

$$1^c = 1 \quad \forall c \quad (6.2b)$$

$$a^c > 0 \quad \forall c \quad (6.2c)$$

$$a^c > 1 \quad \forall a > 1 \quad \text{and} \quad c > 0 \quad (6.2d)$$

$$a^c < 1 \quad \forall a < 1 \quad \text{and} \quad c > 0 \quad (6.2e)$$

$$a^{c+d} = a^c \cdot a^d \quad (6.3a)$$

$$(a \cdot b)^c = a^c \cdot b^c \quad (6.3b)$$

$$(a^b)^c = a^{b \cdot c} \quad (6.3c)$$

$$c < d \Rightarrow a^c < a^d \text{ if } a > 1 \quad (6.4a)$$

$$c < d \Rightarrow a^c > a^d \text{ if } a < 1 \quad (6.4b)$$

$$0 < a \leq b \Rightarrow a^c \leq b^c \quad \forall c > 0 \quad (6.4c)$$

6.1.3 Logarithms

- Consider the equation

$$a^x = y, \quad a > 0$$

with y given and x unknown. First of all, if $a = 1$, it is solvable only if $y = 1$ (and in that case every real number x is a solution). So let $a \neq 1$.

If $y \leq 0$ it has no solution, see property (6.2c).

The following theorem tells us that it has exactly one solution for every $y > 0$:

Theorem 6.2

Let $a > 0$, $a \neq 1$, $y > 0$. There exists a unique real number x such that $a^x = y$.

- This number is called the logarithm to base a of y and is denoted by the symbol:

$$\log_a y$$

- The logarithm is the inverse operation of raising to a power.
- the logarithm of a number to a given base is the exponent to which the base must be raised to obtain the number itself.

Let

$$x, y, a \in \mathbb{R}, \quad x > 0, y > 0, a > 0 \quad (\text{positive reals}) \quad \text{and} \quad a \neq 1,$$

the main properties of logarithms are (they follow from those of exponentials):

$$\log_a(x \cdot y) = \log_a x + \log_a y \quad (6.5a)$$

$$\log_a\left(\frac{x}{y}\right) = \log_a x - \log_a y \quad (6.5b)$$

$$\log_a\left(\frac{1}{y}\right) = -\log_a y \quad (6.5c)$$

$$\log_a x^\alpha = \alpha \log_a x \quad (\forall \alpha \in \mathbb{R}) \quad (6.6a)$$

$$\log_b x = \frac{\log_a x}{\log_a b} \quad (\forall b > 0, b \neq 1) \quad (6.6b)$$

$$\log_a x = \frac{1}{\log_x a} = -\log_{\frac{1}{a}} x \quad (x \neq 1) \quad (6.6c)$$

Notation:

- $\lg n = \log_2 n \rightarrow$ **binary logarithm**
- $\ln n = \log_e n \rightarrow$ **natural logarithm** (where $e \approx 2.718$ is **Euler's number** (Napier's constant)) (sometimes the natural logarithm $\ln$ is also denoted by $\log$)
- $\lg^k n = (\lg n)^k \rightarrow$ **power**
- $\lg \lg n = \lg(\lg n) \rightarrow$ **composition**

6.1.4 Approximations

- a rational number can always be expressed with absolute precision, using either the fractional or the decimal notation (possibly with repeating digits).
- On the other hand, it is of course not possible to write all the decimal digits of an irrational number, since there are infinitely many of them and they follow one another without periodicity.
- What does it mean, then, to “know” or “specify” an irrational number? It means knowing some algorithm that allows us (at least in theory) to write as many exact decimal digits as we wish.

Example 6.4: irrational numbers

- In the (already considered) example of the irrational number:

0.101001000100001...

built according to the rule: write one digit 1, one digit 0, one digit 1, two digits 0, one digit 1, three digits 0, and so on) it is clear that we could write as many digits as we wish.

- In other cases, such as

$$\sqrt{2} \text{ or } \log_2 3,$$

things are more laborious and require iterative computations to determine each successive decimal digit; nevertheless, these computations can actually be carried out.

- At other times, in Mathematical Analysis, a number is specified in a non-constructive way, by denoting it as the unique number that solves a given problem (once it has been proved that such a solution exists and is unique). This is a less satisfactory approach from an operational point of view, but theoretically flawless.

Practical aspects of computing irrational numbers

- Whenever we perform computations with a **pocket calculator**, it will only display numbers with a fixed number of decimal digits after the decimal point (typically 9). This means that we are working only with rational numbers, indeed with a finite subset of the set $\mathbb{Q}$.
- Working with a **computer** things improve somewhat, but we still remain within finite subsets of $\mathbb{Q}$. Of course, one needs to be aware of this.
- Besides the approximation produced by computing tools, sometimes we ourselves are not interested in too many decimal digits; we then deliberately introduce approximations, writing for example

$$\sqrt{2} \approx 1.414$$

Rounding rule: the last digit written is rounded down (up) if the first digit discarded is from 0 to 4 (respectively, from 5 to 9).

Example 6.5: Rounding

For example:

$$2.4138 \approx 2.41$$

$$2.4152 \approx 2.42$$

6.2 Modular arithmetic

Definition 6.3: of integer part

Given a real number $a \in \mathbb{R}$, we denote by $[a]$ or $\lfloor a \rfloor$ the **integer part** (or “**floor**”) of a :

$$[a] = \text{integer } n \text{ such that } n \leq a < n + 1$$

Equivalently, the integer part of a is the **largest** integer less than or equal to a :

$$\lfloor a \rfloor = \max \{n \in \mathbb{Z} : n \leq a\}.$$

- The two descriptions coincide. If $n \leq a < n + 1$, then n belongs to the set $\{k \in \mathbb{Z} : k \leq a\}$ and every other element k of it satisfies $k \leq a < n + 1$, that is, $k \leq n$ (since k and n are integers): hence n is its maximum. Conversely, if n is the maximum of that set, then $n \leq a$ and moreover $a < n + 1$, since otherwise $n + 1$ would belong to the set and n would not be its maximum.

Example 6.6: Integer part

$$[2.38] = 2; \quad [3] = 3; \quad [-1.8] = -2.$$

- While for positive numbers the integer part is obtained simply by “throwing away the digits after the decimal point”, for negative numbers one must take the largest integer $\leq a$, which is different from what is obtained by throwing away the digits after the decimal point (except when a is already an integer)

Definition 6.4: of fractional part

Given a real number $a \in \mathbb{R}$, we denote by (a) the **fractional part** (or decimal part, in Italian “*mantissa*”) of a :

$$(a) = a - [a]$$

Example 6.7: Fractional part

$$(2.38) = 0.38; \quad (3) = 0; \quad (-1.8) = 0.2.$$

- The fractional part is therefore not an integer but a real number, lying in $[0, 1)$. Indeed, setting $n = [a]$, the definition of integer part gives $n \leq a < n + 1$ and, subtracting n ,

$$0 \leq \underbrace{a - n}_{(a)} < 1.$$

- For positive numbers, it is obtained simply by “throwing away the digits before the decimal point”; for negative numbers the fractional part is the **one’s complement** of the number obtained by throwing away the digits before the decimal point.

Definition 6.5: of upper integer part

Given a real number $a \in \mathbb{R}$, we denote by $\lceil a \rceil$ the **upper integer part** (or “**ceil**”) of a :

$$\lceil a \rceil = \text{integer } n \text{ such that } n - 1 < a \leq n$$

Equivalently, the upper integer part of a is the **smallest** integer greater than or equal to a :

$$\lceil a \rceil = \min \{n \in \mathbb{Z} : n \geq a\}.$$

- Here too the two descriptions coincide. If $n - 1 < a \leq n$, then n belongs to the set $\{k \in \mathbb{Z} : k \geq a\}$ and every other element k of it satisfies $k \geq a > n - 1$, that is, $k \geq n$: hence n is its minimum. Conversely, if n is the minimum of that set, then $a \leq n$ and moreover $n - 1 < a$, since otherwise $n - 1$ would belong to the set and n would not be its minimum.

Example 6.8: of ceil

$$\lceil 2.38 \rceil = 3; \quad \lceil 3 \rceil = 3; \quad \lceil -1.8 \rceil = -1.$$

Some properties:

Given a real number $a \in \mathbb{R}$, we have

$$a - 1 < \lfloor a \rfloor \leq a \leq \lceil a \rceil < a + 1$$

- This follows at once from the two definitions: from $\lfloor a \rfloor \leq a < \lfloor a \rfloor + 1$ we get $\lfloor a \rfloor \leq a$ and $a - 1 < \lfloor a \rfloor$; from $\lceil a \rceil - 1 < a \leq \lceil a \rceil$ we get $a \leq \lceil a \rceil$ and $\lceil a \rceil < a + 1$.

Moreover, the following relations between the two functions hold.

Observation 6.6: integer part, upper integer part and change of sign

For every real number $a \in \mathbb{R}$ we have

$$-\lfloor a \rfloor = \lceil -a \rceil \quad \text{and} \quad -\lceil a \rceil = \lfloor -a \rfloor \quad (6.7)$$

and moreover

$$\lfloor a \rfloor = a \iff a \in \mathbb{Z} \iff \lceil a \rceil = a \quad (6.8)$$

Proof. Set $n = \lfloor a \rfloor$, that is, $n \leq a < n + 1$. Multiplying by -1 (and reversing the inequalities) we obtain

$$-n - 1 < -a \leq -n, \quad \text{that is,} \quad (-n) - 1 < -a \leq -n.$$

Since $-n$ is an integer, the definition of upper integer part gives $\lceil -a \rceil = -n = -\lfloor a \rfloor$: this is the first identity in (6.7). Applying the first identity to the number $-a$ we obtain $-\lfloor -a \rfloor = \lceil a \rceil$, that is, the second one.

As for (6.8): if $\lfloor a \rfloor = a$ then a is an integer, because $\lfloor a \rfloor \in \mathbb{Z}$. Conversely, if $a \in \mathbb{Z}$ then $a \leq a < a + 1$ and hence $\lfloor a \rfloor = a$. In the same way, if $\lceil a \rceil = a$ then $a \in \mathbb{Z}$; and if $a \in \mathbb{Z}$ then $a - 1 < a \leq a$, hence $\lceil a \rceil = a$. $\square$

The two functions are completely characterized by the following inequalities.

Observation 6.7: characterizations of the integer part and of the upper integer part

For every real number $a \in \mathbb{R}$ and every integer $n \in \mathbb{Z}$ we have:

$$\lfloor a \rfloor = n \iff n \leq a < n + 1 \quad (6.9a)$$

$$\lfloor a \rfloor = n \iff a - 1 < n \leq a \quad (6.9b)$$

$$\lceil a \rceil = n \iff n - 1 < a \leq n \quad (6.9c)$$

$$\lceil a \rceil = n \iff a \leq n < a + 1 \quad (6.9d)$$

$$a < n \iff \lfloor a \rfloor < n \quad (6.9e)$$

$$n \leq a \iff n \leq \lfloor a \rfloor \quad (6.9f)$$

$$a \leq n \iff \lceil a \rceil \leq n \quad (6.9g)$$

$$n < a \iff n < \lceil a \rceil \quad (6.9h)$$

and moreover

$$\lfloor a + n \rfloor = \lfloor a \rfloor + n \quad \text{and} \quad \lceil a + n \rceil = \lceil a \rceil + n \quad (6.10)$$

Proof. • Equation (6.9a) is the definition of integer part and (6.9c) is the definition of upper integer part.

- Equation (6.9b) is a rewriting of (6.9a): the condition $n \leq a < n + 1$ is equivalent to “ $n \leq a$ and $a < n + 1$ ”, that is, to “ $n \leq a$ and $a - 1 < n$ ”. In the same way (6.9d) is a rewriting of (6.9c): $n - 1 < a \leq n$ is equivalent to “ $a \leq n$ and $n < a + 1$ ”.
- (6.9e): if $a < n$, then $\lfloor a \rfloor \leq a < n$. Conversely, if $\lfloor a \rfloor < n$, then, both being integers, $\lfloor a \rfloor \leq n - 1$ and hence $a < \lfloor a \rfloor + 1 \leq n$.

- (6.9g): if $a \leq n$, then n belongs to the set $\{k \in \mathbb{Z} : k \geq a\}$, whose minimum is $\lceil a \rceil$, hence $\lceil a \rceil \leq n$. Conversely, if $\lceil a \rceil \leq n$, then $a \leq \lceil a \rceil \leq n$.
- Equations (6.9f) and (6.9h) are obtained by negating both sides of (6.9e) and of (6.9g) respectively: the negation of $a < n$ is $n \leq a$ and the negation of $\lfloor a \rfloor < n$ is $n \leq \lfloor a \rfloor$; the negation of $a \leq n$ is $n < a$ and the negation of $\lceil a \rceil \leq n$ is $n < \lceil a \rceil$.
- (6.10): setting $m = \lfloor a \rfloor$, that is, $m \leq a < m + 1$, and adding n we obtain $m + n \leq a + n < (m + n) + 1$ with $m + n \in \mathbb{Z}$, hence $\lfloor a + n \rfloor = m + n = \lfloor a \rfloor + n$. In the same way, setting $m = \lceil a \rceil$, from $m - 1 < a \leq m$ we obtain $(m + n) - 1 < a + n \leq m + n$, hence $\lceil a + n \rceil = m + n = \lceil a \rceil + n$.

□

Given an integer $n \in \mathbb{Z}$, we have

$$\left\lfloor \frac{n}{2} \right\rfloor + \left\lceil \frac{n}{2} \right\rceil = n$$

Proof. We distinguish the two cases.

- If n is **even**, there exists $m \in \mathbb{Z}$ such that $n = 2m$, and since m is an integer (6.8) gives $\lfloor m \rfloor = \lceil m \rceil = m$:

$$\left\lfloor \frac{n}{2} \right\rfloor + \left\lceil \frac{n}{2} \right\rceil = \left\lfloor \frac{2m}{2} \right\rfloor + \left\lceil \frac{2m}{2} \right\rceil = \lfloor m \rfloor + \lceil m \rceil = m + m = 2m = n.$$

- If n is **odd**, there exists $m \in \mathbb{Z}$ such that $n = 2m + 1$. Since $m \leq m + \frac{1}{2} < m + 1$, (6.9a) gives $\lfloor m + \frac{1}{2} \rfloor = m$; since $(m + 1) - 1 < m + \frac{1}{2} \leq m + 1$, (6.9c) gives $\lceil m + \frac{1}{2} \rceil = m + 1$. Hence

$$\left\lfloor \frac{n}{2} \right\rfloor + \left\lceil \frac{n}{2} \right\rceil = \left\lfloor m + \frac{1}{2} \right\rfloor + \left\lceil m + \frac{1}{2} \right\rceil = m + (m + 1) = 2m + 1 = n.$$

□

Given two positive integers $r, s \in \mathbb{Z}$, $r, s > 0$, we have

$$\left\lceil \frac{r}{s} \right\rceil \leq \frac{r + (s - 1)}{s} \quad (6.11a)$$

$$\left\lfloor \frac{r}{s} \right\rfloor \geq \frac{r - (s - 1)}{s} \quad (6.11b)$$

Given, in addition, a non-negative real number $p \in \mathbb{R}$, $p \geq 0$, we have

$$\left\lceil \frac{\left\lfloor \frac{p}{r} \right\rfloor}{s} \right\rceil = \left\lceil \frac{p}{r s} \right\rceil \quad (6.12a)$$

$$\left\lfloor \frac{\left\lceil \frac{p}{r} \right\rceil}{s} \right\rfloor = \left\lfloor \frac{p}{r s} \right\rfloor \quad (6.12b)$$

Definition 6.8: Divisor

A **divisor** of an integer $n \in \mathbb{Z}$, also called a **factor** of n , is an integer $m \in \mathbb{Z}$ that can be multiplied by some integer $q \in \mathbb{Z}$ to obtain n , i.e., if $n = q \cdot m$.

- If m is a divisor of n , n is a **multiple** of m .
- An integer n is **divisible** by another integer m if m is a *divisor* of n .
- For an integer $a \in \mathbb{Z}$ and a positive integer $n \in \mathbb{Z}, n > 0$, the value $a \bmod n$ is the remainder of the division $\frac{a}{n}$.

Definition 6.9: of modulo (remainder)

Given $a \in \mathbb{Z}$ and $n \in \mathbb{Z}, n > 0$,

$$a \bmod n = a - n \left\lfloor \frac{a}{n} \right\rfloor$$

- It follows that $0 \leq a \bmod n < n$

If $(a \bmod n) = (b \bmod n)$, we write

$$a \equiv b \pmod{n}$$

and we say that a is **equivalent** to b , modulo n .

Example 6.9

For example, 23 and 13 are equivalent modulo 5 and we write $23 \equiv 13 \pmod{5}$

- Equivalently, $a \equiv b \pmod{n}$ if a and b have the same remainder when divided by n .
- Equivalently, $a \equiv b \pmod{n}$ if and only if n is a divisor of $|b - a|$.

Summations, geometric and arithmetic progressions

7.1 Summations

Definition 7.1: summation

Let $a_1, a_2, \dots, a_n$ be n real numbers. Their sum

$$a_1 + a_2 + \dots + a_n$$

can be written in compact form with the summation symbol:

$$\sum_{k=1}^n a_k$$

which is read: “sum for k from 1 to n of a_k ”. The symbol k is called the summation index.

- The summation symbol is therefore just a shorthand, which is nevertheless very useful when the terms a_k are defined explicitly as a function of the index k .

Example 7.1: summations

$$\sum_{k=1}^{10} \frac{1}{k} = 1 + \frac{1}{2} + \frac{1}{3} + \frac{1}{4} + \frac{1}{5} + \frac{1}{6} + \frac{1}{7} + \frac{1}{8} + \frac{1}{9} + \frac{1}{10}$$

$$\sum_{k=3}^n k^2 = 3^2 + 4^2 + 5^2 + \dots + n^2$$

- The summation index is a **dummy index**. This means that if we replace k with i , j or any other index (in all its occurrences), the value of the summation does not change.

Example 7.2: dummy index

We have:

$$\sum_{k=1}^n k^2 = \sum_{i=1}^n i^2$$

since both symbols denote the sum of the squares of the first n natural numbers (without zero).

On the other hand, we have:

$$\sum_{k=1}^n k^2 \neq \sum_{k=1}^m k^2$$

since the two symbols denote the sum of, respectively, the first n or the first m squares (if $n \neq m$ the result will be different).

7.1.1 Main properties of summations

Observation 7.2: product by a constant

Given a summation $\sum_{k=1}^n a_k$ and a real number $c \in \mathbb{R}$, we have:

$$\sum_{k=1}^n (c \cdot a_k) = c \sum_{k=1}^n a_k \quad (7.1)$$

Proof. By the distributive property we have:

$$\underbrace{c a_1 + c a_2 + \cdots + c a_n}_{=\sum_{k=1}^n (c \cdot a_k)} = \underbrace{c (a_1 + a_2 + \cdots + a_n)}_{=c \sum_{k=1}^n a_k}$$

□

Observation 7.3: summation with a constant term

For every natural number $n \geq 1$ and every real number $c \in \mathbb{R}$, we have:

$$\sum_{k=1}^n c = c \cdot n \quad (7.2)$$

Proof. We have:

$$\underbrace{c + c + \cdots + c}_{\text{that is, } c \text{ added } n \text{ times}} = c n$$

$$= \sum_{k=1}^n c$$

□

Observation 7.4: union of summations

Given two summations $\sum_{k=1}^n a_k$ and $\sum_{k=1}^n b_k$, we have:

$$\sum_{k=1}^n a_k + \sum_{k=1}^n b_k = \sum_{k=1}^n (a_k + b_k) \quad (7.3)$$

Proof. We have:

$$\underbrace{a_1 + a_2 + \cdots + a_n + b_1 + b_2 + \cdots + b_n}_{=\sum_{k=1}^n a_k + \sum_{k=1}^n b_k} = \underbrace{a_1 + b_1 + a_2 + b_2 + \cdots + a_n + b_n}_{=\sum_{k=1}^n (a_k + b_k)}$$

□

- The three properties that follow (decomposition, index translation and index reflection) are simply different ways of writing and/or ordering the terms of the summations.

Observation 7.5: decomposition

Given two natural numbers $n \geq 1$ and $m \geq 1$, we have:

$$\sum_{k=1}^{n+m} a_k = \sum_{k=1}^n a_k + \sum_{k=n+1}^{n+m} a_k \quad (7.4)$$

Proof. The terms of the summation $\sum_{k=1}^{n+m} a_k$ can be grouped into the first n and the last m ones:

$$\begin{aligned} & \underbrace{a_1 + a_2 + \cdots + a_n}_{=\sum_{k=1}^n a_k} + \underbrace{a_{n+1} + a_{n+2} + \cdots + a_{n+m}}_{=\sum_{k=n+1}^{n+m} a_k} \\ &= \underbrace{a_1 + a_2 + \cdots + a_n + a_{n+1} + \cdots + a_{n+m}}_{=\sum_{k=1}^{n+m} a_k} \end{aligned}$$

□

Observation 7.6: index translation

Given a summation $\sum_{k=1}^n a_k$ and a natural number $m \geq 1$, we have:

$$\sum_{k=1}^n a_k = \sum_{k=1+m}^{n+m} a_{k-m} = \sum_{k=1-m}^{n-m} a_{k+m} \quad (7.5)$$

Proof. In the summation $\sum_{k=1+m}^{n+m} a_{k-m}$ the index k ranges from $1+m$ to $n+m$, and therefore the index $k-m$ ranges from 1 to n :

$$\underbrace{a_{(1+m)-m} + a_{(2+m)-m} + \cdots + a_{(n+m)-m}}_{=\sum_{k=1+m}^{n+m} a_{k-m}} = \underbrace{a_1 + a_2 + \cdots + a_n}_{=\sum_{k=1}^n a_k}$$

In the same way, in the summation $\sum_{k=1-m}^{n-m} a_{k+m}$ the index k ranges from $1-m$ to $n-m$, and therefore the index $k+m$ ranges from 1 to n :

$$\underbrace{a_{(1-m)+m} + a_{(2-m)+m} + \cdots + a_{(n-m)+m}}_{=\sum_{k=1-m}^{n-m} a_{k+m}} = \underbrace{a_1 + a_2 + \cdots + a_n}_{=\sum_{k=1}^n a_k}$$

□

Observation 7.7: index reflection

Given a summation $\sum_{k=1}^n a_k$, we have:

$$\sum_{k=1}^n a_k = \sum_{k=1}^n a_{n-k+1} = \sum_{k=0}^{n-1} a_{n-k} \quad (7.6)$$

Proof. In the summation $\sum_{k=1}^n a_{n-k+1}$ the index k ranges from 1 to n , and therefore the index $n-k+1$ ranges from n to 1, that is, the terms are the same ones listed in reverse order:

$$\underbrace{a_n + a_{n-1} + \cdots + a_1}_{=\sum_{k=1}^n a_{n-k+1}} = \underbrace{a_1 + a_2 + \cdots + a_n}_{=\sum_{k=1}^n a_k}$$

In the same way, in the summation $\sum_{k=0}^{n-1} a_{n-k}$ the index k ranges from 0 to $n-1$, and therefore the index $n-k$ ranges from n to 1:

$$\underbrace{a_n + a_{n-1} + \cdots + a_1}_{=\sum_{k=0}^{n-1} a_{n-k}} = \underbrace{a_1 + a_2 + \cdots + a_n}_{=\sum_{k=1}^n a_k}$$

□

7.1.2 Some important summations

Observation 7.8: sum of the first n natural numbers (without zero)

For every natural number $n \geq 1$, we have:

$$\sum_{k=1}^n k = \frac{n(n+1)}{2}$$

Proof. Using the index reflection (7.6) we have:

$$\begin{aligned} \sum_{k=1}^n k &= \frac{1}{2} \left(\sum_{k=1}^n k + \sum_{k=1}^n k \right) = \frac{1}{2} \left(\sum_{k=1}^n k + \sum_{k=1}^n (n - k + 1) \right) \\ &= \frac{1}{2} \sum_{k=1}^n (k + n - k + 1) = \frac{1}{2} \sum_{k=1}^n (n + 1) = \frac{n(n+1)}{2} \end{aligned}$$

□

Observation 7.9: sum of the first n odd numbers

For every natural number $n \geq 1$, we have:

$$\sum_{k=1}^n (2k-1) = n^2 \quad \text{or equivalently} \quad \sum_{k=0}^{n-1} (2k+1) = n^2$$

Proof. Using the properties of summations we have:

$$\begin{aligned} \sum_{k=1}^n (2k-1) &= 2 \sum_{k=1}^n k - \sum_{k=1}^n 1 \\ &= 2 \left(\frac{n(n+1)}{2} \right) - n = n^2 + n - n = n^2 \\ \\ \sum_{k=0}^{n-1} (2k+1) &= 2 \sum_{k=0}^{n-1} k + \sum_{k=0}^{n-1} 1 \\ &= 2 \sum_{k=1}^n (k-1) + n = 2 \left(\sum_{k=1}^n k - \sum_{k=1}^n 1 \right) + n \\ &= 2 \left(\frac{n(n+1)}{2} - n \right) + n = n^2 + n - 2n + n = n^2 \end{aligned}$$

□

Observation 7.10: sum of the first n even numbers (without zero)

For every natural number $n \geq 1$, we have:

$$\sum_{k=1}^n 2k = n(n+1)$$

Proof.

$$\sum_{k=1}^n 2k = 2 \sum_{k=1}^n k = 2 \left(\frac{n(n+1)}{2} \right) = n(n+1)$$

□

- The three important summations we have just seen can also be proved by induction, in the chapter “Principle of induction”.

7.2 Geometric progressions

Definition 7.11: geometric progression

A sequence of real numbers is a **geometric progression** if the ratio between each term (starting from the second one) and the previous one is constant. This constant is called the common ratio of the progression.

Given the first term $a \in \mathbb{R}$ and the common ratio $q \in \mathbb{R}$, the associated geometric progression is:

$$a, \quad aq, \quad aq^2, \quad aq^3, \quad aq^4, \quad \dots$$

Each term (starting from the second one) is obtained from the previous one by multiplying it by q . The k -th term ($k \in \mathbb{N}$, $k \geq 1$) can be written as aq^{k-1} and we have:

$aq^{1-1} = aq^0 = a$	first term, in position $k = 1$
$aq^{2-1} = aq^1 = aq$	second term, in position $k = 2$
$aq^{3-1} = aq^2$	third term, in position $k = 3$
$aq^{4-1} = aq^3$	fourth term, in position $k = 4$
$\dots$	$\dots$

Observation 7.12: k -th term of a geometric progression

Given the first term $a \in \mathbb{R}$ and the common ratio $q \in \mathbb{R}$, let t_k denote the term in position k of the geometric progression. The recursive relation that defines the progression is:

$$t_1 = a \quad \text{and} \quad t_k = t_{k-1} q \quad \text{for every natural number } k \geq 2 \quad (7.7)$$

and the closed formula of the k -th term is:

$$t_k = a q^{k-1} \quad \text{for every natural number } k \geq 1 \quad (7.8)$$

Proof. The closed formula (7.8) is obtained by applying repeatedly the recursive relation (7.7):


$$\begin{aligned} t_1 &= a = a q^0 \\ t_2 &= t_1 q = a q = a q^1 \\ t_3 &= t_2 q = a q q = a q^2 \\ t_4 &= t_3 q = a q^2 q = a q^3 \\ &\vdots \\ t_k &= t_{k-1} q = a q^{k-2} q = a q^{k-1} \end{aligned}$$

□

Example 7.3: geometric progressions


- With $a = 1$ and $q = \frac{1}{2}$, the first 4 terms are:

$$1, \quad \frac{1}{2}, \quad \frac{1}{4}, \quad \frac{1}{8}$$



- With $a = 1$ and $q = 2$, the first 4 terms are:

$$1, \quad 2, \quad 4, \quad 8$$




Example 7.4: geometric progression with a negative common ratio

- With $a = 1$ and $q = -2$, the first 4 terms are:

$$1, -2, 4, -8$$

With a negative common ratio the terms change sign at every step: those in odd positions are positive, those in even positions are negative.



7.2.1 Sums of the terms of geometric progressions

Observation 7.13: sum of the first n terms of the geometric progression ($a = 1$)

Given $q \in \mathbb{R}$, for every natural number $n \geq 1$ we have:

$$\sum_{k=1}^n q^{k-1} = \begin{cases} \frac{q^n - 1}{q - 1} & \text{if } q \neq 1 \\ n & \text{otherwise} \end{cases} \quad (7.9)$$

Proof. If $q \neq 1$, we prove the equation in the following equivalent form:

$$(q - 1) \sum_{k=1}^n q^{k-1} = q^n - 1$$

Applying the properties of summations, in particular the product by a constant (7.1) and the index translation (7.5), we get:

$$\begin{aligned} (q - 1) \sum_{k=1}^n q^{k-1} &= q \sum_{k=1}^n q^{k-1} - \sum_{k=1}^n q^{k-1} = \\ &= \sum_{k=1}^n q^k - \sum_{k=1}^n q^{k-1} = \sum_{k=1}^n q^k - \sum_{k=0}^{n-1} q^k = \\ &= \sum_{k=1}^{n-1} q^k + q^n - \left(1 + \sum_{k=1}^{n-1} q^k \right) = q^n - 1 \end{aligned}$$

Dividing by $q - 1 \neq 0$ we obtain:

$$\sum_{k=1}^n q^{k-1} = \frac{q^n - 1}{q - 1}$$

If $q = 1$, we have instead:

$$\sum_{k=1}^n q^{k-1} = \sum_{k=1}^n 1 = n$$

□

- The proof never uses the sign of the common ratio: formula (7.9) holds for every $q \in \mathbb{R}$, positive, zero or negative. The only case to be made precise is $q = 0$: the first term of the summation is $q^{1-1} = 0^0$, which here is understood to be equal to 1 (convention $0^0 = 1$). With this convention, for every $n \geq 1$ we have:

$$\sum_{k=1}^n 0^{k-1} = 1 + 0 + \cdots + 0 = 1 \quad \text{and} \quad \frac{0^n - 1}{0 - 1} = \frac{-1}{-1} = 1$$

and therefore formula (7.9) remains true.

- Given $q \in \mathbb{R}, n \in \mathbb{N}, n \geq 1$ and $a \in \mathbb{R}$, formula (7.9) extends as follows:

$$\sum_{k=1}^n a q^{k-1} = \begin{cases} a \left(\frac{q^n - 1}{q - 1} \right) & \text{if } q \neq 1 \\ a n & \text{otherwise} \end{cases} \quad \text{since} \quad \sum_{k=1}^n a q^{k-1} = a \sum_{k=1}^n q^{k-1} \quad (7.10)$$

clearly:

$$\frac{q^n - 1}{q - 1} = \frac{1 - q^n}{1 - q} \quad \text{and hence we also have} \quad \sum_{k=1}^n a q^{k-1} = \begin{cases} a \left(\frac{1 - q^n}{1 - q} \right) & \text{if } q \neq 1 \\ a n & \text{otherwise} \end{cases}$$

- Formula (7.9) can also be proved by induction, in the chapter “Principle of induction”.

Example 7.5: sum of the first n terms of geometric progressions

- With $a = 2$ and $q = \frac{1}{2}$, the first 4 terms are:

$$2, \quad 1, \quad \frac{1}{2}, \quad \frac{1}{4} \quad \text{and} \quad 2 + 1 + \frac{1}{2} + \frac{1}{4} = \frac{8 + 4 + 2 + 1}{4} = \frac{15}{4}$$

the sum of the first $n = 4$ terms is given by the formula:

$$\sum_{k=1}^4 2 \left(\frac{1}{2} \right)^{k-1} = \sum_{k=1}^4 \frac{2}{2^{k-1}} = 2 \frac{1 - \frac{1}{2^4}}{1 - \frac{1}{2}} = 2 \frac{1 - \frac{1}{16}}{\frac{1}{2}} = \frac{15}{16} \cdot 4 = \frac{15}{4}$$

- With $a = 1$ and $q = \frac{1}{2}$, the first 4 terms are:

$$1, \quad \frac{1}{2}, \quad \frac{1}{4}, \quad \frac{1}{8} \quad \text{and} \quad 1 + \frac{1}{2} + \frac{1}{4} + \frac{1}{8} = \frac{8 + 4 + 2 + 1}{8} = \frac{15}{8}$$

the sum of the first $n = 4$ terms is given by the formula:

$$\sum_{k=1}^4 \left(\frac{1}{2} \right)^{k-1} = \sum_{k=1}^4 \frac{1}{2^{k-1}} = \frac{1 - \frac{1}{2^4}}{1 - \frac{1}{2}} = \frac{1 - \frac{1}{16}}{\frac{1}{2}} = \frac{15}{16} \cdot 2 = \frac{15}{8}$$

- With $a = 1$ and $q = 2$, the first 4 terms are:

$$1, \quad 2, \quad 4, \quad 8 \quad \text{and} \quad 1 + 2 + 4 + 8 = 15$$

the sum of the first $n = 4$ terms is given by the formula:

$$\sum_{k=1}^4 2^{k-1} = \frac{2^4 - 1}{2 - 1} = 16 - 1 = 15$$

- With $a = 1$ and $q = -2$, the first 4 terms are:

$$1, \quad -2, \quad 4, \quad -8 \quad \text{and} \quad 1 - 2 + 4 - 8 = -5$$

the sum of the first $n = 4$ terms is given by the formula:

$$\sum_{k=1}^4 (-2)^{k-1} = \frac{(-2)^4 - 1}{-2 - 1} = \frac{16 - 1}{-3} = \frac{15}{-3} = -5$$

7.3 Arithmetic progressions

Definition 7.14: arithmetic progression

A sequence of real numbers is an **arithmetic progression** if the difference between each term (starting from the second one) and the previous one is constant. This constant is called the common difference of the progression.

Given the first term $a \in \mathbb{R}$ and the common difference $d \in \mathbb{R}$, the associated arithmetic progression is:

$$a, \quad a + d, \quad a + 2d, \quad a + 3d, \quad a + 4d, \quad \dots$$

Each term (starting from the second one) is obtained from the previous one by adding d to it. The k -th term ($k \in \mathbb{N}$, $k \geq 1$) can be written as $a + (k - 1)d$ and we have:

$a + (1 - 1)d = a + 0d = a$	first term, in position $k = 1$
$a + (2 - 1)d = a + 1d = a + d$	second term, in position $k = 2$
$a + (3 - 1)d = a + 2d$	third term, in position $k = 3$
$a + (4 - 1)d = a + 3d$	fourth term, in position $k = 4$
$\dots$	$\dots$

- The arithmetic progression is the additive analogue of the geometric progression: in the geometric progression each term is *multiplied* by the common ratio q , in the arithmetic progression the common difference d is *added* to each term.

Observation 7.15: k -th term of an arithmetic progression

Given the first term $a \in \mathbb{R}$ and the common difference $d \in \mathbb{R}$, let t_k denote the term in position k of the arithmetic progression. The recursive relation that defines the progression is:

$$t_1 = a \quad \text{and} \quad t_k = t_{k-1} + d \quad \text{for every natural number } k \geq 2 \quad (7.11)$$

and the closed formula of the k -th term is:

$$t_k = a + (k - 1)d \quad \text{for every natural number } k \geq 1 \quad (7.12)$$

Proof. The closed formula (7.12) is obtained by repeatedly applying the recursive relation (7.11):

$$\begin{aligned}
 t_1 &= a = a + 0 \, d \\
 t_2 &= t_1 + d = a + d = a + 1 \, d \\
 t_3 &= t_2 + d = a + d + d = a + 2 \, d \\
 t_4 &= t_3 + d = a + 2 \, d + d = a + 3 \, d \\
 &\vdots \\
 t_k &= t_{k-1} + d = a + (k-2) \, d + d = a + (k-1) \, d
 \end{aligned}$$

□

- The closed formula (7.12) can also be obtained by the following argument, which does not need the dots. We write the recursive relation (7.11) in the form $t_j - t_{j-1} = d$ for every position j from 2 to k :

$$\begin{aligned}
 t_2 - t_1 &= d \\
 t_3 - t_2 &= d \\
 t_4 - t_3 &= d \\
 &\vdots \\
 t_k - t_{k-1} &= d
 \end{aligned}$$

We now add all these $k-1$ equalities. On the right-hand side we add d to itself $k-1$ times and we obtain $d(k-1)$. On the left-hand side the sum is **telescopic**: every term appears twice, once with the sign $+$ and once with the sign $-$, except t_k and t_1 :

$$\underbrace{(t_2 - t_1) + (t_3 - t_2) + (t_4 - t_3) + \cdots + (t_k - t_{k-1})}_{=\sum_{j=2}^k (t_j - t_{j-1})} = t_k - t_1 = t_k - a$$


Equating the two sides we get $t_k - a = d(k-1)$, that is, the closed formula:

$$t_k = a + (k-1) \, d \quad \text{for every natural number } k \geq 1$$

Example 7.6: arithmetic progressions

- With $a = 1$ and $d = \frac{1}{2}$, the first 4 terms are:


$$1, \quad \frac{3}{2}, \quad 2, \quad \frac{5}{2}$$



- With $a = 2$ and $d = 2$, the first 4 terms are:

$$2, 4, 6, 8$$

that is, the first 4 even numbers (without zero).



7.3.1 Sums of the terms of arithmetic progressions

Observation 7.16: sum of the first n terms of the arithmetic progression ($a = 0$)

Given $d \in \mathbb{R}$, for every natural number $n \geq 1$ we have:

$$\sum_{k=1}^n d(k-1) = d \left(\frac{n(n-1)}{2} \right) \quad (7.13)$$

Proof. Using the product by a constant (7.1) and the index reflection (7.6), with the same argument used for the sum of the first n natural numbers, we have:

$$\begin{aligned}\sum_{k=1}^n d(k-1) &= \frac{d}{2} \left(\sum_{k=1}^n (k-1) + \sum_{k=1}^n (k-1) \right) = \frac{d}{2} \left(\sum_{k=1}^n (k-1) + \sum_{k=1}^n (n-k) \right) \\ &= \frac{d}{2} \sum_{k=1}^n (k-1+n-k) = \frac{d}{2} \sum_{k=1}^n (n-1) = d \left(\frac{n(n-1)}{2} \right)\end{aligned}$$

where in the second step the index of the second summation has been reflected: the term in position k is $a_k = k-1$ and therefore $a_{n-k+1} = (n-k+1)-1 = n-k$. $\square$

- Given $d \in \mathbb{R}, n \in \mathbb{N}, n \geq 1$ and $a \in \mathbb{R}$, formula (7.13) extends as follows:

$$\begin{aligned}\sum_{k=1}^n (a + d(k-1)) &= a n + d \left(\frac{n(n-1)}{2} \right) \quad (7.14) \\ \text{since } \sum_{k=1}^n (a + d(k-1)) &= \underbrace{\sum_{k=1}^n a}_{=a n} + \sum_{k=1}^n d(k-1)\end{aligned}$$

where the union of summations (7.3) and the summation with a constant term (7.2) have been used. This formula computes the sum of the first n terms of the arithmetic progression with a generic first term a .

- The same proof by index reflection, applied directly to the summation of the first n terms, gives the equivalent form:

$$\sum_{k=1}^n (a + d(k-1)) = \frac{n}{2} \left(\underbrace{a}_{=t_1} + \underbrace{a + (n-1)d}_{=t_n} \right) = n \left(\frac{t_1 + t_n}{2} \right)$$

since, reflecting the index, the term in position k and the term in position $n-k+1$ have constant sum $t_k + t_{n-k+1} = 2a + (n-1)d = t_1 + t_n$. The sum of the first n terms of an arithmetic progression is therefore n times the average of the first and the last term.

- With $a = 1$ and $d = 1$ the arithmetic progression is $1, 2, 3, \dots$ and formula (7.14) gives $n + \frac{n(n-1)}{2} = \frac{n(n+1)}{2}$, that is, the sum of the first n natural numbers (without zero) already seen in the section “Some important summations”. With $a = 1$ and $d = 2$ the progression is $1, 3, 5, \dots$ and the formula gives $n + 2 \frac{n(n-1)}{2} = n^2$, that is, the sum of the first n odd numbers.

Example 7.7: sum of the first n terms of arithmetic progressions

- With $a = 1$ and $d = \frac{1}{2}$, the first 4 terms are:

$$1, \quad \frac{3}{2}, \quad 2, \quad \frac{5}{2} \quad \text{and} \quad 1 + \frac{3}{2} + 2 + \frac{5}{2} = \frac{2+3+4+5}{2} = \frac{14}{2} = 7$$

the sum of the first $n = 4$ terms is given by the formula:

$$\sum_{k=1}^4 \left(1 + \frac{1}{2} (k-1) \right) = 1 \cdot 4 + \frac{1}{2} \left(\frac{4 \cdot 3}{2} \right) = 4 + 3 = 7$$

- With $a = 2$ and $d = 2$, the first 4 terms are:

$$2, \quad 4, \quad 6, \quad 8 \quad \text{and} \quad 2 + 4 + 6 + 8 = 20$$

the sum of the first $n = 4$ terms is given by the formula:

$$\sum_{k=1}^4 (2 + 2 (k-1)) = 2 \cdot 4 + 2 \left(\frac{4 \cdot 3}{2} \right) = 8 + 12 = 20$$

and with the equivalent form: $4 \left(\frac{2+8}{2} \right) = 4 \cdot 5 = 20$.

- With $a = 1$ and $d = -\frac{1}{2}$, the first 4 terms are:

$$1, \quad \frac{1}{2}, \quad 0, \quad -\frac{1}{2} \quad \text{and} \quad 1 + \frac{1}{2} + 0 - \frac{1}{2} = 1$$

the sum of the first $n = 4$ terms is given by the formula:

$$\sum_{k=1}^4 \left(1 - \frac{1}{2} (k-1) \right) = 1 \cdot 4 - \frac{1}{2} \left(\frac{4 \cdot 3}{2} \right) = 4 - 3 = 1$$

Principle of mathematical induction

8.1 The principle of mathematical induction

- We now present a proof technique called proof by induction. This procedure can be applied to theorems with the following structure:

“ for every $n \in \mathbb{N}$, $n \geq n_0$, property $p(n)$ holds ”

- The number n_0 is the smallest integer for which we want the property to be true; if $n_0 = 0$ the theorem simply states that the property is true for every $n \in \mathbb{N}$.

A proof by induction consists of the following two steps:

1. We prove that $p(n)$ is true for $n = n_0$ (**base case of the induction**).
2. We prove that, if n is an arbitrary natural number $\geq n_0$, the fact that $p(n)$ is true implies that $p(n + 1)$ is true (**inductive step**).

We can then conclude that for every $n \geq n_0$, $p(n)$ is true.

- The validity of this proof technique, *intuitively*, is based on the following fact:
 1. By point 1, we know that $p(n_0)$ is true. Suppose for example $n_0 = 1$: we therefore know that $p(1)$ is true (this must be proved explicitly).
 2. By point 2, since $p(1)$ is true, $p(2)$ will be true: indeed, we have proved that for any n , if $p(n)$ is true then $p(n + 1)$ is also true. But then, since $p(2)$ is true, $p(3)$ will be true; but then $p(4)$ is true, $\dots$ and so on, hence $p(n)$ is true for every $n \geq 1$.
- In practice, the proof consists of two phases.
 1. Prove $p(n_0)$ directly;

2. Assume $p(n)$ as a hypothesis (**inductive hypothesis**) and prove $p(n+1)$.

- This is the delicate point, often subject to misunderstandings. “Assuming as a hypothesis” $p(n)$ does not mean assuming the thesis as a hypothesis. What must be proved is that:

“ for every $n \geq n_0$, if $p(n)$ is true then $p(n+1)$ is also true ”

and not

“ if $p(n)$ is true for every n , then $p(n+1)$ is also true”

8.2 Proofs based on the principle of mathematical induction

8.2.1 Bernoulli's inequality

Observation 8.1: Bernoulli's inequality

For every integer $n \geq 0$, $x \in \mathbb{R}$, $x \geq -1$, we have:

$$(1+x)^n \geq 1+n x \quad (8.1)$$

Proof. By induction on n .

- **Base case of the induction**

Let $n = 0$. Then the statement becomes:

$$(1+x)^0 \geq 1+0 x \quad \text{i.e.} \quad 1 \geq 1$$

which is clearly true.

- **Inductive step**

Suppose it is true for n , and let us prove it for $(n+1)$. By the inductive hypothesis, we have:

$$(1+x)^n \geq 1+n x$$

and moreover we have

$$(1+x) \geq 0 \quad \text{since} \quad x \geq -1.$$

Then we can write:

$$\begin{aligned} (1+x)^{n+1} &= (1+x) \cdot (1+x)^n \\ &\geq (1+x) \cdot (1+n x) \\ &= 1 + (n+1) x + n x^2 \\ &\geq 1 + (n+1) x \end{aligned}$$

where in the last inequality we used the fact that $n x^2 \geq 0$.


The chain of inequalities shows that, for $n + 1$, we have


$$(1 + x)^{n+1} \geq 1 + (n + 1) x$$

which is exactly the desired statement, for $n + 1$.

□

Example 8.1: Bernoulli's inequality





8.2.2 Some important summations

Observation 8.2: sum of the first n natural numbers

For every integer $n \geq 1$, we have:

$$\sum_{k=1}^n k = \frac{n(n+1)}{2}$$

Proof. By induction on n .

- **Base case of the induction**

Let $n = 1$. Then the statement becomes:

$$\sum_{k=1}^1 k = \frac{1(1+1)}{2} \quad \text{i.e.} \quad 1 = 1$$

which is clearly true.

- **Inductive step**

Suppose it is true for n , and let us prove it for $(n+1)$. By the inductive hypothesis, we have:

$$\sum_{k=1}^n k = \frac{n(n+1)}{2}.$$

Hence we can write:

$$\begin{aligned}
 \sum_{k=1}^{n+1} k &= \sum_{k=1}^n k + (n+1) \\
 &= \frac{n(n+1)}{2} + (n+1) \\
 &= \frac{n(n+1) + 2(n+1)}{2} \\
 &= \frac{(n+1)(n+2)}{2} = \frac{(n+1)((n+1)+1)}{2}
 \end{aligned}$$

which is exactly the desired statement, for $n+1$.

□

Observation 8.3: sum of the first n odd numbers

For every integer $n \geq 1$, we have:

$$\sum_{k=0}^{n-1} (2k+1) = n^2 \quad \text{or equivalently} \quad \sum_{k=1}^n (2k-1) = n^2$$

Proof. By induction on n .

- **Base case of the induction**

Let $n = 1$. Then the statement becomes:

$$\sum_{k=0}^0 (2k+1) = 1^2 \quad \text{i.e.} \quad 1 = 1$$

which is clearly true.

- **Passo induttivo**

Suppose it is true for n , and let us prove it for $(n+1)$. By the inductive hypothesis, we have:

$$\sum_{k=0}^{n-1} (2k+1) = n^2$$

Hence we can write:

$$\sum_{k=0}^{(n+1)-1} (2k+1) = \sum_{k=0}^{n-1} (2k+1) + 2n+1 = n^2 + 2n+1 = (n+1)^2$$

which is exactly the desired statement, for $n+1$.

The second sum coincides with the first one, by an index shift ($k \rightarrow k + 1$):

$$\sum_{k=1}^n (2k - 1) = \sum_{k=0}^{n-1} (2(k + 1) - 1) = \sum_{k=0}^{n-1} (2k + 1)$$

□

Observation 8.4: sum of the first n even numbers

For every integer $n \geq 1$, we have:

$$\sum_{k=1}^n 2k = n(n + 1)$$

Proof. By induction on n .

- **Base case of the induction**

Let $n = 1$. Then the statement becomes:

$$\sum_{k=1}^1 2k = 1 \cdot 2 \quad \text{i.e.} \quad 2 = 2$$

which is clearly true.

- **Passo induttivo**

Suppose it is true for n , and let us prove it for $(n + 1)$. By the inductive hypothesis, we have:

$$\sum_{k=1}^n 2k = n(n + 1)$$

Hence we can write:

$$\begin{aligned} \sum_{k=1}^{n+1} 2k &= \sum_{k=1}^n 2k + 2(n + 1) = n^2 + n + 2(n + 1) \\ &= n^2 + 2n + 1 + n + 1 = (n + 1)^2 + (n + 1) = (n + 1)((n + 1) + 1) \end{aligned}$$

which is exactly the desired statement, for $n + 1$.

□

8.2.3 Sum of the terms of the geometric progression

Observation 8.5: sum of the first n terms of the geometric progression ($a = 1$)

Given $q \in \mathbb{R}$, for every integer $n \geq 1$ we have (with the convention $0^0 = 1$ for the case $q = 0$):

$$\sum_{k=1}^n q^{k-1} = \begin{cases} \frac{q^n - 1}{q - 1} & \text{if } q \neq 1 \\ n & \text{otherwise} \end{cases} \quad (8.2)$$

Proof. By induction on n .

- **Base case of the induction**

Let $n = 1$. Then the statement becomes:

$$\sum_{k=1}^1 q^{k-1} = \frac{q^1 - 1}{q - 1} \quad \text{i.e.} \quad 1 = 1$$

which is clearly true.

- **Inductive step**

Suppose it is true for n , and let us prove it for $(n+1)$. By the inductive hypothesis, we have:

$$\sum_{k=1}^n q^{k-1} = \frac{q^n - 1}{q - 1}$$

Hence we can write

$$\begin{aligned} \sum_{k=1}^{n+1} q^{k-1} &= \sum_{k=1}^n q^{k-1} + q^n = \frac{q^n - 1}{q - 1} + q^n \\ &= \frac{q^n - 1 + q^{n+1} - q^n}{q - 1} = \frac{q^{n+1} - 1}{q - 1} \end{aligned}$$

which is exactly the desired statement, for $n + 1$.

□

8.2.4 Inequality of arithmetic and geometric means

Observation 8.6: inequality of arithmetic and geometric means

For every integer $n \geq 2$ and every vector $\mathbf{a} = (a_1, a_2, \dots, a_n) \in \mathbb{R}_{\geq 0}^n$, we have:

$$\frac{\sum_{i=1}^n a_i}{n} \geq \sqrt[n]{\prod_{i=1}^n a_i} \quad (8.3)$$

- The inequality (8.3) states that the *arithmetic mean* is greater than or equal to the *geometric mean*.
- If $a_i = a$ for every $i \in \{1, 2, \dots, n\}$, we have:

$$\frac{\sum_{i=1}^n a_i}{n} = \frac{n a}{n} = a \quad \text{and} \quad \sqrt[n]{\prod_{i=1}^n a_i} = \sqrt[n]{a^n} = a$$

and the arithmetic mean and the geometric mean are equal (this is the only case in which equality holds).

Proof. By induction on n .

- **Base case of the induction**

Let $n = 2$. Since both sides are non-negative, we can square them:

$$\begin{aligned} \frac{a_1 + a_2}{2} \geq \sqrt{a_1 a_2} &\iff \left(\frac{a_1 + a_2}{2} \right)^2 \geq a_1 a_2 \\ &\iff \frac{a_1^2 + 2 a_1 a_2 + a_2^2}{4} \geq a_1 a_2 \\ &\iff a_1^2 - 2 a_1 a_2 + a_2^2 \geq 0 \\ &\iff (a_1 - a_2)^2 \geq 0 \end{aligned}$$

which is clearly true.

- **Inductive step**

Suppose it is true for n , and let us prove it for $(n+1)$. By the inductive hypothesis, the arithmetic mean of n non-negative real numbers is greater than or equal to their geometric mean. We need to prove that:

$$\frac{\sum_{i=1}^{n+1} a_i}{n+1} \geq \sqrt[n+1]{\prod_{i=1}^{n+1} a_i} \quad (\text{A})$$

Let α be the arithmetic mean of the $n + 1$ non-negative real numbers:

$$\alpha = \frac{\sum_{i=1}^{n+1} a_i}{n+1} \quad (\text{B})$$

If $a_i = \alpha$ for every $i \in \{1, 2, \dots, n+1\}$, then (A) holds with equality. Otherwise, there is at least one value greater than α and at least one value smaller than α . Without loss of generality, we reorder the values so that:

$$a_n > \alpha \quad \text{and} \quad a_{n+1} < \alpha$$

Then we have:

$$a_n - \alpha > 0 \quad \text{and} \quad \alpha - a_{n+1} > 0 \implies (a_n - \alpha)(\alpha - a_{n+1}) > 0 \quad (\text{C})$$

From (B) we have:

$$(n+1) \alpha = \sum_{i=1}^{n+1} a_i \iff n \alpha = \sum_{i=1}^{n-1} a_i + \underbrace{a_n + a_{n+1} - \alpha}_{=y} \iff \alpha = \frac{\sum_{i=1}^{n-1} a_i + y}{n}$$

where $y = a_n + a_{n+1} - \alpha \geq a_n - \alpha > 0$, since $a_{n+1} \geq 0$. Hence α is also the arithmetic mean of the n non-negative numbers $a_1, a_2, \dots, a_{n-1}$ and y . By the inductive hypothesis, $\alpha^n \geq \left(\prod_{i=1}^{n-1} a_i\right) y$, and therefore:

$$\alpha^{n+1} = \alpha^n \alpha \geq \left(\prod_{i=1}^{n-1} a_i\right) y \alpha \quad (\text{D})$$

From (C) it follows that:

$$(a_n - \alpha)(\alpha - a_{n+1}) = \underbrace{(a_n + a_{n+1} - \alpha)}_{=y} \alpha - a_n a_{n+1} > 0$$

and therefore:

$$y \alpha > a_n a_{n+1} \quad (\text{E})$$

Substituting (E) into (D) (the product $\prod_{i=1}^{n-1} a_i$ is non-negative), we obtain:

$$\alpha^{n+1} \geq \prod_{i=1}^{n+1} a_i \iff \frac{\sum_{i=1}^{n+1} a_i}{n+1} \geq \sqrt[n+1]{\prod_{i=1}^{n+1} a_i}$$

that is, (A), which is exactly the desired statement, for $n + 1$.

□

8.2.5 Fibonacci sequence

- The **Fibonacci sequence** is defined by the following two initial values and recurrence relation:

$$F_0 = 0, \quad F_1 = 1, \quad F_n = F_{n-1} + F_{n-2} \quad \text{for every integer } n \geq 2. \quad (8.4)$$

- The first terms are 0, 1, 1, 2, 3, 5, 8, 13, 21, 34, ...: each term is the sum of the two preceding ones.
- The two solutions of the equation $x^2 - x - 1 = 0$ are called the **golden ratio** ϕ and the **conjugate of the golden ratio** $\hat{\phi}$:

$$\phi = \frac{1 + \sqrt{5}}{2} \quad \text{and} \quad \hat{\phi} = \frac{1 - \sqrt{5}}{2}$$

Since they are solutions of $x^2 = x + 1$, the following two equalities hold:

$$\phi^2 = \phi + 1 \quad \text{and} \quad \hat{\phi}^2 = \hat{\phi} + 1 \quad (8.5)$$

- The following proposition gives a closed formula for F_n : the n -th term can be computed directly, without going through all the preceding terms.

Observation 8.7: closed formula of the Fibonacci sequence

For every integer $n \geq 0$, we have:

$$F_n = \frac{\phi^n - \hat{\phi}^n}{\sqrt{5}} \quad (8.6)$$

Proof. By induction on n . Since the recurrence relation links F_{n+1} to the **two** preceding terms, the base case of the induction checks the statement for $n = 0$ and for $n = 1$, and the inductive step assumes it true for n and for $n - 1$.

- Base case of the induction**

Let $n = 0$ and $n = 1$. Then the statement becomes:

$$F_0 = \frac{\phi^0 - \hat{\phi}^0}{\sqrt{5}} = \frac{1 - 1}{\sqrt{5}} = 0$$

$$\text{and} \quad F_1 = \frac{\phi^1 - \hat{\phi}^1}{\sqrt{5}} = \frac{\frac{1+\sqrt{5}}{2} - \frac{1-\sqrt{5}}{2}}{\sqrt{5}} = \frac{\sqrt{5}}{\sqrt{5}} = 1$$

which is clearly true.

- Inductive step**

Suppose it is true for n and for $n - 1$, with $n \geq 1$, and let us prove it for $(n + 1)$. By the definition of the Fibonacci sequence we have $F_{n+1} = F_n + F_{n-1}$, while by

the inductive hypothesis we have:

$$F_n = \frac{\phi^n - \hat{\phi}^n}{\sqrt{5}} \quad \text{and} \quad F_{n-1} = \frac{\phi^{n-1} - \hat{\phi}^{n-1}}{\sqrt{5}}$$

Hence we can write:

$$\begin{aligned} F_{n+1} &= F_n + F_{n-1} = \frac{\phi^n - \hat{\phi}^n}{\sqrt{5}} + \frac{\phi^{n-1} - \hat{\phi}^{n-1}}{\sqrt{5}} \\ &= \frac{(\phi^n + \phi^{n-1}) - (\hat{\phi}^n + \hat{\phi}^{n-1})}{\sqrt{5}} = \frac{\phi^{n-1}(\phi + 1) - \hat{\phi}^{n-1}(\hat{\phi} + 1)}{\sqrt{5}} \\ &= \frac{\phi^{n-1} \phi^2 - \hat{\phi}^{n-1} \hat{\phi}^2}{\sqrt{5}} = \frac{\phi^{n+1} - \hat{\phi}^{n+1}}{\sqrt{5}} \end{aligned}$$

where in the second-to-last equality we used the relations (8.5). This is exactly the desired statement, for $n + 1$.

□

- The Fibonacci sequence is studied in more detail in the chapter “Fibonacci sequence”.

Factorials, binomial coefficients and triangle inequality

9.1 Factorials

Definition 9.1: factorial of n

The factorial of n is the product of the first n positive integers. It is denoted by $n!$ and is read “ n factorial”. In formulas:

$$n! = \prod_{k=1}^n k = 1 \cdot 2 \cdot 3 \cdot \dots \cdot (n-1) \cdot n$$

By definition, we set $0! = 1$.

- The number $n!$ grows very rapidly as n increases. The first values are:

n	0	1	2	3	4	5	6	7	8	9	10
$n!$	1	1	2	6	24	120	720	5,040	40,320	362,880	3,628,800

- Some properties of the factorial, which are immediate to verify, are:

$$n! = n \cdot (n-1)!$$

$$\frac{n!}{(n-k)!} = n \cdot (n-1) \cdot (n-2) \cdot \dots \cdot (n-k+1), \quad \text{with } 1 \leq k \leq n \quad (9.1)$$

With $1 \leq k \leq n$, it becomes the product of k factors, starting from n and decreasing by one unit at a time.

Example 9.1: Computing the factorial

$$\frac{100!}{95!} = \frac{100!}{(100-5)!} = 100 \cdot 99 \cdot 98 \cdot 97 \cdot 96 = 9,034,502,400$$

It is always convenient to simplify expressions containing the factorial as much as possible, before computing them!

9.2 Binomial coefficients**Definition 9.2: binomial coefficient**

The **binomial coefficient** is defined as the number:

$$c_{n,k} = \frac{n!}{k! (n-k)!} \quad \text{with } 0 \leq k \leq n. \quad (9.2)$$

The binomial coefficient $c_{n,k}$ is usually denoted by the symbol: $\binom{n}{k}$ which is read “ n choose k ”.

- Given rule (9.1), we have:

$$c_{n,k} = \frac{n \cdot (n-1) \cdot (n-2) \cdot \dots \cdot (n-k+1)}{k!} \quad (9.3)$$

with $k \geq 1$, an expression that is more convenient for computing the binomial coefficient.

We have:

$$\binom{n}{n-k} = \binom{n}{k} \quad (9.4)$$

Since:

$$\binom{n}{n-k} = \frac{n!}{(n-k)! (n-(n-k))!} = \frac{n!}{k! (n-k)!} = \binom{n}{k}$$

We have:

$$\binom{n-1}{k-1} + \binom{n-1}{k} = \binom{n}{k} \quad \text{for every } n \geq 2 \text{ and } 1 \leq k \leq n-1 \quad (9.5)$$

(the proof below needs $(k-1)!$ and $(n-k-1)!$, hence $k \geq 1$ and $k \leq n-1$). Since:

$$\begin{aligned}
 \binom{n-1}{k-1} + \binom{n-1}{k} &= \frac{(n-1)!}{(k-1)! \underbrace{(n-1-(k-1))!}_{(n-k)!(n-k-1)!}} + \frac{(n-1)!}{k!(n-k-1)!} \\
 &= \frac{(n-1)!}{(k-1)!(n-k)(n-k-1)!} + \frac{(n-1)!}{k(k-1)!(n-k-1)!} \\
 &= \frac{k(n-1)! + (n-k)(n-1)!}{k(k-1)!(n-k)(n-k-1)!} = \frac{\overbrace{(n-1)!}^{=n!} n}{\underbrace{k(k-1)!}_{k!} \underbrace{(n-k)(n-k-1)!}_{(n-k)!}} \\
 &= \frac{n!}{k!(n-k)!} = \binom{n}{k}
 \end{aligned}$$

It also follows that:

$$\binom{n}{k-1} + \binom{n}{k} = \binom{n+1}{k} \quad \text{for every } n \geq 1 \text{ and } 1 \leq k \leq n$$

9.2.1 Newton's formula

The n -th power of a binomial $(a+b)$ can be computed with the following formula (from which the name binomial coefficient derives):

Observation 9.3: Newton's formula

For every integer $n \geq 0$, with $a, b \in \mathbb{R}$, we have:

$$(a+b)^n = \sum_{k=0}^n \binom{n}{k} a^{n-k} b^k \quad (9.6)$$

Proof. By induction on n .

- **Base case of the induction**

Let $n = 0$. Then the statement becomes: $(a+b)^0 = \binom{0}{0} a^0 b^0$ i.e. $1 = 1$ which is clearly true (with the convention $0^0 = 1$, which also covers the case $a = b = 0$).

- **Inductive step**

Suppose it is true for n , and let us prove it for $(n+1)$. By the inductive hypothesis,

we have: $(a+b)^n = \sum_{k=0}^n \binom{n}{k} a^{n-k} b^k$. Then:

$$\begin{aligned}
 (a+b)^{n+1} &= (a+b) \cdot (a+b)^n = (a+b) \sum_{k=0}^n \binom{n}{k} a^{n-k} b^k \\
 &= \sum_{k=0}^n \binom{n}{k} a^{n-k+1} b^k + \underbrace{\sum_{k=0}^n \binom{n}{k} a^{n-k} b^{k+1}}_{= \sum_{k=1}^{n+1} \binom{n}{k-1} a^{n-k+1} b^k} \\
 &= a^{n+1} + \sum_{k=1}^n \binom{n}{k} a^{n-k+1} b^k + b^{n+1} + \sum_{k=1}^n \binom{n}{k-1} a^{n-k+1} b^k \\
 &= a^{n+1} + b^{n+1} + \sum_{k=1}^n \underbrace{\left(\binom{n}{k} + \binom{n}{k-1} \right)}_{= \binom{n+1}{k}} a^{n-k+1} b^k \\
 &= \sum_{k=0}^{n+1} \binom{n+1}{k} a^{n+1-k} b^k
 \end{aligned}$$

which is exactly the desired statement, for $n+1$. □

Observation 9.4

For every integer $n \geq 0$ we have:

$$\sum_{k=0}^n \binom{n}{k} = 2^n \quad (9.7)$$

Proof. We write:

$$2^n = (1+1)^n$$

We now apply Newton's formula (9.6):

$$(1+1)^n = \sum_{k=0}^n \binom{n}{k} 1^{n-k} 1^k = \sum_{k=0}^n \binom{n}{k}$$

□

9.2.2 Recursive computation of binomial coefficients

Relation (9.5) allows us to compute the binomial coefficients $\binom{n}{k}$ by means of the so-called **Pascal's triangle** (or Tartaglia's triangle).

The rules for building the triangle are:

1. At the top of the triangle we place the number $\binom{0}{0} = 1$ (by definition).
2. On the sides we place the numbers $\binom{n}{0} = \binom{n}{n} = 1$ for every $n \geq 1$.
3. For $0 < k < n$, the number $\binom{n}{k}$ is written at the intersection of the n -th row and the k -th column.
4. Again for $0 < k < n$, the number $\binom{n}{k}$ is the sum of the two numbers located in the previous row, the one in the same column and the one in the previous column.

Example 9.2: Pascal's triangle (or Tartaglia's triangle)

Pascal's triangle with $n \leq 10$ and $k \leq 10$ is:

	$k = 0$	$k = 1$	$k = 2$	$k = 3$	$k = 4$	$k = 5$	$k = 6$	$k = 7$	$k = 8$	$k = 9$	$k = 10$
$n = 0$	1										
$n = 1$	1	1									
$n = 2$	1	2	1								
$n = 3$	1	3	3	1							
$n = 4$	1	4	6	4	1						
$n = 5$	1	5	10	10	5	1					
$n = 6$	1	6	15	20	15	6	1				
$n = 7$	1	7	21	35	35	21	7	1			
$n = 8$	1	8	28	56	70	56	28	8	1		
$n = 9$	1	9	36	84	126	126	84	36	9	1	
$n = 10$	1	10	45	120	210	252	210	120	45	10	1

To compute the binomial coefficient $\binom{5}{3}$, corresponding to the blue cell, we can use relation (9.5) and add the two binomial coefficients $\binom{4}{2}$ and $\binom{4}{3}$, corresponding to the red cells:

$$\binom{5}{3} = \binom{4}{2} + \binom{4}{3} = 6 + 4 = 10.$$

Example 9.3: computing the power of a binomial using Pascal's triangle

Thanks to the previous Pascal's triangle and to Newton's formula (9.6) we can compute:

$$(a + b)^5 = a^5 + 5 a^4 b + 10 a^3 b^2 + 10 a^2 b^3 + 5 a b^4 + b^5$$

9.3 Absolute value

Definition 9.5: absolute value

The **absolute value** of a real number $a \in \mathbb{R}$ (or **modulus** of a) is the non-negative number defined as follows:

$$|a| = \begin{cases} a & \text{if } a \geq 0 \\ -a & \text{if } a < 0 \end{cases} \quad (9.8)$$

From the definition of absolute value it immediately follows that:

$$\forall \varepsilon \geq 0, a \in \mathbb{R}, \quad |a| \leq \varepsilon \iff -\varepsilon \leq a \leq \varepsilon \quad (9.9)$$

9.3.1 Triangle inequality in $\mathbb{R}$

Observation 9.6: triangle inequality in $\mathbb{R}$

$$|b + c| \leq |b| + |c| \quad \forall b, c \in \mathbb{R} \quad (9.10)$$

Proof. We write the two relations:

$$-|b| \leq b \leq |b|, \quad -|c| \leq c \leq |c|$$

and add them side by side:

$$-(|b| + |c|) \leq b + c \leq |b| + |c|$$

Hence, by (9.9), with $\varepsilon = |b| + |c| \geq 0$ and $a = b + c$, (9.10) follows. $\square$

- The triangle inequality is also used in the following form:

$$|d - e| \leq |d - f| + |e - f| \quad \forall d, e, f \in \mathbb{R} \quad (9.11)$$

To obtain it, it suffices to set in (9.10):

$$b = d - f, \quad c = f - e$$

we obtain:

$$|d - f + f - e| = |d - e| \leq |d - f| + |f - e| = |d - f| + |e - f|$$

since:

$$|f - e| = |e - f| \quad \forall e, f \in \mathbb{R}$$

- Moreover, the triangle inequality can also be written in the following form:

$$|g| \leq |g - h| + |h| \quad \text{i.e.} \quad |g| - |h| \leq |g - h| \quad \forall g, h \in \mathbb{R} \quad (9.12)$$

To obtain it, it suffices to set in (9.10):

$$b = g - h, \quad c = h$$

Observation 9.7: reverse triangle inequality in $\mathbb{R}$

$$||g| - |h|| \leq |g - h|, \quad \forall g, h \in \mathbb{R}. \quad (9.13)$$

Proof. From (9.12) we have

$$|g| - |h| \leq |g - h| \quad \forall g, h \in \mathbb{R}$$

Similarly, swapping g and h in (9.12) we obtain:

$$|h| - |g| \leq |h - g| = |g - h| \quad \text{that is} \quad |g| - |h| \geq -|g - h|$$

hence we have

$$-(|g - h|) \leq |g| - |h| \leq |g - h| \quad \forall g, h \in \mathbb{R}$$

Hence, by (9.9), with $\varepsilon = |g - h| \geq 0$ and $a = |g| - |h|$, the reverse triangle inequality follows. $\square$

- Inequality (9.10) can easily be extended to the case of k terms:

$$\left| \sum_{i=1}^k b_i \right| \leq \sum_{i=1}^k |b_i|. \quad (9.14)$$

- The following immediate properties also hold:

$$\begin{aligned} |bc| &= |b| |c|, & |-b| &= |b| \quad \forall b, c \in \mathbb{R}, \\ \left| \frac{b}{c} \right| &= \frac{|b|}{|c|} & \forall b, c \in \mathbb{R}, \quad c &\neq 0. \end{aligned} \quad (9.15)$$

The Fibonacci sequence

10.1 The Fibonacci sequence

Definition 10.1: Fibonacci sequence

$$\begin{aligned} F_0 &= 0, \\ F_1 &= 1, \\ F_i &= F_{i-1} + F_{i-2}, \quad \text{with } i \geq 2, i \in \mathbb{N} \end{aligned} \quad (10.1)$$

- Therefore, each Fibonacci number is the sum of the two previous ones, which gives the sequence

$$0, 1, 1, 2, 3, 5, 8, 13, 21, 34, 55, \dots$$

- The Fibonacci sequence is related to the **golden ratio** ϕ and to the **conjugate of the golden ratio** $\hat{\phi}$,

Definition 10.2: golden ratio


The **golden ratio** ϕ and the **conjugate of the golden ratio** $\hat{\phi}$ are the two roots of the equation:

$$x^2 = x + 1$$

and are given by the formulas:

$$\begin{aligned} \phi &= \frac{1 + \sqrt{5}}{2} & \hat{\phi} &= \frac{1 - \sqrt{5}}{2} \\ &\approx 1.61803 & &\approx -0.61803 \end{aligned}$$

The following figure shows the graph of the function $f(x) = x^2 - x - 1$ on the interval $[-3, 3]$; the red dots correspond to the roots of the equation $x^2 = x + 1$.



The golden ratio ϕ and the conjugate of the golden ratio $\hat{\phi}$ clearly satisfy the equation $x^2 = x + 1$ since:

$$\phi^2 = \left(\frac{1 + \sqrt{5}}{2} \right)^2 = \frac{1 + 2\sqrt{5} + 5}{4} = \frac{3 + \sqrt{5}}{2} = \frac{1 + \sqrt{5}}{2} + 1 = \phi + 1$$

$$\hat{\phi}^2 = \left(\frac{1 - \sqrt{5}}{2} \right)^2 = \frac{1 - 2\sqrt{5} + 5}{4} = \frac{3 - \sqrt{5}}{2} = \frac{1 - \sqrt{5}}{2} + 1 = \hat{\phi} + 1$$

Observation 10.3: closed formula of the Fibonacci sequence

$$F_i = \frac{\phi^i - \hat{\phi}^i}{\sqrt{5}}, \quad i = 0, 1, 2, \dots$$

Proof. By induction on i .

- **Base case of the induction**

We prove that the formula holds for $i = 0$ and $i = 1$:

$$F_0 = \frac{\phi^0 - \hat{\phi}^0}{\sqrt{5}} = \frac{1 - 1}{\sqrt{5}} = 0, \quad F_1 = \frac{\phi^1 - \hat{\phi}^1}{\sqrt{5}} = \frac{\sqrt{5}}{\sqrt{5}} = 1.$$

- **Inductive step**

Suppose it is true for $i = k$ and $i = k - 1$ with $k \geq 1$, and let us prove it for $i = k + 1$. By the definition of the Fibonacci sequence, we have $F_{k+1} = F_k + F_{k-1}$, while by the inductive hypothesis we have:

$$F_k = \frac{\phi^k - \hat{\phi}^k}{\sqrt{5}} \quad \text{and} \quad F_{k-1} = \frac{\phi^{k-1} - \hat{\phi}^{k-1}}{\sqrt{5}}$$

Hence we can write

$$\begin{aligned} F_{k+1} &= F_k + F_{k-1} \\ &= \frac{\phi^k - \hat{\phi}^k}{\sqrt{5}} + \frac{\phi^{k-1} - \hat{\phi}^{k-1}}{\sqrt{5}} = \frac{(\phi^k - \hat{\phi}^k) + (\phi^{k-1} - \hat{\phi}^{k-1})}{\sqrt{5}} \\ &= \frac{(\phi^k + \phi^{k-1}) - (\hat{\phi}^k + \hat{\phi}^{k-1})}{\sqrt{5}} = \frac{\phi^{k-1}(\phi + 1) - \hat{\phi}^{k-1}(\hat{\phi} + 1)}{\sqrt{5}} \\ &= \frac{\phi^{k-1}(\phi^2) - \hat{\phi}^{k-1}(\hat{\phi}^2)}{\sqrt{5}} = \frac{\phi^{k+1} - \hat{\phi}^{k+1}}{\sqrt{5}} \end{aligned}$$

which is exactly the desired statement, for $i = k + 1$. □

Observation 10.4: computing F_i by rounding

$$F_i = \left\lfloor \frac{\phi^i}{\sqrt{5}} + \frac{1}{2} \right\rfloor, \quad i = 0, 1, 2, \dots$$

Proof. Since $|\hat{\phi}| < 1$, we have

$$\frac{|\hat{\phi}^i|}{\sqrt{5}} \leq \frac{1}{\sqrt{5}} < \frac{1}{2} \quad \text{and} \quad -\frac{1}{2} < \frac{\hat{\phi}^i}{\sqrt{5}} < \frac{1}{2}, \quad i = 0, 1, 2, \dots$$


Since $F_i \in \mathbb{N}$ and $F_i = \frac{\phi^i}{\sqrt{5}} - \frac{\hat{\phi}^i}{\sqrt{5}}$, the i -th Fibonacci number F_i is equal to $\frac{\phi^i}{\sqrt{5}}$ rounded to the nearest integer:

$$F_i = \left\lfloor \frac{\phi^i}{\sqrt{5}} + \frac{1}{2} \right\rfloor, \quad i = 0, 1, 2, \dots$$




- The values of

$$\frac{\hat{\phi}^i}{\sqrt{5}} \text{ with } i = 0, 1, \dots, 10 \text{ are :}$$



- The values of


$$\frac{\phi^i}{\sqrt{5}} - \left\lfloor \frac{\phi^i}{\sqrt{5}} \right\rfloor \text{ with } i = 0, 1, \dots, 10 \text{ are :}$$



- Since $F_i \in \mathbb{N}$, we moreover have:

$$\left(\frac{\phi^i}{\sqrt{5}} - \left\lfloor \frac{\phi^i}{\sqrt{5}} \right\rfloor \right) - \frac{\hat{\phi}^i}{\sqrt{5}} \in \{0, 1\}, \quad i = 0, 1, 2, \dots$$

- The values of F_i , with $i = 0, 1, \dots, 10$, are:



Cardinality of infinite sets

11.1 Countable cardinality

Is it possible to compare the “size” of infinite sets?

- Let us approach the question starting from the important number sets that we have introduced: $\mathbb{N}$, $\mathbb{Z}$, $\mathbb{Q}$, $\mathbb{R}$, and ask ourselves: how many elements does each of these sets have?
- Intuitively, the answer seems obvious: each of these sets has infinitely many elements; however, the rational numbers are more numerous than the integers, since $\mathbb{Z} \subset \mathbb{Q}$, and for the same reason the real numbers are more numerous than the rational numbers, since $\mathbb{Q} \subset \mathbb{R}$.
- How can we state, at the same time, that two sets are both infinite, but one is more numerous than the other?

How is it possible, in general, to compare the size of infinite sets?

- To give meaning to these questions, even before being able to answer them, we must define what we mean by two sets having the same size.
- Abstracting from the experience of counting the elements of a finite set, the idea of **same size** has been identified with that of **one-to-one correspondence** (bijection):

Definition 11.1: equal cardinality of two sets

Two sets A , B are said to have **equal cardinality** (or **power**) if they can be put in one-to-one correspondence with each other, i.e., if there exists a rule that associates with each element of A one and only one element of B , and vice versa.

- The cardinality (or power) of a set expresses the intuitive idea of size

Observation 11.2

The set of integers $\mathbb{Z}$ and the set of natural numbers $\mathbb{N}$ have the same cardinality

Proof. The following rule puts the integers and the natural numbers in one-to-one correspondence:

$\mathbb{Z}$	0	1	-1	2	-2	...	n	$-n$	...
	$\updownarrow$	$\updownarrow$	$\updownarrow$	$\updownarrow$	$\updownarrow$		$\updownarrow$	$\updownarrow$	
$\mathbb{N}$	0	1	2	3	4	...	$2n-1$	$2n$	...

□

- Therefore, even though from the point of view of inclusion $\mathbb{Z}$ has “more elements” than $\mathbb{N}$ (in the sense that it has all the elements of $\mathbb{N}$ plus others), the two sets have the same cardinality.

Two sets that have the same cardinality are also called **equipotent** and should be thought of as equally numerous.

Definition 11.3: countable set

A set that has the same cardinality as $\mathbb{N}$ is called **countable**.

- The term countable indicates that the elements of the set can be enumerated, that is, arranged in a numbered list (place 1, place 2, place 3, ...)

The cardinality of $\mathbb{N}$ is called the **cardinality of the countable** (countable infinity).

Theorem 11.4

The set $\mathbb{Q}$ is countable

Proof. • We begin by proving that the set of positive rational numbers is countable.

- To do this, we represent the positive rational numbers as fractions n/m , with n, m positive integers, and we arrange these fractions in an **infinite triangular table**, in the following way:

$\frac{n}{m}$, with $n + m = \dots$				
2	$\frac{1}{1}$			
3	$\frac{1}{2}$	$\frac{2}{1}$		
4	$\frac{1}{3}$	$\frac{2}{2}$	$\frac{3}{1}$	
5	$\frac{1}{4}$	$\frac{2}{3}$	$\frac{3}{2}$	$\frac{4}{1}$
...	...			

Note that: all the fractions representing positive rational numbers appear in this table at least once; some numbers are repeated several times (e.g., $\frac{1}{1} = \frac{2}{2}$). Each row has finite length.

- Hence we can put $\mathbb{N}$ in one-to-one correspondence with the set of positive rationals, by going through the table row by row (and skipping an element when it is equal to one already encountered). For example:

1	2	3	4	5	6	7	8	9	...
$\frac{1}{1}$	$\frac{1}{2}$	$\frac{2}{1}$	$\frac{1}{3}$	$\frac{3}{1}$	$\frac{1}{4}$	$\frac{2}{3}$	$\frac{3}{2}$	$\frac{4}{1}$	...

(note that we skipped $\frac{2}{2}$ because it is equal to $\frac{1}{1}$, already encountered).

- This proves that the set of positive rationals is countable.
- Then $\mathbb{Q}$ is also countable, and this can be proved with an argument similar to the one we used to prove that $\mathbb{Z}$ is countable: denoting by $q_1, q_2, q_3, \dots$ the positive rationals, we set:

$\mathbb{N}$	0	1	2	3	4	5	6	...
	$\updownarrow$	$\updownarrow$	$\updownarrow$	$\updownarrow$	$\updownarrow$	$\updownarrow$	$\updownarrow$	
$\mathbb{Q}$	0	q_1	$-q_1$	q_2	$-q_2$	q_3	$-q_3$	...

which gives a one-to-one correspondence between $\mathbb{N}$ and $\mathbb{Q}$.

□

- Having discovered that several infinite sets, each properly contained in the next ($\mathbb{N}, \mathbb{Z}, \mathbb{Q}$), have the same cardinality, might suggest that this is true for all infinite sets. This is not true, as shown in the next section.

11.2 Cardinality of the continuum

Theorem 11.5

The set of real numbers $\mathbb{R}$ is not countable

- The proof technique is called **Cantor's diagonal argument**

Proof. • We will prove that the interval $[0, 1]$ has uncountable cardinality, from which the uncountability of $\mathbb{R}$ obviously follows

- Suppose therefore, **by contradiction**, that $[0, 1]$ is countable, and arrange **all** the real numbers of the interval $[0, 1]$ in a list $r_1, r_2, r_3, \dots$
- We write each number r_i in decimal form:

$$0.a_1a_2a_3\dots$$

where the a_i are digits from 0 to 9; if the digits are all zero we have $r_i = 0$ and if they are all nine we have $r_i = 0.\overline{9} = 1$:

$$r_1 = 0.a_{11}a_{12}a_{13}\dots$$

$$r_2 = 0.a_{21}a_{22}a_{23}\dots$$

$$r_3 = 0.a_{31}a_{32}a_{33}\dots$$

...

- We now define the following decimal number:

$$r = 0.b_1b_2b_3\dots$$

where the digits b_i are defined by the following rule:

$$b_i = \begin{cases} 5 & \text{if } a_{ii} \text{ is one of the digits } 0, 1, 2, 3 \text{ or } 4 \\ 4 & \text{if } a_{ii} \text{ is one of the digits } 5, 6, 7, 8 \text{ or } 9 \end{cases}$$

With this definition we have $b_i \neq a_{ii}$ for every i . Note that the number r has been constructed by reasoning on the diagonal of the infinite table whose rows are the numbers r_i , hence the name of the argument. □

Proof. • We now observe that r is a real number belonging to the interval $[0, 1]$ (because it is of the form $0.b_1b_2b_3\dots$ with all digits equal to 4 or 5) and on the other hand it is not equal to any of the numbers r_i in the list, since:

$\rightarrow r \neq r_1$: the first digit of r is different from the first digit of r_1 ($b_1 \neq a_{11}$);


$\rightarrow r \neq r_2$: the second digit of r is different from the second digit of r_2 ($b_2 \neq a_{22}$);
 $\rightarrow \dots$

- But this leads to a **contradiction**, since we had assumed that the r_i completely exhausted the set of real numbers of the interval $[0, 1]$. Therefore $[0, 1]$ is not countable.

□

Since $\mathbb{N}$ cannot be put in one-to-one correspondence with $\mathbb{R}$, but it can be put in one-to-one correspondence with a proper subset of $\mathbb{R}$ ($\mathbb{N}$ itself!), we say that $\mathbb{N}$ has **smaller cardinality** than $\mathbb{R}$.

- We also note that the interval $[0, 1]$, and any interval of $\mathbb{R}$ (open or closed, bounded or unbounded), has the same cardinality as $\mathbb{R}$.
- This fact can be proved, for example, with an elementary geometric construction that gives a one-to-one correspondence between the points of the line and those of a segment $[A, B]$; as shown in the following figure, by drawing suitable segments from the two fixed points P_1, P_2 to the line itself:



- This implies that, for example, the points of a line have the same cardinality as the points of a segment.

The cardinality of $\mathbb{R}$ is called the **cardinality of the continuum**.

- It is not only every interval of $\mathbb{R}$ that has this same cardinality. It can be proved that the same holds for the plane, for three-dimensional space and for their “continuous” subsets: for example, a plane and a sphere both have the cardinality of the continuum.

- So far we have encountered only two hierarchical levels of infinity: the cardinality of the countable and that of the continuum. One should not believe that only these two exist! For example, the set of all subsets of $\mathbb{R}$ (its power set) is even more numerous than $\mathbb{R}$; and with this procedure we can always construct a set more numerous than a given set.
- Therefore the hierarchical levels of infinite cardinalities are themselves infinite.

Complex numbers


12.1 Definition of $\mathbb{C}$ and field structure

- We have denoted by $\mathbb{R}^2$ (short for $\mathbb{R} \times \mathbb{R}$) the set of ordered pairs (a, b) of real numbers.
- On these pairs we directly define the operations of sum and product with the following rules:


$$(a, b) + (c, d) = (a + c, b + d) \quad (12.1)$$

$$(a, b) \cdot (c, d) = (ac - bd, ad + bc) \quad (12.2)$$

Example 12.1: sum



Example 12.2: product



- This “sum” and this “product” satisfy the **commutative**, **associative** and **distributive** properties
- We also observe that, $\forall (a, b) \in \mathbb{R}^2$:

$$(a, b) + (0, 0) = (0, 0) + (a, b) = (a, b)$$

hence the pair $(0, 0)$ is the **identity element for the sum**. Moreover:

$$(a, b) \cdot (1, 0) = (1, 0) \cdot (a, b) = (a, b)$$

hence the pair $(1, 0)$ is the **identity element for the product**. We have:

$$(a, b) + (-a, -b) = (0, 0)$$

hence $(-a, -b)$ is the **opposite** of (a, b) . Moreover, if $(a, b) \neq (0, 0)$ then:

$$(a, b) \cdot \left(\frac{a}{a^2 + b^2}, \frac{-b}{a^2 + b^2} \right) = (1, 0)$$

hence the pair $(a/(a^2 + b^2), -b/(a^2 + b^2))$ is the **reciprocal** of (a, b) .

Definition 12.1: field of complex numbers

Properties R_1, R_2 are satisfied by the sum and the product defined above, and therefore the set $\mathbb{R}^2$ with this structure is a field, which we will call the **field of complex numbers** and denote by $\mathbb{C}$

- We now observe that $\mathbb{C}$ contains the subset $\mathbb{C}_0$ of pairs of the form $(a, 0)$; it is a subfield of $\mathbb{C}$, since the sum and product of pairs of this form are again pairs of the same form; indeed we have:

$$(a, 0) + (b, 0) = (a + b, 0) \quad \text{and} \quad (a, 0) \cdot (b, 0) = (a \cdot b, 0)$$

Moreover, $\mathbb{C}_0$ can be ordered by setting $(a, 0) < (b, 0)$ if $a < b$.

- If we then put the set of real numbers $\mathbb{R}$ in one-to-one correspondence with $\mathbb{C}_0$, by setting

$$(a, 0) \longleftrightarrow a$$

we can identify the real numbers $a \in \mathbb{R}$ with the complex numbers of the form $(a, 0) \in \mathbb{R}^2$. In this sense the field of complex numbers $\mathbb{C}$ is an **extension** of the field of real numbers $\mathbb{R}$.

Let us now consider the number $(0, 1) \in \mathbb{C}$. It has the remarkable property that:

$$(0, 1) \cdot (0, 1) = (-1, 0)$$

i.e., its square coincides with the real number -1

Definition 12.2: imaginary unit

The pair $(0, 1) \in \mathbb{C}$ is denoted by the letter “ i ” and is called the **imaginary unit**

12.2 Algebraic form of complex numbers

We observe that, if we write any complex number $(c, 0)$ simply as c , we have:

$$(a, b) = (a, 0) + \underbrace{(0, 1)}_{=i} \cdot (b, 0) = a + ib$$

With this notation, rules (12.1) and (12.2) are the ordinary rules of algebraic calculation, keeping in mind that $i^2 = -1$:

$$(a + ib) + (c + id) = (a + c) + i(b + d) \quad (12.3)$$

$$(a + ib) \cdot (c + id) = (ac - bd) + i(ad + bc) \quad (12.4)$$

Definition 12.3: algebraic form, real part and imaginary part

The expression:


$$z = a + ib \quad (12.5)$$

is called the **algebraic form of complex numbers**; a is called the **real part** of z and is denoted by $\Re(z)$ (or $\text{Re}(z)$) while b is called the **imaginary part** and is denoted by $\Im(z)$ (or $\text{Im}(z)$).

Complex plane

- In a Cartesian plane, the complex numbers $a + ib$ can be represented as points with coordinates (a, b) . In this context:

- the plane is called the **complex plane** or **Gauss plane**
- the x , y axes are called the **real axis** and the **imaginary axis**
- the points on the real axis are the **real numbers**
- the points on the imaginary axis are the **purely imaginary numbers** (i.e., of the form ib)
- The **sum of two complex numbers** is the complex number whose coordinates are the sums of the coordinates: the geometric meaning of this fact is that the point $z + t$ is constructed from the points z , t according to the “**parallelogram rule**”, illustrated in the following figure:



Observation 12.4

The set of complex numbers $\mathbb{C}$ is not an ordered field

Proof. • We have seen that $\mathbb{C}$ satisfies the field axioms; however, it does not satisfy those of an ordered field, that is, it is not possible to define a relation $\leq$ between complex numbers in such a way that properties $R3$ hold

- It can be proved that properties $R3$ imply that the square of any number is never negative, and on the other hand, if a number is positive its opposite is negative. Now, in $\mathbb{C}$ we have:

$$1^2 = 1 \quad \text{and} \quad i^2 = -1$$

We therefore have two squares, each the opposite of the other. Neither of them, however, can be negative (because they are squares), and this is absurd (because

between a and $-a$ one must be negative, if $a \neq 0$). We conclude that $\mathbb{C}$ is not an ordered field. $\square$

Definition 12.5: conjugate

The complex number $a - ib$ is called the **complex conjugate** of $z = a + ib$ and is denoted by $\bar{z}$.

We have:

$$z + \bar{z} = 2a = (a + ib) + (a - ib) = 2a = 2 \Re(z) \quad (12.6)$$

$$z - \bar{z} = 2ib = (a + ib) - (a - ib) = 2ib = 2i \Im(z) \quad (12.7)$$

$$z \cdot \bar{z} = (a + ib) \cdot (a - ib) = a^2 - aib + iba - \underbrace{i^2}_{=-1} b^2 = a^2 + b^2 \geq 0 \quad (12.8)$$

- The conjugation operation has the following elementary properties with respect to sum and product:

$$\overline{(z_1 + z_2)} = \bar{z}_1 + \bar{z}_2 \quad (12.9)$$

$$\overline{(z_1 \cdot z_2)} = \bar{z}_1 \cdot \bar{z}_2 \quad (12.10)$$

$$\overline{\left(\frac{1}{z}\right)} = \frac{1}{\bar{z}} \quad (12.11)$$

Definition 12.6: modulus

The **modulus** of $z = a + ib$ is the non-negative real number $\sqrt{a^2 + b^2}$; it is denoted by $|z|$.

- If $z = a$ is real, its modulus is called the absolute value and is still denoted by $|a|$. The following properties hold:
 - $|z| = 0 \iff z = 0$, moreover $|z| \geq 0$
 - $|z| = |\bar{z}|$
 - $\Re(z) \leq |z| \quad \Im(z) \leq |z| \quad |z| \leq |\Re(z)| + |\Im(z)|$
 - $|z_1 + z_2| \leq |z_1| + |z_2|$ **triangle inequality**
 - $|z_1 + z_2| \geq ||z_1| - |z_2||$
- Properties a), b), c) are immediately verified.

Proof. • Let us prove properties d) and e). They are equivalent to the following:

$$(|z_1| - |z_2|)^2 \leq |z_1 + z_2|^2 \leq (|z_1| + |z_2|)^2$$

- Setting $z_1 = a + ib$, $z_2 = c + id$ we obtain:

$$\begin{aligned} |z_1 + z_2|^2 &= |(a + ib) + (c + id)|^2 \\ &= |(a + c) + i(b + d)|^2 \\ &= \left(\sqrt{(a + c)^2 + (b + d)^2} \right)^2 = (a + c)^2 + (b + d)^2 \end{aligned}$$

and hence we have:

$$\begin{aligned} \underbrace{\left(\sqrt{a^2 + b^2} - \sqrt{c^2 + d^2} \right)^2}_{=(a^2+b^2)+(c^2+d^2)-2\sqrt{a^2+b^2}\cdot\sqrt{c^2+d^2}} &\leq \underbrace{(a + c)^2 + (b + d)^2}_{=a^2+2ac+c^2+b^2+2bd+d^2} \\ &\leq \underbrace{\left(\sqrt{a^2 + b^2} + \sqrt{c^2 + d^2} \right)^2}_{=(a^2+b^2)+(c^2+d^2)+2\sqrt{a^2+b^2}\cdot\sqrt{c^2+d^2}} \end{aligned}$$

After simplification, this double inequality reduces to:

$$-\sqrt{a^2 + b^2} \cdot \sqrt{c^2 + d^2} \leq ac + bd \leq \sqrt{a^2 + b^2} \cdot \sqrt{c^2 + d^2}$$

which is equivalent to the following:

$$|ac + db| \leq \sqrt{a^2 + b^2} \cdot \sqrt{c^2 + d^2}$$

Squaring both sides we arrive at:

$$\underbrace{(ac + db)^2}_{=a^2c^2+d^2b^2+2acbd} \leq \underbrace{(a^2 + b^2) \cdot (c^2 + d^2)}_{=a^2c^2+a^2d^2+b^2c^2+b^2d^2}$$


that is,

$$0 \leq -2acbd + a^2d^2 + b^2c^2 = (ad - bc)^2$$

which is true for every $a, b, c, d \in \mathbb{R}$. □

- Geometrically, $|z|$ represents the distance of the point (or complex number) z from the origin; $|z_1 - z_2|$ represents the distance between the two points z_1 and z_2 ; inequalities d) and e) express the well-known theorem on the **lengths of the sides of a triangle**¹:

¹In a non-degenerate triangle, the sum of the lengths of two sides is greater than the length of the third one.



- Using the concepts just introduced, we can write the quotient of two complex numbers in algebraic form:

$$\frac{a + ib}{c + id}$$

it suffices to multiply the numerator and the denominator by $c - id$, and we obtain:

$$\frac{a + ib}{c + id} = \frac{(a + ib)(c - id)}{\underbrace{(c + id)(c - id)}_{=c^2+d^2}} = \frac{ac - aid + ibc - i^2bd}{c^2 + d^2} = \frac{(ac + bd)}{c^2 + d^2} + i \frac{(bc - ad)}{c^2 + d^2}$$

12.2.1 Equations in the complex field

- Let us see how to solve an equation in the complex field, when it involves the unknown $z = x + iy$ also through $\Re(z)$, $\Im(z)$, z , $|z|$.
- We illustrate, with the following example, the procedure of transforming the equation in one complex unknown into a system of two equations in two real unknowns. The method consists in taking the real and imaginary parts of the equation.

Example 12.3: equations in the complex field (algebraic method)

- We want to solve:

$$z^2 + i \Im z + 2 \bar{z} = 0$$

We set $z = x + iy$, with x, y real unknowns, and rewrite the equation:

$$\begin{cases} z^2 = (x + iy)^2 = x^2 - y^2 + 2ixy \\ i \Im z = iy \\ 2\bar{z} = 2(x - iy) = 2x - 2iy \end{cases}$$

Substituting, we therefore obtain:

$$(x^2 - y^2 + 2ixy) + (iy) + (2x - 2iy) = 0$$

- Now, a complex number is zero if and only if its real part and imaginary part are zero. Therefore we separate the real part and the imaginary part of the left-hand side:

$$(x^2 - y^2 + 2x) + i(2xy + y - 2y) = 0$$

and set both equal to zero:

$$\begin{cases} x^2 - y^2 + 2x = 0 \\ 2xy - y = 0 \end{cases}$$

We have thus transformed the equation in one complex unknown into a system of two equations in two real unknowns.

- We solve the system. The second equation gives:

$$y = 0 \quad \text{or} \quad x = \frac{1}{2}$$

1. For $y = 0$ the first equation becomes

$$x^2 + 2x = 0 \quad \text{which gives} \quad x = 0 \quad \text{or} \quad x = -2$$

2. For $x = \frac{1}{2}$ the first equation becomes

$$-y^2 + \frac{5}{4} = 0 \quad \text{which gives} \quad y = \pm \frac{\sqrt{5}}{2}$$


- Hence the equation has the following 4 solutions:

$$z = 0, \quad z = -2, \quad z = \frac{1}{2} + i \frac{\sqrt{5}}{2}, \quad z = \frac{1}{2} - i \frac{\sqrt{5}}{2}$$

The method seen in this example can in principle be applied to any equation in $\mathbb{C}$, but a generic system of two equations in two unknowns is almost always impossible to solve algebraically.

12.3 Trigonometric form of complex numbers

- As is known from Geometry, the points of the plane can be identified not only by their Cartesian coordinates, but also by their **polar coordinates**:
 - $\varrho \rightarrow$ **polar radius**, i.e., the distance of the point from the origin
 - $\vartheta \rightarrow$ **polar angle**, i.e., the angle that the line joining the point to the origin forms with the positive x -axis, measured counterclockwise.
- Clearly, a pair ϱ, ϑ , with $\varrho > 0$, identifies a well-defined point of the plane; conversely, a point of the plane uniquely determines the coordinate ϱ , but the angle ϑ , measured in radians, is determined only up to multiples of 2π .



- Given a complex number z , its **modulus** $|z|$ coincides with the **polar radius** ϱ of the point representing it in the complex plane.
- We call **argument** of z , and denote by $\arg(z)$, any of the angles ϑ associated with the point z . In this way the argument of z is not uniquely determined. Often this indeterminacy does not cause any problem. At other times, however, it is preferable to assign a well-defined argument to a complex number. This can be done in infinitely many ways, by fixing any interval of length 2π within which the angle ϑ is allowed to vary
- The intervals most commonly used for this purpose are $[0, 2\pi)$ and $(-\pi, \pi]$; the argument of z is then called the **principal argument**.

Example 12.4: arguments of complex numbers

- the number $-i$ has argument $-\pi/2$ or $3\pi/2$ or any other value of the form $-\pi/2 + 2k\pi$ with $k \in \mathbb{Z}$. Its principal argument will be $3\pi/2$ if we adopt the convention $\vartheta \in [0, 2\pi)$, and $-\pi/2$ with the convention $\vartheta \in (-\pi, \pi]$.
- Positive real numbers have principal argument 0 and negative ones π , with both conventions.

Given the number $z = a + ib$, from trigonometry we immediately obtain the relations between the Cartesian coordinates a, b and the polar ones ϱ, ϑ :

$$a = \varrho \cos \vartheta \quad \text{and} \quad b = \varrho \sin \vartheta \quad (12.12)$$

The inverse relations are:

$$\varrho = \sqrt{a^2 + b^2}, \quad \cos \vartheta = \frac{a}{\sqrt{a^2 + b^2}} \quad \text{and} \quad \sin \vartheta = \frac{b}{\sqrt{a^2 + b^2}} \quad (12.13)$$

Definition 12.7: trigonometric form

A complex number $z = a + ib$ can also be written in the form

$$z = \varrho (\cos \vartheta + i \sin \vartheta) \quad (12.14)$$

which is called the **trigonometric form** of complex numbers.

Example 12.5: trigonometric form

Let us write the following complex number in trigonometric form:

$$z = \sqrt{3} + i$$

We have $\varrho = \sqrt{a^2 + b^2} = \sqrt{3 + 1} = 2$, hence:

$$\sqrt{3} + i = 2 \left(\frac{\sqrt{3}}{2} + i \frac{1}{2} \right) = 2 \left(\cos \frac{\pi}{6} + i \sin \frac{\pi}{6} \right)$$

12.3.1 De Moivre's formulas

- The trigonometric form is convenient for expressing products and quotients of complex numbers. Indeed, if we have:

$$z_1 = \varrho_1 (\cos \vartheta_1 + i \sin \vartheta_1) \quad z_2 = \varrho_2 (\cos \vartheta_2 + i \sin \vartheta_2)$$

for the **product** we obtain:

$$\begin{aligned} z_1 z_2 &= \varrho_1 \varrho_2 \cdot \left\{ \cos \vartheta_1 \cos \vartheta_2 - \sin \vartheta_1 \sin \vartheta_2 + i (\sin \vartheta_1 \cos \vartheta_2 + \cos \vartheta_1 \sin \vartheta_2) \right\} \\ &= \varrho_1 \varrho_2 \cdot \left\{ \cos (\vartheta_1 + \vartheta_2) + i \sin (\vartheta_1 + \vartheta_2) \right\} \end{aligned} \quad (12.15)$$

for the **quotient**, if $z_2 \neq 0$, we have:

$$\frac{z_1}{z_2} = \frac{\varrho_1}{\varrho_2} \cdot \frac{\cos \vartheta_1 + i \sin \vartheta_1}{\cos \vartheta_2 + i \sin \vartheta_2},$$

multiplying numerator and denominator by $(\cos \vartheta_2 - i \sin \vartheta_2)$ and, taking into account that $(\cos \vartheta_2)^2 + (\sin \vartheta_2)^2 = 1$, we obtain:

$$\begin{aligned}\frac{z_1}{z_2} &= \frac{\varrho_1}{\varrho_2} \cdot \left\{ (\cos \vartheta_1 + i \sin \vartheta_1) \cdot (\cos \vartheta_2 - i \sin \vartheta_2) \right\} \\ &= \frac{\varrho_1}{\varrho_2} \cdot \left\{ \cos(\vartheta_1 - \vartheta_2) + i \sin(\vartheta_1 - \vartheta_2) \right\}\end{aligned}\quad (12.16)$$

- Therefore the modulus of the product and of the quotient of two complex numbers is, respectively, the product and the quotient of the moduli; the argument is, respectively, the sum and the difference of the arguments:

$$|z_1 \cdot z_2| = |z_1| \cdot |z_2| \quad \text{and} \quad \arg(z_1 \cdot z_2) = \arg(z_1) + \arg(z_2) \quad (12.17)$$

$$\left| \frac{z_1}{z_2} \right| = \frac{|z_1|}{|z_2|} \quad \text{and} \quad \arg\left(\frac{z_1}{z_2}\right) = \arg(z_1) - \arg(z_2) \quad (12.18)$$

- Formula (12.15) generalizes to the case of any number of factors $z_1, z_2, \dots, z_n$:

$$z_1 z_2 \dots z_n = \varrho_1 \varrho_2 \dots \varrho_n \cdot \left\{ \cos(\vartheta_1 + \vartheta_2 + \dots + \vartheta_n) + i \sin(\vartheta_1 + \vartheta_2 + \dots + \vartheta_n) \right\} \quad (12.19)$$

If, moreover, all the factors are equal, we obtain:

$$z^n = \varrho^n \cdot \left\{ \cos(n\vartheta) + i \sin(n\vartheta) \right\} \quad (12.20)$$

- These relations on products and quotients of complex numbers are known as De Moivre's formulas.

Example 12.6: powers of complex numbers with De Moivre's formulas

Write in algebraic form:

$$z = (1 + i)^7$$

We determine the modulus and argument of $(1 + i)$ and then apply De Moivre's formula.

$$|1 + i| = \sqrt{2} \quad \text{and} \quad \arg(1 + i) = \frac{\pi}{4} \quad \left(\cos \vartheta = \frac{1}{\sqrt{2}}, \quad \sin \vartheta = \frac{1}{\sqrt{2}} \right)$$

Hence we have:

$$|(1 + i)^7| = (\sqrt{2})^7 = 2^{\frac{7}{2}} = 2^{3+\frac{1}{2}} = 8\sqrt{2} \quad \text{and} \quad \arg(1 + i)^7 = \frac{7}{4} \pi$$

Consequently:

$$(1 + i)^7 = 8\sqrt{2} \left(\underbrace{\cos \frac{7}{4} \pi}_{=\frac{1}{\sqrt{2}}} + i \underbrace{\sin \frac{7}{4} \pi}_{=-\frac{1}{\sqrt{2}}} \right) = 8\sqrt{2} \left(\frac{1}{\sqrt{2}} - i \frac{1}{\sqrt{2}} \right) = 8 - 8i$$

- De Moivre's formulas allow us to give a geometric interpretation of the product of complex numbers.
- Let z , to begin with, be a complex number with modulus 1, hence of the form $(\cos \vartheta + i \sin \vartheta)$. Then, multiplying a number by z means adding ϑ to its argument, i.e., performing a **rotation by the angle ϑ** .
- If z has modulus ϱ instead of 1, in addition to a rotation we perform a **dilation by the factor ϱ** .

Example 12.7: geometric interpretation of the product of complex numbers

- multiplying by i means performing a rotation by $\frac{\pi}{2}$;
- multiplying by -1 means performing a rotation by π ;
- multiplying by $(1 + i)$ means performing a dilation by the factor $\sqrt{2}$ and a rotation by $\frac{\pi}{4}$

Example 12.8: equations in the complex field (trigonometric method)

- We want to solve:

$$z^3 - |z| = 0$$

We rewrite the equation in the form

$$z^3 = |z|$$

and we set $z = \varrho (\cos \vartheta + i \sin \vartheta)$ (trigonometric form) and rewrite the equation:

$$\begin{cases} z^3 = \varrho^3 (\cos 3\vartheta + i \sin 3\vartheta) \\ |z| = \varrho \end{cases}$$

Substituting, we therefore obtain:

$$\varrho^3 (\cos 3\vartheta + i \sin 3\vartheta) = \varrho$$

The equation is satisfied if and only if the two sides have equal moduli and arguments that differ by multiples of 2π (the right-hand side has argument 0), that is:

$$\begin{cases} \varrho^3 = \varrho \\ 3\vartheta = 2k\pi \quad \text{with } k \in \mathbb{Z} \end{cases}$$

The first equation gives $\varrho = 0$ and $\varrho = 1$ (careful: ϱ must be ≥ 0 because it is the modulus of the complex number; therefore $\varrho = -1$ is not acceptable); the second one gives $\vartheta = \frac{2k\pi}{3}, k \in \mathbb{Z}$. Hence:

$$z = 0, \quad z = \cos \frac{2k\pi}{3} + i \sin \frac{2k\pi}{3} \quad \text{with } k \in \mathbb{Z}$$

Explicitly:

$$z = 0, \quad z = 1, \quad z = -\frac{1}{2} + i \frac{\sqrt{3}}{2}, \quad z = -\frac{1}{2} - i \frac{\sqrt{3}}{2}$$

12.3.2 n -th roots of complex numbers

Definition 12.8: n -th root of a complex number

Given a complex number w , we say that z is an n -th (complex) root of w if $z^n = w$.

Theorem 12.9

Let $w \in \mathbb{C}$, $w \neq 0$, and n an integer ≥ 1 . There exist exactly n complex n -th roots $z_0, z_1, \dots, z_{n-1}$ of w ; setting

$$w = r (\cos \varphi + i \sin \varphi) \quad \text{and} \quad z_k = \varrho_k (\cos \vartheta_k + i \sin \vartheta_k)$$

we have

$$\begin{cases} \varrho_k = r^{1/n} \\ \vartheta_k = \frac{\varphi + 2k\pi}{n} \end{cases} \quad k = 0, 1, 2, \dots, n-1 \quad (12.21)$$

Proof. • The numbers z_k are clearly roots of w , as can be seen by computing z_k^n with De Moivre's formula. Let us show that there are no others.

- If a number $R(\cos \psi + i \sin \psi)$ is an n -th root of w , we must have

$$R^n = r \quad \text{and} \quad n\psi = \varphi + 2h\pi \quad \text{with} \quad h \in \mathbb{Z}$$

or equivalently:

$$R = r^{1/n} \quad \text{and} \quad \psi = \varphi/n + 2h\pi/n \quad \text{with} \quad h \in \mathbb{Z}$$

- Giving h the values $0, 1, \dots, n-1$ we find exactly the numbers z_k .
- Giving h any other value $\bar{h}$ different from the previous ones, it can be written in the form $\bar{h} = k + mn$ ($m \in \mathbb{Z}$ is the quotient and k is the remainder of the division of $\bar{h}$ by n), so that we would have

$$\psi = \frac{\varphi}{n} + \frac{2k\pi}{n} + 2m\pi = \vartheta_k + 2m\pi$$

and we would find again the same z_k as before. □

Example 12.9: fifth root of a complex number

- Let us compute $\sqrt[5]{1+i}$: the complex number $(1+i)$ has modulus $\sqrt{2}$ and argument $\pi/4$. The fifth roots will therefore be:

$$\sqrt[5]{1+i} = \sqrt[5]{\sqrt{2}} \left[\cos \left(\frac{\frac{\pi}{4} + 2k\pi}{5} \right) + i \sin \left(\frac{\frac{\pi}{4} + 2k\pi}{5} \right) \right] =$$

$$= \sqrt[10]{2} \left[\cos \left(\frac{\pi}{20} + \frac{2}{5}k\pi \right) + i \sin \left(\frac{\pi}{20} + \frac{2}{5}k\pi \right) \right] \quad \text{with } k = 0, 1, 2, 3, 4$$

- The angles found are not standard angles, but if desired, their sines and cosines can be computed approximately using a calculator.

- Unfortunately, a somewhat ambiguous notation is used for complex roots, the same one used to denote the arithmetic root; that is, $\sqrt[n]{z}$ or $z^{1/n}$ denotes the set of the n complex roots of z .
- This can create confusion when z is real. Indeed, the symbol $\sqrt{4}$, understood as the arithmetic root of 4, is 2; understood as the complex root of 4, it is the set of the two numbers $+2$ and -2 .

Example 12.10: cube root of a complex number

- Let us compute $\sqrt[3]{-1}$: the complex number (-1) has modulus 1 and argument π . The cube roots will therefore be:

$$\sqrt[3]{-1} = \underbrace{\sqrt[3]{1}}_{=1} \left[\cos \left(\frac{\pi}{3} + \frac{2}{3}k\pi \right) + i \sin \left(\frac{\pi}{3} + \frac{2}{3}k\pi \right) \right] \quad \text{with } k = 0, 1, 2$$

that is, the numbers z_k of the form:

$$\cos \vartheta_k + i \sin \vartheta_k \quad \text{with } \vartheta_k = \frac{\pi}{3} + \frac{2}{3}k\pi \quad \text{and } k = 0, 1, 2$$


- Explicitly we have:

$$\begin{cases} z_0 = \cos \left(\frac{\pi}{3} \right) + i \sin \left(\frac{\pi}{3} \right) = \frac{1}{2} + i \frac{\sqrt{3}}{2} = \frac{1}{2}(1 + i\sqrt{3}) \\ z_1 = \cos \left(\frac{\pi}{3} + \frac{2\pi}{3} \right) + i \sin \left(\frac{\pi}{3} + \frac{2\pi}{3} \right) = \cos(\pi) + i \sin(\pi) = -1 \\ z_2 = \cos \left(\frac{\pi}{3} + \frac{4\pi}{3} \right) + i \sin \left(\frac{\pi}{3} + \frac{4\pi}{3} \right) \\ \quad = \cos \left(\frac{5\pi}{3} \right) + i \sin \left(\frac{5\pi}{3} \right) = \frac{1}{2} - i \frac{\sqrt{3}}{2} = \frac{1}{2}(1 - i\sqrt{3}) \end{cases}$$

- The arrangement of the roots of complex numbers in the Gauss plane is not random.
- Indeed, if $w = r(\cos \varphi + i \sin \varphi)$, the n -th roots $z_0, z_1, \dots, z_{n-1}$ of w lie at the vertices of the regular polygon with n sides inscribed in the circle with center 0 and radius $r^{1/n}$, with the vertex z_0 located at the point with argument $\vartheta = \varphi/n$.

Example 12.11: roots in the Gauss plane

- The figure shows the cube roots of -1 : z_0, z_1, z_2 from the previous exercise:




- The figure shows the sixth roots of i : $z_0, z_1, z_2, z_3, z_4, z_5$,

$$\sqrt[6]{i} = \underbrace{\sqrt[6]{1}}_{=1} \left[\cos \left(\frac{\pi}{2 \cdot 6} + \frac{2}{6} k \pi \right) + i \sin \left(\frac{\pi}{2 \cdot 6} + \frac{2}{6} k \pi \right) \right] \quad \text{with } k = 0, 1, 2, 3, 4, 5$$

that is, the numbers z_k of the form:

$$\cos \vartheta_k + i \sin \vartheta_k \quad \text{with } \vartheta_k = \frac{\pi}{12} + \frac{1}{3} k \pi \quad \text{and } k = 0, 1, 2, 3, 4, 5$$




12.3.3 Exponential form of complex numbers

- The following notation is useful:

$$e^{i\varphi} = \cos \varphi + i \sin \varphi \quad \text{with } \varphi \in \mathbb{R}$$

where e is Euler's number (Napier's constant).



With this notation, the *trigonometric form* can be equivalently rewritten as:

$$z = |z| (\cos \varphi + i \sin \varphi) = |z| e^{i\varphi}, \quad \varphi = \arg(z) \quad \text{with } z \neq 0$$

This notation is called the **exponential form** of complex numbers.

- This form is useful since we have:

$$e^{i\varphi_1} \cdot e^{i\varphi_2} = e^{i(\varphi_1 + \varphi_2)}$$

that is, it is very easy to compute products of complex numbers.

De Moivre's formulas become:

$$(e^{i\varphi})^n = e^{in\varphi}$$

The exponential notation is widely used in applications because it greatly simplifies manipulations involving trigonometric quantities.

We have:

$$e^{2k\pi i} = 1 \quad \text{and} \quad e^{i(\varphi + 2k\pi)} = e^{i\varphi} \quad \forall k \in \mathbb{Z}, \quad \text{moreover} \quad |e^{i\varphi}| = 1$$

We also have:

$$e^{i\pi} = -1, \quad e^{i\frac{\pi}{2}} = i, \quad e^{i\frac{3\pi}{2}} = -i, \quad e^{i\frac{3\pi}{4}} = \frac{-1+i}{\sqrt{2}}$$

Rewriting, we obtain **Euler's formula**:

$$e^{i\pi} + 1 = 0$$

which links in a simple way five of the most important constants: 0, 1, i , π , e .

The exponential notation can be generalized consistently with the properties of powers, by defining the **complex exponential**:

$$e^z = e^{x+iy} = e^x e^{iy} = e^x (\cos y + i \sin y), \quad \forall z = x + iy \in \mathbb{C}$$

- From the definition of exponential form it follows that

$$e^{-i\varphi} = \cos(-\varphi) + i \sin(-\varphi) = \cos \varphi - i \sin \varphi$$

and we easily obtain:

$$\cos \varphi = \frac{e^{i\varphi} + e^{-i\varphi}}{2}, \quad \sin \varphi = \frac{e^{i\varphi} - e^{-i\varphi}}{2i} \quad \text{with } \varphi \in \mathbb{R}$$

These formulas allow us to express sine and cosine as combinations of complex exponentials.

12.4 Quadratic equations

- A **second-degree equation** or **quadratic equation** in one **real** unknown x is an algebraic equation in which the highest power of the unknown is 2, and it can always be written in the form:

$$a x^2 + b x + c = 0 \quad (a \neq 0) \quad (12.22)$$

where a, b and c are real numbers.

The solutions are also called **roots** or **zeros** of the equation.

Observation 12.10: quadratic formula

Given a quadratic equation $a x^2 + b x + c = 0$ ($a \neq 0$), its zeros or roots are:

$$x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a} \quad (12.23)$$

- A quadratic equation is called a “complete quadratic equation” when all its coefficients are different from 0. It is solved with the so-called **method of completing the square**, so called because the equation is modified until its left-hand side becomes the square of a binomial.

Proof. Isolating the constant term, we obtain:

$$a x^2 + b x = -c$$

$$\underbrace{4 a^2 x^2}_{=(2 a x)^2} + \underbrace{4 a b x}_{=2 (2 a x) b} = -4 a c$$

$$(2 a x)^2 + 2 (2 a x) b + b^2 = b^2 - 4 a c$$

$$(2 a x + b)^2 = \underbrace{b^2 - 4 a c}_{:=\Delta \text{ (discriminant)}}$$

$$2 a x + b = \pm \sqrt{b^2 - 4 a c}$$

$$x = \frac{-b \pm \sqrt{b^2 - 4 a c}}{2 a}$$

□

- If the discriminant Δ is negative there are no real solutions.
- If $\Delta = 0$, the quadratic formula becomes:

$$x = -\frac{b}{2a}$$

therefore there is only one root, with *multiplicity two*.

- Finally, if $\Delta < 0$, the equation has no real solutions. In particular, there are always two solutions, but they belong to the field of complex numbers: they are **two complex conjugate numbers** and are computed with the two formulas:

$$x_+ = \frac{-b}{2a} + i \left(\frac{\sqrt{4ac - b^2}}{2a} \right)$$

$$x_- = \frac{-b}{2a} - i \left(\frac{\sqrt{4ac - b^2}}{2a} \right)$$

where i is the **imaginary unit** ($i^2 = -1$).

A quadratic equation in one **complex** unknown z has the following form

$$a z^2 + b z + c = 0 \quad (a \neq 0) \quad (12.24)$$

where a, b and c are complex numbers. It is solved with the same formula as in the case of a real unknown:

$$z = \frac{-b \pm \sqrt{b^2 - 4 a c}}{2 a} \quad (12.25)$$

where the square root is understood in the complex sense (the $\pm$ sign is superfluous, because in the complex field the root denotes two numbers, each the opposite of the other)

Example 12.12: quadratic equations in the complex field

- We want to solve:

$$z^2 + 2i z - \sqrt{3}i = 0 \quad a = 1, b = 2i, c = -\sqrt{3}i$$

we have (switching to the trigonometric form to compute the root):

$$\begin{aligned} z &= \frac{-2i \pm \sqrt{2^2 i^2 + 2^2 \sqrt{3}i}}{2} = \frac{-2i \pm 2\sqrt{-1 + \sqrt{3}i}}{2} = -i \pm \sqrt{-1 + \sqrt{3}i} \\ &= -i \pm \sqrt{2 \left(-\frac{1}{2} + i \frac{\sqrt{3}}{2} \right)} = -i \pm \sqrt{2 \left(\cos \frac{2}{3}\pi + i \sin \frac{2}{3}\pi \right)} \\ &= -i \pm \sqrt{2} \left(\underbrace{\cos \frac{\pi}{3}}_{=\frac{1}{2}} + i \underbrace{\sin \frac{\pi}{3}}_{=\frac{\sqrt{3}}{2}} \right) = \pm \frac{\sqrt{2}}{2} + i \left(-1 \pm \frac{\sqrt{6}}{2} \right) \end{aligned}$$

- The previous theorem tells us that a polynomial of the form $z^n + a$ (with a complex) has exactly n roots in $\mathbb{C}$; in the real field, instead, the equation $x^n + a = 0$ may have two, one, or no roots (examples: $x^2 - 1 = 0, x^3 - 1 = 0, x^2 + 1 = 0$); we now know that such an equation always has n roots in $\mathbb{C}$, but only occasionally one or two of them lie in $\mathbb{R}$.
- The result is much more general, as stated by the following theorem, whose proof we do not report.

Theorem 12.11: fundamental theorem of algebra


A polynomial equation of the form

$$a_0 + a_1 z + \cdots + a_n z^n = 0 \quad (a_n \neq 0)$$

with arbitrary complex coefficients has exactly n roots in $\mathbb{C}$, if each of them is counted with its multiplicity ^a.

^aIf $P(z)$ is a polynomial in z of degree n and z_0 is one of its roots, we say that z_0 has multiplicity k (k integer, ≥ 1) if the formula $P(z) = (z - z_0)^k Q(z)$ holds, where Q is a polynomial such that $Q(z_0) \neq 0$.

Cosines and sines of the main angles



Exercises

The sheets of worked exercises on the topics of this volume, one per section, with the solution right after each statement. The sheets are by Fabio Furini and Gianluca Priori; many exercises are taken from M. Bramanti, *Esercitazioni di Analisi matematica 1*, Esculapio.

13.1 Maxima, minima, suprema and infima

Exercise 13.1

Determine whether the set

$$A = [0, \sqrt{2}] \cap \mathbb{Q}$$

has a maximum, a minimum, a supremum, an infimum and, if so, determine these elements.

Solution

The minimum of A is 0. A has no maximum; its supremum is $\sqrt{2}$.

Exercise 13.2

Determine whether the set

$$A = [\sqrt{2}, \sqrt{3}] \cap \mathbb{Q}$$

has a maximum, a minimum, a supremum, an infimum and, if so, determine these elements.

Solution

A has neither a minimum nor a maximum. The infimum is $\sqrt{2}$, the supremum is $\sqrt{3}$.

Exercise 13.3

Determine whether the set

$$A = [0, \sqrt{2}] \cap (\mathbb{R} - \mathbb{Q})$$

has a maximum, a minimum, a supremum, an infimum and, if so, determine these elements.

Solution

A has no minimum; the infimum is 0. The maximum of A is $\sqrt{2}$.

Exercise 13.4

In each of the following cases, say whether the set

$$A = (-\infty, \sqrt{2}] \cap \mathbb{Q}$$

has a maximum, a minimum, a supremum, an infimum and, if so, determine these elements.

Solution

A is not bounded below, hence it has no infimum in $\mathbb{R}$. A has no maximum; the supremum is $\sqrt{2}$.

Exercise 13.5

Say whether the following set $A \subset \mathbb{R}$ has a maximum, a minimum, a supremum, an infimum, and whether it is bounded:

$$A = \left\{ \frac{2n+1}{n} : n \in \mathbb{N} \setminus \{0\} \right\}$$

Solution

From $(2n+1)/n = 2 + (1/n)$ we have that the maximum of A is attained for $n = 1$ and is equal to 3. A has no minimum; the infimum is 2. In particular, A is bounded.

Exercise 13.6

Say whether the following set $A \subset \mathbb{R}$ has a maximum, a minimum, a supremum, an infimum, and whether it is bounded:

$$A = \left\{ \frac{n-1}{n} : n \in \mathbb{N} \setminus \{0\} \right\}$$

Solution

From $(n-1)/n = 1 - (1/n)$ we have that the minimum of A is attained for $n = 1$ and is equal to 0. A has no maximum; the supremum is 1. In particular, A is bounded.

Exercise 13.7

Say whether the following set $A \subset \mathbb{R}$ has a maximum, a minimum, a supremum, an infimum, and whether it is bounded:

$$A = \left\{ \frac{3n^2 + 1}{n^2} : n \in \mathbb{N} \setminus \{0\} \right\}$$

Solution

From $(3n^2 + 1)/n^2 = 3 + (1/n^2)$ we have that the maximum of A is attained for $n = 1$ and is equal to 4. A has no minimum; the infimum is 3. In particular, A is bounded.

Exercise 13.8

Say whether the following set $A \subset \mathbb{R}$ has a maximum, a minimum, a supremum, an infimum, and whether it is bounded:

$$A = \left\{ \frac{1}{n^2 + 1} : n \in \mathbb{N} \setminus \{0\} \right\}$$

Solution

The maximum value of A is attained for $n = 1$ and is equal to $1/2$. A has no minimum; the infimum is 0. In particular, A is bounded.

Exercise 13.9

Let

$$A = \left\{ \left| \frac{3x}{x+1} \right| : -\frac{1}{2} < x \leq 2 \right\}.$$

Choose the correct statements among the following:

- (a) $\sup A \notin A$ (b) $\inf A = \min A = 2$
 (c) $\max A = 3$ (d) none of the other answers is correct

Solution

A is the set of values of $|f(x)|$ on the interval $(-1/2, 2]$, with $f(x) = \frac{3x}{x+1}$. From


$$\frac{3x}{x+1} = 3 - \frac{3}{x+1}$$

we have that the function f is strictly increasing on the given interval $(-1/2, 2]$, negative on $(-1/2, 0)$, zero for $x = 0$, and positive on $(0, 2]$. It follows that

$$|f(x)| = \left| \frac{3x}{x+1} \right|$$

is strictly decreasing on $(-1/2, 0]$, where it takes all the values in $[0, 3)$, and strictly increasing on $[0, 2]$, where it takes all the values in $[0, 2]$. The minimum value is therefore attained

for $x = 0$ and is equal to 0. The supremum is 3; there is no maximum value. The correct answer is (a).



Exercise 13.10

For $I = [1/3, +\infty)$ consider the function

$$f : I \rightarrow \mathbb{R}, \quad f(x) = \exp\left(\left|\frac{2x-1}{x}\right|\right).$$

Determine, among the following intervals, the set $J = f(I)$ of the values taken by f .

- | | | |
|--------------------|-------------------|----------------------|
| (a) $[e, e^2)$ | (b) $(e^{-2}, e]$ | (c) $[1, e^2)$ |
| (d) $(e, +\infty)$ | (e) $(0, e^2)$ | (f) another interval |


Solution

The function f has the same monotonicity behavior as

$$|g(x)| = \left|\frac{2x-1}{x}\right|$$

with

$$g(x) = \frac{2x-1}{x} = 2 - \frac{1}{x}.$$




Solution

The function g is strictly increasing on the interval $I = [1/3, +\infty)$, negative on $[1/3, 1/2)$, zero for $x = 1/2$, and positive on $(1/2, +\infty)$. It follows that $|g(x)|$ is strictly decreasing on $[1/3, 1/2]$, where it takes all the values in $[0, 1]$, and strictly increasing on $[1/2, +\infty)$, where it takes all the values in $[0, 2)$. The set of values $|g|(I)$ is $[0, 2)$, hence for $f(x) = e^{|g(x)|}$ we have

$$f(I) = [1, e^2).$$

The correct answer is (c).



More precisely, f is strictly decreasing on $[1/3, 1/2]$, where it takes all the values in $[1, e]$, and strictly increasing on $[1/2, +\infty)$, where it takes all the values in $[1, e^2]$.

Exercise 13.11

Let $A = A_+ \cup A_-$ with

$$A_+ = \left\{ x + \frac{2}{x} : x > 0 \right\}, \quad A_- = \left\{ x + \frac{2}{x} : x < 0 \right\}.$$

Choose the correct statements among the following:

- (a) $\sup A = +\infty$ (b) $\inf A_+ = 2\sqrt{2}$ (c) A_+ has no minimum
- (d) A_- has no maximum (e) none of the other answers is correct

Solution

Clearly, A_+ is not bounded above, hence statement (a) is correct (and statement (e) is false). For $x \neq 0$, we consider the equation


$$x + \frac{2}{x} = y,$$

$$x^2 - yx + 2 = 0.$$

There are solutions $x \in \mathbb{R}$ if and only if $y \in (-\infty, -2\sqrt{2}] \cup [2\sqrt{2}, +\infty)$. Moreover, for every $y \in (-\infty, -2\sqrt{2}]$ the solutions x are negative, while for every $y \in [2\sqrt{2}, +\infty)$ the solutions x are positive. This means

$$A_+ = [2\sqrt{2}, +\infty), \quad A_- = (-\infty, -2\sqrt{2}].$$

Therefore (b) is true, (c) is false, (d) is false.



13.2 Equations and inequalities in one real variable

Exercise 13.12

Solve the following inequality in the real variable x :

$$\frac{x^2 - 2x}{x^2 - 4x + 3} > 0$$

Solution

The numerator $x^2 - 2x$ is positive for $x < 0$ or for $x > 2$, zero for $x = 0$, $x = 2$, and negative for $0 < x < 2$. The denominator $x^2 - 4x + 3$ is positive for $x < 1$ or for $x > 3$, and negative for $1 < x < 3$. The quotient is defined and has the required positive sign for

$$x \in (-\infty, 0) \cup (1, 2) \cup (3, +\infty).$$

Exercise 13.13

Solve the following inequality in the real variable x :

$$\log(x-1)^2 - \log(x-2) > 0$$

Solution

The function $f(x) = \log(x-1)^2 - \log(x-2)$ is defined for $x > 2$, since for such x , and only for such x , we have $x-2 > 0$, $(x-1)^2 > 0$. In its domain, we have, as required, $f(x) > 0$ if and only if

$$\log(x-1)^2 > \log(x-2), \quad x > 2$$

hence if and only if

$$\begin{cases} (x-1)^2 > x-2 \\ x > 2 \end{cases}$$

We obtain

$$\begin{cases} x^2 - 3x + 3 > 0 \\ x > 2 \end{cases}$$

which is equivalent to

$$x > 2$$

since $x^2 - 3x + 3 > 0$ is satisfied for every x . On its whole domain $(2, +\infty)$ the function $f(x)$ takes positive values.

Exercise 13.14

Solve the following inequality in the real variable x :

$$e^x + e^{-x} < \frac{10}{3}$$

Solution

For every $x \in \mathbb{R}$

$$e^x + e^{-x} < \frac{10}{3}$$

is equivalent to

$$3e^{2x} - 10e^x + 3 < 0$$

hence to

$$\frac{1}{3} < e^x < 3.$$

It follows that the inequality is solved by

$$-\log 3 < x < \log 3.$$

Exercise 13.15

Solve the following inequality in the real variable x :

$$8^{x+1} \geq 2^{x^2}$$

Solution

For every $x \in \mathbb{R}$

$$8^{x+1} \geq 2^{x^2}$$

is equivalent to

$$2^{3x+3} \geq 2^{x^2}$$

hence to

$$3x + 3 \geq x^2$$

since 2^x is a strictly increasing function on $\mathbb{R}$. It follows that the inequality is solved by

$$\frac{3 - \sqrt{21}}{2} \leq x \leq \frac{3 + \sqrt{21}}{2}.$$

Exercise 13.16

Solve the following inequality in the real variable x :

$$\log(1 - \sin x) \geq 0$$

Solution

The function $f(x) = \log(1 - \sin x)$ is defined for $\sin x < 1$, hence for

$$x \neq \frac{\pi}{2} + 2k\pi, \quad k \in \mathbf{Z}.$$

For such x we have $f(x) \geq 0$ if and only if

$$1 - \sin x \geq 1$$

hence for

$$\sin x \leq 0.$$

The given inequality is solved for

$$\pi + 2k\pi \leq x \leq 2\pi + 2k\pi, \quad k \in \mathbf{Z}$$

i.e., for

$$x \in \bigcup_{k \in \mathbf{Z}} [\pi + 2k\pi, 2\pi + 2k\pi].$$

Exercise 13.17

Solve the following inequality in the real variable x :

$$|x|\sqrt{1 - 2x^2} > 2x^2 - 1$$

Solution

$\sqrt{1 - 2x^2}$ is defined for $1 - 2x^2 \geq 0$. When $1 - 2x^2 = 0$ the inequality is not satisfied, since it reduces to $0 > 0$. When $1 - 2x^2 > 0$ the inequality is satisfied, since the left-hand side $|x|\sqrt{1 - 2x^2}$ is the product of two positive numbers while the right-hand side $2x^2 - 1 = -(1 - 2x^2)$ is negative. Therefore the inequality is solved for

$$-\sqrt{\frac{1}{2}} < x < \sqrt{\frac{1}{2}}.$$

Exercise 13.18

Solve the following equation in the real variable x :

$$\sin^2 x = 2 \cos^2 x - \frac{1}{2}$$

Solution

The equation

$$\sin^2 x = 2 \cos^2 x - \frac{1}{2}$$

is equivalent to

$$2 \sin^2 x = 4(1 - \sin^2 x) - 1$$

$$6 \sin^2 x = 3$$

$$\sin x = \pm \sqrt{\frac{1}{2}}.$$

The solutions are given by

$$x = \frac{\pi}{4} + \frac{k\pi}{2}, \quad k \in \mathbf{Z}.$$

Exercise 13.19

Solve the following inequality in the real variable x :

$$e^{\sin^2 x - \sin x} \leq 1$$

Solution

For every $x \in \mathbb{R}$, $e^{\sin^2 x - \sin x} \leq 1$ is equivalent to

$$\sin^2 x - \sin x \leq 0$$

hence to

$$0 \leq \sin x \leq 1.$$

The solutions are given by

$$2k\pi \leq x \leq \pi + 2k\pi, \quad k \in \mathbf{Z}$$

i.e., by

$$x \in \bigcup_{k \in \mathbf{Z}} [2k\pi, \pi + 2k\pi].$$

Exercise 13.20

Solve the following inequality in the real variable x :

$$2|x^2 - x| > |x|$$

Solution

The inequality $2|x^2 - x| > |x|$ is not satisfied for $x = 0$. For $x \neq 0$, dividing by the positive term $|x|$, it is equivalent to

$$2|x - 1| > 1$$

hence to

$$|x - 1| > 1/2.$$

The solutions are given by

$$(x - 1 < -1/2, x \neq 0) \vee (x - 1 > 1/2),$$

$$(x < 1/2, x \neq 0) \vee x > 3/2$$

hence by

$$x \in (-\infty, 0) \cup (0, 1/2) \cup (3/2, +\infty).$$

Exercise 13.21

Solve the following inequality in the real variable x :

$$(x + 1)^{x^2 - 1} > 1$$

Solution

The function $f(x) = (x + 1)^{x^2 - 1}$ is defined for $x > -1$. For such x , $f(x) > 1$ is equivalent to

$$e^{(x^2 - 1)\log(x + 1)} > 1$$

hence to

$$(x^2 - 1)\log(x + 1) > 0, \quad x > -1.$$

Dividing by the positive term $x + 1$ we obtain

$$(x - 1)\log(x + 1) > 0, \quad x > -1.$$

The factor $x - 1$ is positive for $x > 1$, zero for $x = 1$, and negative for $-1 < x < 1$. The factor $\log(x + 1)$ is positive for $x > 0$, zero for $x = 0$, and negative for $-1 < x < 0$. By the rule of signs, the solutions are given by

$$x \in (-1, 0) \cup (1, +\infty).$$

Exercise 13.22

Solve the following equation in the real variable x :

$$3^{|x^2 - 4|} = 0$$

Solution

$3^{|x^2 - 4|} = 0$ has no solutions, since $3^y > 0$ for every $y \in \mathbb{R}$.

Exercise 13.23

Solve the following inequality in the real variable x :

$$\frac{\log_a(4x-3)}{\log_a(2x-1)} > 1 \quad 0 < a < 1$$

Solution

Because of the existence conditions of the logarithm and of the denominator, we must impose the system

$$\begin{cases} 4x-3 > 0 \\ 2x-1 > 0 \\ \log_a(2x-1) \neq 0 \end{cases}$$

which gives $\frac{3}{4} < x < 1, x > 1$. Now moving everything to the left-hand side, finding the least common denominator and using the properties of logarithms, we obtain

$$\frac{\log_a(\frac{4x-3}{2x-1})}{\log_a(2x-1)} > 0$$

Studying the sign of the numerator and of the denominator, and keeping in mind that $a < 1$, for the numerator we obtain

$$\log_a\left(\frac{4x-3}{2x-1}\right) > 0 \iff \frac{4x-3}{2x-1} < 1 \iff \frac{1}{2} < x < 1$$

while for the denominator we obtain

$$\log_a(2x-1) > 0 \iff x < 1$$

Combining the signs, the inequality is satisfied for $\frac{1}{2} < x < 1, x > 1$. Intersecting this set of solutions with the existence conditions studied above, we obtain that the inequality is satisfied for every x belonging to the set $(\frac{3}{4}; 1) \cup (1, +\infty)$.

Exercise 13.24

Solve the following inequality in the real variable x :

$$\sqrt{x^2 - 3x + 5} \leq x + 3$$

Solution

This irrational inequality has the form $\sqrt{A(x)} \leq B(x)$, with $A(x)$ and $B(x)$ polynomials in x . First of all, we must impose the existence condition of the root, i.e., $x^2 - 3x + 5 \geq 0$. Moreover, since a square root is always positive or zero in $\mathbb{R}$, we impose $x + 3 \geq 0$. We therefore solve the system:

$$\begin{cases} x^2 - 3x + 5 \geq 0 \\ x + 3 \geq 0 \\ x^2 - 3x + 5 \leq (x + 3)^2 \end{cases}$$

where, in the last inequality, we were allowed to square both sides, since they are positive. The system reduces to:

$$\begin{cases} \forall x \in \mathbb{R} \\ x \geq -3 \\ x \geq -\frac{4}{9} \end{cases}$$

whose set of solutions is the interval $[-\frac{4}{9}, +\infty)$.

Exercise 13.25

Solve the following rational inequality in the real variable x :

$$\frac{x^3 - 3x^2 + 2x - 6}{x^2 - 2x - 3} > \frac{2}{x + 1}$$

Solution

First of all, we must write the inequality in canonical form (0 on the right-hand side). The existence conditions (denominators different from zero) will be automatically included in the sign study. We obtain

$$\frac{x^3 - 3x^2 + 2x - 6}{x^2 - 2x - 3} - \frac{2}{x + 1} > 0$$

We factor the numerator and the denominator of the first fraction, obtaining:

$$\frac{(x^2 + 2)(x - 3)}{(x - 3)(x + 1)} - \frac{2}{x + 1} > 0$$

Since we can cancel $x - 3$ and obtain an equivalent fraction, provided that $x \neq 3$ by the existence condition, we have

$$\frac{x^2 + 2}{x + 1} - \frac{2}{x + 1} > 0$$

that is,

$$\frac{x^2}{x + 1} > 0$$

The numerator is positive $\forall x \in \mathbb{R}, x \neq 0$, and the denominator is positive for $x > -1$. Combining the signs, and taking into account the condition $x \neq 3$, we obtain the solution set $(-1, 0) \cup (0, 3) \cup (3, +\infty)$.

13.3 Summations

Exercise 13.26

Prove, using the properties of summations, that for every integer $n \geq 1$ we have:

$$\sum_{k=1}^n k^2 = \frac{n(n+1)(2n+1)}{6}$$

(sum of the squares of the first n natural numbers, excluding zero).

Solution

We start by writing the sum $\sum_{k=0}^n (k+1)^3$ in two different ways:

$$\begin{aligned} 1) \quad \sum_{k=0}^n (k+1)^3 &= \sum_{k=0}^n (k^3 + 3k^2 + 3k + 1) \\ &= \left(\sum_{k=1}^n k^3 + 3 \sum_{k=1}^n k^2 + 3 \sum_{k=1}^n k + \sum_{k=1}^n 1 \right) + 1 \\ &= \sum_{k=1}^n k^3 + 3 \sum_{k=1}^n k^2 + \frac{3n(n+1)}{2} + n + 1 \\ 2) \quad \sum_{k=0}^n (k+1)^3 &= \sum_{k=1}^{n+1} k^3 = \sum_{k=1}^n k^3 + (n+1)^3 \end{aligned}$$

Equating the two expressions and canceling the sum with k^3 , we obtain:

$$3 \sum_{k=1}^n k^2 + \frac{3n(n+1)}{2} + n + 1 = (n+1)^3$$

Isolating the sum with k^2 , we obtain:

$$\begin{aligned} 3 \sum_{k=1}^n k^2 &= (n+1)^3 - \frac{3n(n+1)}{2} - n - 1 \\ &= n^3 + 3n^2 + 3n + 1 - \frac{3n^2 + 3n}{2} - n - 1 \\ &= \frac{2n^3 + 3n^2 + n}{2} \end{aligned}$$

Hence:

$$\sum_{k=1}^n k^2 = \frac{2n^3 + 3n^2 + n}{6} = \frac{n(n+1)(2n+1)}{6}$$

Exercise 13.27

Prove, using the properties of summations, that for every integer $n \geq 1$ we have:

$$\sum_{k=1}^n k^3 = \frac{n^2 (n+1)^2}{4}$$

(sum of the cubes of the first n natural numbers, excluding zero). That is, we have:

$$\sum_{k=1}^n k^3 = \frac{n^2 (n+1)^2}{4} = \left(\frac{n(n+1)}{2} \right)^2 = \left(\sum_{k=1}^n k \right)^2$$

and hence the sum of the cubes is equal to the square of the sum of the natural numbers.

Solution

We start by writing the following sum in two different ways: $\sum_{k=0}^n (k+1)^4$.

$$\begin{aligned} 1) \quad \sum_{k=0}^n (k+1)^4 &= \sum_{k=0}^n (k^4 + 4k^3 + 6k^2 + 4k + 1) \\ &= \left(\sum_{k=1}^n k^4 + 4 \sum_{k=1}^n k^3 + 6 \sum_{k=1}^n k^2 + 4 \sum_{k=1}^n k + \sum_{k=1}^n 1 \right) + 1 \\ &= \sum_{k=1}^n k^4 + 4 \sum_{k=1}^n k^3 + n(n+1)(2n+1) + 2n(n+1) + n + 1 \\ &= \sum_{k=1}^n k^4 + 4 \sum_{k=1}^n k^3 + 2n^3 + 5n^2 + 4n + 1 \\ 2) \quad \sum_{k=0}^n (k+1)^4 &= \sum_{k=1}^{n+1} k^4 = \sum_{k=1}^n k^4 + (n+1)^4 \end{aligned}$$

Equating the two expressions and canceling the sum with k^4 , we obtain:

$$4 \sum_{k=1}^n k^3 + 2n^3 + 5n^2 + 4n + 1 = (n+1)^4$$

Isolating the sum with k^3 , we obtain:

$$4 \sum_{k=1}^n k^3 = n^4 + 4n^3 + 6n^2 + 4n + 1 - 2n^3 - 5n^2 - 4n - 1 = n^4 + 2n^3 + n^2$$

Hence:

$$\sum_{k=1}^n k^3 = \frac{n^4 + 2n^3 + n^2}{4} = \frac{n^2 (n+1)^2}{4}$$

13.4 Principle of mathematical induction

Exercise 13.28

Prove by induction that for every integer $n \geq 1$ the following holds:

$$\sum_{k=1}^n k^2 = \frac{n(n+1)(2n+1)}{6}$$

(sum of the squares of the first n natural numbers)

Solution

By induction on n .

- **Base case**

Let $n = 1$. Then the statement becomes:

$$\sum_{k=1}^1 k^2 = \frac{1(1+1)(2 \cdot 1 + 1)}{6} \quad \text{i.e.} \quad 1 = 1$$

which is clearly true.

- **Inductive step**

Assume that it is true for n , and let us prove it for $(n+1)$. By the inductive hypothesis, we have:

$$\sum_{k=1}^n k^2 = \frac{n(n+1)(2n+1)}{6}.$$

Hence we can write:

$$\begin{aligned} \sum_{k=1}^{n+1} k^2 &= \sum_{k=1}^n k^2 + (n+1)^2 \\ &= \frac{n(n+1)(2n+1)}{6} + (n+1)^2 = \frac{n(n+1)(2n+1) + 6(n+1)^2}{6} \\ &= \frac{(n+1)(n(2n+1) + 6(n+1))}{6} = \frac{(n+1)(2n^2 + n + 6n + 6)}{6} \\ &= \frac{(n+1)(2n^2 + 4n + 3n + 6)}{6} = \frac{(n+1)((2n)(n+2) + 3(n+2))}{6} \\ &= \frac{(n+1)(n+2)(2n+3)}{6} = \frac{(n+1)(n+1+1)(2(n+1)+1)}{6} \end{aligned}$$

which is exactly the desired statement for $n+1$.

Exercise 13.29

Prove by induction that for every integer $n \geq 1$ the following holds:

$$\sum_{k=0}^{n-1} (2k+1) = n^2 \quad \text{or} \quad \sum_{k=1}^n (2k-1) = n^2$$

(sum of the first n odd numbers)

Solution

By induction on n .

- **Base case**

Let $n = 1$. Then the statement becomes:

$$\sum_{k=0}^0 (2k+1) = 1^2 \quad \text{i.e.} \quad 1 = 1$$

which is clearly true.

- **Inductive step**

Assume that it is true for n , and let us prove it for $(n+1)$. By the inductive hypothesis, we have:

$$\sum_{k=0}^{n-1} (2k+1) = n^2$$

Hence we can write:

$$\begin{aligned} \sum_{k=0}^{n-1+1} (2k+1) &= \sum_{k=0}^{n-1} (2k+1) + 2n+1 \\ &= n^2 + 2n+1 \\ &= (n+1)^2 \end{aligned}$$

which is exactly the desired statement for $n+1$.

Exercise 13.30

Prove by induction that for every integer $n \geq 1$ the following holds:

$$\sum_{k=1}^n 2k = n(n+1)$$

(sum of the first n even numbers)

Solution

By induction on n .

- **Base case**

Let $n = 1$. Then the statement becomes:

$$\sum_{k=1}^1 2k = 1 \cdot 2 \quad \text{i.e.} \quad 2 = 2$$

which is clearly true.

- **Inductive step**

Assume that it is true for n , and let us prove it for $(n+1)$. By the inductive hypothesis, we have:

$$\sum_{k=1}^n 2k = n(n+1)$$

Hence we can write:

$$\begin{aligned} \sum_{k=1}^{n+1} 2k &= \sum_{k=1}^n 2k + 2(n+1) \\ &= 2 \left(\sum_{k=1}^n k + (n+1) \right) \\ &= 2 \left(\frac{n(n+1)}{2} + (n+1) \right) \\ &= 2 \left(\frac{n(n+1) + 2(n+1)}{2} \right) \\ &= (n+1)(n+2) = (n+1)(n+1+1) \end{aligned}$$

which is exactly the desired statement for $n+1$.

Exercise 13.31

Prove by induction that for every integer $n \geq 1$ the following holds:

$$\sum_{k=1}^n \frac{k}{2^k} = 2 - \frac{n+2}{2^n}$$

Solution

By induction on n .

- **Base case**

Let $n = 1$. Then the statement becomes:

$$\sum_{k=1}^1 \frac{k}{2^k} = 2 - \frac{1+2}{2^1} \quad \text{i.e.} \quad \frac{1}{2} = \frac{1}{2}$$

which is clearly true.

- **Inductive step**

Assume that it is true for n , and let us prove it for $(n+1)$. By the inductive hypothesis, we have:

$$\sum_{k=1}^n \frac{k}{2^k} = 2 - \frac{n+2}{2^n}$$

Hence we can write:

$$\begin{aligned} \sum_{k=1}^{n+1} \frac{k}{2^k} &= \sum_{k=1}^n \frac{k}{2^k} + \frac{n+1}{2^{n+1}} \\ &= 2 - \frac{n+2}{2^n} + \frac{n+1}{2^{n+1}} \\ &= 2 - \frac{2(n+2)}{2^{n+1}} + \frac{n+1}{2^{n+1}} \\ &= 2 - \left(\frac{2(n+2) - n - 1}{2^{n+1}} \right) \\ &= 2 - \left(\frac{n+3}{2^{n+1}} \right) \\ &= 2 - \frac{(n+1)+2}{2^{n+1}} \end{aligned}$$

which is exactly the desired statement for $n+1$.

Exercise 13.32

Given a set X with n elements, prove by induction that the power set $\mathcal{P}(X)$ has 2^n elements, that is,

$$|\mathcal{P}(X)| = 2^n \quad \text{for } n \geq 0$$

Solution

By induction on n .

- **Base case**

Let $n = 0$. A set with 0 elements is the empty set $(\emptyset)$. The power set of the empty set contains only the empty set as an element. Then the statement becomes:

$$|\mathcal{P}(\emptyset)| = 2^0 \quad \text{i.e.} \quad 1 = 1$$

which is clearly true.

• **Inductive step**

Assume the statement is true for a set of n elements, and let us prove it for a set of $(n + 1)$ elements. By the inductive hypothesis, the power set of a set of n elements has 2^n elements.

A set Y of $(n + 1)$ elements can be written as:

$$Y = \{a_1, a_2, \dots, a_n, a_{n+1}\}$$

hence:

$$\mathcal{P}(Y) = \{S : S \subseteq Y\} = \{S : S \subseteq \{a_1, a_2, \dots, a_n, a_{n+1}\}\}$$

The subsets of Y either contain the element a_{n+1} or do not contain it. We therefore define the set Z_a formed by the first n elements of Y :

$$Z_a = \{a_1, a_2, \dots, a_n\} \quad \text{hence} \quad |\mathcal{P}(Z_a)| = 2^n$$

We now define the following set Z_b :

$$Z_b = \{S \cup \{a_{n+1}\} : S \subseteq \underbrace{\{a_1, a_2, \dots, a_n\}}_{=Z_a}\} = \{S \cup \{a_{n+1}\} : S \subseteq Z_a\} \quad \text{and} \quad |Z_b| = 2^n$$

since Z_b has one element for each element of $\mathcal{P}(Z_a)$.

In other words, $\mathcal{P}(Z_a)$ contains all the subsets of Y in which the element a_{n+1} is not present, while Z_b contains all the subsets of Y in which the element a_{n+1} is present.

We also note that the sets $\mathcal{P}(Z_a)$ and Z_b are disjoint, that is: $\mathcal{P}(Z_a) \cap Z_b = \emptyset$. Hence $|\mathcal{P}(Z_a) \cup Z_b| = |\mathcal{P}(Z_a)| + |Z_b|$. Finally, we have:

$$|\mathcal{P}(Y)| = |\mathcal{P}(Z_a) \cup Z_b| = 2^n + 2^n = 2^{n+1}$$

which is exactly the desired statement for $n + 1$.

Exercise 13.33

Prove by induction that for every integer $n \geq 1$ the following holds:

$$\sum_{k=1}^n k^3 = \frac{n^4 + 2n^3 + n^2}{4} = \frac{n^2(n+1)^2}{4}$$

(sum of the cubes of the first n natural numbers). Then compute the sum of the cubes of the first 10, 100 and 1000 natural numbers (without zero) and prove that the sum of the cubes of the first n natural numbers equals the square of the sum of the first n natural numbers:

$$\sum_{k=1}^n k^3 = \left(\sum_{k=1}^n k \right)^2$$

Solution

By induction on n .

- **Base case**

Let $n = 1$. Then the statement becomes:

$$\sum_{k=1}^1 k^3 = \frac{1+2+1}{4} \quad \text{i.e.} \quad 1 = 1$$

which is clearly true.

- **Passo induttivo**

Suppose it is true for n , and let us prove it for $(n+1)$. By the inductive hypothesis, we have:

$$\sum_{k=1}^n k^3 = \frac{n^4 + 2n^3 + n^2}{4}.$$

Hence we can write:

$$\begin{aligned} \sum_{k=1}^{n+1} k^3 &= \sum_{k=1}^n k^3 + (n+1)^3 = \frac{n^4 + 2n^3 + n^2}{4} + (n+1)^3 \\ &= \frac{n^4 + 2n^3 + n^2 + 4(n+1)^3}{4} = \frac{n^4 + 2n^3 + n^2 + 4n^3 + 12n^2 + 12n + 4}{4} \\ &= \frac{n^4 + 6n^3 + 13n^2 + 12n + 4}{4} \\ &= \frac{(n^4 + 4n^3 + 6n^2 + 4n + 1) + 2(n^3 + 3n^2 + 3n + 1) + (n^2 + 2n + 1)}{4} \\ &= \frac{(n+1)^4 + 2(n+1)^3 + (n+1)^2}{4} \end{aligned}$$

which is exactly the desired statement for $n+1$.

The sum of the cubes of the first 10, 100 and 1000 natural numbers (without zero) is:

$$\begin{aligned} \sum_{k=1}^{10} k^3 &= \frac{10^4 + 2 \cdot 10^3 + 10^2}{4} = 3025, & \sum_{k=1}^{100} k^3 &= \frac{100^4 + 2 \cdot 100^3 + 100^2}{4} = 25,502,500 \\ \sum_{k=1}^{1000} k^3 &= \frac{1000^4 + 2 \cdot 1000^3 + 1000^2}{4} = 250,500,250,000 \end{aligned}$$

Finally, the sum of the cubes of the first n natural numbers equals the square of the sum of the first n natural numbers, since:

$$\sum_{k=1}^n k^3 = \frac{n^4 + 2n^3 + n^2}{4} = \frac{n^2(n^2 + 2n + 1)}{4} = \frac{n^2(n+1)^2}{4} = \left(\frac{n(n+1)}{2}\right)^2 = \left(\sum_{k=1}^n k\right)^2$$

Exercise 13.34

Prove by induction that for every integer $n \geq 1$ the following holds:

$$\sum_{k=1}^n \frac{1}{k(k+1)} = \frac{n}{n+1} = 1 - \frac{1}{n+1}$$

Then compute the value of the sum for $n = 10$, $n = 100$ and $n = 1000$.

Solution

By induction on n .

- **Base case**

Let $n = 1$. Then the statement becomes:

$$\sum_{k=1}^1 \frac{1}{k(k+1)} = \frac{1}{1+1} \quad \text{i.e.} \quad \frac{1}{2} = \frac{1}{2}$$

which is clearly true.

- **Passo induttivo**

Suppose it is true for n , and let us prove it for $(n+1)$. By the inductive hypothesis, we have:

$$\sum_{k=1}^n \frac{1}{k(k+1)} = \frac{n}{n+1}.$$

Hence we can write:

$$\begin{aligned} \sum_{k=1}^{n+1} \frac{1}{k(k+1)} &= \sum_{k=1}^n \frac{1}{k(k+1)} + \frac{1}{(n+1)(n+2)} = \frac{n}{n+1} + \frac{1}{(n+1)(n+2)} \\ &= \frac{n(n+2) + 1}{(n+1)(n+2)} = \frac{n^2 + 2n + 1}{(n+1)(n+2)} = \frac{(n+1)^2}{(n+1)(n+2)} = \frac{n+1}{(n+1)+1} \end{aligned}$$

which is exactly the desired statement for $n+1$.

The value of the sum for $n = 10$, $n = 100$ and $n = 1000$ is:

$$\sum_{k=1}^{10} \frac{1}{k(k+1)} = \frac{10}{11}, \quad \sum_{k=1}^{100} \frac{1}{k(k+1)} = \frac{100}{101}, \quad \sum_{k=1}^{1000} \frac{1}{k(k+1)} = \frac{1000}{1001}$$

Exercise 13.35

Prove by induction that for every integer $n \geq 1$ the following holds:

$$\sum_{k=1}^n k(k+1) = \frac{2n+3}{3} \frac{n^2+n^3}{3} = \frac{n(n+1)(n+2)}{3}$$

Then compute the value of the sum for $n = 10$, $n = 100$ and $n = 1000$.

Solution

By induction on n .

- **Base case**

Let $n = 1$. Then the statement becomes:

$$\sum_{k=1}^1 k(k+1) = \frac{2+3+1}{3} \quad \text{i.e.} \quad 2 = 2$$

which is clearly true.

- **Passo induttivo**

Suppose it is true for n , and let us prove it for $(n+1)$. By the inductive hypothesis, we have:

$$\sum_{k=1}^n k(k+1) = \frac{2n+3n^2+n^3}{3}.$$

Hence we can write:

$$\begin{aligned} & \sum_{k=1}^{n+1} k(k+1) \\ &= \sum_{k=1}^n k(k+1) + (n+1)((n+1)+1) = \frac{2n+3n^2+n^3}{3} + (n+1)(n+2) \\ &= \frac{2n+3n^2+n^3+3(n+1)(n+2)}{3} = \frac{n^3+6n^2+11n+6}{3} \\ &= \frac{2(n+1)+3(n^2+2n+1)+(n^3+3n^2+3n+1)}{3} \\ &= \frac{2(n+1)+3(n+1)^2+(n+1)^3}{3} \end{aligned}$$

which is exactly the desired statement for $n+1$.

The value of the sum for $n = 10$, $n = 100$ and $n = 1000$ is:

$$\begin{aligned} \sum_{k=1}^{10} k(k+1) &= \frac{10 \cdot 11 \cdot 12}{3} = 440, & \sum_{k=1}^{100} k(k+1) &= \frac{100 \cdot 101 \cdot 102}{3} = 343,400 \\ \sum_{k=1}^{1000} k(k+1) &= \frac{1000 \cdot 1001 \cdot 1002}{3} = 334,334,000 \end{aligned}$$

Exercise 13.36

Prove by induction that for every integer $n \geq 1$ the following holds:

$$n! \geq 2^{n-1}$$

Solution

By induction on n .

- **Base case**

Let $n = 1$. Then the statement becomes:

$$1! \geq 2^{1-1} \quad \text{i.e.} \quad 1 \geq 1$$

which is clearly true.

- **Passo induttivo**

Suppose it is true for n , and let us prove it for $(n + 1)$. By the inductive hypothesis, we have $n! \geq 2^{n-1}$. Moreover, for every $n \geq 1$:

$$(n + 1) 2^{n-1} \geq 2^n \iff n + 1 \geq \frac{2^n}{2^{n-1}} = 2 \iff n \geq 1$$

Hence we can write:

$$(n + 1)! = (n + 1) n! \geq (n + 1) 2^{n-1} \geq 2^n = 2^{(n+1)-1}$$

which is exactly the desired statement for $n + 1$.

Exercise 13.37

Prove by induction that for every integer $n \geq 1$ the following holds:

$$n^n \geq 2^{n-1} n!$$

Solution

By induction on n .

- **Base case**

Let $n = 1$. Then the statement becomes:

$$1^1 \geq 2^{1-1} \cdot 1! \quad \text{i.e.} \quad 1 \geq 1$$

which is clearly true.

- **Passo induttivo**

Suppose it is true for n , and let us prove it for $(n + 1)$. By the inductive hypothesis, we have $n^n \geq 2^{n-1} n!$. Moreover, for every $n \geq 1$:

$$(n + 1)^n \geq 2 n^n \iff \left(\frac{n + 1}{n} \right)^n \geq 2 \iff \left(1 + \frac{1}{n} \right)^n \geq 2$$

and the last inequality holds by Bernoulli's inequality, $(1+x)^m \geq 1+m x$ for every integer $m \geq 0$ and $x \geq -1$: with $x = \frac{1}{n}$ and $m = n$ we get

$$\left(1 + \frac{1}{n}\right)^n \geq 1 + n \frac{1}{n} = 2$$

Hence we can write:

$$\begin{aligned} (n+1)^{n+1} &= (n+1) (n+1)^n \geq (n+1) 2 n^n \geq (n+1) 2 \cdot 2^{n-1} n! \\ &= 2^n (n+1)! = 2^{(n+1)-1} (n+1)! \end{aligned}$$

which is exactly the desired statement for $n+1$.

Exercise 13.38

Prove by induction that for every integer $n \geq 1$ the following holds:

$$n! \leq n^n$$

Solution

By induction on n .

- **Base case**

Let $n = 1$. Then the statement becomes:

$$1! \leq 1^1 \quad \text{i.e.} \quad 1 \leq 1$$

which is clearly true.

- **Passo induttivo**

Suppose it is true for n , and let us prove it for $(n+1)$. By the inductive hypothesis, we have $n! \leq n^n$. Moreover, for every $n \geq 1$:

$$(n+1)^n \geq n^n \iff \left(\frac{n+1}{n}\right)^n \geq 1 \iff \left(1 + \frac{1}{n}\right)^n \geq 1$$

and the last inequality is clearly true, since $1 + \frac{1}{n} > 1$. Hence we can write:

$$(n+1)! = (n+1) n! \leq (n+1) n^n \leq (n+1) (n+1)^n = (n+1)^{n+1}$$

which is exactly the desired statement for $n+1$.

Exercise 13.39

Prove by induction that for every integer $n \geq 0$ the natural number $10^n - 1$ is divisible by 9, that is:

$$\forall n \in \mathbb{N}, \exists m \in \mathbb{N} : 10^n - 1 = 9 m$$

Solution

By induction on n .

- **Base case**

Let $n = 0$. Then:

$$10^0 - 1 = 0 = 9 \cdot 0$$

and, since 0 is a natural number, the statement is true.

- **Passo induttivo**

Suppose it is true for n , and let us prove it for $(n + 1)$. By the inductive hypothesis, there exists $m \in \mathbb{N}$ such that $10^n - 1 = 9m$. Hence we can write:

$$\begin{aligned} 10^{n+1} - 1 &= 10 \cdot 10^n - 1 = 10 \cdot 10^n - 10 + 9 = 10(10^n - 1) + 9 \\ &= 10(9m) + 9 = 9(10m + 1) \end{aligned}$$

and, since $10m + 1$ is a natural number, the statement is true for $n + 1$.

Exercise 13.40

Prove by induction that for every integer $n \geq 1$ the natural number $n(n + 1)$ is divisible by 2, that is:

$$\forall n \in \mathbb{N}, n \geq 1, \exists m \in \mathbb{N} : n(n + 1) = 2m$$

Solution

By induction on n .

- **Base case**

Let $n = 1$. Then:

$$1(1 + 1) = 2 = 2 \cdot 1$$

and, since 1 is a natural number, the statement is true.

- **Passo induttivo**

Suppose it is true for n , and let us prove it for $(n + 1)$. By the inductive hypothesis, there exists $m \in \mathbb{N}$ such that $n(n + 1) = 2m$. Hence we can write:

$$\begin{aligned} (n + 1)((n + 1) + 1) &= (n + 1)(n + 2) = n^2 + n + 2n + 2 = n(n + 1) + 2n + 2 \\ &= 2m + 2n + 2 = 2(m + n + 1) \end{aligned}$$

and, since $m + n + 1$ is a natural number, the statement is true for $n + 1$.

Exercise 13.41

Prove by induction that for every integer $n \geq 2$ the natural number $n^3 - n$ is divisible by 6, that is:

$$\forall n \in \mathbb{N}, n \geq 2, \exists m \in \mathbb{N} : n^3 - n = 6m$$

(hint: use the previous exercise).

Solution

By induction on n .

- **Base case**

Let $n = 2$. Then:

$$2^3 - 2 = 6 = 6 \cdot 1$$

and, since 1 is a natural number, the statement is true.

- **Passo induttivo**

Suppose it is true for n , and let us prove it for $(n + 1)$. By the inductive hypothesis, there exists $m \in \mathbb{N}$ such that $n^3 - n = 6m$. Moreover, by the previous exercise, $n(n + 1)$ is even: there exists $h \in \mathbb{N}$ such that $n(n + 1) = 2h$. Hence we can write:

$$\begin{aligned} (n + 1)^3 - (n + 1) &= n^3 + 3n^2 + 3n + 1 - n - 1 = (n^3 - n) + (3n^2 + 3n) \\ &= 6m + 3n(n + 1) = 6m + 3 \cdot 2h = 6(m + h) \end{aligned}$$

and, since $m + h$ is a natural number, the statement is true for $n + 1$.

Exercise 13.42

Prove by induction that for every integer $n \geq 0$ and every $x \in \mathbb{R}$, $x \geq -\frac{2}{3}$, the following holds:

$$(1 + x)e^n \geq \frac{1}{n + 3}$$

Solution

By induction on n .

- **Base case**

Let $n = 0$. Then the statement becomes:

$$(1 + x)e^0 \geq \frac{1}{0 + 3} \quad \text{i.e.} \quad x \geq -\frac{2}{3}$$

which is true by hypothesis.

- **Passo induttivo**

Suppose it is true for n , and let us prove it for $(n + 1)$. By the inductive hypothesis,

we have $(1+x)e^n \geq \frac{1}{n+3}$. Hence, since $e > 1$, we can write:

$$(1+x)e^{n+1} = e((1+x)e^n) \geq \frac{e}{n+3} \geq \frac{1}{n+3} \geq \frac{1}{(n+1)+3}$$

which is exactly the desired statement for $n+1$.

13.5 Complex numbers

Exercise 13.43

Write the following complex numbers z in algebraic form, identifying their real part and imaginary part. Then, for each of them, find the modulus, the conjugate and the reciprocal.

$$z = \frac{2+5i}{1+i} \qquad z = \frac{2+i}{1-3i}$$

Solution

$$z = \frac{2+5i}{1+i} \cdot \frac{1-i}{1-i} = \frac{7+3i}{2} = \frac{7}{2} + i\frac{3}{2}$$

from which

$$\Re z = \frac{7}{2}, \Im z = \frac{3}{2}, |z| = \sqrt{\frac{49}{4} + \frac{9}{4}} = \frac{\sqrt{58}}{2}, \bar{z} = \frac{7}{2} - i\frac{3}{2}.$$

Finally

$$z^{-1} = \frac{1}{z} \cdot \frac{\bar{z}}{\bar{z}} = \frac{\bar{z}}{|z|^2} = \frac{7-3i}{2} \cdot \frac{4}{58} = \frac{7}{29} - i\frac{3}{29}.$$

Solution

$$z = \frac{2+i}{1-3i} \cdot \frac{1+3i}{1+3i} = \frac{-1+7i}{10} = -\frac{1}{10} + i\frac{7}{10}$$

from which

$$\Re z = -\frac{1}{10}, \Im z = \frac{7}{10}, |z| = \sqrt{\frac{1}{100} + \frac{49}{100}} = \frac{\sqrt{2}}{2}, \bar{z} = -\frac{1}{10} - i\frac{7}{10}.$$

Finally

$$z^{-1} = \frac{1}{z} \cdot \frac{\bar{z}}{\bar{z}} = \frac{\bar{z}}{|z|^2} = -\frac{1+7i}{10} \cdot 2 = -\frac{1}{5} - i\frac{7}{5}.$$

Exercise 13.44

Determine all the complex numbers z that satisfy:

$$|z| \leq |z-2i|$$

Solution

It is convenient to use the form $z = x + iy$, $x, y \in \mathbb{R}$.

$$|x + iy| \leq |x + i(y - 2)|$$

$$|x + iy|^2 \leq |x + i(y - 2)|^2$$

$$x^2 + y^2 \leq x^2 + (y - 2)^2$$

$$0 \leq -4y + 4$$

$$y \leq 1.$$

Representing the solutions in the Cartesian plane, we obtain a closed half-plane.

Exercise 13.45

Determine all the complex numbers z that satisfy:

$$|z - (3 + i)| \leq 2$$

Solution

This inequality is satisfied by those complex numbers $z = x + iy$ which, in the Gauss plane (complex plane), are represented by the points $P(x, y)$ whose distance from the point $(3, 1)$ is less than or equal to 2.

That is, all and only the points of the **closed disk** with center $C(3, 1)$ and radius 2 (careful: it is not the circle, but the disk, i.e., the interior together with the boundary). The circle has equation:

$$\gamma: (x - 3)^2 + (y - 1)^2 = 4$$

while the closed disk has equation:

$$(x - 3)^2 + (y - 1)^2 \leq 4$$

Alternatively, the inequality can be handled through the substitution $z = x + iy$, obtaining

$$|x + iy - 3 - i| \leq 2$$

in which, factoring out the imaginary unit,

$$|x - 3 + i(y - 1)| \leq 2$$

Applying the definition of the modulus of a complex number, we have

$$\sqrt{(x - 3)^2 + (y - 1)^2} \leq 2$$

Since we can square both non-negative sides, we finally obtain:

$$(x - 3)^2 + (y - 1)^2 \leq 4$$

Exercise 13.46

Determine all the complex numbers z that satisfy:

$$(1 - i)z - (1 + i)\bar{z} = i$$

Solution

$$(1 - i)(x + iy) - (1 + i)(x - iy) = i$$

$$x + y - ix + iy - x - y - ix + iy = i$$

$$i(2y - 2x) = i$$

$$2y - 2x = 1.$$

Representing the solutions in the Cartesian plane, we obtain a line.

Exercise 13.47

Determine all the complex numbers z that satisfy:

$$|z + \bar{z}| + |z - \bar{z}| \leq 2$$

Solution

$$|z + \bar{z}| + |z - \bar{z}| \leq 2$$

$$2|\Re z| + 2|\Im z| \leq 2$$

$$|x| + |y| \leq 1.$$

Representing the solutions in the Cartesian plane, we obtain a closed square with vertices $\pm 1, \pm i$.

Exercise 13.48

Determine all the complex numbers z that satisfy:

$$\Im(z) - |z + \bar{z}|^2 < 1$$

Solution

$$\Im(z) - |z + \bar{z}|^2 < 1$$

$$\Im(z) - 4|\Re z|^2 < 1$$

$$y < 4x^2 + 1.$$

Representing the solutions in the Cartesian plane, we obtain the open region below the parabola $y = 4x^2 + 1$.

Exercise 13.49

Determine all the complex numbers z that satisfy:

$$\begin{cases} z\bar{z} \leq 2 \\ z + \bar{z} \leq 2 \end{cases}$$

Solution

$$\begin{cases} |z|^2 \leq 2 \\ 2\Re z \leq 2 \end{cases} \quad \begin{cases} x^2 + y^2 \leq 2 \\ x \leq 1 \end{cases}$$

Representing the solutions in the Cartesian plane, we obtain the part of the plane common to the closed disk with center at the origin and radius $\sqrt{2}$ and to the closed half-plane to the left of the vertical line $x = 1$.

Exercise 13.50

The system in $\mathbb{C}$

$$\begin{cases} z^2 + \bar{z}^2 = 0 \\ |\Re(z)| + |\Im(z)| = 1 \end{cases}$$

has:

- (a) two solutions (b) three solutions
(c) four solutions (d) none of the other answers is correct

Solution

Setting $z = x + iy$, $x, y \in \mathbb{R}$, we have

$$\begin{cases} (x + iy)^2 + (x - iy)^2 = 0 \\ |x| + |y| = 1 \end{cases}$$

$$\begin{cases} x^2 - y^2 + 2ixy + x^2 - y^2 - 2ixy = 0 \\ |x| + |y| = 1 \end{cases}$$

$$\begin{cases} x^2 = y^2 \\ |x| + |y| = 1 \end{cases} \quad \begin{cases} |x| = |y| \\ |x| + |y| = 1 \end{cases} \quad \begin{cases} |x| = 1/2 \\ |y| = 1/2 \end{cases}$$

We obtain the four points $(1 + i)/2$, $(-1 + i)/2$, $(1 - i)/2$, $(-1 - i)/2$. The correct answer is (c).

Exercise 13.51

The solutions $z = x + iy$ of the complex equation

$$z^2 - z\bar{z} + iz = -3 + 2i$$

satisfy

$$\begin{aligned} \text{(a)} \quad \left(x + \frac{1}{6}\right)^2 + \left(y + \frac{1}{4}\right)^2 &= 1 & \text{(b)} \quad y &= \frac{2}{3}x + 1 \\ \text{(c)} \quad \left(x - \frac{1}{6}\right)^2 + \left(y - \frac{1}{4}\right)^2 &= 1 & \text{(d)} \quad y &= \frac{3}{2}x \end{aligned}$$

Solution

$$(x + iy)^2 - (x^2 + y^2) + ix - y = -3 + 2i$$

$$-2y^2 - y + i(x + 2xy) = -3 + 2i$$

$$\begin{cases} 2y^2 + y - 3 = 0 \\ x + 2xy = 2 \end{cases} \quad \begin{cases} y = 1 \vee y = -3/2 \\ x = 2/(1 + 2y) \end{cases}$$

We obtain the two points $(2/3) + i$, $-1 - i(3/2)$, which lie on the line $y = (3/2)x$. The correct answer is (d).

Exercise 13.52

Find the modulus and the principal argument of

$$\text{(a)} \quad z = -1 - i\sqrt{3} \quad , \quad \text{(b)} \quad z = -4i.$$

Solution

(a) We have $|z| = \sqrt{1+3} = 2$. The principal argument $\text{Arg } z$ is the unique $\vartheta \in [-\pi, \pi)$ such that

$$\cos \vartheta = \Re z / |z| = -1/2, \quad \sin \vartheta = \Im z / |z| = -\sqrt{3}/2$$

hence $\text{Arg } z = -2\pi/3$.

Solution

(b) We have $|z| = 4$. The principal argument $\text{Arg } z$ is the unique $\vartheta \in [-\pi, \pi)$ such that

$$\cos \vartheta = \Re z / |z| = 0, \quad \sin \vartheta = \Im z / |z| = -1$$

hence $\text{Arg } z = -\pi/2$.

Exercise 13.53

Let

$$w = \frac{z + 1 - i}{z + i}$$

Determine, locating them in the Gauss plane, the complex numbers z for which w is a purely imaginary number.

Solution

The complex number w is defined for $z \neq -i$. Moreover,

$$w = \frac{z + 1 - i}{z + i} \cdot \frac{z - i}{z - i} = \frac{z^2 + z - 1 + i(-2z - 1)}{z^2 + 1}$$

That is, separating the real part of w from its imaginary part:

$$w = \frac{z^2 + z - 1}{z^2 + 1} + i \frac{-2z - 1}{z^2 + 1}$$

The complex number w is purely imaginary when $\Re(w) = 0$, i.e.,

$$\Re(w) = \frac{z^2 + z - 1}{z^2 + 1} = 0$$

This rational equation, under the condition $z \neq \pm\sqrt{-1}$, i.e., $z \neq \pm i$, gives the real numbers

$$z_1 = -\frac{1}{2} + \frac{\sqrt{5}}{2}, \quad z_2 = -\frac{1}{2} - \frac{\sqrt{5}}{2}$$

which, in the Gauss plane, can be represented as two points on the real axis. In conclusion, w is purely imaginary for the two values of z found.

Exercise 13.54

Let $z = \sqrt{3} - 3i$. The number of solutions w with $\Im(w) < 0$ of the equation

$$w^6 = |z|(z + \bar{z})$$

is:

- (a) two (b) three
(c) zero (d) none of the other answers is correct

Solution

We have $|z| = \sqrt{12}$, $z + \bar{z} = 2\sqrt{3}$, hence

$$w^6 = 12.$$

The complex sixth roots of a positive real number are six, pairwise opposite, and two of them are real. Of the remaining four, two have positive imaginary part and two have negative imaginary part. The correct answer is (a). We nevertheless want to determine their values.

Setting $w = r(\cos \vartheta + i \sin \vartheta)$ and taking into account that $|12| = 12$, $\arg 12 = 2k\pi$, $k \in \mathbb{Z}$, we obtain

$$\begin{cases} r^6 = 12 \\ 6\vartheta = 2k\pi \end{cases} \quad \begin{cases} r = 12^{1/6} \\ \vartheta = k\pi/3 \end{cases}$$

We therefore have the six roots

$$w_k = 12^{1/6}(\cos k\pi/3 + i \sin k\pi/3), \quad k = 0, 1, 2, 3, 4, 5,$$

i.e.,

$$w_0 = 12^{1/6}, \quad w_1 = 12^{1/6}(1/2 + i\sqrt{3}/2), \quad w_2 = 12^{1/6}(-1/2 + i\sqrt{3}/2)$$

$$w_3 = -12^{1/6}, \quad w_4 = 12^{1/6}(-1/2 - i\sqrt{3}/2), \quad w_5 = 12^{1/6}(1/2 - i\sqrt{3}/2).$$

The two with negative imaginary part are w_4 and w_5 .

Exercise 13.55

Solve the following system in the unknown $z \in \mathbb{C}$.

$$\begin{cases} z^6 = i \\ \Re(z)\Im(z) < 0 \end{cases}$$

Solution

We first solve the equation $z^6 = i$. Setting $z = r(\cos \vartheta + i \sin \vartheta)$ and taking into account that $|i| = 1$, $\arg(i) = \pi/2 + 2k\pi$, $k \in \mathbb{Z}$, we obtain

$$\begin{cases} r^6 = 1 \\ 6\vartheta = \pi/2 + 2k\pi \end{cases} \quad \begin{cases} r = 1 \\ \vartheta = \pi/12 + k\pi/3. \end{cases}$$

We therefore have the six roots

$$z_k = \cos((\pi + 4k\pi)/12) + i \sin((\pi + 4k\pi)/12), \quad k = 0, 1, 2, 3, 4, 5.$$

Those satisfying the condition $\Re(z)\Im(z) < 0$ (points in the second or fourth quadrant) are:

$$z_2 = -\sqrt{1/2} + i\sqrt{1/2}, \quad z_5 = \sqrt{1/2} - i\sqrt{1/2}.$$

Exercise 13.56

Solve the following equation in the complex variable z .

$$z^3 + 8 = 0$$

Solution

The equation to be solved is, clearly,

$$z^3 = -8$$

Setting $z = r(\cos \vartheta + i \sin \vartheta)$ and taking into account that $|-8| = 8$, $\arg(-8) = \pi + 2k\pi$, $k \in \mathbb{Z}$, we obtain

$$\begin{cases} r^3 = 8 \\ 3\vartheta = \pi + 2k\pi \end{cases} \quad \begin{cases} r = 2 \\ \vartheta = (2k+1)\pi/3. \end{cases}$$

We therefore have the three roots

$$z_k = 2 \cos((2k+1)\pi/3) + 2i \sin((2k+1)\pi/3), \quad k = 0, 1, 2,$$

i.e.,

$$z_0 = 1 + \sqrt{3}i, \quad z_1 = -2, \quad z_2 = 1 - \sqrt{3}i.$$

Exercise 13.57

Solve the following equation in the complex variable z .

$$z^3 = -27i$$

Solution

Setting $z = r(\cos \vartheta + i \sin \vartheta)$ and taking into account that $|-27i| = 27$, $\arg(-27i) = -\pi/2 + 2k\pi$, $k \in \mathbb{Z}$, we obtain

$$\begin{cases} r^3 = 27 \\ 3\vartheta = -\pi/2 + 2k\pi \end{cases} \quad \begin{cases} r = 3 \\ \vartheta = (4k-1)\pi/6. \end{cases}$$

We therefore have the three roots

$$z_k = 3 \cos((4k-1)\pi/6) + 3i \sin((4k-1)\pi/6), \quad k = 0, 1, 2,$$

i.e.,

$$z_0 = 3\sqrt{3}/2 - 3i/2, \quad z_1 = 3i, \quad z_2 = -3\sqrt{3}/2 - 3i/2.$$

Exercise 13.58

Solve the following equation in the complex variable z .

$$z^5 + (3 + i\sqrt{3})z = 0$$

Solution

We have

$$z(z^4 + 3 + i\sqrt{3}) = 0$$

hence one solution is $z = 0$. The other four are found by solving

$$z^4 = -3 - i\sqrt{3}.$$

Setting $z = r(\cos \vartheta + i \sin \vartheta)$ and taking into account that

$$|-3 - i\sqrt{3}| = 2\sqrt{3}, \quad \arg(-3 - i\sqrt{3}) = -5\pi/6 + 2k\pi, k \in \mathbb{Z}$$

we obtain

$$\begin{cases} r^4 = 12^{1/2} \\ 4\vartheta = -5\pi/6 + 2k\pi \end{cases} \quad \begin{cases} r = 12^{1/8} \\ \vartheta = -5\pi/24 + k\pi/2. \end{cases}$$

Besides the solution $z = 0$, we therefore have four more roots

$$z_k = 12^{1/8} \cos((12k - 5)\pi/24) + 12^{1/8}i \sin((12k - 5)\pi/24), \quad k = 0, 1, 2, 3.$$

Exercise 13.59

Solve the following equation in the complex variable z .

$$\left(\frac{z+i}{1+i}\right)^3 = -8i$$

Solution

Setting $w = (z+i)/(1+i)$, we solve $w^3 = -8i$. Writing $w = r(\cos \vartheta + i \sin \vartheta)$ and taking into account that $|-8i| = 8$, $\arg(-8i) = -\pi/2 + 2k\pi$, $k \in \mathbb{Z}$, we obtain

$$\begin{cases} r^3 = 8 \\ 3\vartheta = -\pi/2 + 2k\pi \end{cases} \quad \begin{cases} r = 2 \\ \vartheta = (4k - 1)\pi/6. \end{cases}$$

We therefore have the three roots

$$w_k = 2 \cos((4k - 1)\pi/6) + 2i \sin((4k - 1)\pi/6), \quad k = 0, 1, 2,$$

i.e.,

$$w_0 = \sqrt{3} - i, \quad w_1 = 2i, \quad w_2 = -\sqrt{3} - i.$$

Going back to the variable z , from $z = (1+i)w - i$, the three solutions of the given equation are

$$z_0 = \sqrt{3} + 1 + i\sqrt{3} - 2i, \quad z_1 = -2 + i, \quad z_2 = 1 - \sqrt{3} - i\sqrt{3} - 2i.$$

Exercise 13.60

Solve the following equation in the complex variable z .

$$\left(\frac{z-i}{i}\right)^4 = -16$$

Solution

From $i^4 = 1$ we have $(z - i)^4 = -16$. Setting $w = z - i$, we solve $w^4 = -16$. Writing $w = r(\cos \vartheta + i \sin \vartheta)$ and taking into account that $|-16| = 16$, $\arg(-16) = -\pi + 2k\pi$, $k \in \mathbb{Z}$, we obtain

$$\begin{cases} r^4 = 16 \\ 4\vartheta = -\pi + 2k\pi \end{cases} \quad \begin{cases} r = 2 \\ \vartheta = -\pi/4 + k\pi/2. \end{cases}$$

We therefore have the four roots

$$w_k = 2 \cos((2k - 1)\pi/4) + 2i \sin((2k - 1)\pi/4), \quad k = 0, 1, 2, 3,$$

i.e.,

$$w_0 = \sqrt{2} - i\sqrt{2}, \quad w_1 = \sqrt{2} + i\sqrt{2}, \quad w_2 = -\sqrt{2} + i\sqrt{2}, \quad w_3 = -\sqrt{2} - i\sqrt{2}.$$

Going back to the variable z , from $z = w + i$, the four solutions of the given equation are

$$z_0 = \sqrt{2} - i\sqrt{2} + i, \quad z_1 = \sqrt{2} + i\sqrt{2} + i, \quad z_2 = -\sqrt{2} + i\sqrt{2} + i, \quad z_3 = -\sqrt{2} - i\sqrt{2} + i.$$

Exercise 13.61

Solve the following equation in the complex variable z .

$$z^5 = -\bar{z}$$

Solution

The equation can be seen as

$$z^5 = (-1)\bar{z}$$

Setting $z = r(\cos \vartheta + i \sin \vartheta)$ and taking into account that $|\bar{z}| = r$, $\arg(\bar{z}) = -\vartheta + 2k\pi$, $k \in \mathbb{Z}$, and recalling that $-1 = e^{i\pi}$, we obtain

$$\begin{cases} r^5 = r \\ 5\vartheta = \pi - \vartheta + 2k\pi \end{cases} \quad \begin{cases} r = 0, r = 1 \\ \vartheta = (2k + 1)\pi/6. \end{cases}$$

At this point,

- for $r = 0$, we obtain $z_0 = 0$;
- for $r = 1$, we obtain the roots $z = \cos((2k + 1)\pi/6) + i \sin((2k + 1)\pi/6)$, $k = 0, 1, 2, 3, 4, 5$.

Exercise 13.62

The equation

$$z^4 - 2z^3 + 4z^2 - 2z + 3 = 0$$

has $z_1 = i$ as a solution. Denoting by z_2, z_3, z_4 the other solutions, we have:

$$(a) \quad \Im(z_1 z_2 z_3 z_4) = 0 \quad (b) \quad \Re(z_1 + z_2 + z_3 + z_4) = 0$$

- (c) $\Re((z_1 + z_2 + z_3 + z_4)^{-1}) = 1/2$ (d) none of the other answers is correct

Solution

The equation has real coefficients, hence the solutions come in conjugate pairs; in particular, another solution is $z_2 = -i$. It follows that the polynomial

$$p(z) = z^4 - 2z^3 + 4z^2 - 2z + 3$$

is divisible by $(z - i)(z + i)$, i.e., by $z^2 + 1$. Performing the division we obtain

$$p(z) = (z^2 + 1)(z^2 - 2z + 3)$$

hence among the solutions we also have the roots of

$$z^2 - 2z + 3 = 0$$

which are $z_3 = 1 - i\sqrt{2}$, $z_4 = 1 + i\sqrt{2}$.

Summarizing, the solutions of the given equation are

$$z_1 = i, \quad z_2 = -i, \quad z_3 = 1 - i\sqrt{2}, \quad z_4 = 1 + i\sqrt{2}.$$

Since $z_1 + z_2 + z_3 + z_4 = 2$ and $z_1 z_2 z_3 z_4 = 3$, the correct answers are (a) and (c).

Exercise 13.63

Let z_h , $h = 1, \dots, 6$ be the solutions of

$$z^6 - (8 - i)z^3 - 8i = 0.$$

We have:

$$(a) \sum_{h=1}^6 \Re(z_h) = -1 \quad (b) \sum_{h=1}^6 \Im(z_h) = -1$$

$$(c) \sum_{h=1}^6 \Re(z_h) = 1 \quad (d) \sum_{h=1}^6 \Im(z_h) = 0$$

$$(e) \sum_{h=1}^6 \Re(z_h) = 2 \quad (f) \text{ none of the other answers is correct}$$

Solution

It is convenient to factor it as

$$(z^3 - 8)(z^3 + i) = 0$$

and then solve the two equations

$$z^3 = 8, \quad z^3 = -i$$

with the usual method illustrated several times in the solutions of the previous exercises.

We obtain the values:

$$z_1 = 2, \quad z_2 = -1 + i\sqrt{3}, \quad z_3 = -1 - i\sqrt{3},$$

$$z_4 = \sqrt{3}/2 - i/2, \quad z_5 = i, \quad z_6 = -\sqrt{3}/2 - i/2.$$

The correct answer is (d).