

L'Hôpital's rule and differentiability



Outline

L'Hôpital's rule

Limit of the derivative and differentiability

L'Hôpital's rule

Introduction: L'Hôpital's rule

- ▶ In this section we see how differential calculus helps to compute limits that appear in indeterminate form
- ▶ L'Hôpital's rule allows us to replace the quotient f/g with the quotient of the derivatives f'/g'
- ▶ We will see when it is convenient to use it, when it does not help and which precautions to take



- ▶ A remarkable application of differential calculus is the computation of limits that appear in the indeterminate forms:

$$\left[\frac{0}{0} \right] \quad \text{and} \quad \left[\frac{\infty}{\infty} \right]$$

Theorem 1: L'Hôpital's rule

Let f, g be differentiable functions on an interval (a, b) with $g, g' \neq 0$ in (a, b) . If

$$\lim_{x \rightarrow a^+} f(x) = \lim_{x \rightarrow a^+} g(x) = 0 \quad (\text{or } \pm \infty) \quad \text{and} \quad \lim_{x \rightarrow a^+} \frac{f'(x)}{g'(x)} = \ell \in \mathbb{R}^*$$

then:

$$\lim_{x \rightarrow a^+} \frac{f(x)}{g(x)} = \ell$$

The theorem still holds if $a = -\infty$ or if we consider the limit as $x \rightarrow b^-$, with $b \leq +\infty$ (a case we do not prove). Hence, if the hypotheses hold for a^- and a^+ , it can clearly be applied also at an interior point of an interval.

Example

We want to compute:

$$\lim_{x \rightarrow 0} \frac{\sin x}{x} = \left[\frac{0}{0} \right]$$

$$\text{and } f(x) = \sin x, f'(x) = \cos x, g(x) = x, g'(x) = 1$$

applying L'Hôpital's rule, for example with $x \in (-1, 0) \cup (0, 1)$, we obtain:

$$\lim_{x \rightarrow 0} \frac{\sin x}{x} \stackrel{H}{=} \lim_{x \rightarrow 0} \frac{\cos x}{1} = 1$$

Example

We want to compute:

$$\lim_{x \rightarrow \frac{\pi}{2}} \frac{1 - \sin x}{\cos x} = \left[\frac{0}{0} \right]$$

and $f(x) = 1 - \sin x$, $f'(x) = -\cos x$, $g(x) = \cos x$, $g'(x) = -\sin x$

applying L'Hôpital's rule, for example with $x \in (\frac{\pi}{4}, \frac{\pi}{2}) \cup (\frac{\pi}{2}, \frac{3\pi}{4})$, we obtain:

$$\lim_{x \rightarrow \frac{\pi}{2}} \frac{1 - \sin x}{\cos x} \stackrel{H}{=} \lim_{x \rightarrow \frac{\pi}{2}} \frac{-\cos x}{-\sin x} = \lim_{x \rightarrow \frac{\pi}{2}} \frac{\cos x}{\sin x} = \frac{0}{1} = 0$$

- ▶ We now prove the theorem for the form $\begin{bmatrix} 0 \\ 0 \end{bmatrix}$ as $x \rightarrow a^+$
- ▶ The idea is to apply Lagrange's theorem to an auxiliary function h built from f and g

Proof (1/2)

We consider the case:

$$f(x), g(x) \rightarrow 0, \quad \text{as } x \rightarrow a^+$$

Let $\{x_n\}$ be a sequence such that $x_n \rightarrow a^+$ as $n \rightarrow +\infty$ and $x_n \neq a, \forall n$, and let us extend f and g by continuity at a by setting $f(a) = g(a) = 0$.

We now define the following function:

$$h(x) = f(x_n) g(x) - g(x_n) f(x)$$

The function h satisfies the hypotheses of Lagrange's theorem on the interval $[a, x_n]$, hence there exists $t_n \in (a, x_n)$ such that

$$h'(t_n) = \frac{h(x_n) - h(a)}{x_n - a} = 0 \quad \text{since} \quad h(a) = h(x_n) = 0$$

Proof (2/2)

Computing

$$h'(x) = f(x_n) g'(x) - g(x_n) f'(x),$$

$$\text{we have } \underbrace{f(x_n) g'(t_n) - g(x_n) f'(t_n)}_{=h'(t_n)} = 0$$

Hence for every x_n there exists a point $t_n \in (a, x_n)$ such that:

$$\frac{f(x_n)}{g(x_n)} = \frac{f'(t_n)}{g'(t_n)} \quad (1)$$

As $n \rightarrow +\infty$ we have $t_n \rightarrow a^+$, since $a^+ < t_n < x_n$ and $x_n \rightarrow a^+$ as $n \rightarrow +\infty$, therefore by the hypothesis of the theorem we have:

$$\frac{f'(t_n)}{g'(t_n)} \rightarrow \ell \quad \text{and consequently} \quad \frac{f(x_n)}{g(x_n)} \rightarrow \ell$$

which is what we wanted to prove. □

With L'Hôpital's rule we can easily prove the theorem on the hierarchy of infinities for functions.

Proof (1/2)

We prove:

$$\lim_{x \rightarrow +\infty} \frac{x^\alpha}{b^{\lambda x}} = 0, \quad \text{with } \alpha > 0, \lambda > 0, b > 1$$

Setting $f(x) = x$, $g(x) = b^{\mu x}$ ($\mu > 0$), we have $f'(x) = 1$, $g'(x) = \mu \log b \, b^{\mu x}$ and since:

$$\lim_{x \rightarrow +\infty} \frac{f'(x)}{g'(x)} = \lim_{x \rightarrow +\infty} \frac{1}{\mu \log b \, b^{\mu x}} = 0 \quad \text{then} \quad \lim_{x \rightarrow +\infty} \frac{x}{b^{\mu x}} = 0$$

Observing now that

$$\frac{x^\alpha}{b^{\lambda x}} = \left(\frac{x}{b^{(\lambda/\alpha) x}} \right)^\alpha = \left(\frac{x}{b^{\mu x}} \right)^\alpha$$

$$\text{with } \mu = \frac{\lambda}{\alpha} > 0 \quad \text{then} \quad \lim_{x \rightarrow +\infty} \frac{x^\alpha}{b^{\lambda x}} = \left(\lim_{x \rightarrow +\infty} \frac{x}{b^{\mu x}} \right)^\alpha = 0$$

Proof (2/2)

We prove:

$$\lim_{x \rightarrow +\infty} \frac{(\log_a x)^\beta}{x^\alpha} = 0, \quad \text{with } \beta > 0, \alpha > 0, a > 1$$

it suffices to perform the change of variable

$$t = \log_a x, \quad x = a^t, \quad \text{we have that as } x \rightarrow +\infty \text{ also } t \rightarrow +\infty$$

hence

$$\lim_{x \rightarrow +\infty} \frac{(\log_a x)^\beta}{x^\alpha} = \lim_{t \rightarrow +\infty} \frac{t^\beta}{a^{\alpha t}} = 0$$

since the second limit has the same form as the one just proved.



- ▶ L'Hôpital's rule can be useful for limits that cannot be solved using only fundamental limits or the asymptotic expansions derived from them.



Example

We compute

$$\lim_{x \rightarrow 0} \frac{x - \sin x}{x^3} = \left[\frac{0}{0} \right]$$

Fundamental limits are not sufficient to resolve the indeterminate form, since:

$$\frac{x - \sin x}{x^3} = \frac{1}{x^2} \left(1 - \frac{\sin x}{x} \right)$$

$$\text{and } \lim_{x \rightarrow 0} \frac{1}{x^2} \underbrace{\left(1 - \frac{\sin x}{x} \right)}_{\rightarrow 1} = [\infty \cdot 0]$$

L'Hôpital's rule gives:

$$\lim_{x \rightarrow 0} \frac{x - \sin x}{x^3} \stackrel{H}{=} \lim_{x \rightarrow 0} \frac{(x - \sin x)'}{(x^3)'} = \lim_{x \rightarrow 0} \frac{1 - \cos x}{3x^2} = \frac{1}{3} \cdot \lim_{x \rightarrow 0} \underbrace{\frac{1 - \cos x}{x^2}}_{\rightarrow \frac{1}{2}} = \frac{1}{6}$$

Example (1/2)

We now treat the same limit with the first-order asymptotic expansion:

$$\sin x = x + o(x) \quad \text{as } x \rightarrow 0$$

we obtain:

$$\lim_{x \rightarrow 0} \frac{x - \sin x}{x^3} = \lim_{x \rightarrow 0} \frac{x - (x + o(x))}{x^3} = \lim_{x \rightarrow 0} \frac{o(x)}{x^3}$$

That is, we still have an indeterminate form, since:

$$\lim_{x \rightarrow 0} \frac{o(x)}{x^3} = \lim_{x \rightarrow 0} \frac{1}{x^2} \frac{o(x)}{x} = \lim_{x \rightarrow 0} \frac{1}{x^2} o(1) = [+\infty \cdot 0]$$

where $o(1)$ denotes a generic function that tends to 0 as $x \rightarrow 0$.

Example (2/2)

We reach the same conclusion also with the following reasoning. The symbol $o(x)$ as $x \rightarrow 0$ denotes the set of functions that, divided by x , tend to 0 as $x \rightarrow 0$. Hence, for example, $5x^2 = o(x)$ but also $x^3 = o(x)$, and the value of the limit cannot be determined with the first-order asymptotic expansion. As we will see later, to determine this limit with asymptotic expansions we need an asymptotic expansion of order higher than the first.

Example (1/2)

We compute

$$\lim_{x \rightarrow +\infty} \frac{x - \sin x}{x + \sin x} = \left[\frac{+\infty}{+\infty} \right]$$

L'Hôpital's rule gives:

$$\lim_{x \rightarrow +\infty} \frac{x - \sin x}{x + \sin x} \stackrel{H}{=} \lim_{x \rightarrow +\infty} \frac{(x - \sin x)'}{(x + \sin x)'} = \lim_{x \rightarrow +\infty} \frac{1 - \cos x}{1 + \cos x} \quad \text{does not exist}$$

the fact that this limit does not exist does not allow us to conclude that the limit we are looking for does not exist.

Example (2/2)

Let us proceed instead with asymptotic estimates:

$$\frac{x - \sin x}{x + \sin x} \sim \frac{x}{x} \quad \text{as } x \rightarrow +\infty$$

Since

$$\lim_{x \rightarrow +\infty} \frac{x - \sin x}{x} = \lim_{x \rightarrow +\infty} 1 - \underbrace{\frac{\sin x}{x}}_{\rightarrow 0}$$

$$\text{and } \lim_{x \rightarrow +\infty} \frac{x + \sin x}{x} = \lim_{x \rightarrow +\infty} 1 + \underbrace{\frac{\sin x}{x}}_{\rightarrow 0}$$

hence:

$$\lim_{x \rightarrow +\infty} \frac{x - \sin x}{x + \sin x} = \lim_{x \rightarrow +\infty} \frac{x}{x} = 1$$

- ▶ The theorem is useful when its application simplifies the limit instead of complicating it, that is, when the order of the infinitesimal (or of the infinity) in the numerator or in the denominator decreases.

Example

We compute

$$\lim_{x \rightarrow 0^+} \frac{e^{-1/x^2}}{x} = \left[\frac{0}{0} \right]$$

L'Hôpital's rule gives:

$$\lim_{x \rightarrow 0^+} \frac{e^{-1/x^2}}{x} \stackrel{H}{=} \lim_{x \rightarrow 0^+} \frac{(e^{-1/x^2})'}{(x)'} = \lim_{x \rightarrow 0^+} \frac{2 e^{-1/x^2}}{x^3} = \left[\frac{0}{0} \right]$$

Applying instead the change of variable:

$$x^2 = \frac{1}{t}, \quad x \rightarrow 0^+ \text{ is equivalent to } t \rightarrow +\infty$$

substituting we have:

$$\lim_{x \rightarrow 0^+} \frac{e^{-1/x^2}}{(x^2)^{1/2}} = \lim_{t \rightarrow +\infty} \frac{e^{-t}}{\left(\frac{1}{t}\right)^{1/2}} = \lim_{t \rightarrow +\infty} \frac{e^{-t}}{t^{-1/2}} = \lim_{t \rightarrow +\infty} \frac{\sqrt{t}}{e^t} = 0$$

where the last limit is obtained thanks to the theorem on the hierarchy of infinities.

- Sometimes the theorem must be applied several times consecutively to resolve the indeterminate form. Also in this case, already after the first application, one should notice that the order of the infinitesimal (or of the infinity) in the numerator or in the denominator has decreased.

Example

We compute

$$\lim_{x \rightarrow 0} \frac{1 - \cos^3 x}{x^3 - x^2} = \begin{bmatrix} 0 \\ 0 \end{bmatrix}$$

L'Hôpital's rule gives:

$$\lim_{x \rightarrow 0} \frac{1 - \cos^3 x}{x^3 - x^2} \stackrel{H}{=} \lim_{x \rightarrow 0} \frac{3 \sin x \cos^2 x}{3x^2 - 2x} = \begin{bmatrix} 0 \\ 0 \end{bmatrix}$$

L'Hôpital's rule applied twice gives:

$$\lim_{x \rightarrow 0} \frac{1 - \cos^3 x}{x^3 - x^2} \stackrel{H}{=} \lim_{x \rightarrow 0} \frac{3 \sin x \cos^2 x}{3x^2 - 2x} \stackrel{H}{=} \lim_{x \rightarrow 0} \frac{3 \overset{\rightarrow 1}{\cos x} \left(\overset{\rightarrow 1}{\cos^2 x} - \overset{\rightarrow 0}{2 \sin^2 x} \right)}{\underbrace{6x}_{\rightarrow 0} - 2} = -\frac{3}{2}$$

Precautions for use

- ▶ The theorem is used for quotients, not for products
- ▶ The theorem is used for quotients that are actual indeterminate forms:

$$\left[\frac{0}{0} \right] \quad \text{and} \quad \left[\frac{\infty}{\infty} \right]$$

- ▶ The theorem prescribes computing the quotient of the derivatives, not the derivative of the quotient
- ▶ If the limit of f'/g' does not exist, nothing can be said about the limit of f/g .

Summary

- ▶ **L'Hôpital's rule:** for the forms $\left[\frac{0}{0}\right]$ and $\left[\frac{\infty}{\infty}\right]$, if $\lim f'/g' = \ell$ exists then $\lim f/g = \ell$
- ▶ The proof uses Lagrange's theorem applied to $h(x) = f(x_n)g(x) - g(x_n)f(x)$
- ▶ It allows us to prove the hierarchy of infinities and to solve limits that fundamental limits cannot handle
- ▶ It is convenient when it lowers the order of the infinitesimal (or infinity); it can be applied several times
- ▶ If $\lim f'/g'$ does not exist, nothing can be said about $\lim f/g$

Limit of the derivative and differentiability



Introduction: Limit of the derivative

- ▶ In this section we see a way to establish the differentiability of a function at a point without computing the limit of the difference quotient
- ▶ The theorem on the limit of the derivative links the right (left) derivative to the right (left) limit of f'
- ▶ It is useful to recognize corner points, but care is needed when the limit of f' does not exist

Theorem 2: limit of the derivative

If $f : [a, b) \rightarrow \mathbb{R}$ is continuous at a , differentiable in (a, b) , and $\lim_{x \rightarrow a^+} f'(x) = m \in \mathbb{R}^$, then $f'_+(a) = m$.*

If the function is continuous at a and the right-hand limit of the derivative exists, then the right derivative exists and coincides with that limit. An analogous statement of the theorem holds for the left derivative and hence for the derivative.

Proof

Let $h > 0$. Applying Lagrange's theorem to f on the interval $[a, a + h]$, we obtain that there exists $t_h \in [a, a + h]$ such that

$$\frac{f(a + h) - f(a)}{h} = f'(t_h)$$

As $h \rightarrow 0^+$ we have $t_h \rightarrow a^+$, hence by the hypothesis of the theorem $f'(t_h) \rightarrow m$ as $t_h \rightarrow a^+$. Consequently, there exists

$$f'_+(a) = \lim_{h \rightarrow 0^+} \frac{f(a + h) - f(a)}{h} = m$$



Under the hypotheses of the theorem it is therefore possible to compute the right derivative, the left derivative or the derivative without using the limit of the difference quotient



Example (1/4)


Suppose we want to study whether or not the following function is differentiable:

$$f(x) = x \cdot |\log x|$$

defined for $x > 0$. For $x \neq 1$, where $\log x \neq 0$, we have:

$$f(x) = \begin{cases} x \cdot \log x, & x > 1 \\ -x \cdot \log x, & 0 < x < 1 \end{cases} \quad \text{hence} \quad f'(x) = \begin{cases} \log x + 1, & x > 1 \\ -(\log x + 1), & 0 < x < 1 \end{cases}$$

Example (2/4)



Example (3/4)

At $x = 1$ the function $f(x)$ is continuous, and we have:

$$\lim_{x \rightarrow 1^+} f'(x) = \lim_{x \rightarrow 1^+} (\log x + 1) = 1$$


$$\text{and } \lim_{x \rightarrow 1^-} f'(x) = \lim_{x \rightarrow 1^-} (-\log x - 1) = -1$$

hence the theorem is applicable (from the right and from the left)
and we have:

$$f'_+(1) = 1, f'_-(1) = -1, \text{ so that } x = 1$$

is a corner point and $f'(1)$ does not exist

Example (4/4)



Example (1/4)

Suppose we want to study whether or not the following function is differentiable:

$$f(x) = x^2 \cdot \sin\left(\frac{1}{x}\right)$$


defined for $x \neq 0$. We know that

$$\lim_{x \rightarrow 0} x^2 \cdot \sin\left(\frac{1}{x}\right) = 0$$

and we can extend f by continuity by defining $f(0) = 0$, and f turns out to be continuous also at 0. For $x \neq 0$, we have:

$$f'(x) = 2x \cdot \sin\left(\frac{1}{x}\right) + x^2 \cdot \cos\left(\frac{1}{x}\right) \cdot \left(-\frac{1}{x^2}\right) = \underbrace{2x \cdot \sin\left(\frac{1}{x}\right)}_{\rightarrow 0 \text{ as } x \rightarrow 0} - \cos\left(\frac{1}{x}\right)$$

Example (2/4)



Near zero, the function $\cos \frac{1}{x}$, like the function $\sin \frac{1}{x}$, cannot be drawn since it has infinitely many oscillations.

Example (3/4)

Hence the hypotheses of the theorem are not satisfied, since

$$\lim_{x \rightarrow 0} f'(x) \text{ does not exist, } \lim_{x \rightarrow 0^-} f'(x)$$


$$\text{does not exist, } \lim_{x \rightarrow 0^+} f'(x) \text{ does not exist}$$

However, this does not allow us to conclude that the function is not differentiable at $x_0 = 0$. To compute the derivative we need to use the limit of the difference quotient. We have

$$f'(0) = \lim_{h \rightarrow 0} \frac{f(h) - 0}{h} = \lim_{h \rightarrow 0} h \cdot \sin\left(\frac{1}{h}\right) = 0$$

hence the function is differentiable at $x = 0$

Example (4/4)



Summary

- ▶ **Limit of the derivative theorem:** if f is continuous at a , differentiable in (a, b) and $\lim_{x \rightarrow a^+} f'(x) = m$, then $f'_+(a) = m$
- ▶ An analogous statement holds for the left derivative and hence for the derivative
- ▶ Example $x |\log x|$: $f'_+(1) = 1 \neq f'_-(1) = -1$, hence $x = 1$ is a corner point
- ▶ Example $x^2 \sin \frac{1}{x}$: the limit of f' does not exist, but $f'(0) = 0$ via the difference quotient

Final summary

- ▶ L'Hôpital's rule resolves the indeterminate forms $\left[\frac{0}{0}\right]$ and $\left[\frac{\infty}{\infty}\right]$ by passing to the quotient of the derivatives
- ▶ It must be used with care: only for quotients, only for actual indeterminate forms, and the non-existence of $\lim f'/g'$ says nothing
- ▶ The limit of the derivative allows us to compute right and left derivatives without the difference quotient, when it exists