

Second derivative

Outline

Second derivative and second derivative function

Second derivative, concavity and convexity

Convexity and derivatives

Convexity and tangent lines

Inflection points

Second derivative and second derivative function



Introduction: Second derivative

- ▶ In this section we ask whether the derivative function f' is itself differentiable
- ▶ We define the second derivative, the second derivative function and the derivatives of order n , with their notations
- ▶ We will then see the geometric meaning of the second derivative: the curvature of the graph

- ▶ We can now ask whether the function $f'(x)$ is in turn differentiable (at a point or on an interval).

Definition 1: second derivative

Let $f : (a, b) \rightarrow \mathbb{R}$, $x_0 \in (a, b)$ and $f' : (x_0 - \delta, x_0 + \delta) \rightarrow \mathbb{R}$ with $\delta > 0$; f is said to be twice differentiable at x_0 if the following limit exists and is finite

$$\lim_{h \rightarrow 0} \frac{f'(x_0 + h) - f'(x_0)}{h}$$

this limit is called the second derivative of f at x_0 .

Notation

- To denote the second derivative function the following notations are used:

$$\underbrace{f''(x_0)}$$

Lagrange's notation

$$\underbrace{\ddot{f}(x_0)}$$

Newton's notation

$$\underbrace{\left. \frac{d^2 f}{dx^2} \right|_{x=x_0} \quad \text{and} \quad \left. \frac{d^2 y}{dx^2} \right|_{x=x_0}}_{\text{Leibniz's notation}}$$

Definition 2: second derivative function

If a derivative function f' is differentiable at every point of the interval (a, b) , the function

$$f'' : (a, b) \rightarrow \mathbb{R}, \quad f'' : x \mapsto f''(x)$$

*is called the **second derivative function** of f .*

Notation

- To denote the second derivative function the following notations are used:

$$\underbrace{f''(x)}$$

Lagrange's notation

$$\underbrace{\ddot{f}(x)}$$

Newton's notation

$$\underbrace{\frac{d^2 f}{dx^2}, \quad \frac{d^2 f(x)}{dx^2} \quad \text{and} \quad \frac{d^2 y}{dx^2}}_{\text{Leibniz's notation}}$$

Derivative of order n

- In a completely analogous way one defines the derivative of order n , or n -th derivative, and the n -th derivative function, which are denoted by the symbols:

$$\underbrace{f^{(n)}(x_0), \quad f^{(n)}(x)}_{\text{Lagrange's notation}} \quad \underbrace{\left. \frac{d^n f}{dx^n} \right|_{x=x_0} \quad \text{and} \quad \left. \frac{d^n y}{dx^n} \right|_{x=x_0}, \quad \frac{d^n f}{dx^n}, \quad \frac{d^n f(x)}{dx^n} \quad \text{and} \quad \frac{d^n y}{dx^n}}_{\text{Leibniz's notation}}$$

Geometric meaning of the second derivative

While the first derivative has, as its geometric meaning, the slope of the graph, the second derivative represents the **rate of change of this slope** and therefore is a measure of the **degree to which the graph deviates from a straight line**.



Geometric meaning of the second derivative

- We start by considering the family of functions that satisfy the conditions:

$$f(0) = f'(0) = 0, \quad f''(0) \geq 0$$


and the family of semicircles with center on the y -axis, tangent to the graph of f at the origin

$$c_r(x) = r - \sqrt{r^2 - x^2}, \quad \text{where } r > 0 \text{ is the radius}$$

For every $r \in \mathbb{R}_+$, we have:

$$c'_r(0) = c_r(0) = 0 \quad \text{since} \quad c'_r(x) = \frac{x}{\sqrt{r^2 - x^2}}$$

For example, the function $f(x) = 1 - \cos x$ with $f(0) = f'(0) = 0$ and the circles with radii $r \in \{0.5, 1, 1.5\}$ are:



- Among these semicircles we want to select the one that not only has the same tangent at $x = 0$ but also the same rate of change of the slope at $x = 0$. That is, we want to choose r so that:

$$c_r''(0) = f''(0) \quad (1)$$

We have:

$$c_r'(x) = \frac{x}{\sqrt{r^2 - x^2}} \quad \text{and} \quad c_r''(x) = \frac{r^2}{(r^2 - x^2)^{3/2}}$$

hence, if we want (1) to be satisfied, we need to choose r so that:

$$\underbrace{\frac{1}{r}}_{=c_r''(0)} = f''(0) \quad (2)$$


Equation (2) expresses the geometric meaning of the second derivative at $x = 0$ for the family of functions that satisfy the conditions $f(0) = f'(0) = 0$, $f''(0) \geq 0$. That is, $f''(0)$ represents the reciprocal of the radius of the semicircle that best approximates f at $x = 0$.

► Let us go back to the function $f(x) = 1 - \cos x$; we have

$$f'(x) = \sin x \quad \text{and} \quad f''(x) = \cos x$$

hence

$$f''(0) = 1 \quad \text{and} \quad \frac{1}{r} = 1 \quad \text{hence} \quad r = 1$$



Curvature

Repeating an analogous construction at a generic point $(x, f(x))$ of the graph of f (without assuming a priori $f'(x) = 0$) one finds that the **circle that best approximates the graph of the function at the point has radius $r(x)$** given by:

$$\frac{1}{r(x)} = \frac{|f''(x)|}{\left(1 + (f'(x))^2\right)^{3/2}} \quad (3)$$

The value $\frac{1}{r(x)}$ is called the **curvature** (of the graph) of f at x and the value $r(x)$ is the **radius of curvature**.

Example (1/3)

Consider the (parametric) parabola:

$$f(x) = a x^2 \quad \text{with } a \in \mathbb{R}$$

$$\text{we have } f'(x) = 2 a x \quad \text{and} \quad f''(x) = 2 a$$

and the following curvature:

$$\frac{1}{r(x)} = \frac{2 |a|}{(1 + 4 a^2 x^2)^{3/2}}$$

Note that the curvature is maximum at $x = 0$, i.e., at the vertex.


Example (2/3)

Let us now consider the parabola

$$y = 2x^2$$

and compute the radius $r(x)$ of the circle that best approximates the graph at the points with abscissa $x = 0$ and $x = \frac{1}{2}$. We have:

$$\frac{1}{r(0)} = \frac{2|2|}{(1 + 4 \cdot 2^2 \cdot 0^2)^{3/2}} = 4 \quad \text{hence} \quad r(0) = \frac{1}{4}$$



Example (3/3)

$$\frac{1}{r\left(\frac{1}{2}\right)} = \frac{2|2|}{\left(1 + 4 \cdot 2^2 \left(\frac{1}{2}\right)^2\right)^{3/2}} = \frac{4}{5^{3/2}} \approx 0.3577$$

$$\text{hence } r\left(\frac{1}{2}\right) \approx 2.7950$$

Summary

- ▶ **Second derivative:** derivative of the derivative function,
$$f''(x_0) = \lim_{h \rightarrow 0} \frac{f'(x_0+h) - f'(x_0)}{h}$$
- ▶ Lagrange, Newton and Leibniz notations; derivatives of order n are defined analogously
- ▶ **Geometric meaning:** f'' measures the rate of change of the slope
- ▶ **Curvature:** $\frac{1}{r(x)} = \frac{|f''(x)|}{(1+(f'(x))^2)^{3/2}}$, where $r(x)$ is the radius of curvature

Second derivative, concavity and convexity



Introduction: Concavity and convexity

- ▶ In this section we make the idea of curvature of the graph precise through the concept of convexity
- ▶ We start from convex sets in the plane and arrive at convex and concave functions via the epigraph
- ▶ Convexity is a fundamental property in analysis and optimization

- ▶ We will now see how to make this idea of curvature of the graph more precise, qualitatively and quantitatively, through the concept of **convexity**.

Definition 3: convex set

*A set $F \subseteq \mathbb{R}^2$ (a portion of the Euclidean space) is **convex** if for every pair of points $P_a, P_b \in F$ the segment joining P_a to P_b (called a **chord**) is entirely contained in F .*

Convex combination

Given two points P_a and P_b and a value λ between 0 and 1, i.e., $0 \leq \lambda \leq 1$, the point:

$$P_c = \lambda P_a + (1 - \lambda) P_b$$

is called a **convex (linear) combination** of the points P_a and P_b . As λ varies, the points P_c move along the segment joining P_a to P_b .

Definition 4: convex set – equivalent definition

A set $F \subseteq \mathbb{R}^2$ is *convex* if:


$$\forall P_a, P_b \in F, \quad 0 \leq \lambda \leq 1, \quad P_c = \lambda P_a + (1 - \lambda) P_b \in F$$

Example (1/2)

Consider, for example, the following convex set given by the intersection of 4 half-planes:


$$F = \left\{ (x, y) \in \mathbb{R}^2 : \begin{array}{ll} x + y \geq 3, & x + y \leq 9, \\ -x + y \leq 3, & -x + y \geq -3 \end{array} \right\}$$

Example (2/2)



Example (1/2)

Let us now consider the following set:



Example (2/2)

The set is not convex since, for example, the convex combination of the points $(5, 3)$ and $(3, 5)$ with $\lambda = \frac{1}{2}$, that is, the point:

$$\left(\frac{1}{2} \cdot 5 + \frac{1}{2} \cdot 3, \frac{1}{2} \cdot 3 + \frac{1}{2} \cdot 5 \right) = (4, 4)$$


does not belong to the set.

Definition 5: epigraph

Consider a function $f : I \rightarrow \mathbb{R}$. The **epigraph** of f is the set:

$$\text{epi } f = \{(x, y) \in \mathbb{R}^2 : x \in I \text{ and } y \geq f(x)\}$$


Example




Definition 6: convex (concave) function

*A function $f : I \rightarrow \mathbb{R}$ is **convex** on I if its epigraph is a convex set. A function is **concave** on I if $-f$ is convex on I .*

Example



Example




Definition 7: convex (concave) function – equivalent definition

*A function $f : I \rightarrow \mathbb{R}$ is **convex** (**concave**) on I if for every pair of points $x_1, x_2 \in I$ the segment (“chord”) with endpoints $(x_1, f(x_1))$, $(x_2, f(x_2))$ has no points below (above) the graph of f .*

The definition of convex function translates into the analytic inequality:

$$f(\lambda x_1 + (1 - \lambda) x_2) \leq \lambda f(x_1) + (1 - \lambda) f(x_2), \quad \forall x_1, x_2 \in I, 0 \leq \lambda \leq 1$$



- ▶ If in the previous inequality the strict $<$ always holds (with $\lambda \neq 0, 1$), the function is called **strictly convex**. For concave (or strictly concave) functions an analogous inequality holds with the sign $\geq$ (or $>$).

- Observe that as $0 \leq \lambda \leq 1$ varies: the point $\lambda x_1 + (1 - \lambda) x_2$ runs along the segment $[x_1, x_2]$ on the x -axis; the point $\lambda f(x_1) + (1 - \lambda) f(x_2)$ runs along the segment $[f(x_1), f(x_2)]$ on the y -axis; the point:

$$\left(\lambda x_1 + (1 - \lambda) x_2, f(\lambda x_1 + (1 - \lambda) x_2) \right)$$

runs along the graph of the function; the point:

$$(\lambda x_1 + (1 - \lambda) x_2, \lambda f(x_1) + (1 - \lambda) f(x_2))$$


runs along the segment with endpoints $(x_1, f(x_1))$ and $(x_2, f(x_2))$.

Example (1/2)

Consider the convex function $f(x) = (x - 2)^2 + 1$ with domain $[\frac{1}{2}, 3]$ and take, for example, $x_1 = 1$ and $x_2 = \frac{5}{2}$ and $\lambda = \frac{1}{3}$. We have the following convex (linear) combination:

$$\left(\frac{1}{3} \cdot 1 + \frac{2}{3} \cdot \frac{5}{2} , \frac{1}{3} \cdot 2 + \frac{2}{3} \cdot \frac{5}{4} \right) = \left(2 , \frac{3}{2} \right)$$

Example (2/2)



- Note that the definition of convex function does not require a priori that the function be continuous or differentiable on an interval.

Theorem 1


A convex (or concave) function on an interval I is continuous on I , except possibly at the endpoints of the interval. Moreover, it has a right and a left derivative at every interior point of the interval.


Proof

Omitted



- ▶ Corner points inside the interval and points of discontinuity at the endpoints of the interval are the only irregular behaviors allowed for a convex or concave function, as shown by the following examples:





Summary

- ▶ **Convex set:** it contains every chord between two of its points, i.e., every convex combination $\lambda P_a + (1 - \lambda)P_b$
- ▶ **Convex function:** convex epigraph; equivalently $f(\lambda x_1 + (1 - \lambda)x_2) \leq \lambda f(x_1) + (1 - \lambda)f(x_2)$
- ▶ f is concave if $-f$ is convex
- ▶ A convex function on an interval is continuous in the interior and has right and left derivatives at interior points

Convexity and derivatives



Introduction: Convexity and derivatives

- ▶ In this section we link convexity to the first and second derivatives
- ▶ The sign of f'' tells us where a function is convex or concave
- ▶ We apply the criterion to exponentials, logarithms and powers

- ▶ If we know a priori that the function is differentiable once or twice on the interval considered, then convexity is related to the first and second derivative of the function.

Theorem 2

Let $f : (a, b) \rightarrow \mathbb{R}$.

- a) If f is differentiable in (a, b) , then f is convex (concave) on (a, b) if and only if f' is increasing (decreasing) on (a, b) .
- b) If f is twice differentiable in (a, b) , then f is convex (concave) on (a, b) if and only if:


$$f''(x) \geq 0 \ (\leq 0), \quad \forall x \in (a, b)$$

- ▶ The theorem is modified in the obvious way for strictly convex or concave functions.



Proof

We do not prove part (a). Part (b) follows from (a) by the monotonicity test applied to f' . □


As a consequence of this theorem, the study of the sign of the second derivative allows us to decide on the convexity or concavity of a function.



$$f''(x_0) = 0$$



$$f''(x_0) = 0$$



Example

The exponential functions

$$f(x) = a^x$$

are convex on $\mathbb{R}$, for any base $a > 0, a \neq 1$, since:

$$f'(x) = a^x \log a; \quad f''(x) = a^x \log^2 a > 0, \quad \forall x \in \mathbb{R}, \forall a > 0, a \neq 1$$

Example

The logarithmic functions

$$f(x) = \log_a x$$

are concave on $(0, +\infty)$ if $a > 1$ and convex on $(0, +\infty)$ if $0 < a < 1$, since:

$$f'(x) = \frac{1}{x \log a}; \quad f''(x) = -\frac{1}{x^2 \log a} \quad \begin{cases} < 0, \forall x > 0, & \text{if } a > 1 \\ > 0, \forall x > 0, & \text{if } 0 < a < 1 \end{cases}$$

Example (1/2)

The power functions with real exponent α :


$$f(x) = x^\alpha$$

are convex on $(0, +\infty)$ if $\alpha > 1$ or $\alpha < 0$, and concave on $(0, +\infty)$ if $0 < \alpha < 1$, since:

$$f'(x) = \alpha x^{\alpha-1}; \quad f''(x) = \alpha(\alpha-1)x^{\alpha-2}$$

which has, for every $x > 0$, the sign of $\alpha(\alpha-1)$.

Example (2/2)



Summary

- ▶ If f is differentiable: f convex (concave) $\iff f'$ increasing (decreasing)
- ▶ If f is twice differentiable: f convex (concave) $\iff f'' \geq 0$ (≤ 0)
- ▶ a^x is convex; $\log_a x$ is concave for $a > 1$ and convex for $0 < a < 1$; x^α is convex for $\alpha > 1$ or $\alpha < 0$, concave for $0 < \alpha < 1$

Convexity and tangent lines



Introduction: Convexity and tangent lines

- ▶ In this section we see a geometric characterization of convexity through tangent lines
- ▶ A convex function lies above its tangent lines and below its chords
- ▶ This provides useful inequalities, such as $e^x \geq 1 + x$

- ▶ A useful geometric characterization of convexity involves the tangent lines to the graph of the function

Theorem 3

A function $f : (a, b) \rightarrow \mathbb{R}$, differentiable in (a, b) , is convex (concave) on (a, b) if and only if, for any choice of a point $x_0 \in (a, b)$, the graph of f lies on the whole of (a, b) above (below) the graph of its tangent line at $(x_0, f(x_0))$.

Proof

Omitted



Example (1/2)

Consider the function:

$$f(x) = e^x, \quad f'(x) = e^x, \quad f''(x) = e^x \quad \text{is convex on all of } \mathbb{R}$$


The tangent line to the graph of f , for example at $x = 0$, that is, at the point $(0, 1)$, is:

$$y = 1 + x \quad \text{hence} \quad e^x \geq 1 + x, \quad x \in \mathbb{R}$$

The tangent line to the graph of f , for example at $x = 1$, that is, at the point $(1, e)$, is:

$$y = e + e(x - 1) \quad \text{hence} \quad e^x \geq ex, \quad x \in \mathbb{R}$$

Example (2/2)



A (differentiable) convex function lies above its tangent lines and, at the same time, below its chords (definition of convexity).

- ▶ This allows us to conclude that, given any two points on the graph of a convex function, the graph between those two points lies entirely in the **triangle** whose sides are the chord joining them and the tangent lines to the graph at the two points.

Example (1/2)


Consider the function:

$$f(x) = x^2, \quad f'(x) = 2x, \quad f''(x) = 2 \quad \text{is convex on all of } \mathbb{R}$$

The tangent lines to the graph of f , for example at the points $x_1 = -0.5$ and $x_2 = 1$, are:

$$y = \frac{1}{4} - \left(x + \frac{1}{2}\right) \quad \text{and} \quad y = 1 + 2(x - 1)$$

Example (2/2)



Summary

- ▶ A differentiable f is convex (concave) $\iff$ its graph lies above (below) each of its tangent lines
- ▶ Examples: $e^x \geq 1 + x$ and $e^x \geq ex$ for every $x \in \mathbb{R}$
- ▶ Between two points of the graph, a convex function lies between the tangent lines and the chord

Inflection points

Introduction: Inflection points

- ▶ In this section we study the points where a function changes concavity: the inflection points
- ▶ We will see that at an inflection point, if it exists, f'' vanishes, but the converse is false
- ▶ We will see inflection points with horizontal, oblique and vertical tangent

- ▶ The direction of concavity of a function (that is, whether it is convex or concave) can change within its domain; this leads us to the concept of inflection point.

Definition 8: inflection point

*Let $f : (a, b) \rightarrow \mathbb{R}$ be a function and let $x_0 \in (a, b)$ be a point of differentiability for f , or let $f'(x_0) = \pm\infty$. The point x_0 is called an **inflection point** of f if there exist a right neighborhood $(x_0, x_0 + h)$, $h > 0$, in which f is convex (concave) and a left neighborhood $(x_0 - h, x_0)$, $h > 0$, in which f is concave (convex).*

- ▶ When crossing an inflection point, the second derivative of f (if it exists) changes sign. We then expect f'' to vanish at this point.

Theorem 4

Let x_0 be an inflection point of f ; if $f''(x_0)$ exists, then $f''(x_0) = 0$.

Proof

Note that, if we knew that f'' exists in a neighborhood of x_0 **and is continuous** at x_0 , then the thesis of the theorem would follow from the intermediate value theorem for continuous functions (applied to f''). The theorem can also be proved without these additional hypotheses (omitted). □

The converse of the implication stated by the theorem is not true: a point where the second derivative vanishes may not be an inflection point.


Example (1/2)

Consider the function:

$$f(x) = x^4, \quad f'(x) = 4x^3, \quad f''(x) = 12x^2 \quad \text{is convex on all of } \mathbb{R}$$

Since $f'(x) > 0$ for $x > 0$, the function is increasing for $x > 0$, decreasing for $x < 0$ and has a minimum point at $x = 0$. We have $f''(0) = 0$, but the function is convex on all of $\mathbb{R}$, hence $x = 0$ is **not an inflection point**.

Example (2/2)



- ▶ The geometric meaning of inflection points is clarified by the following theorem.

Theorem 5

If $f : (a, b) \rightarrow \mathbb{R}$ is differentiable in (a, b) and $x_0 \in (a, b)$ is an inflection point, then the graph of $f(x)$ crosses its tangent line at $(x_0, f(x_0))$.

Proof

Let us draw the tangent line to the graph of $f(x)$ at the point with abscissa x_0 . If f is (for example) concave on (a, x_0) , since f is differentiable, the graph of f lies below the line on (a, x_0) ; on the other hand f is convex on (x_0, b) , so its graph lies above the line on (x_0, b) . Consequently, at x_0 the graph crosses the tangent line. $\square$

Example (1/2)


Consider the function:

$$f(x) = x^3, \quad f'(x) = 3x^2, \quad f''(x) = 6x$$

Since $f'(x) > 0$ for $x \neq 0$, the function is increasing on all of $\mathbb{R}$. It has a stationary point at $x = 0$, which, however, is not a maximum or minimum point, because the function is always increasing.

We have $f''(0) = 0$. Moreover, $f''(x) < 0$ for $x < 0$, hence it is concave for $x < 0$, and $f''(x) > 0$ for $x > 0$, hence it is convex for $x > 0$. Consequently, the point x_0 is **an inflection point with horizontal tangent**, and the graph of the function crosses its tangent line at $(0, 0)$, that is, the line $y = 0$.

Example (2/2)



Example (1/3)

Consider the function:

$$f(x) = e^{-x^2}, \quad f'(x) = -2x e^{-x^2}, \quad f''(x) = e^{-x^2} (4x^2 - 2)$$

We have $f'(x) > 0$ for $x < 0$, hence the function increases for $x \leq 0$; and $f'(x) < 0$ for $x > 0$, hence it decreases for $x \geq 0$, and moreover $f'(0) = 0$. Therefore it has a relative maximum point at $x = 0$.

We have:

$$f''(x) = e^{-x^2} (4x^2 - 2) \geq 0 \quad \text{for } x^2 \geq \frac{1}{2}$$

$$\text{that is } x \in \left(-\infty, -\frac{1}{\sqrt{2}}\right] \cup \left[\frac{1}{\sqrt{2}}, +\infty\right)$$

The function is convex for these values, and concave for $-\frac{1}{\sqrt{2}} \leq x \leq \frac{1}{\sqrt{2}}$. Hence it has inflection points at $x = \pm \frac{1}{\sqrt{2}}$, with tangent line of slope:

Example (2/3)


$$f' \left(\pm \frac{1}{\sqrt{2}} \right) = \mp \sqrt{\frac{2}{e}}$$

$$\text{since } -2 \left(\pm \frac{1}{\sqrt{2}} \right) e^{-\left(\pm \frac{1}{\sqrt{2}} \right)^2} = \mp \sqrt{2} \cdot \frac{1}{\sqrt{e}}$$

The tangent lines are:

$$y = \frac{1}{\sqrt{e}} + \sqrt{\frac{2}{e}} \left(x + \frac{1}{\sqrt{2}} \right) \quad \text{and} \quad y = \frac{1}{\sqrt{e}} - \sqrt{\frac{2}{e}} \left(x - \frac{1}{\sqrt{2}} \right)$$

Example (3/3)



Example (1/2)


Consider the function:

$$f(x) = x^{1/3}, \quad f'(x) = \frac{1}{3x^{2/3}},$$

$$\text{for } x > 0, \quad f''(x) = -\frac{2}{9x^{5/3}}, \quad \text{for } x > 0$$

For $x_0 = 0$, we have seen that $f'(0) = +\infty$. In this case, however, $f''(0)$ does not exist and the function has an inflection point with vertical tangent.

Example (2/2)




Example (1/2)

Consider the function:

$$f(x) = x|x|, \text{ with } x \neq 0, \quad f(x) = \begin{cases} x^2, & x > 0 \\ -x^2, & x < 0 \end{cases} \quad f'(x) = \begin{cases} 2x, & x > 0 \\ -2x, & x < 0 \end{cases} \quad f''(x) = \begin{cases} 2, & x > 0 \\ -2, & x < 0 \end{cases}$$

Hence, for $x > 0$ the function is convex and for $x < 0$ it is concave, and $f'(0) = 0$; therefore x_0 is an inflection point (but $f''(0)$ does not exist).

Example (2/2)



Summary

- ▶ **Inflection point:** point where f changes from convex to concave (or vice versa)
- ▶ If x_0 is an inflection point and $f''(x_0)$ exists, then $f''(x_0) = 0$; the converse is false (x^4 at $x = 0$)
- ▶ At an inflection point the graph crosses its tangent line
- ▶ Examples: x^3 (horizontal tangent), e^{-x^2} (oblique tangent), $x^{1/3}$ (vertical tangent), $x|x|$ ($f''(0)$ does not exist)

Final summary

- ▶ The second derivative measures the change of the slope and is related to the curvature of the graph
- ▶ Convexity and concavity are defined via chords and the epigraph and are studied through the sign of f''
- ▶ A convex function lies above its tangent lines and below its chords
- ▶ At inflection points the concavity changes and the graph crosses its tangent line