

Invertible functions and inverse functions



Outline

Invertible functions and inverse functions

Graph of the inverse function

Inverse power functions

Inverse trigonometric functions

Inverse hyperbolic functions

Invertible functions and inverse functions



Introduction: Invertible functions and inverse functions

- ▶ A function associates to each input x a unique output $y = f(x)$: we ask when it is possible to “go back”, that is, to recover from y the unique x that produces it
- ▶ In this section we define **invertible functions** and the **inverse function**, with a graphical criterion to recognize them
- ▶ We will prove that **strictly monotone** functions are invertible, and we will see that this is not a necessary condition

Invertible functions

- ▶ Given a real function of a real variable $f : D \rightarrow \mathbb{R}$, for every input x in the domain D there exists a unique output $y = f(x)$ in the image of the domain $f(D)$.
- ▶ If, in addition, for every output $y = f(x) \in f(D)$ there exists a unique input $x \in D$, then f is called **invertible**, and it establishes a **one-to-one correspondence** between the domain D of f and the image of the domain $f(D)$.

Definition 1: of invertible function

A function $f : D \rightarrow \mathbb{R}$ is **invertible** on the domain D if one of the following equivalent conditions holds:

$$\forall x_1, x_2 \in D, \quad x_1 \neq x_2 \implies f(x_1) \neq f(x_2)$$

$$\forall x_1, x_2 \in D, \quad f(x_1) = f(x_2) \implies x_1 = x_2$$

$$\forall y \in f(D), \quad \exists! x \in D : y = f(x)$$

that is, if f is **injective**.

Warning

- ▶ **WARNING:** the definition of invertible function given here for real functions of a real variable is different from the usual one given for functions between arbitrary sets, which also requires surjectivity.
- ▶ In this case *injectivity is enough* because we define the inverse on the image of f and not on its whole codomain.

Definition 2: of inverse function


*Given an invertible function $f : D \rightarrow \mathbb{R}$, the function that associates to each output $y \in f(D)$ the unique input $x \in D$ such that $f(x) = y$ is called the **inverse function** of f and is denoted by the symbol f^{-1} .*

The pair f and f^{-1}

The pair f and f^{-1} is written as:

$$f : \begin{cases} y = f(x) \\ x \in D \end{cases} \quad f^{-1} : \begin{cases} x = f^{-1}(y) \\ y \in f(D) \end{cases}$$


The black box of f^{-1} works backwards with respect to that of f , according to the following scheme:



Graphical criterion for invertibility


- ▶ The invertibility condition is equivalent to requiring that the graph of f be intersected in at most one point by every line parallel to the x -axis.
- ▶ We now see an example of the graph of an invertible function and one of a non-invertible function

Example



Graph of a function that is **invertible** on $[a, b]$, since every line parallel to the x -axis either does not intersect the graph of f or intersects it in exactly one point.

Example



Graph of a function that is **not invertible** on $[a, b]$, since for the indicated value $\tilde{y}$ there are several values x (precisely x_1 , x_2 and x_3) whose image is $\tilde{y}$.


Strictly monotone functions

Theorem 1


If a function $f : D \rightarrow \mathbb{R}$ is strictly increasing (decreasing) on D , then it is invertible on D . Moreover, its inverse function is strictly increasing (decreasing).

Graphical interpretation (1/2)

For the strictly increasing case, graphically we have:



Graphical interpretation (2/2)



Proof (1/2)

Consider the case of a function that is strictly increasing on D and two arbitrary values $x_1, x_2 \in D$ (the strictly decreasing case is analogous).

If $x_1 \neq x_2$, then either $x_1 < x_2$ or $x_1 > x_2$. Since f is strictly increasing, we have:

$$\text{if } x_1 < x_2, \text{ then } f(x_1) < f(x_2)$$

$$\text{if } x_1 > x_2, \text{ then } f(x_1) > f(x_2)$$

In both cases $f(x_1) \neq f(x_2)$, therefore f is invertible.

Proof (2/2)

Consider the inverse function f^{-1} , that is, $x = f^{-1}(y)$, and let us prove that it is strictly increasing. Consider two arbitrary values $y_1, y_2 \in f(D)$, with $y_1 < y_2$.

If we had $f^{-1}(y_1) = x_1 \geq x_2 = f^{-1}(y_2)$, since f is strictly increasing we would have $y_1 = f(x_1) \geq f(x_2) = y_2$, that is, $y_1 \geq y_2$. This cannot happen, since it contradicts $y_1 < y_2$ (**contradiction**).

Consequently, we have $f^{-1}(y_1) = x_1 < x_2 = f^{-1}(y_2)$, that is, $f^{-1}(y_1) < f^{-1}(y_2)$, and hence f^{-1} is strictly increasing. □


Strict monotonicity is not necessary

- ▶ However, a function can be invertible even without being strictly increasing or decreasing.

Example

For example, the following piecewise-defined function is invertible:

$$f(x) = \begin{cases} \frac{1}{2}x & \text{if } -1 \leq x < 0, \\ \frac{1}{2} - \frac{1}{2}x & \text{if } 0 \leq x \leq \frac{2}{3} \end{cases}$$



Non-invertible functions

- ▶ Two classes of functions that are certainly not invertible are:
 1. even (symmetric) functions, since:

$$f(-x) = f(x)$$

2. periodic functions, since:

$$f(x + T) = f(x)$$

Summary

Invertible and inverse functions:

- ▶ f is invertible on D if it is injective: equal outputs correspond to equal inputs
- ▶ The inverse f^{-1} is defined on the image $f(D)$ and associates to each y the unique x with $f(x) = y$
- ▶ Graphical criterion: every horizontal line intersects the graph in at most one point

Monotonicity:

- ▶ Strictly monotone $\Rightarrow$ invertible, with inverse monotone of the same type (the converse does not hold)
- ▶ Even functions and periodic functions are not invertible

Graph of the inverse function



Introduction: Graph of the inverse function

- ▶ In this section we see how to obtain the **graph** of f^{-1} from that of f : a symmetry with respect to the bisector $y = x$ is enough
- ▶ We then see how to find the **analytic expression** of the inverse by solving the equation $f(x) = y$ for x
- ▶ Examples: affine function, x^2 , exponential; and a case in which the inverse exists but cannot be written explicitly

Symmetry with respect to the bisector


- ▶ The relations between an invertible function f and its inverse function f^{-1} :

$$f : \begin{cases} y = f(x) \\ x \in D \end{cases} \qquad f^{-1} : \begin{cases} x = f^{-1}(y) \\ y \in f(D) \end{cases}$$

indicate that if the point (x_0, y_0) lies on the graph of f , then the point (y_0, x_0) lies on the graph of f^{-1} .

- ▶ Since the points (x_0, y_0) and (y_0, x_0) are **symmetric with respect to the bisector** with equation $y = x$, the graph of f^{-1} is obtained from that of f by symmetry with respect to the bisector.

Symmetry with respect to the bisector



Analytic expression of the inverse

- ▶ If the analytic expression of f is known and f is invertible, the **analytic expression** of f^{-1} is found by trying to solve for x the equation:

$f(x) = y$ that is, find the input x that produces the output y

- ▶ Note that it is not always possible to find it, even in cases where the inverse function exists!

Inverse of an affine function

Given $a, b \in \mathbb{R}$, $a \neq 0$, we have:

$$f : \begin{cases} y = ax + b \\ x \in \mathbb{R} \end{cases}$$

$$f^{-1} : \begin{cases} x = \frac{y-b}{a} \\ y \in \mathbb{R} \end{cases}$$

Example (1/3)

The function $f : \mathbb{R} \rightarrow \mathbb{R}, x \mapsto 2x + 3$ is strictly increasing, hence invertible on $\mathbb{R}$, and the equation


$$2x + 3 = y \text{ solved for } x, \quad \text{that is } x = \frac{y - 3}{2}$$

gives for f^{-1} the analytic expression:

$$f^{-1}(y) = \frac{y - 3}{2}$$


that is, the function: $f^{-1} : \mathbb{R} \rightarrow \mathbb{R}, y \mapsto \frac{y-3}{2}$.

Example (2/3)



Example (3/3)

The graphs $y = 2x + 3$ and $y = \frac{x-3}{2}$ are symmetric with respect to the bisector $y = x$



Inverse of x^2

We have:

$$f : \begin{cases} y = x^2 \\ x \geq 0 \end{cases}$$

$$f^{-1} : \begin{cases} x = \sqrt{y} \\ y \geq 0 \end{cases}$$

Example (1/2)

The function $f : \mathbb{R} \rightarrow \mathbb{R}, x \mapsto x^2$ is strictly increasing on the interval $[0, +\infty)$, hence invertible there; the equation

$$x^2 = y \text{ solved for } x, \text{ that is } x = \sqrt{y}$$


gives for f^{-1} the analytic expression

$$f^{-1}(y) = \sqrt{y}$$

that is, the function: $f^{-1} : [0, +\infty) \rightarrow \mathbb{R}, y \mapsto \sqrt{y}$.

Example (2/2)

The graphs $y = x^2$ and $y = \sqrt{x}$ are symmetric with respect to the bisector $y = x$



Inverse of the exponential

Given $a \in \mathbb{R}_+$, $a \neq 1$, we have:

$$f : \begin{cases} y = a^x \\ x \in \mathbb{R} \end{cases}$$

$$f^{-1} : \begin{cases} x = \log_a y \\ y > 0 \end{cases}$$

Example (1/2)

The function $f : \mathbb{R} \rightarrow \mathbb{R}, x \mapsto e^x$ is strictly increasing on all of $\mathbb{R}$ and hence invertible; the equation

$$e^x = y \text{ solved for } x, \text{ that is } x = \log y$$


gives for f^{-1} the analytic expression

$$f^{-1}(y) = \log y$$

that is, the function: $f^{-1} : (0, +\infty) \rightarrow \mathbb{R}, y \mapsto \log y$.

Example (2/2)

The graphs $y = e^x$ and $y = \log x$ are symmetric with respect to the bisector $y = x$



An inverse that cannot be written explicitly

- ▶ Now consider the function:

$$f(x) = x + e^x$$

Being the sum of two functions that are strictly increasing on all of $\mathbb{R}$, $f(x)$ is strictly increasing and hence invertible on all of $\mathbb{R}$ (see the Theorem on strictly monotone functions).

- ▶ However, we would try in vain to solve the equation $x + e^x = y$ for x . In other words, f^{-1} exists, but we cannot write it explicitly.

Summary

Graph and expression of the inverse:

- ▶ (x_0, y_0) lies on the graph of f if and only if (y_0, x_0) lies on the graph of f^{-1} : the graphs are symmetric with respect to the bisector $y = x$
- ▶ The expression of f^{-1} is found by solving $f(x) = y$ for x

Examples:

- ▶ $y = ax + b \Leftrightarrow x = \frac{y-b}{a}$; $y = x^2 (x \geq 0) \Leftrightarrow x = \sqrt{y}$;
 $y = a^x \Leftrightarrow x = \log_a y$
- ▶ $f(x) = x + e^x$ is invertible, but f^{-1} cannot be written explicitly

Inverse power functions



Introduction: Inverse power functions

- ▶ In this section we study the inverse of the **power** functions $y = x^\alpha$
- ▶ The inverse of a power is again a power, with exponent $\frac{1}{\alpha}$ (the **roots**)
- ▶ The domain on which the power is invertible depends on the exponent: even powers must be restricted to $x \geq 0$, odd powers are invertible on all of $\mathbb{R}$

Powers and their inverses (1/2)

Given $\alpha \in \mathbb{R}, \alpha \neq 0$, we have:

$$f : \begin{cases} y = x^\alpha \\ x > 0 \end{cases} \qquad f^{-1} : \begin{cases} x = y^{\frac{1}{\alpha}} \\ y > 0 \end{cases}$$

If $\alpha > 0$ we have:

$$f : \begin{cases} y = x^\alpha \\ x \geq 0 \end{cases} \qquad f^{-1} : \begin{cases} x = y^{\frac{1}{\alpha}} \\ y \geq 0 \end{cases}$$

Powers and their inverses (2/2)

If $\alpha = \frac{m}{n} \in \mathbb{Q}$, with $m \in \mathbb{Z}$, $n \in \mathbb{N}_+$ odd and coprime, we have:


$$f : \begin{cases} y = x^{\frac{m}{n}} \\ x \in \mathbb{R} \end{cases} \qquad f^{-1} : \begin{cases} x = y^{\frac{n}{m}} \\ y \in \mathbb{R} \end{cases}$$

Even powers

The **even powers**:


$$x^{2n} \quad \text{with } n = 1, 2, \dots$$

are not invertible on the whole real line, but only on the half-line $x \geq 0$.



Odd powers

The **odd powers** x^{2n+1} (with $n = 1, 2, \dots$) and the **powers with rational exponent** $x^{\frac{m}{n}}$ (with n, m positive odd integers), being strictly increasing monotone functions, are invertible from $-\infty$ to $+\infty$.



Summary

Inverse powers:

- ▶ The inverse of $y = x^\alpha$ ($x > 0$) is $x = y^{\frac{1}{\alpha}}$; if $\alpha > 0$ this also holds on $[0, +\infty)$
- ▶ Even powers x^{2n} : invertible only on $x \geq 0$ (inverses: $x^{\frac{1}{2}}, x^{\frac{1}{4}}, \dots$)
- ▶ Odd powers x^{2n+1} and rational exponents with m, n odd: strictly increasing, invertible on all of $\mathbb{R}$ (inverses: $x^{\frac{1}{3}}, x^{\frac{1}{5}}, \dots$)

Inverse trigonometric functions

Introduction: Inverse trigonometric functions

- ▶ The trigonometric functions are periodic and hence **not invertible**: to invert them we restrict them to intervals on which they are strictly monotone
- ▶ In this way we obtain the **arcsine**, the **arccosine** and the **arctangent**, with their graphs
- ▶ These functions are used to write the solutions of trigonometric equations and inequalities with non-standard angles

Trigonometric functions are not invertible

- ▶ Being periodic, the trigonometric functions cannot be invertible. Indeed, for example, the equation

$$y = \sin x$$

has infinitely many solutions if $-1 \leq y \leq 1$ (the output y corresponds to infinitely many inputs), or has no real solutions if $|y| > 1$.

- ▶ To speak of inverse functions of sine, cosine and tangent, we will need to restrict ourselves to intervals on which these functions are strictly monotone and therefore invertible.

Arcsine

- An interval on which the sine function is invertible is $[-\frac{\pi}{2}, \frac{\pi}{2}]$.

Definition 3: of arcsine

*The inverse function of the sine on the interval $[-\frac{\pi}{2}, \frac{\pi}{2}]$ is the **arcsine**:*


$$f : [-1, 1] \rightarrow \left[-\frac{\pi}{2}, \frac{\pi}{2}\right], \quad f : y \mapsto \arcsin y$$

We have:

$$f : \begin{cases} y = \sin x \\ x \in [-\frac{\pi}{2}, \frac{\pi}{2}] \end{cases} \qquad f^{-1} : \begin{cases} x = \arcsin y \\ y \in [-1, 1] \end{cases}$$

Graph of the arcsine

The graph of the arcsine is obtained from the arc of the sine curve restricted to the interval $[-\frac{\pi}{2}, \frac{\pi}{2}]$ (strictly monotone), by symmetry with respect to the bisector $y = x$:



Arccosine

- An interval on which the cosine function is invertible is $[0, \pi]$.

Definition 4: of arccosine

*The inverse function of the cosine on the interval $[0, \pi]$ is the **arccosine**:*


$$f : [-1, 1] \rightarrow [0, \pi], \quad f : y \mapsto \arccos y$$

We have:

$$f : \begin{cases} y = \cos x \\ x \in [0, \pi] \end{cases} \qquad f^{-1} : \begin{cases} x = \arccos y \\ y \in [-1, 1] \end{cases}$$

Graph of the arccosine

The graph of the arccosine is obtained from the arc of the cosine curve restricted to the interval $[0, \pi]$ (strictly monotone), by symmetry with respect to the bisector $y = x$:



Arctangent

- Note that on the interval $(-\frac{\pi}{2}, \frac{\pi}{2})$ the tangent is strictly monotone and hence invertible. Its inverse function is called the arctangent ($\arctan$), and it is defined on $\mathbb{R}$.

Definition 5: of arctangent

*The inverse function of the tangent on the interval $(-\frac{\pi}{2}, \frac{\pi}{2})$ is the **arctangent**:*

$$f : \mathbb{R} \rightarrow \left(-\frac{\pi}{2}, \frac{\pi}{2}\right), \quad f : y \mapsto \arctan y$$

Arctangent


We have:

$$f : \begin{cases} y = \tan x \\ x \in \left(-\frac{\pi}{2}, \frac{\pi}{2}\right) \end{cases}$$

$$f^{-1} : \begin{cases} x = \arctan y \\ y \in \mathbb{R} \end{cases}$$

- ▶ The graph of the arctangent is obtained from that of the tangent restricted to the interval $\left(-\frac{\pi}{2}, \frac{\pi}{2}\right)$ (strictly monotone), by symmetry with respect to the bisector $y = x$.

Graph of the arctangent



Trigonometric equations and inequalities

- ▶ By means of the inverse trigonometric functions, we can express the solutions of a **trigonometric equation or inequality** when it involves angles that are not standard angles.


Example (1/3)

The solutions of:

$$\sin x = \frac{1}{4}$$

are:

$$(i) \ x = \arcsin \frac{1}{4} + 2k\pi; \quad (ii) \ x = \pi - \arcsin \frac{1}{4} + 2k\pi \quad (k \in \mathbb{Z})$$




Example (2/3)

The solutions of:

$$\cos x < \frac{1}{5}$$

are:

$$\arccos \frac{1}{5} + 2k\pi < x < 2\pi - \arccos \frac{1}{5} + 2k\pi \quad (k \in \mathbb{Z})$$



Example (3/3)

The solutions of:

$$\tan x \geq 3$$

are:

$$\arctan 3 + k\pi \leq x < \frac{\pi}{2} + k\pi \quad (k \in \mathbb{Z})$$

Summary

Inverse trigonometric functions:

- ▶ $\arcsin : [-1, 1] \rightarrow [-\frac{\pi}{2}, \frac{\pi}{2}]$, inverse of the sine restricted to $[-\frac{\pi}{2}, \frac{\pi}{2}]$
- ▶ $\arccos : [-1, 1] \rightarrow [0, \pi]$, inverse of the cosine restricted to $[0, \pi]$
- ▶ $\arctan : \mathbb{R} \rightarrow (-\frac{\pi}{2}, \frac{\pi}{2})$, inverse of the tangent restricted to $(-\frac{\pi}{2}, \frac{\pi}{2})$

Application:

- ▶ They allow us to write the solutions of trigonometric equations and inequalities with non-standard angles

Inverse hyperbolic functions



Introduction: Inverse hyperbolic functions

- ▶ In this section we invert the **hyperbolic sine** and the **hyperbolic cosine** (the latter restricted to $x \geq 0$)
- ▶ Unlike the trigonometric functions, here we can find the **analytic expression** of the inverse, by solving a quadratic equation in e^x
- ▶ We obtain the **inverse hyperbolic sine** and the **inverse hyperbolic cosine**, expressed through the logarithm

Inverse hyperbolic sine (1/3)

- Consider the **hyperbolic sine** function:

$$y = \operatorname{Sh} x = \frac{e^x - e^{-x}}{2}$$

It is defined and strictly increasing on all of $\mathbb{R}$, therefore it is invertible.

- To solve the equation for x , multiplying both sides by e^x we obtain:

$$\begin{aligned} 0 &= e^x y - e^x \frac{e^x - e^{-x}}{2} \\ &= 2 e^x y - (e^x e^x - \underbrace{e^x e^{-x}}_{=e^{x-x}=e^0}) \\ &= 2 y e^x - e^{2x} + 1 \\ &= e^{2x} - 2 y e^x - 1 \end{aligned}$$

Inverse hyperbolic sine (2/3)

- ▶ This is a quadratic equation in the unknown e^x . Setting $t = e^x$ we obtain

$$0 = t^2 - 2 y t - 1$$

We obtain:

$$t = y \pm \sqrt{y^2 + 1} \quad \text{and substituting back} \quad e^x = y \pm \sqrt{y^2 + 1}$$

since $e^x > 0$, the solution with the minus sign must be discarded. Hence we are left with:

$$e^x = \underbrace{y + \sqrt{y^2 + 1}}_{>0, \forall y \in \mathbb{R}} \quad \text{and hence} \quad x = \log \left(y + \sqrt{y^2 + 1} \right)$$

Inverse hyperbolic sine (3/3)

- This is the analytic expression of the inverse function of $\text{Sh } x$, which is called the **inverse hyperbolic sine** (area hyperbolic sine), and is also denoted by SettSh . It is defined for every real y .

Definition 6: of inverse hyperbolic sine

*The inverse function of the hyperbolic sine is the **inverse hyperbolic sine**:*


$$f : \mathbb{R} \rightarrow \mathbb{R}, \quad f : y \mapsto \text{SettSh } y$$

We have:

$$f : \begin{cases} y = \text{Sh } x \\ x \in \mathbb{R} \end{cases} \qquad f^{-1} : \begin{cases} x = \text{SettSh } y \\ y \in \mathbb{R} \end{cases}$$

Graph of the inverse hyperbolic sine

The graph of the inverse hyperbolic sine is obtained from that of the hyperbolic sine on $\mathbb{R}$ (strictly monotone), by symmetry with respect to the bisector $y = x$:



Inverse hyperbolic cosine (1/3)

- Consider the **hyperbolic cosine** function:

$$y = \operatorname{Ch} x = \frac{e^x + e^{-x}}{2}$$

It is defined on all of $\mathbb{R}$, strictly increasing for $x \geq 0$, decreasing for $x \leq 0$. Therefore it is not invertible on all of $\mathbb{R}$.

- Its restriction to $x \geq 0$, however, is invertible. We want to determine the analytic expression of the inverse function of this restriction.
- Proceeding as before, we obtain:

$$e^x = y \pm \sqrt{y^2 - 1}$$

Inverse hyperbolic cosine (2/3)

- This time both numbers $y \pm \sqrt{y^2 - 1}$ are positive; recall, however, that we are reasoning only for $x \geq 0$, which is equivalent to choosing the plus sign. Therefore:

$$x = \log \left(y + \sqrt{y^2 - 1} \right)$$

This is the analytic expression of the inverse function of $\text{Ch } x$, which is called the inverse hyperbolic cosine (area hyperbolic cosine), and is also denoted by SettCh . Note that it is defined for $y \geq 1$.

Inverse hyperbolic cosine (3/3)

Definition 7: of inverse hyperbolic cosine

*The inverse function of the hyperbolic cosine on the interval $[0, +\infty)$ is the **inverse hyperbolic cosine**:*

$$f : [1, +\infty) \rightarrow [0, +\infty), \quad f : y \mapsto \operatorname{SettCh} y$$


We have:

$$f : \begin{cases} y = \operatorname{Ch} x \\ x \in [0, +\infty) \end{cases}$$

$$f^{-1} : \begin{cases} x = \operatorname{SettCh} y \\ y \in [1, +\infty) \end{cases}$$

Graph of the inverse hyperbolic cosine

The graph of the inverse hyperbolic cosine is obtained from that of the hyperbolic cosine on $[0, +\infty)$ (strictly monotone), by symmetry with respect to the bisector $y = x$.



Example (1/2)

- ▶ The equation:

$$\operatorname{Sh} x = 2$$

has the unique solution:

$$x = \operatorname{SettSh} 2 = \log(2 + \sqrt{5})$$


- ▶ The equation:

$$\operatorname{Ch} x = 3$$

has two solutions:

$$x = \pm \operatorname{SettCh} 3 = \pm \log(3 + 2\sqrt{2})$$

Example (2/2)



Inverse hyperbolic functions:

- ▶ Sh is strictly increasing on $\mathbb{R}$: its inverse is the inverse hyperbolic sine $\text{SettSh } y = \log \left(y + \sqrt{y^2 + 1} \right)$, defined for every $y \in \mathbb{R}$
- ▶ Ch is invertible only if restricted to $x \geq 0$: its inverse is the inverse hyperbolic cosine $\text{SettCh } y = \log \left(y + \sqrt{y^2 - 1} \right)$, defined for $y \geq 1$
- ▶ The expressions are obtained by solving a quadratic equation in $t = e^x$

Summary (1/2)

Invertible functions:

- ▶ f is invertible if it is injective; the inverse f^{-1} is defined on $f(D)$
- ▶ Strictly monotone $\Rightarrow$ invertible; even or periodic $\Rightarrow$ not invertible
- ▶ The graph of f^{-1} is symmetric to that of f with respect to the bisector $y = x$
- ▶ The expression of f^{-1} is obtained by solving $f(x) = y$ for x (not always possible)

Summary (2/2)

Inverses of the elementary functions:

- ▶ Powers: the inverse of x^α is $y^{\frac{1}{\alpha}}$ (even powers only for $x \geq 0$)
- ▶ Exponential and logarithm: $y = a^x \leftrightarrow x = \log_a y$
- ▶ Trigonometric (on intervals of monotonicity): arcsin, arccos, arctan
- ▶ Hyperbolic: $\text{SettSh } y = \log \left(y + \sqrt{y^2 + 1} \right),$
 $\text{SettCh } y = \log \left(y + \sqrt{y^2 - 1} \right)$