


Comparisons and asymptotic estimates



- **Author:** Fabio Furini.
- **Book:** “Analisi matematica 1”, Marco Bramanti, Carlo D. Pagani and Sandro Salsa, Zanichelli.

Contents

1 Comparisons and asymptotic estimates

2

1 Comparisons and asymptotic estimates

- We have seen that a sequence tending to 0 is an **infinitesimal**; a sequence that diverges (to $+\infty$, to $-\infty$) is called an **infinity** (infinite quantity).
- When two sequences are both infinitesimals or both infinities, it is useful to be able to compare them, to understand which of the two tends “**more rapidly**” to 0 or to infinity.

Example 1: Infinities

Examples of infinities are the following sequences:

$$\{\log n\}, \quad \{\sqrt{n}\}, \quad \{n^2\}, \quad \{2^n\}$$

Example 2: Infinitesimals

Examples of infinitesimals are the following sequences:

$$\left\{\frac{1}{\log n}\right\}, \quad \left\{\frac{1}{\sqrt{n}}\right\}, \quad \left\{\frac{1}{n^2}\right\}, \quad \left\{\frac{1}{2^n}\right\}$$

- Let $\{a_n\}$ and $\{b_n\}$ be **two infinities**; considering the limit of their ratio we have 4 cases:

$$\lim_{n \rightarrow +\infty} \frac{a_n}{b_n} = \begin{cases} 0 & \text{case 1): } \{a_n\} \text{ is an infinity of } \mathbf{lower\ order} \text{ than } \{b_n\} \\ l \in \mathbb{R}, l \neq 0 & \text{case 2): } \{a_n\} \text{ and } \{b_n\} \text{ are infinities of the } \mathbf{same\ order} \\ \pm\infty & \text{case 3): } \{a_n\} \text{ is an infinity of } \mathbf{higher\ order} \text{ than } \{b_n\} \\ \text{nonexistent} & \text{case 4): } \{a_n\} \text{ and } \{b_n\} \text{ are not comparable} \end{cases}$$

- Let $\{a_n\}$ and $\{b_n\}$ be **two infinitesimals** (with b_n eventually different from zero); considering the limit of their ratio we have 4 cases:

$$\lim_{n \rightarrow +\infty} \frac{a_n}{b_n} = \begin{cases} 0 & \text{case 1): } \{a_n\} \text{ is an infinitesimal of } \mathbf{higher\ order} \text{ than } \{b_n\} \\ l \in \mathbb{R}, l \neq 0 & \text{case 2): } \{a_n\} \text{ and } \{b_n\} \text{ are infinitesimals of the } \mathbf{same\ order} \\ \pm\infty & \text{case 3): } \{a_n\} \text{ is an infinitesimal of } \mathbf{lower\ order} \text{ than } \{b_n\} \\ \text{nonexistent} & \text{case 4): } \{a_n\} \text{ and } \{b_n\} \text{ are not comparable} \end{cases}$$

- The case

$$\frac{a_n}{b_n} \rightarrow 1$$

is particularly important: in this case we say that the two sequences $\{a_n\}$ and $\{b_n\}$ are **asymptotic** (asymptotically equivalent).

- To indicate this, we write

$$a_n \sim b_n$$

(read: a_n is asymptotic to b_n)

- The asymptotic symbol is very useful in computing limits, thanks to the following **properties**:

Proposition 1: Asymptotic behavior

1. If $a_n \sim b_n$, the two sequences have the same behavior:

- either they converge to the same limit,
- or they both diverge to $\pm\infty$,
- or neither of them has a limit.

2. We can write chains of asymptotic relations, i.e.:

$$\text{if } a_n \sim b_n \sim \dots \sim c_n \text{ then } a_n \sim c_n$$

3. An expression made of a product or quotient of several factors can be estimated factor by factor:

$$\text{if } a_n \sim a'_n, b_n \sim b'_n, c_n \sim c'_n \text{ then } \frac{a_n b_n}{c_n} \sim \frac{a'_n b'_n}{c'_n}$$

- **Warning:** the same does not hold for sums or for exponentials.

Proof. 1. We prove the first statement

if $a_n \sim b_n$ then $\{a_n\}$ and $\{b_n\}$ have the same behavior

- If $a_n \rightarrow l \in \mathbb{R}$, since

$$b_n = \frac{b_n}{a_n} \cdot a_n \text{ and } \frac{b_n}{a_n} \rightarrow 1 \text{ (by definition of asymptotic),}$$

then by the theorem on the algebra of limits we have

$$b_n \rightarrow 1 \cdot l = l.$$

- With the same steps, the theorem on the partial arithmetization of the infinity symbol allows us to conclude that

$$\text{if } a_n \rightarrow \pm\infty \text{ and } a_n \sim b_n \text{ then } b_n \rightarrow \pm\infty.$$

Observing that the asymptotic relation is symmetric, what we have just proved also shows that if $\{b_n\}$ converges (diverges), then $\{a_n\}$ converges (diverges) as well.

- We conclude that if $\{a_n\}$ is irregular, then $\{b_n\}$ is irregular as well, because if, by contradiction, it were not, then by what we have just proved $\{a_n\}$ would also be convergent or divergent.

2. We prove the transitivity of the asymptotic relation:

$$\text{if } a_n \sim b_n \sim c_n \text{ then } a_n \sim c_n.$$

The hypotheses mean that

$$\frac{a_n}{b_n} \rightarrow 1 \text{ and } \frac{b_n}{c_n} \rightarrow 1$$

Then by the theorem on the algebra of limits we have

$$\frac{a_n}{c_n} = \frac{a_n}{b_n} \cdot \frac{b_n}{c_n} \rightarrow 1.$$

3. The third property is proved analogously.

□

Example 3: Asymptotic sequences by applying the definition

We prove that:

$$\log n \sim \log(n+1)$$

Factoring out n in the second sequence we obtain

$$\log(n+1) = \log\left(n\left(1 + \frac{1}{n}\right)\right) = \log n + \log\left(1 + \frac{1}{n}\right)$$

Hence

$$\lim_{n \rightarrow +\infty} \frac{\log n}{\log(n+1)} = \lim_{n \rightarrow +\infty} \frac{\log n}{\underbrace{\log n + \log\left(1 + \frac{1}{n}\right)}_{\rightarrow 0}} = 1$$

- A typical way to show that

$$a_n \sim b_n$$

consists in writing

$$a_n = b_n c_n \text{ with } c_n \rightarrow 1.$$

That is, decomposing $\{a_n\}$ into the product of a sequence $\{b_n\}$ and a sequence $\{c_n\}$ that tends to 1.

Example 4: Asymptotic sequences via the decomposition method

Example:

$$\underbrace{2n^2 + 3n + 1}_{a_n} = \underbrace{2n^2}_{b_n} \underbrace{\left(1 + \frac{3}{2n} + \frac{1}{2n^2}\right)}_{c_n} \sim 2n^2 \quad \text{hence } a_n \sim b_n$$

since

$$\underbrace{\left(1 + \frac{3}{2n} + \frac{1}{2n^2}\right)}_{c_n} \rightarrow 1$$

Example 5: Computing limits with asymptotic estimates

We compute the limit

$$\lim_{n \rightarrow +\infty} \frac{2n^3 + 4n + 1}{5(n+1)^3} = \left[\frac{\infty}{\infty} \right] \quad (\text{indeterminate form})$$

Proceeding as before we can write

$$\begin{aligned} \underbrace{2n^3 + 4n + 1}_{a_n} &= \underbrace{2n^3}_{a'_n} \underbrace{\left(1 + \frac{2}{n^2} + \frac{1}{2n^3}\right)}_{\rightarrow 1} \sim 2n^3 \\ 5(n+1)^3 &= \underbrace{5(n^3 + 3n^2 + 3n + 1)}_{c_n} = \underbrace{5n^3}_{c'_n} \underbrace{\left(1 + \frac{3}{n} + \frac{3}{n^2} + \frac{1}{n^3}\right)}_{\rightarrow 1} \sim 5n^3 \end{aligned}$$

Using point 3 of Proposition (??) on asymptotic behavior we can write:

$$\text{if } a_n \sim a'_n, c_n \sim c'_n \quad \text{then} \quad \frac{a_n}{c_n} \sim \frac{a'_n}{c'_n}$$

and obtain

$$\frac{2n^3 + 4n + 1}{5(n+1)^3} \sim \frac{2n^3}{5n^3}$$

that is, the two sequences have the same behavior. Hence

$$\lim_{n \rightarrow +\infty} \frac{2n^3 + 4n + 1}{5(n+1)^3} = \lim_{n \rightarrow +\infty} \frac{2n^3}{5n^3} = \frac{2}{5}$$

- Another way to show that

$$a_n \sim b_n$$

is to use the **substitution principle**.

- For example, knowing that

$$\lim_{n \rightarrow +\infty} \underbrace{\log n}_{a_n} \rightarrow +\infty$$

we can state

$$\lim_{n \rightarrow +\infty} \log c_n \rightarrow +\infty$$

where $\{c_n\}$ is any sequence diverging to $+\infty$, hence

$$\underbrace{\log n}_{a_n} \sim \underbrace{\log c_n}_{b_n}$$

Example 6: Computing limits with the substitution principle and asymptotic estimates

We compute the limit

$$\lim_{n \rightarrow +\infty} \log_3(n^2 + 4n + 1)$$

using the substitution principle and defining

$$c_n = n^2 + 4n + 1 \quad \text{we have} \quad \lim_{n \rightarrow +\infty} c_n = +\infty$$

then

$$\log_3 n \sim \log_3(n^2 + 4n + 1)$$

and hence

$$\lim_{n \rightarrow +\infty} \log_3(n^2 + 4n + 1) = \lim_{n \rightarrow +\infty} \log_3 n = +\infty$$

The fact that the asymptotic relation satisfies the 3 properties:

1. *Reflexive*: $a_n \sim a_n$
2. *Symmetric*: if $a_n \sim b_n$ then $b_n \sim a_n$
3. *Transitive*: if $a_n \sim b_n$ and $b_n \sim c_n$ then $a_n \sim c_n$

means that “**asymptotic**” is an **equivalence relation**.